

# Longitudinal Renal Function Degradation Among Florida Agricultural Workers

Roxana C. Chicas, PhD, Lisa Elon, MS, MPH, Nezahualcoyotl Xiuhotecutli, PhD, Donghai Liang, PhD, Madelyn C. Houser, PhD, Tuzo Mwarumba, BA, Liris Berra, MPH, Vicki Hertzberg, PhD, Jeff M. Sands, MD, and Linda McCauley, PhD

**Objective:** This longitudinal study evaluated renal function and acute kidney injury (AKI) over time in US agricultural workers. **Methods:** We followed Florida agricultural workers from January 2020 to August 2022, collecting blood and urine preworkday and postworkday during five visits. **Results:** Preworkday estimated glomerular filtration rate function in all participants was lower in summers but relatively consistent over time. In participants who worked almost exclusively in fernery operations (piece-rate compensation), we observed a high incidence of postworkday AKI in 2020 (21%) that increased to 43% by the end of the study. In comparison, 11% of nursery workers (hourly compensation) had AKI, and this rate was fairly stable. **Conclusion:** AKI risk over time differs according to the type of agricultural work. Piece rate workers who are incentivized to forgo rest breaks and hydration to earn higher wages demonstrate steadily increasing rates of AKI.

**Keywords:** agricultural workers, piece rate, acute kidney injury, dehydration, heat

Renal dysfunction among agricultural workers has been an area of intense scientific debate and investigation, with the role of heat stress being a prominent area of inquiry.<sup>1-6</sup> Most studies to date have documented chronic kidney disease of unknown etiology (CKDu) (also referred to as chronic kidney disease of nontraditional origin [CKDnt]<sup>7</sup> and Mesoamerican nephropathy<sup>8</sup>) among workers in agricultural regions in Central America. The causes of CKDu remain elusive, but several global studies have documented the degradation of renal function longitudinally in agricultural workers,<sup>9-16</sup> suggesting that repeated episodes of AKI among agricultural workers may increase the risk of CKDu development.<sup>7,17-20</sup>

From the Nell Hodgson Woodruff School of Nursing, Emory University, Atlanta, Georgia (R.C.C., N.X., M.C.H., V.H., L.M.C.); Rollins School of Public Health, Department of Biostatistics and Bioinformatics, Emory University, Atlanta, Georgia (L.E.); Farmworker Association of Florida, Apopka, Florida (N.X.); Rollins School of Public Health, Emory University, Atlanta, Georgia (D.L., L.B.); Department of Medicine, Emory University, Atlanta, Georgia (T.M.); and Division of Renal Medicine, Department of Medicine, Emory University, Atlanta, Georgia (J.M.S.).

Funding sources: This study was funded by a research grant from the National Institute for Occupational Safety & Health (grant number CDC/NIOSH R01OH011782) and the National Institute of Nursing Research (NIH/NINR K23NR020356).

Chicas, Elon, Xiuhotecutli, Liang, Houser, Mwarumba, Berra, Hertzberg, Sands, and McCauley have no relationships/conditions/circumstances that present potential conflict of interest.

The JOEM editorial board and planners have no financial interest related to this research. Author contributions:

The authors confirm contribution to the paper as follows:

Study conception and design: RCC, NX, VH, JMS, LM

Data collection: RCC, NX, LB

Analysis and interpretation of results: RCC, LE, NX, DL, MCH, VH, JMS, LM

Draft manuscript preparation: RCC, LE, MCH, TM, LB

All authors reviewed the results and approved the final version of the manuscript.

Data availability: Data available on request from the authors.

Ethical considerations & disclosure: The Institutional Review Board at Emory University provided approval (IRB00112681) for the study, and all participants provided informed consent.

Supplemental digital contents are available for this article. Direct URL citation appears in the printed text and is provided in the HTML and PDF versions of this article on the journal's Web site ([www.joem.org](http://www.joem.org)).

Address correspondence to: Roxana C. Chicas, PhD, 1520 Clifton Road, Atlanta, GA 30322 ([rchicas@emory.edu](mailto:rchicas@emory.edu)).

Copyright © 2024 American College of Occupational and Environmental Medicine  
DOI: 10.1097/JOM.00000000000003142

## CME Learning Objectives

After completing this enduring educational activity, the learner will be better able to:

- Differentiate risk factors for AKI and renal function between fernery workers (piece-rate compensation) and nursery (hourly compensation).
- Discuss how and why piece-rate workers may be at increased risk of AKI and renal dysfunction.
- Outline the renal function and AKI risk over time in U.S. agricultural workers.

Cross-sectional studies that measured creatinine prework and postwork shift have reported acute kidney injury (AKI) among agricultural workers both in the United States (US)<sup>21,22</sup> and internationally.<sup>14,23,24</sup> Agricultural worker studies of heat stress and renal dysfunction have determined AKI using the Kidney Disease: Improving Global Outcomes (KDIGO) criteria that defines AKI as either an increase in the creatinine level by  $\geq 0.3$  mg/dL or an increase in the creatinine level to  $\geq 1.5$  times the preshift level.<sup>14,21-25</sup>

While the prevalence of AKI has been reported in both agricultural workers in Central America and in the US, few studies have measured renal function longitudinally over the course of a harvest period or over multiple harvesting periods. Renal function for longer than a harvest season has been reported by including retrospective clinical data from pre-employment screenings before previous harvest seasons. Wesseling et al (2016)<sup>11</sup> documented a decrease in estimated glomerular filtration rate (eGFR) after only 9 weeks of sugarcane harvesting, and Laws et al (2016)<sup>10</sup> indicated an association of decreases in renal function with biomarkers suggestive of tubular injury. Sorenson et al (2020) studied Guatemalan sugarcane workers over a 4-year period by retrospectively analyzing data from the preharvest medical employment screenings (years 2012–2016) for the study participants they prospectively monitored in 2017, and they described degradation in eGFR.<sup>9</sup>

There is a clear need for longitudinal tracking of renal function and interventions to interrupt this decline in renal function. The US is also susceptible to rising global temperature, and AKI in agricultural workers has been reported in cross-sectional studies in Florida and California.<sup>21,22</sup> Furthermore, two studies have reported 'hot spots' of unexplained end-stage renal disease in agricultural areas in the US.<sup>26,27</sup> Thus, incidence of AKI and progressive degradation in renal function may be increasing among US agricultural workers. To our knowledge, no published studies in the US have examined renal function and injury of agricultural workers longitudinally. This paucity of research is often influenced by the inability to recruit and follow workers because of worker immigration status even though they may have worked in agriculture for years.<sup>28-32</sup>

This paper describes the results of a community-engaged research program conducted in partnership with the Farmworker Association of Florida (FWAF), which supports and advocates for this marginalized population that is at increased risk of climate-related health

impacts. Through this partnership, we have been able to follow research participants engaged in different types of agricultural work over multiple harvesting seasons to characterize differences in renal function when working in both cool and hot environments and to study degradation of renal function and AKI incidence over time. In addition to the influence of heat exposure on renal dysfunction and decline among these workers, we examined risks falling under the following three categories: health factors, working conditions and habits, and hydration practices and indicators.

## METHODS

Trained community workers from the FWF recruited a convenience sample of agricultural workers in two central Florida locations in January 2020 for a longitudinal study of heat exposure and renal function. Workers were eligible for the study if: (a) they were 18 to 49 years of age at the time of study, (b) self-identified as Hispanic/Latino, and (c) had worked in agricultural settings for at least 1 month before study participation. Exclusions included a history of type 1 diabetes, pregnancy at the time of enrollment, under treatment for hypertension, or reported history of glomerulonephritis, pyelonephritis, renal calculi, or snake bite. This study was approved by the institutional review board of Emory University (IRB00112681), and informed consent was obtained.

### Data Collection Procedures and Variables

At each visit, participants met with study personnel in the FWF field offices for clinical and nonclinical data collection; all data were saved in the REDCap (Vanderbilt University Medical Center, Nashville, Tennessee) platform.<sup>28</sup> At the first visit, participants met on a nonworkday to have their height, weight, and blood pressure recorded and a fingerstick to collect blood for analysis of hemoglobin A1C (A1C NOW®, pts Diagnostics, Whitestown, IN); weight, A1C, and blood pressure were measured again at each visit. Study personnel orally queried participants in the participant's primary language; information reported here included sociodemographics (age, sex, country of origin, marital status, and years of education), health history (frequency of nonsteroidal anti-inflammatory drugs use [NSAID; sometimes/daily or rarely/never], history of gout or kidney stones, family history of gout or kidney stones, family history of diabetes, hypertension, or kidney disease), and general work characteristics (payment method [by-the-piece/hourly or salary/other], number of years in US agriculture, training in heat-related illness prevention, use of pesticides at work). We also inquired if participants had relatives or household members that were also participating, using this information to adjust for possible clustering (nonindependence) effects in statistical analyses.

Before and after their work shift on the observation day, blood and urine were collected. Blood was analyzed with the iSTAT Handheld Blood Analyzer Point of Care system with a Chem8+ cartridge (Abbott Laboratories, Abbott Park, IL), measuring sodium, potassium, chloride, BUN (blood urea nitrogen), and creatinine. Urine specific gravity was measured with an ATAGO meter (Atago PAL-10S digital refractometer, Bellevue, WA). At visits 1 through 4, before work, study staff fitted participants with a monitor worn around the chest (Zephyr™, Boulder, CO) that collected heart rate and physical activity measures every 30 seconds; the monitor was removed when participants returned to the study office after work. Average heart rate and physical activity were calculated for the period between each participant's self-reported work start and stop times.

After work, a FWF community health worker queried participants (one-to-one) by administering a structured survey that included questions on their work start and stop times, beverage consumption (type and quantities), and heat-related illness symptoms (heavy sweating, headache, nausea, dizziness, confusion, muscle cramps, fainting, dysuria) experienced during that workday. The sum of sports drinks, energy drinks, juice, and soda were reported as "ounces of sugary beverages."

The Florida Automated Weather Network (FAWN) is managed by the University of Florida's Institute of Food and Agricultural Sciences,

which provides rural weather information to agricultural communities. The FAWN has 35 monitoring stations, which collect weather data every 15 minutes throughout Florida's agricultural communities. The FAWN data were obtained from the monitoring station located within each study area (Pierson and Apopka) in which the participants were working. Using the Lu and Romps extended heat index (HI) algorithm based on ambient temperature and relative humidity,<sup>27</sup> mean HI, max HI, and hours spent at HI >=100°F were calculated for each participant's work hours.

The primary kidney function outcomes were prework and postwork shift eGFR and acute kidney injury (AKI). We estimated the glomerular filtration rate using the 2021 CKD-EPI Creatinine Equation,<sup>33</sup> as recommended for US adults by the joint National Kidney Foundation/American Society of Nephrology Task Force.<sup>34</sup> Incident AKI was defined as an increase in creatinine from prework to postwork shift of at least 50% or ≥0.3 mg/dL in accordance with KDIGO guidelines<sup>25</sup> and as recommended by the First International Research Workshop on Mesoamerican Nephropathy.<sup>35</sup>

In addition to age, sex, work type, and time in study, risk factors examined for the primary outcomes of low eGFR and AKI incidence fall into three *a priori* categories: health factors, work conditions, and hydration practices and indicators. Health risk factors included body mass index, blood pressure (normal [ $<120$  systolic and  $<80$  diastolic], elevated 120–139 [systolic or 80–89 diastolic], high [ $\geq 140$  systolic or  $\geq 90$  diastolic]), A1C  $\geq 5.7$  (ie, prediabetes or diabetes), resting heart rate, sometimes/daily use of NSAIDs, and family history of kidney risk factors (any diabetes/hypertension/kidney disease). Work risk factors included training in HRI prevention (yes/no), occupation as a pesticide applicator (years), workday duration, shade breaks taken, heat exposure (mean HI, max HI, hours at HI  $\geq 100^\circ\text{F}$ ), and average heart rate and physical activity (Vector magnitude units (VMU) units) during workday. Hydration practices and indicators included total self-reported beverage consumption and average beverage consumption per work hour, sugary beverage consumption, dehydration (urine specific gravity  $\geq 1.020$  both before and after work), and serum osmolality (mOsm/kg).

Additionally, the effects of primary work type were examined. The two locations were selected because the nature of agricultural work in these two Florida locations is quite distinct. In Apopka, 95% of the recruited participants worked at a nursery while in Pierson, and 90% worked in a fernery. Common tasks that nursery workers performed were planting; cutting; trimming; packing; loading and unloading; watering and pruning trees, shrubs, and plants; and moving potted plants and shrubs. Fernery workers mainly cut fern and bundled it into bunches, which they loaded onto a truck. While correspondence between location and primary work type were not absolute, for simplicity, workers from the Apopka cohort will henceforth be referred to as "nursery workers" and the Pierson cohort as "fernery workers." Payment method was not examined as a separate risk factor as it was colinear with primary work type with fernery workers likely to be paid by piece work. This practice is not observed in nursery work.

### Statistical Analysis

Descriptive statistics were presented as % (*n*) or median (first quartile, third quartile). Simple differences in characteristics between primary work types at baseline were examined with methods that adjusted for the random effect of household cluster: the R package "lqmm" (linear quantile mixed models)<sup>36,37</sup> for median values of continuous variables, generalized linear mixed models for categorical variables.

Before attempting multivariate modeling of the longitudinal data, we used multiple imputation for risk factors having more than 10% missing values. Missing values were assumed missing at random as they were mainly due to machine malfunction. The imputation model included all risk factors, outcomes, and intended interaction terms that would be included in analysis models, plus demographic and clinical auxiliary variables. We conducted the imputation using the R package "mice,"<sup>38</sup> which uses fully conditional specification,

and employed predictive mean matching (numeric data), logistic regression imputation (binary data, factor with 2 levels), or proportional odds model for (ordered, >2 levels), as appropriate; 35 imputations and 15 iterations were specified. After examining convergence plots and imputed values, the originally missing *outcome* values were reset to missing.

Because workday heart rate and physical activity were not measured at the final visit, we examined how variable a participant's measurements were over time by calculating the intraclass correlation coefficients (ICC). ICC for mean heart rate was 0.60 for winter and 0.62 for summer (0.55 overall); for mean physical activity, it was 0.81 for winter and 0.86 for summer (0.75 overall), indicating low variability. We thus chose to use visit 1 values as the predictor rather than employing a less than optimal imputation of all final visit values. To account for the nested data structure, linear mixed modeling and generalized linear mixed modeling with a random intercept for ID were used for simple and multivariate inference. For predicting prework and postwork eGFR, a base model included time, primary work, and their interaction plus covariates age and sex. This simple model was run with all five visits, and again for summer and winter visits separately, dropping the interaction term if it was not significant in the base model. To the summer and winter models, risk factors were added one by one. Risk factors measured at visit 1 were also examined for interaction with time to examine whether they influenced the course of eGFR over time. Models that included imputed risk factors used SAS MIANALYZE<sup>39</sup> (Cary, NC) to calculate estimates and statistical significance. A similar analytical plan was executed for predicting AKI incidence, except that no winter model was run due to the limited number of cases of AKI. Additionally, the base model included only time, primary work, age, and sex as the interaction term of time by primary work was not significant. The assumptions of logit linearity for continuous predictors and absence of multicollinearity were confirmed.

Analyses were performed with SAS version 9.4 software (Cary, NC) and R 4.3.0. Statistical significance was evaluated using  $\alpha = 0.05$ .

## RESULTS

### Longitudinal Study Retention

This study was conducted over 32 months from January 2020 to August 2022, with study visits and workday monitoring conducted in two winters and in three summers during that time. Sixty-seven participants in Apopka, FL, primarily working in nurseries, and forty-eight participants in Pierson, FL, primarily working in ferneries, were initially recruited. As previously reported,<sup>40</sup> more than three quarters of participants were retained over the first two years of the study, with retention decreasing for the final summer visit which occurred more than a year after the previous study visit. Retention at the last visit was higher among Pierson fernery workers (79%) compared with Apopka nursery workers (52%).

### Demographic Characteristics

Several differences were identified in the nursery and fernery worker groups (Table 1). While more than two-thirds of the recruited participants working in either setting identified as female, there was a higher percentage (33%) of males among the fernery workers compared to the nursery workers (16%,  $P = 0.03$ ). There were no significant differences in age or in the age at which participants started working in agriculture, but there was a significant difference ( $P = 0.003$ ) in the number of years participants had been working in US agriculture (median 6 in nurseries vs. median 16 in ferneries). Additionally, nearly all (94%) fernery workers claimed Mexican nationality with the remainder being from the US, while nursery workers were a less homogeneous group ( $P < 0.001$ ) with roots in Mexico (57%), Guatemala (24%), the US (3%), and other countries (16%). A lower percentage (21%) of fernery workers were also single compared to nursery workers (40%,  $P = 0.04$ ). Educational level was similarly low in the two cohorts (median of 8 years,  $P = 0.8$ ).

### Health Characteristics

At the initial study visit (January 2020), the median body mass index among fernery workers (30.7) was higher than that of nursery workers (28.5,  $P = 0.04$ ), and this pattern remained consistent across study visits (Table S1, <http://links.lww.com/JOM/B598>). Fernery workers also had higher blood pressure (BP;  $P = 0.049$ ); at the initial study visit, 41% of nursery workers had BP readings classified as elevated or hypertensive (>120/80) compared to 59% of fernery workers (Table 1). By the final study visit, the percentage of participants with elevated or hypertensive BP readings had increased in both groups (48% nursery workers, 66% fernery workers), but the distinction between the groups remained (Table S1, <http://links.lww.com/JOM/B598>). No differences were found between the worker groups in diabetes status (based on A1C percent), family or personal history of kidney disease-related medical issues (gout, kidney stones, hypertension, or diabetes), NSAID use, or resting heart rate (Table 1). A seasonal pattern was observed for A1C percent, with slightly higher values in the summers in both worker cohorts (Table S1, <http://links.lww.com/JOM/B598>).

### Work Environment and Practices

Nearly all (96%) nursery workers were paid by the hour and none by the piece, whereas most fernery workers (73%) were compensated per piece or by a combination of piece and hours (6%) (Table 1). This difference in compensation structure likely contributed to the differences in participants' reports of their workday break practices ( $P = 0.03$ ); among fernery workers, 80% reported taking no breaks or just one break during workdays; 46% of nursery workers reported taking two or more breaks. Differences in work hours may also have contributed to these differences in breaks. Participants affiliated with nurseries worked longer hours (median 9.0) than did participants affiliated with ferneries (median 7.2) ( $P < 0.001$ ). There were no significant differences between the agricultural settings in participants' reports of working with pesticides or recent training to prevent heat-related illness.

Participants' physical activity also differed between the agricultural settings (Table 1). Fernery workers had a higher average heart rate (median 96.3 bpm) and activity level (median 0.20 VMU) compared with nursery workers (median 91.5 bpm,  $P = 0.04$ ; median 0.11 VMU,  $P < 0.001$ ). A median of 0.20 VMU is the activity equivalent of someone continuously walking during their whole work shift.<sup>41</sup> Fernery workers were also bending more during their work shifts ( $P < 0.001$ ). These activity differences remained consistent at all timepoints measured (Table 2).

Heat indices were calculated in each location for each study visit day (Fig. 1). While there was greater variation in heat indices across study visit days for fernery workers, the mean heat index as well as the number of hours in which the heat index was over 100°F were comparable for both groups. The max heat index did not differ for the first three study visits but was slightly higher for fernery workers at the last two summer study visits.

### Dehydration Prevalent Among Fernery Workers

While overall self-reported workday beverage consumption did not differ between fernery and nursery workers, choices of beverages did, with more fernery workers drinking sugary beverages than nursery workers ( $P = 0.002$ ) (Table 1). Sugary beverage intake during work was more common in both groups in the summers than in the winters, but the greater prevalence and higher intake among fernery workers persisted at all study visits (Table S2, <http://links.lww.com/JOM/B598>). Coffee consumption was higher among nursery than fernery workers, but overall, the numbers of participants reporting drinking coffee during work were lower than those reporting drinking sugary beverages or water (Table S2, <http://links.lww.com/JOM/B598>).

Over the course of the study, there was greater evidence of dehydration among fernery workers compared to nursery workers (Table 3). At all but one study timepoint (August 2020 preworkday),

**TABLE 1.** Participant Personal and Work Characteristics at Initial Visit, by Primary Work. OHEaRD Study Jan2020

Characteristic	n	Overall <sup>a</sup> , N = 115	Nursery <sup>a</sup> , n = 67	Fernery <sup>a</sup> , n = 48	P <sup>b</sup>
<b>Demographics</b>					
Age, yrs	115	38.9 (33.4, 44.5)	37.6 (33.4, 43.2)	41.1 (33.7, 46.0)	0.1
Sex	115				0.03
Male		23% (27)	16% (11)	33% (16)	
Female		77% (88)	84% (56)	67% (32)	
Nationality	115				<0.001
Mexico		72% (83)	57% (38)	94% (45)	
Guatemala		14% (16)	24% (16)	0% (0)	
USA		4% (5)	3% (2)	6% (3)	
Other		10% (11)	16% (11)	0% (0)	
Education, yrs	114	8.0 (6.0, 10.0)	8.0 (6.0, 11.0)	8.0 (6.0, 9.0)	0.8
Marital status	115				0.04
Coupled		68% (78)	60% (40)	79% (38)	
Single		32% (37)	40% (27)	21% (10)	
<b>Health-related</b>					
BMI	115	29.1 (25.7, 33.4)	28.5 (24.7, 32.2)	30.7 (27.0, 33.8)	0.04
Heart rate, resting, bpm	115	73.0 (66.5, 81.0)	73.0 (66.0, 82.5)	74.0 (68.0, 79.2)	0.5
Blood pressure	107				0.049
<120/80		52% (56)	59% (38)	42% (18)	
120–139 or 80–89		34% (36)	25% (16)	47% (20)	
140+ or 90+		14% (15)	16% (10)	12% (5)	
A1C	114				0.1
Normal		84% (96)	79% (52)	92% (44)	
Prediabetic		13% (15)	18% (12)	6% (3)	
Diabetic		3% (3)	3% (2)	2% (1)	
NSAID, sometimes/daily	115	14% (16)	13% (9)	15% (7)	0.9
Hx gout/kidney stones	114	4% (4)	4% (3)	2% (1)	0.5
Family Hx gout/kidney stones	108	13% (14)	15% (9)	11% (5)	0.5
Family Hx DB/HTN/KD	110	52% (57)	52% (33)	51% (24)	0.9
<b>Work-related</b>					
Work type	113				
Fernery		38% (43)	0% (0)	90% (43)	<0.001
Nursery		56% (63)	95% (62)	2% (1)	
Other		6% (7)	5% (3)	8% (4)	
Work hours	113	8.5 (7.0, 9.5)	9.0 (8.5, 9.5)	7.2 (6.0, 8.5)	<0.001
Age started in agriculture	115	20.0 (17.0, 26.0)	21.0 (15.5, 29.0)	19.0 (17.0, 25.0)	0.4
Working in US agriculture, yrs	114	11.0 (4.0, 18.0)	6.0 (2.5, 14.0)	16.0 (10.0, 22.5)	0.003
Payment	115				<0.001
By the piece		30% (35)	0% (0)	73% (35)	
By the hour		64% (74)	96% (64)	21% (10)	
Combination piece/hour		3% (4)	1% (1)	6% (3)	
Salary		2% (2)	3% (2)	0% (0)	
Took breaks	113				0.03
None		35% (39)	31% (20)	40% (19)	
One		30% (34)	23% (15)	40% (19)	
Two		33% (37)	43% (28)	19% (9)	
Three or more		3% (3)	3% (2)	2% (1)	
Took shade breaks		58% (65)	63% (41)	50% (24)	0.2
HRI training in past year	115	33% (38)	31% (21)	35% (17)	0.6
Works with pesticides	115	8% (9)	12% (8)	2% (1)	0.1
Heart rate, avg during work, bpm	105	94.5 (87.9, 100.8)	91.5 (86.4, 99.4)	96.3 (90.2, 104.4)	0.04
Activity, avg during work, VMU in g	108	0.13 (0.10, 0.19)	0.11 (0.09, 0.13)	0.20 (0.15, 0.24)	<0.001
Posture, avg during work, degrees from vertical	108	7 (–3, 27)	0 (–4, 8)	26 (10, 37)	<0.001
<b>Hydration</b>					
Beverages, oz	113	44.0 (30.0, 64.0)	40.0 (24.0, 64.0)	48.0 (36.5, 64.0)	0.2
Sugary beverages, oz	113	0.0 (0.0, 12.0)	0.0 (0.0, 8.0)	12.0 (0.0, 16.0)	0.002

<sup>a</sup>% (n) or median (first quartile, third quartile).

<sup>b</sup>Differences tested using linear quantile mixed models (continuous variables) or generalized linear mixed models (categorical variables), unless the random effect of household cluster was negligible. In the latter case, simple linear regression or chi-squared tests were conducted. DB, Diabetes; HTN, Hypertension; KD, Kidney Disease.

a higher percentage of fernery workers than nursery workers had urine specific gravity (USG) ≥ 1.020, and at all timepoints, more fernery than nursery workers were classified as dehydrated by USG ≥ 1.020 both before and after work shifts (odds ratio [OR] 2.1, 95% confidence interval

[CI] 1.1, 3.8). In the summers, the percentage of workers with USG ≥ 1.020 was higher postworkday than it was preworkday in both agricultural settings, but the increases in the percentage of dehydrated workers preworkday to postworkday were higher among fernery

Downloaded from http://journals.lww.com/jeem by BNDMfsePHKav1ZEoum1tQIN4a+kJLHEZgbsHh04XM10hCwCX1AW nYQp/llQH333D000dRy/7TVSFI4C3Vc1y0abgqZQZxGg12MwIzlel= on 09/06/2024

**TABLE 2.** Workday Physical Activity Summaries Based on 30-Second Epoch Continuous Monitoring

	Jan2020 <sup>a</sup>	July2020 <sup>a</sup>	Jan2021 <sup>a</sup>	July2021 <sup>a</sup>
<b>Fernery</b>	<b>n = 45<sup>b</sup></b>	<b>n = 28</b>	<b>n = 35</b>	<b>n = 14</b>
Activity, mean	0.20 (0.15, 0.24)	0.15 (0.13, 0.22)	0.20 (0.14, 0.25)	0.20 (0.16, 0.24)
Posture, mean	26 (10, 37)	10 (0, 24)	26 (-3, 41)	31 (14, 40)
Steps, walking	3,407 (2,681, 6,086)	3,180 (1,906, 5,442)	4,141 (2,381, 6,892)	2,381 (2,057, 3,830)
Steps, running	15 (6, 25)	30 (8, 78)	15 (6, 52)	10 (4, 29)
Steps, walking + running	3,463 (2,700, 6,107)	3,190 (1,910, 6,110)	4,152 (2,384, 7,046)	2,422 (2,074, 3,848)
Heart rate, mean	96 (90, 104)	95 (87, 108)	96 (90, 104)	104 (95, 116)
<b>Nursery</b>	<b>n = 63<sup>b</sup></b>	<b>n = 46</b>	<b>n = 47</b>	<b>n = 41</b>
Activity, mean	0.11 (0.09, 0.13)	0.10 (0.08, 0.14)	0.10 (0.08, 0.13)	0.11 (0.08, 0.13)
Posture, mean	0 (-4, 8)	-1 (-4, 4)	-2 (-5, 4)	0 (-5, 5)
Steps, walking	7,574 (4,586, 10,052)	5,712 (4,415, 10,370)	5,380 (4,279, 9,794)	6,672 (3,871, 9,907)
Steps, running	58 (28, 156)	58 (14, 126)	48 (26, 112)	77 (29, 134)
Steps, walking + running	7,779 (4,655, 10,361)	5,784 (4,425, 10,424)	5,609 (4,308, 9,864)	6,934 (3,903, 9,984)
Heart rate, mean	92 (86, 99)	94 (87, 102)	91 (83, 98)	91 (84, 100)

Activity was not measured during the 2022 season.

Summer data are shaded.

<sup>a</sup>Median (IQR)

<sup>b</sup>Monitor files were included in calculations if at least 80% of the workday was captured. Additional missingness of m<sub>Fernery</sub> = 19, m<sub>Nursery</sub> = 27 records for steps at visit 1 due to technical issues.

workers than nursery workers. Furthermore, in the summers, markedly higher percentages of fernery workers compared to nursery workers were classified as severely dehydrated (USG ≥ 1.030) postworkday (25%, 11%, and 18% vs 2%, 4%, and 9%, respectively [OR 4.9, 95% CI 1.6, 14.8]). At all but one timepoint, higher percentages of fernery workers compared to nursery workers also met the criteria for hyperosmolality after workday.

**Preworkday eGFR Slightly Higher in Fernery Versus Nursery Workers**

Preworkday eGFR values adjusted for sex and age of participants remained more consistent across the study than did postworkday eGFR values (Fig. 2). There was no significant interaction between work season and primary work type, as there was with postworkday eGFR, but there were significant main effects, with preworkday eGFR values slightly lower in the summers (beta -1.5, 95% CI -2.7, -0.3) and slightly higher in fernery compared with nursery workers (beta 4.1, 95% CI 1.4, 6.9). Prework eGFR <90 was only seen in a single participant during the study (Table 3).


**Increasing AKI and Declining Postworkday eGFR in Fernery Workers**

Patterns of AKI across the study differed between nursery and fernery workers (Fig. 3, Table 3). At all but one timepoint (July 2021), about 10% of nursery workers met the criteria for AKI postwork shift. At all but one timepoint (Aug 2020), more than 20% of fernery workers met the criteria for AKI postwork shift, and this percentage increased progressively over the last three study timepoints from 21% in

January 2021 to 43% in August 2022 (OR 8.1 for fernery workers in summer 2021 and 12.6 in summer 2022 relative to summer 2020 compared with 2.4 and 1.1 for nursery workers, respectively, Table S3, <http://links.lww.com/JOM/B598>). Besides agricultural setting, none of the demographic factors, health factors, work conditions, or hydration measures collected in this study were significantly associated with summer AKI (Table S3, <http://links.lww.com/JOM/B598>).

Estimated GFR levels calculated prework and postwork shift also differed across the study for workers in the two agricultural settings (Table 3, Fig. 2). After adjustment for sex and age of the participants, postworkday eGFR values for fernery workers were higher than for nursery workers at the first three study timepoints (winter 2020, summer 2020, winter 2021), but, in alignment with AKI results, postworkday eGFR values for fernery workers dropped below nursery workers at the last two summer timepoints (Fig. 3, Table 3). For fernery workers, comparing base summer 2020 to the two following summers showed substantial decline in eGFR (2021 beta -9.3, 95% CI -16.9, -1.7; 2022 beta -10.5, 95% CI -18.1, -2.9) that was not observed for nursery workers (2021 beta -2.6, 95% CI -7.5, 2.2; 2022 beta 2.8, 95% CI -2.3, 7.9) (Table 4). At these timepoints, we also observed higher percentages of fernery workers with postworkday eGFR <90 than had been seen at previous timepoints or were ever observed among nursery workers in this study (Table 3).

Our primary predictor, heat exposure, significantly impacted summer postworkday eGFR; for every degree increase in average heat index, postworkday eGFR declined by 0.5 units (P = 0.036), and controlling for it in the model revealed significant declines in eGFR from summer 2020 to summer 2021 in nursery workers (beta -6.0, 95% CI -11.7, -0.3) while deattenuating the effect in fernery workers for the



**FIGURE 1.** Heat indices largely comparable in both agricultural settings. At each study visit, the distribution of participants' work shift heat exposure: mean and max heat index experienced during their work shift and number of hours in which the heat index was above 100°F. Boxplots summarize values for all participants, with the line inside the box representing the median and the symbol inside the box representing the mean. Weather data obtained from the FAWN monitoring stations. Heat index calculated using Lu and Romps extended heat index algorithm based on ambient temperature and relative humidity.

Downloaded from http://journals.lww.com/joem by BMDMSEPHKav1ZEoum1tQIN4a+kLHEZgbsHio4XMl0hCwCX1AW nYQp/llQHd313D00dRv7ITVSFI4C3VC1y0abg9QZxdgGj2mWlZel= on 09/06/2024

Downloaded from http://journals.lww.com/jeem by BNDMSEPHKAV1ZEoum1IQIN44+kLLHEZgbsH04XM10hCwCX1AW on 09/06/2024

**TABLE 3. Urine and Serum Characteristics, OHEARD Study Jan2020–Aug2022**

Characteristic	Nursery										Fernery					
	Jan2020 <sup>a</sup> n = 67	Aug2020 <sup>a</sup> n = 60	Jan2021 <sup>a</sup> n = 50	Jul2021 <sup>a</sup> n = 52	Aug2022 <sup>a</sup> n = 35	Jan2020 <sup>a</sup> n = 48	Aug2020 <sup>a</sup> n = 39	Jan2021 <sup>a</sup> n = 37	Jul2021 <sup>a</sup> n = 36	Aug2022 <sup>a</sup> n = 38						
USG > =1.020, am	30% (19)	45% (25)	24% (12)	39% (20)	37% (13)	40% (17)	44% (16)	49% (18)	53% (19)	50% (19)						
USG > =1.020, pm	28% (17)	68% (40)	28% (14)	44% (22)	49% (17)	43% (18)	78% (28)	41% (14)	67% (24)	66% (25)						
USG > =1.020, AM and PM	14% (8)	33% (18)	10% (5)	27% (13)	23% (8)	23% (9)	39% (13)	32% (11)	44% (16)	39% (15)						
USG > =1.030, am	3% (2)	0% (0)	0% (0)	2% (1)	6% (2)	2% (1)	3% (1)	5% (2)	3% (1)	5% (2)						
USG > =1.030, pm	7% (4)	2% (1)	4% (2)	4% (2)	9% (3)	2% (1)	25% (9)	6% (2)	11% (4)	18% (7)						
Serum Osm, am																
Hyperosmolality <sup>b</sup>	41% (25)	33% (18)	27% (12)	61% (27)	57% (20)	49% (23)	32% (11)	21% (7)	50% (15)	43% (16)						
Serum Osm, pm																
Hyperosmolality <sup>b</sup>	14% (9)	30% (17)	14% (6)	49% (20)	40% (14)	40% (19)	33% (11)	18% (6)	48% (15)	46% (17)						
AKI	10% (6)	11% (6)	10% (4)	24% (9)	11% (4)	21% (10)	6% (2)	21% (6)	33% (9)	43% (16)						
eGFR, am																
Median (IQR)	121 (117, 128)	121 (115, 126)	122 (117, 126)	118 (113, 125)	119 (114, 126)	124 (116, 128)	119 (115, 126)	124 (118, 134)	121 (114, 128)	124 (115, 131)						
eGFR, pm																
Median (IQR)	117 (112, 125)	117 (108, 123)	119 (112, 125)	115 (105, 118)	117 (110, 121)	117 (111, 127)	116 (110, 121)	118 (114, 126)	114 (82, 122)	113 (92, 124)						
eGFR < 90, am	0% (0)	0% (0)	0% (0)	2% (1)	0% (0)	0% (0)	0% (0)	0% (0)	0% (0)	0% (0)						
eGFR < 90, pm	3% (2)	5% (3)	5% (2)	12% (5)	0% (0)	8% (4)	0% (0)	0% (0)	28% (9)	14% (5)						

Summer values are shaded.  
<sup>a</sup>% (n) or median (first quartile, third quartile).  
<sup>b</sup>>295 mOsm/kg.  
eGFR, estimated glomerular filtration rate.

same period (beta -5.8, 95% CI -14, 2.3) (Table 4). Similar effects were observed when accounting for the number of work hours in which the heat index was above 100°F. A significant negative association ( $P = 0.024$ ) was identified between summer postworkday eGFR and the volume of beverage consumption during the workday independent of agricultural setting. Besides age, no other risk factors were significantly associated with this outcome, nor were they confounders of the relationship between work type and time.

Postworkday eGFR in the winters appeared to have a different relationship with predictors evaluated in this study (Table 5). There was not a clear effect of agricultural setting nor evidence of decline between winter 2020 and winter 2021. Other than age, the only predictor that was significantly associated with winter postworkday eGFR was serum osmolality, with eGFR declining 1 unit for every unit increase in serum osmolality (95% CI -1.5, -0.5).


There were no appreciable differences longitudinally across the summers or the winters (Tables 6 and 7, respectively). Resting heart rate, NSAID use, and work applying pesticides were significant predictors of summertime prework eGFR. Every beat per minute increase in resting heart rate was associated with an increase of 0.1 units in preworkday eGFR (95% CI 0.1, 0.2,  $P = 0.001$ ). Participants who reported using NSAIDs more frequently had higher preworkday eGFR (beta 2.2, 95% CI 0, 4.3,  $P = 0.049$ ). Each year of occupational pesticide use was associated with a decline of 0.4 units in preworkday eGFR (95% CI -0.9, 0,  $P = 0.042$ ). Beyond age and sex, no other risk factors assessed were significantly associated with summertime eGFR, nor were they confounders of the relationship between preworkday eGFR and work type (Table 6). Additionally, other than age, none of the risk factors assessed were significantly associated with wintertime preworkday eGFR (Table 7).

**DISCUSSION**

In this longitudinal study of two agricultural communities in Florida, we found differences in their levels of renal dysfunction. Preworkday eGFR in both communities was slightly lower in the summers. The study participants in Pierson, who worked almost exclusively in fernery operations, had slightly higher preworkday eGFR compared with participants from Apopka, who worked primarily in agricultural nurseries, and this did not vary greatly over time. In contrast, postworkday eGFR for fernery workers, which had been slightly higher than for nursery workers in previous study seasons, dropped below that for nursery workers at the last two summer timepoints. Fernery workers also had a high rate of AKI postworkday when they were first measured in 2020 (21%), and the rate had more than doubled by the end of the study (43%). In contrast, in Apopka, AKI developed in approximately 10% to 11% of participants, and the rate was stable over time.

Fernery workers (Pierson, FL) have several risk factors that may place them at increased risk of AKI and lower postwork eGFR compared to nursery workers (Apopka, FL). Fernery workers are primarily compensated by the piece rate system that likely contributes to them taking fewer rest breaks (0–1 rest break) and have higher physical activity, whereas the nursery workers are primarily hourly workers, more of them took two rest breaks, and their physical activity was lower. Furthermore, the fernery workers consumed more sugary drinks, specifically in the summers, compared to the nursery workers.


We observed a significant negative association ( $P = 0.024$ ) between summer postworkday eGFR and the volume of fluid consumption during the workday independent of agricultural setting. Although workers in both locations were consuming more fluid in the summer, the average fluid consumption per hour was between 8 and 12 ounces, whereas the recommended amount for those working strenuously in the heat is 24 to 32 ounces per hour. Workers who were drinking more were probably under more heat stress, but the amount of fluid they were drinking was not enough to keep them hydrated. Heat index



**FIGURE 2.** Longitudinal and seasonal dynamics of prework and postwork estimated kidney function. Average prework and postwork eGFR calculated from serum creatinine values at each timepoint in each agricultural setting as predicted by linear mixed model containing the following variables: primary work type, timepoint, the interaction of primary work type and timepoint, age, and sex. Error bars reflect 95% confidence limits.

was a significant predictor of postwork eGFR, and the fernery workers experienced slightly higher heat index with some differences in max heat index; however, the mean was quite similar in both groups. Including heat index as a covariate in analytical models with AKI or postworkday eGFR as outcomes did not markedly impact the significant relationship with work type, indicating that the difference between the worker groups was not all attributable to heat index.

The fernery workers in Florida have similar risk factors as agricultural workers with CKDu in Mesoamerica, such as being compensated by the piece-rate system, intense physical exertion, high ambient temperature, dehydration, and increased consumption of sugary drinks.<sup>7,18–20,42</sup> Degradation of renal function over time has been documented among Mesoamerican sugarcane workers.<sup>9–11</sup> While our assessments of preworkday eGFR suggest that Florida fernery workers did not experience irreversible



**FIGURE 3.** Acute kidney injury more prevalent among fernery workers. Percentage of participants meeting the criteria for AKI (an increase in serum creatinine from before to after work shift of at least 50% or  $\geq 0.3$  mg/dL) at each timepoint in each agricultural setting as predicted by generalized linear mixed model containing the following variables: primary work type, timepoint, the interaction of primary work type and timepoint, age, and sex. Error bars reflect 95% confidence limits.

Downloaded from http://journals.lww.com/jeem by BNDMSEPHKAV1ZEUMT1QIN44KJLHEZG9BSH04XMI0h0CwCX1AVW n/Op/llQHHD3E3D00dRv7ITV5F14C3VVC1y0ab9g9QZxqG12MwIzLEl= on 09/06/2024

**TABLE 4. Summertime Predictors of Postwork eGFR Evaluated Using Mixed Models With the Base Model Containing Primary Work Type, Timepoint, and Their Interaction, Controlling for Sex and Age**

Model <sup>a</sup>	Intercept	Primary Work Type		Nursery x Time		Femery x Time		Sex (Ref = Female)	Age (cent40 <sup>d</sup> )	Additional Covariate	P <sup>e</sup>
		Summer20 (Ref = Nursery)	Summer21 (Ref = Summer20)	Summer20 (Ref = Summer20)	Summer21 (Ref = Summer20)	Summer22 (Ref = Summer20)	Summer22 (Ref = Summer20)				
		Estimated Beta (95% Confidence Interval) <sup>b</sup>									
<b>Base model</b>	107.9 (102.5, 113.4)	5.8 (0.2, 11.4)	-2.6 (-7.5, 2.2)	2.8 (-2.3, 7.9)	-9.3 (-16.9, -1.7)	-10.5 (-18.1, -2.9)	-1.1 (-1.3, -0.8)	4 (-1, 9)	-1.1 (-1.3, -0.7)	-0.2 (-0.6, 0.1)	0.22
<b>+ Health factors</b>	107.7 (102.2, 113.1)	6.3 (0.7, 12)	-2.5 (-7.4, 2.3)	2.9 (-2.2, 8)	-9.6 (-17.2, -2)	-10.6 (-18.2, -3)	-1 (-1.3, -0.7)	4.2 (-0.8, 9.1)	-1 (-1.3, -0.7)	0.8 (-3.2, 4.9)	0.687
BMI cent30 <sup>d</sup>	107.4 (101.5, 113.3)	5.7 (0, 11.4)	-2.8 (-7.8, 2.1)	2.8 (-2.4, 7.9)	-8.7 (-16.5, -0.9)	-10.4 (-18.1, -2.8)	-1.1 (-1.4, -0.8)	4.2 (-0.9, 9.3)	-1.1 (-1.4, -0.8)	0.8 (-3.2, 4.9)	0.687
BP 120-139 or 80-89 (ref = <120/80)	107.2 (101.5, 112.9)	6.3 (0.6, 12)	-2.6 (-7.5, 2.3)	3 (-2.2, 8.1)	-9.3 (-16.9, -1.7)	-10.7 (-18.3, -3)	-1.1 (-1.3, -0.8)	4.1 (-0.9, 9.1)	-1.1 (-1.3, -0.8)	0.7 (-4.7, 6.1)	0.798
AIC >= 5.7 at V1 <sup>e</sup>	108.6 (95.6, 121.6)	5.8 (0.2, 11.5)	-2.7 (-7.6, 2.2)	2.7 (-2.6, 8)	-9.3 (-16.9, -1.6)	-10.1 (-17.8, -2.4)	-1.1 (-1.3, -0.8)	4.2 (-0.9, 9.2)	-1.1 (-1.3, -0.8)	2.0 (-3.4, 7.3)	0.461
Resting heart rate	108.1 (102.5, 113.4)	5.8 (0.2, 11.4)	-2.7 (-7.6, 2.2)	2.8 (-2.3, 7.9)	-9.3 (-16.9, -1.6)	-10.5 (-18.1, -2.9)	-1.1 (-1.3, -0.8)	4.1 (-0.9, 9)	-1.1 (-1.3, -0.8)	0 (-0.2, 0.2)	0.899
NSAID use, sometimes or daily	107.3 (101.1, 113.6)	6 (0.1, 11.8)	-3.2 (-8.3, 1.9)	2.7 (-2.6, 8)	-9.2 (-17.1, -1.4)	-10.1 (-17.9, -2.3)	-1.1 (-1.4, -0.8)	4.0 (-1.2, 9.2)	-1.1 (-1.4, -0.8)	-0.4 (-5.2, 4.4)	0.874
Family hx kidney risk factors <sup>f</sup>	107.3 (101.8, 112.8)	5.5 (-0.2, 11.1)	-2.7 (-7.6, 2.1)	2.6 (-2.5, 7.8)	-9.3 (-16.9, -1.8)	-10.3 (-17.9, -2.7)	-1.1 (-1.3, -0.8)	3.8 (-1.1, 8.8)	-1.1 (-1.3, -0.8)	2.7 (-1.4, 6.8)	0.194
<b>+ Work conditions</b>	110 (104, 116)	5.4 (-0.3, 11)	-2.7 (-7.6, 2.1)	2.7 (-2.5, 7.8)	-9.3 (-16.8, -1.7)	-10.4 (-18, -2.9)	-1.0 (-1.3, -0.8)	2.2 (-3.2, 7.7)	-1.0 (-1.3, -0.8)	-0.5 (-1.2, 0.1)	0.127
Training in HRI prevention	108.1 (102.5, 113.7)	5.6 (-0.5, 11.6)	-2.7 (-7.5, 2.2)	2.8 (-2.3, 7.9)	-9.3 (-16.9, -1.7)	-10.5 (-18.1, -2.8)	-1.1 (-1.3, -0.8)	3.9 (-1.1, 9)	-1.1 (-1.3, -0.8)	-0.1 (-1.1, 0.9)	0.812
Pesticide use, years	106.8 (101, 112.6)	5.9 (0.4, 11.5)	-2.8 (-7.6, 2)	2.6 (-2.4, 7.6)	-8.1 (-15.6, -0.6)	-10.2 (-17.7, -2.8)	-1.0 (-1.3, -0.8)	3.4 (-1.6, 8.4)	-1.0 (-1.3, -0.8)	2.8 (-0.7, 6.3)	0.115
Workday duration, hours	109.8 (104.1, 115.5)	5.1 (-0.5, 10.7)	-2.8 (-7.6, 2)	2.1 (-2.9, 7.2)	-5.8 (-14, 2.3)	-9.1 (-16.7, -1.6)	-1.0 (-1.3, -0.8)	4.6 (-0.4, 9.7)	-1.0 (-1.3, -0.8)	-0.5 (-0.9, 0)	0.036
Shade breaks taken	111.3 (105.2, 117.5)	5.5 (-0.1, 11.0)	-5.7 (-11.3, -0.1)	1.8 (-3.5, 7.0)	-6.0 (-14.1, 2.1)	-8.7 (-16.5, -1.0)	-1.0 (-1.3, -0.8)	3.8 (-1.1, 8.7)	-1.0 (-1.3, -0.8)	-1.3 (-2.5, -0.1)	0.030
Heat index mean, cent90 <sup>d</sup>	107.5 (102.1, 113)	6.3 (0.6, 12)	-2.6 (-7.5, 2.2)	2.8 (-2.3, 7.9)	-9.4 (-17, -1.8)	-10.5 (-18.1, -2.9)	-1.1 (-1.4, -0.8)	4.2 (-0.8, 9.1)	-1.1 (-1.4, -0.8)	-0.1 (-0.3, 0.1)	0.436
Heat index, hrs > = 100°F	109.5 (100.7, 118.3)	6.7 (-0.1, 13.5)	-2.7 (-7.5, 2.2)	2.8 (-2.3, 8)	-9.2 (-16.8, -1.6)	-10.5 (-18.1, -2.9)	-1.1 (-1.4, -0.8)	3.6 (-1.5, 8.7)	-1.1 (-1.4, -0.8)	-11.6 (-62.5, 39.4)	0.652
HR mean at V1 <sup>e</sup> , cent95 <sup>d</sup>	108.5 (103, 114.1)	5.6 (-0.1, 11.2)	-2.8 (-7.6, 2.1)	2.8 (-2.3, 7.9)	-8.8 (-16.4, -1.2)	-10.6 (-18.1, -3)	-1.1 (-1.3, -0.8)	4.2 (-0.8, 9.3)	-1.1 (-1.3, -0.8)	-1.0 (-3, 1)	0.307
Workday physical activity mean at V1 <sup>e</sup>	110.5 (104.7, 116.4)	5.6 (0, 11.1)	-2.8 (-7.6, 2)	3.3 (-1.8, 8.4)	-8.8 (-16.3, -1.3)	-10.1 (-17.6, -2.6)	-1.1 (-1.3, -0.8)	1.5 (-3.9, 6.9)	-1.1 (-1.3, -0.8)	-0.06 (-0.1, 0)	0.024
<b>+ Hydration</b>	108.1 (102, 114.2)	5.6 (-0.5, 11.6)	-2.7 (-7.6, 2.3)	2.8 (-2.4, 8)	-9.3 (-17, -1.7)	-10.5 (-18.1, -2.8)	-1.1 (-1.3, -0.8)	3.9 (-1.2, 9)	-1.1 (-1.3, -0.8)	0 (-0.1, 0.1)	0.999
Beverage consumption, oz, cent64 <sup>d</sup>	109.2 (103.4, 114.9)	6.1 (0.5, 11.8)	-2.7 (-7.5, 2.1)	2.6 (-2.5, 7.7)	-9.2 (-16.7, -1.7)	-10.4 (-17.9, -2.9)	-1.1 (-1.3, -0.8)	3.5 (-1.6, 8.5)	-1.1 (-1.3, -0.8)	-2.6 (-6.3, 1.1)	0.174
Sugary drinks consumption, oz	108.3 (102.9, 113.6)	5.7 (0.2, 11.3)	-2.1 (-7, 2.8)	3.3 (-1.8, 8.5)	-9.4 (-17, -1.8)	-10.7 (-18.3, -3.1)	-1.0 (-1.3, -0.8)	2.8 (-2.3, 7.8)	-1.0 (-1.3, -0.8)	-0.4 (-0.8, 0)	0.065
Dehydration, both AM and PM <sup>g</sup>											
Serum osmolality, cent295 <sup>d</sup>											

<sup>a</sup>Additional covariates of interest were added to the base model one at a time.

<sup>b</sup>Bold text indicates relationship with associated P < 0.05.

<sup>c</sup>P values associated with the added covariate.

<sup>d</sup>Variables noted with "cent#%" were centered at the indicated value, thus allowing the intercept interpretation to be the estimated average value of after-work eGFR for a 40-year-old (at visit 1), female nursery worker in the summer of 2020 at the centered value of the additional covariate.

<sup>e</sup>V1 indicates "at visit 1, Winter 2020".

<sup>f</sup>Family history of diabetes, hypertension, or kidney disease.

<sup>g</sup>Urine specific gravity > = 1.020 both before and after work.

HRI, heat-related illness; NSAID, nonsteroidal anti-inflammatory drugs use.

**TABLE 5.** Wintertime Predictors of Postwork eGFR Evaluated Using Mixed Models With the Base Model Containing Primary Work Type and Timepoint, Controlling for Sex and Age

Model <sup>a</sup>	Intercept	Primary Work Type	Winter21	Sex	Age	Additional Covariate	P <sup>e</sup>
		(Ref = Nursery)	(Ref = Winter20)	(Ref = female)	(cent40) <sup>d</sup>		
	Estimated Beta (95% Confidence Interval) <sup>b</sup>						
<b>Base model</b>	112.8 (107.9, 117.6)	2 (-2.1, 6)	2.3 (-1, 5.5)	1.2 (-3.5, 6)	<b>-1.1 (-1.3, -0.8)</b>		
<b>+ Health factors</b>							
BMI (cent30) <sup>d</sup>	112.8 (107.9, 117.8)	1.9 (-2.2, 6)	2.3 (-1, 5.5)	1.2 (-3.6, 6)	<b>-1.1 (-1.4, -0.8)</b>	0 (-0.4, 0.4)	0.8951
BP 120–139 or 80–89 (ref = <120/80)	110.6 (105.2, 116.0)	2.8 (-1.3, 6.9)	1.2 (-2, 4.3)	2.1 (-2.7, 6.9)	<b>-1.1 (-1.3, -0.8)</b>	4.0 (-1.4, 9.5)	0.1473
bp140+ or 90+ (ref = <120/80)						2.2 (-2.1, 6.4)	0.3201
A1C >=5.7 at V1 <sup>e</sup>	112.1 (107.0, 117.2)	2.4 (-1.8, 6.5)	2.2 (-1.1, 5.5)	1.4 (-3.4, 6.2)	<b>-1.1 (-1.4, -0.8)</b>	2.6 (-2.7, 8)	0.3334
Resting heart rate	108.9 (95.3, 122.4)	2.0 (-2.1, 6)	2.4 (-0.9, 5.6)	1.0 (-3.9, 5.8)	<b>-1.1 (-1.3, -0.8)</b>	0.1 (-0.1, 0.2)	0.5426
NSAID use, sometimes/daily	113.0 (108.1, 117.9)	2.0 (-2.1, 6)	2.4 (-0.9, 5.7)	1.3 (-3.5, 6)	<b>-1.1 (-1.3, -0.8)</b>	-1.3 (-6.1, 3.4)	0.5845
Family hx kidney risk factors <sup>f</sup>	111.1 (105.6, 116.6)	2.1 (-2.1, 6.2)	2.1 (-1.2, 5.4)	1.3 (-3.7, 6.2)	<b>-1.1 (-1.4, -0.8)</b>	3.1 (-0.9, 7.1)	0.1296
<b>+ Work conditions</b>							
Training in HRI prevention	111.8 (106.9, 116.8)	1.7 (-2.3, 5.8)	2.3 (-1, 5.5)	1.1 (-3.6, 5.9)	<b>-1.1 (-1.4, -0.8)</b>	3.4 (-0.8, 7.5)	0.1118
Pesticide use, years	112.4 (107.0, 117.7)	2.1 (-2, 6.2)	2.3 (-1, 5.5)	1.6 (-3.6, 6.8)	<b>-1.1 (-1.4, -0.8)</b>	0.1 (-0.6, 0.8)	0.723
Workday duration	111.9 (106.9, 116.8)	3.2 (-1.1, 7.5)	2.3 (-1, 5.5)	1.5 (-3.3, 6.3)	<b>-1.1 (-1.3, -0.8)</b>	0.9 (-0.2, 1.9)	0.107
Shade breaks taken	112.1 (106.8, 117.3)	2.1 (-1.9, 6.2)	2.3 (-1, 5.5)	1.1 (-3.7, 5.9)	<b>-1.1 (-1.4, -0.8)</b>	1.3 (-2.4, 5.1)	0.483
Heat index mean (cent90) <sup>d</sup>	111.1 (103.8, 118.5)	2.1 (-2, 6.2)	1.8 (-1.7, 5.4)	1.5 (-3.4, 6.3)	<b>-1.1 (-1.3, -0.8)</b>	-0.1 (-0.3, 0.1)	0.5608
HR mean at V1 <sup>e</sup> (cent95) <sup>d</sup>	112.6 (107.7, 117.6)	2.1 (-2, 6.3)	2.3 (-1, 5.5)	1.3 (-3.5, 6.1)	<b>-1.1 (-1.4, -0.8)</b>	0 (-0.3, 0.2)	0.7682
Workday physical activity mean at V1 <sup>e</sup>	110.9 (103.9, 118.0)	0.9 (-4.2, 5.9)	2.3 (-1, 5.5)	1.6 (-3.3, 6.4)	<b>-1.1 (-1.3, -0.8)</b>	13.8 (-24.9, 52.5)	0.484
HRI symptoms	113.6 (108.5, 118.7)	1.8 (-2.3, 5.8)	1.9 (-1.4, 5.3)	1.1 (-3.7, 5.9)	<b>-1.1 (-1.4, -0.8)</b>	-1.4 (-4.4, 1.6)	0.3462
<b>+ Hydration</b>							
Beverage consumption, oz. cent64 <sup>d</sup>	113 (108.1, 117.9)	1.9 (-2.2, 5.9)	2.2 (-1.1, 5.4)	1.4 (-3.4, 6.3)	<b>-1.1 (-1.3, -0.8)</b>	0 (0, 0.1)	0.5398
Sugary drinks consumption, oz	111.4 (106.3, 116.5)	2.7 (-1.7, 7.2)	2.3 (-0.9, 5.5)	1.8 (-3, 6.6)	<b>-1.1 (-1.3, -0.8)</b>	0.1 (-0.1, 0.2)	0.4541
Dehydration both AM and PM <sup>g</sup>	113.7 (108.8, 118.7)	2.5 (-1.5, 6.5)	2.4 (-0.9, 5.7)	0.6 (-4.2, 5.3)	<b>-1.1 (-1.4, -0.9)</b>	-4.1 (-9.5, 1.4)	0.1417
Serum osmolality (cent295) <sup>d</sup>	112.5 (107.9, 117.1)	3.0 (-0.9, 6.8)	0.9 (-2.4, 4.2)	-1.7 (-6.4, 3.1)	<b>-1.0 (-1.3, -0.8)</b>	<b>-1.0 (-1.5, -0.5)</b>	<b>0.0001</b>

<sup>a</sup>Additional covariates of interest were added to the base model one at a time. A model with ‘Heat Index, hrs >=100°F’ was not calculated for winter data.

<sup>b</sup>Bold text indicates relationship with associated P < 0.05.

<sup>c</sup>P values associated with the added covariate.

<sup>d</sup>Variables noted with ‘cent#’ were centered at the indicated value, thus allowing the intercept interpretation to be the estimated average value of after-work eGFR for a 40-year-old (at visit 1), female nursery worker in the summer of 2020 at the centered value of the additional covariate.

<sup>e</sup>V1 indicates ‘at visit 1, Winter 2020’.

<sup>f</sup>Family history of diabetes, hypertension, or kidney disease.

<sup>g</sup>Urine specific gravity >= 1.020 both before and after work.

BMI, body mass index; HRI, heat-related illness; NSAID, nonsteroidal anti-inflammatory drugs use; BP, blood pressure.

declines in kidney function over the 32-month duration of this study, the marked drop in postwork eGFR at the latest study timepoints and the progressive increase in the percentage of fernery workers meeting the criteria for AKI after work raise concern. AKI and chronic degradation of renal function have been described as risk factors for the development of CKD.<sup>17,43,44</sup> However, most of these studies have focused on patients developing AKI in the hospital. One study in Guatemalan sugarcane workers, however, found that workers who experienced the greatest changes in serum creatinine levels—the marker used to calculate eGFR and define AKI in this study—over the course of their work shifts had more pronounced declines in eGFR over a harvest season.<sup>20</sup> To our knowledge, this is the first study of U.S. agricultural workers evaluating renal function over time, and we observed an increase in cross-shift variation in serum creatinine among fernery workers over the course of several harvest seasons. It is possible that recurrent episodes of AKI could contribute to the subsequent development of CKD, and eventually to end-stage renal disease. A longer period of follow-up will be necessary to determine whether the participants in our study who developed AKI progress to CKD. Established CKD inexorably progresses to worsening renal function and ultimately to the need for renal replacement therapy.

The cause of renal dysfunction, both CKDu and AKI, among agricultural workers remains unclear, but factors including exposure to high ambient temperatures, intense physical exertion, elevated core body temperature, and dehydration continue to be implicated.<sup>7,18–20</sup> It has also been postulated that the cause of renal dysfunction among agricultural workers may be multifactorial and that pesticides may have an important role to play.<sup>18,45</sup> A recent study in Mexico assessed kidney function over a 6-month harvest season in 50 agricultural

workers who labored in organic fields, 51 agricultural workers in conventional fields, and 50 office workers within the same region.<sup>18</sup> Although the organic and conventional groups had declines in eGFR, the conventional group with more exposure to pesticides had a more significant decline.<sup>18</sup> The office worker group did not show a decline in eGFR.<sup>18</sup> We have previously reported that while fernery workers perceive that they have high levels of pesticide exposure, urinary markers of organophosphate pesticides are actually higher among nursery workers.<sup>46</sup> In this study, we identified a significant inverse association between years of occupational pesticide exposure and preworkday eGFR, and preworkday eGFR was lower among nursery workers, supporting the idea that pesticides could negatively impact kidney function. Additionally, we identified a positive association between NSAID use and preworkday eGFR, with participants who reported taking NSAIDs more frequently having higher preworkday eGFR. NSAIDs have the potential for nephrotoxicity and have been proposed as a risk factor for the development of CKDu among agricultural workers, but significant associations have rarely been identified. Our results in fact suggest that NSAIDs could have some benefit for baseline kidney function, perhaps combatting inflammatory mechanisms contributing to AKI.<sup>47</sup> Further studies are needed to validate these findings and clarify the role of these risk factors in renal degradation.

In the meantime, minimization of risk factors during the workday to reduce the risk of AKI and the future development of CKD seems prudent. Hydrating with electrolyte solutions has been reported as a potential measure to reduce the development of AKI and renal dysfunction in agricultural workers.<sup>48,49</sup> The piece-rate compensation system seems to be the factor that incentivizes workers to eschew breaks

**TABLE 6.** Summertime Predictors of Prework eGFR Evaluated Using Mixed Models With the Base Model Containing Primary Work Type and Timepoint, Controlling for Sex and Age

Model <sup>a</sup>	Intercept	Primary Work Type	Time		Sex	Age	Additional Covariate	P <sup>c</sup>
		(Ref = Nursery)	Summer21	Summer22	(Ref = Female)	(cent40) <sup>d</sup>		
			(Ref = Summer20)	(Ref = Summer20)				
<b>Base model</b>	<b>114.4 (111.1, 117.8)</b>	<b>3.6 (0.7, 6.5)</b>	-0.7 (-2.2, 0.8)	0.8 (-0.7, 2.3)	<b>4.6 (1.3, 7.8)</b>	<b>-1.0 (-1.2, -0.9)</b>		
<b>+ Health factors</b>								
BMI cent30 <sup>d</sup>	<b>114.4 (111.0, 117.7)</b>	<b>3.6 (0.7, 6.6)</b>	-0.7 (-2.2, 0.8)	0.8 (-0.7, 2.3)	<b>4.6 (1.3, 7.8)</b>	<b>-1.0 (-1.2, -0.8)</b>	0 (-0.3, 0.2)	0.782
BP 120–139 or 80–89 (ref = <120/80)	<b>115.5 (111.7, 119.2)</b>	<b>3.7 (0.8, 6.6)</b>	-0.8 (-2.3, 0.8)	0.7 (-0.9, 2.2)	<b>5.0 (1.7, 8.2)</b>	<b>-1.0 (-1.2, -0.8)</b>	-1.6 (-4.3, 1.0)	0.221
BP140+ or 90+ (ref = <120/80)							-1.5 (-4.0, 1.0)	0.242
A1C > =5.7 at V1 <sup>e</sup>	<b>114.0 (110.6, 117.5)</b>	<b>3.8 (0.9, 6.8)</b>	-0.7 (-2.2, 0.8)	0.8 (-0.7, 2.3)	<b>4.6 (1.4, 7.8)</b>	<b>-1.0 (-1.2, -0.9)</b>	1.4 (-2.3, 5.0)	0.456
Resting heart rate	<b>115.2 (111.9, 118.5)</b>	<b>3.5 (0.8, 6.3)</b>	-0.1 (-1.6, 1.3)	1.8 (0.2, 3.4)	<b>3.8 (0.6, 7.0)</b>	<b>-1.0 (-1.2, -0.9)</b>	<b>0.1 (0.1, 0.2)</b>	<b>0.001</b>
NSAID use, sometimes or daily	<b>114.3 (111.0, 117.6)</b>	<b>3.5 (0.6, 6.4)</b>	-0.7 (-2.1, 0.8)	0.8 (-0.7, 2.3)	<b>4.3 (1.1, 7.5)</b>	<b>-1.1 (-1.2, -0.9)</b>	<b>2.2 (0, 4.3)</b>	<b>0.049</b>
Family hx kidney risk factors <sup>f</sup>	<b>113.9 (110.2, 117.6)</b>	<b>3.6 (0.7, 6.5)</b>	-0.8 (-2.3, 0.7)	0.6 (-0.9, 2.2)	<b>3.9 (0.7, 7.1)</b>	<b>-1.0 (-1.2, -0.8)</b>	2.5 (-0.2, 5.1)	0.069
Urine specific gravity, prework, cent1.020 <sup>d</sup>	<b>114.4 (111.0, 117.8)</b>	<b>3.5 (0.6, 6.5)</b>	-0.7 (-2.3, 0.8)	0.7 (-0.9, 2.3)	<b>5.0 (1.8, 8.3)</b>	<b>-1.0 (-1.2, -0.8)</b>	1.2 (0, 2.5)	0.053
<b>+ Work conditions</b>								
Training in HRI prevention	<b>114.0 (110.6, 117.4)</b>	<b>3.4 (0.6, 6.3)</b>	-0.7 (-2.2, 0.8)	0.8 (-0.7, 2.3)	<b>4.5 (1.3, 7.7)</b>	<b>-1.1 (-1.2, -0.9)</b>	1.5 (-1.4, 4.3)	0.313
Pesticide use, years	<b>116.2 (112.5, 119.8)</b>	<b>3.3 (0.4, 6.1)</b>	-0.7 (-2.2, 0.8)	0.7 (-0.8, 2.3)	3.0 (-0.5, 6.5)	<b>-1.0 (-1.2, -0.9)</b>	<b>-0.4 (-0.9, 0)</b>	<b>0.042</b>

<sup>a</sup>Additional covariates of interest were added to the base model one at a time.

<sup>b</sup>Bold text indicates relationship with associated P < 0.05.

<sup>c</sup>P values associated with the added covariate.

<sup>d</sup>Variables noted with 'cent#' were centered at the indicated value, thus allowing the intercept interpretation to be the estimated average value of after-work eGFR for a 40-year-old (at visit 1), female nursery worker in the summer of 2020 at the centered value of the additional covariate.

<sup>e</sup>V1 indicates 'at visit 1, Winter 2020'.

<sup>f</sup>Family history of diabetes, hypertension, or kidney disease.

BMI, body mass index; HRI, heat-related illness; NSAID, nonsteroidal anti-inflammatory drugs use.

and better hydration with the goal of earning more money, which appears to increase risk of renal dysfunction. Thus, it is necessary to re-evaluate the piece-rate system and integrate safeguards to protect

workers or eliminate the system altogether. Further research is also needed to find more evidence-based solutions to protect workers from kidney dysfunction.

**TABLE 7.** Wintertime Predictors of Prework eGFR Evaluated Using Mixed Models With the Base Model Containing Primary Work Type and Timepoint, Controlling for Sex and Age

Model <sup>a</sup>	Intercept	Primary Work Type	Winter21	Sex	Age	Additional Covariate	P <sup>c</sup>
		(Ref = Nursery)	Winter21	(Ref = female)	(cent40) <sup>d</sup>		
			(Ref = Winter20)				
<b>Base model</b>	<b>117.8 (115, 120.6)</b>	<b>4.0 (1.4, 6.6)</b>	-0.3 (-1.7, 1.1)	2.5 (-0.3, 5.2)	<b>-1.0 (-1.2, -0.8)</b>		
<b>+ Health factors</b>							
BMI cent30 <sup>d</sup>	<b>118 (115.1, 120.8)</b>	<b>3.8 (1.2, 6.5)</b>	-0.3 (-1.7, 1.0)	2.3 (-0.5, 5.1)	<b>-1.0 (-1.2, -0.8)</b>	0.1 (-0.1, 0.3)	0.390
BP 120–139 or 80–89 (ref = <120/80)	<b>118 (114.8, 121.1)</b>	<b>3.8 (1, 6.6)</b>	-0.7 (-2.1, 0.7)	2.6 (-0.2, 5.5)	<b>-1.0 (-1.2, -0.8)</b>	-1.1 (-3.5, 1.2)	0.330
BP140+ or 90+ (ref = <120/80)						0.7 (-2.1, 3.5)	0.613
A1C > =5.7 at V1 <sup>e</sup>	<b>117.4 (114.5, 120.3)</b>	<b>4.2 (1.5, 6.8)</b>	-0.4 (-1.8, 1.0)	2.5 (-0.2, 5.3)	<b>-1.0 (-1.2, -0.8)</b>	1.7 (-1.5, 4.9)	0.296
Resting heart rate	<b>118 (115.1, 120.8)</b>	<b>4.0 (1.4, 6.6)</b>	-0.3 (-1.7, 1.1)	2.2 (-0.6, 5.0)	<b>-1.0 (-1.1, -0.8)</b>	0 (0, 0.1)	0.312
NSAID use, sometimes or daily	<b>117.8 (114.9, 120.6)</b>	<b>4.0 (1.4, 6.6)</b>	-0.4 (-1.8, 1.1)	2.5 (-0.3, 5.2)	<b>-1.0 (-1.2, -0.8)</b>	0.2 (-2.1, 2.5)	0.847
Family hx kidney risk factors <sup>f</sup>	<b>116.8 (113.7, 120)</b>	<b>3.9 (1.2, 6.5)</b>	-0.3 (-1.8, 1.1)	2.1 (-0.7, 4.9)	<b>-1.0 (-1.2, -0.9)</b>	2.3 (-0.1, 4.7)	0.055
Urine specific gravity, prework, cent1.020	<b>117.5 (114.7, 120.4)</b>	<b>3.9 (1.3, 6.5)</b>	-0.5 (-2.0, 0.9)	<b>3.0 (0.2, 5.9)</b>	<b>-1.0 (-1.2, -0.8)</b>	0 (-1.4, 1.5)	0.949
<b>+ Work conditions</b>							
Training in HRI prevention	<b>118.1 (115.2, 121)</b>	<b>4.0 (1.4, 6.7)</b>	-0.3 (-1.7, 1.0)	2.5 (-0.2, 5.2)	<b>-1.0 (-1.1, -0.8)</b>	-0.9 (-3.5, 1.6)	0.468
Pesticide use, years	<b>118.0 (114.9, 121.1)</b>	<b>3.9 (1.3, 6.6)</b>	-0.3 (-1.7, 1.1)	2.3 (-0.7, 5.2)	<b>-1.0 (-1.2, -0.8)</b>	-0.1 (-0.5, 0.3)	0.716

<sup>a</sup>Additional covariates of interest were added to the base model one at a time.

<sup>b</sup>Bold text indicates relationship with associated P < 0.05.

<sup>c</sup>P values associated with the added covariate.

<sup>d</sup>Variables noted with 'cent#' were centered at the indicated value, thus allowing the intercept interpretation to be the estimated average value of after-work eGFR for a 40-year-old (at visit 1), female nursery worker in the summer of 2020 at the centered value of the additional covariate.

<sup>e</sup>V1 indicates 'at visit 1, Winter 2020'.

<sup>f</sup>Family history of diabetes, hypertension, or kidney disease.

BMI, body mass index; HRI, heat-related illness; NSAID, nonsteroidal anti-inflammatory drugs use.

Downloaded from http://journals.lww.com/joem by BMDMSEPHKav1ZEoum1tQIN4e+kLHEZgbsHio4XM10hCwCX1AW nYQp/llQHD33D00dRv7TVSfI4C3VC1y0abgqZQZdgG12MwIzlel= on 09/06/2024

## Strengths and Limitations

A few workers from each location reported working in an agricultural environment other than a nursery or fernery. Furthermore, it is not uncommon for agricultural workers to assume different responsibilities and work in different environments over time. This is also a reflection of the different economic strategies that agricultural workers rely on over time to secure a livelihood under precarious job security situations. Understanding the dynamic nature of this occupation, we decided not to restrict this study to participants only working at a fernery or nursery, even though this necessitated an imperfect comparison of these agricultural settings.

Strengths of this study are that we recorded beverage consumption with a postworkday structured interview using props to demonstrate sizes; we followed workers for 32 months, had good retention both in summer and winter visits, and had a good range of heat indices that strengthened power to detect effect. Although we were able to examine many of the risk factors proposed in the literature, we do not know the types of pesticides or the level of exposure. There also was not much power to detect effects for AKI because of the relatively small number of events. It should also be noted that the classification of AKI was based on changes in serum creatinine, consistent with standard clinical practice, and not confirmed by additional clinical testing such as imaging or biopsies. Temperature and humidity are not from the work sites themselves and may not account for the micro-environment of the worker (inside greenhouse, under shade cloth, etc), nor was core body temperature assessed in this study. This is a limitation because it is a combination of environmental heat exposure as well as physical exertion, which results in hyperthermia and the physiological responses that increase the risk of AKI.

## CONCLUSIONS

To our knowledge, this is the first study of agricultural workers in the US evaluating renal function over time. We found relatively stable preworkday eGFR over the course of the study, with slight decreases in summers, but found increasing incidence of AKI and lower eGFR postworkday as time progressed among the piece-rate workers who are incentivized to forgo rest breaks and hydration, even with high ambient temperature. Workers became more dehydrated in summer despite increased fluid consumption because the increase was not commensurate with their need. Further research is needed to find evidence-based solutions to improve hydration and protect workers from kidney dysfunction.

## REFERENCES

- Glaser J, Lemery J, Rajagopalan B, et al. Climate change and the emergent epidemic of CKD from heat stress in rural communities: the case for heat stress nephropathy. *Clin J Am Soc Nephrol* 2016;11:1472–1483.
- Johnson RJ, Sánchez-Lozada LG, Newman LS, et al. Climate change and the kidney. *Ann Nutr Metab* 2019;74(Suppl 3):38–44.
- Smith DJ, Pius LM, Plantinga LC, Thompson LM, Mac V, Hertzberg VS. Heat stress and kidney function in farmworkers in the US: a scoping review. *J Agromedicine* 2022;27:183–192.
- Sasai F, Roncal-Jimenez C, Rogers K, et al. Climate change and nephrology. *Nephrol Dial Transplant* 2023;38:41–48.
- Chapman CL, Hess HW, Lucas RAI, et al. Occupational heat exposure and the risk of chronic kidney disease of nontraditional origin in the United States. *Am J Physiol Regul Integr Comp Physiol* 2021;321:R141–r151.
- Schlader ZJ, Hostler D, Parker MD, et al. The potential for renal injury elicited by physical work in the heat. *Nutrients* 2019;11:2087.
- Wesseling C, Glaser J, Rodríguez-Guzmán J, et al. Chronic kidney disease of non-traditional origin in Mesoamerica: a disease primarily driven by occupational heat stress. *Rev Panam Salud Publica* 2020;44:e15.
- Sanchez Polo V, Garcia-Trabanino R, Rodriguez G, Madero M. Mesoamerican nephropathy (MeN): what we know so far. *Int J Nephrol Renovasc Dis* 2020;13: 261–272.
- Sorensen CJ, Krisher L, Butler-Dawson J, et al. Workplace screening identifies clinically significant and potentially reversible kidney injury in heat-exposed sugarcane workers. *Int J Environ Res Public Health* 2020;17:8552.

- Laws RL, Brooks DR, Amador JJ, et al. Biomarkers of kidney injury among Nicaraguan sugarcane workers. *Am J Kidney Dis* 2016;67:209–217.
- Wesseling C, Aragon A, Gonzalez M, et al. Kidney function in sugarcane cutters in Nicaragua—a longitudinal study of workers at risk of Mesoamerican nephropathy. *Environ Res* 2016;147:125–132.
- Hansson E, Glaser J, Jakobsson K, et al. Pathophysiological mechanisms by which heat stress potentially induces kidney inflammation and chronic kidney disease in sugarcane workers. *Nutrients* 2020;12:1639.
- Butler-Dawson J, Dally M, Johnson RJ, et al. Association of copeptin, a surrogate marker of arginine vasopressin, with decreased kidney function in sugarcane workers in Guatemala. *Ann Nutr Metab* 2020;76:30–36.
- Butler-Dawson J, Krisher L, Yoder H, et al. Evaluation of heat stress and cumulative incidence of acute kidney injury in sugarcane workers in Guatemala. *Int Arch Occup Environ Health* 2019;92:977–990.
- Dally M, Butler-Dawson J, Cruz A, et al. Longitudinal trends in renal function among first time sugarcane harvesters in Guatemala. *PLoS One* 2020;15:e0229413.
- Hansson E, Glaser J, Weiss I, et al. Workload and cross-harvest kidney injury in a Nicaraguan sugarcane worker cohort. *Occup Environ Med* 2019;76:818–826.
- Kurzhaagen JT, Dellepiane S, Cantaluppi V, Rabb H. AKI: an increasingly recognized risk factor for CKD development and progression. *J Nephrol* 2020; 33:1171–1187.
- López-Gálvez N, Wagoner R, Canales RA, et al. Longitudinal assessment of kidney function in migrant farm workers. *Environ Res* 2021;202:111686.
- Nagai K. Environment and chronic kidney disease in farmers. *Renal Replacement Therapy* 2021;7:55.
- Dally M, Butler-Dawson J, Johnson RJ, et al. Creatinine fluctuations forecast cross-harvest kidney function decline among sugarcane workers in Guatemala. *Kidney Int Rep* 2020;5:1558–1566.
- Mix J, Elon L, Thein Mac VV, et al. Hydration status, kidney function, and kidney injury in Florida agricultural workers. *J Occup Environ Med* 2018;60: e253–e260.
- Moyce S, Armitage T, Mitchell D, Schenker M. Acute kidney injury and workload in a sample of California agricultural workers. *Am J Ind Med* 2020; 63:258–268.
- Kupferman J, Ramirez-Rubio O, Amador JJ, et al. Acute kidney injury in sugarcane workers at risk for Mesoamerican nephropathy. *Am J Kidney Dis* 2018;72:475–482.
- Garcia-Trabanino R, Jarquin E, Wesseling C, et al. Heat stress, dehydration, and kidney function in sugarcane cutters in El Salvador - a cross-shift study of workers at risk of Mesoamerican nephropathy. *Environ Res* 2015;142:746–755.
- Khwaja A. KDIGO clinical practice guidelines for acute kidney injury. *Nephron Clin Pract* 2012;120:c179–c184.
- Anand S, Staniec A, Montez-Rath M, Vlahos P. Using GIS Mapping to Track Hot Spots of Kidney Disease in California. *N Engl J Med* 2020;382:2265–2267.
- Bragg-Gresham J, Morgenstern H, Shahinian V, Robinson B, Abbott K, Saran R. An analysis of hot spots of ESRD in the United States: Potential presence of CKD of unknown origin in the USA? *Clinical nephrology* 2020;93:113–119.
- Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research Electronic Data Capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–381.
- Lu Y-C, Roms DM. Extending the heat index. *J Appl Meteorol Climatol* 2022; 61:1367–1383.
- Aktas E, Bergbom B, Godderis L, et al. Migrant workers occupational health research: an OMEGA-NET working group position paper. *Int Arch Occup Environ Health* 2022;95:765–777.
- Arcury TA, Quandt SA. Delivery of health services to migrant and seasonal farmworkers. *Annu Rev Public Health* 2007;28:345–363.
- Bloss JE, LePrevost CE, Zahra AG, et al. Advancing the health of migrant and seasonal farmworkers in the United States: identifying gaps in the existing literature, 2021. *Health Promot Pract* 2022;23:432–444.
- Inker LA, Eneanya ND, Coresh J, et al. New creatinine- and cystatin C-based equations to estimate GFR without race. *N Engl J Med* 2021;385:1737–1749.
- Delgado C, Baweja M, Crews DC, et al. A unifying approach for GFR estimation: recommendations of the NKF-ASN Task Force on reassessing the inclusion of race in diagnosing kidney disease. *Am J Kidney Dis* 2022;79:268–288.e1.
- Wesseling C, Crowe J, Hogstedt C, Jakobsson K, Lucas R, Wegman DH. Resolving the enigma of the Mesoamerican nephropathy: a research workshop summary. *Am J Kidney Dis* 2014;63:396–404.
- Geraci M. Linear quantile mixed models: the lqmm package for Laplace quantile regression. *J Stat Softw* 2014;57:1–29.
- Geraci M, Bottai M. Linear quantile mixed models. *Stat Comput* 2014;24:461–479.
- van Buuren S, Groothuis-Oudshoorn K. Mice: multivariate imputation by chained equations in R. *J Stat Softw* 2011;45:1–67.
- SAS Institute Inc. *SAS/STAT 13.1 User's Guide*. SAS Institute Inc, Cary, NC; 2013.

40. Abu I, Elon L, Xiuhtecutli N, McCauley L, Chicas R. Retention of agricultural workers participating in a renal longitudinal study. *J Agromedicine* 2024;29:26–33.
41. Medtronic. Measure performance in a new way. Medtronic; 2016. Available at: <https://www.zephyranywhere.com/media/download/zephyr-performance-biopatch-hp-brochure.pdf>. Accessed August 12, 2023.
42. García-Arroyo FE, Cristóbal M, Arellano-Buendía AS, et al. Rehydration with soft drink-like beverages exacerbates dehydration and worsens dehydration-associated renal injury. *Am J Physiol Regul Integr Comp Physiol* 2016;311:R57–R65.
43. Coca SG, Singanamala S, Parikh CR. Chronic kidney disease after acute kidney injury: a systematic review and meta-analysis. *Kidney Int* 2012;81:442–448.
44. Goldstein SL, Jaber BL, Faubel S, Chawla LS. AKI transition of care: a potential opportunity to detect and prevent CKD. *Clin J Am Soc Nephrol* 2013;8:476–483.
45. Valcke M, Levasseur ME, Soares da Silva A, Wesseling C. Pesticide exposures and chronic kidney disease of unknown etiology: an epidemiologic review. *Environ Health* 2017;16:49.
46. Runkle J, Tovar-Aguilar JA, Economos E, et al. Pesticide risk perception and biomarkers of exposure in Florida female farmworkers. *J Occup Environ Med* 2013;55:1286–1292.
47. Houser MC, Mac V, Smith DJ, et al. Inflammation-related factors identified as biomarkers of dehydration and subsequent acute kidney injury in agricultural workers. *Biol Res Nurs* 2021;23:676–688.
48. Chicas R, Suarez J, Elon L, et al. Hydration interventions among agricultural workers: a pilot study. *J Occup Environ Med* 2022;64:e357–e359.
49. Krisher L, Butler-Dawson J, Yoder H, et al. Electrolyte beverage intake to promote hydration and maintain kidney function in Guatemalan sugarcane workers laboring in hot conditions. *J Occup Environ Med* 2020;62:e696–e703.