CDC INFLUENZA REPORT NO. 12 AUGUST 16, 1957

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE Public Health Service Bureau of State Services

> Communicable Disease Center Robert J. Anderson, Chief

Keith E. Jensen, Ph. D. CDC Virus and Rickettsia Section^{*} P. O. Box 61 Montgomery 1, Alabama Telephone No. AMherst 3-4468

*Serving as WHO International Influenza Center for the Americas Robert H. Drachman, M. D. Frederick L. Dunn, M. D. Influenza Surveillance Unit Communicable Disease Center 50 Seventh Street, N. E. Atlanta 23, Georgia Telephone No. TRinity 6-3311 Extension 5455

SPECIAL NOTE

Information contained in this report is a summary of data reported to CDC by State Health Departments, Epidemic Intelligence Service Officers, collaborating influenza diagnostic laboratories, and other pertinent sources. Much of it is preliminary in nature and is primarily intended for those involved in influenza control activities. It is understood that the contents of these reports will not be released to the press, except by the Office of the Surgeon General, Public Health Service, U. S. Department of Health, Education, and Welfare. State Health Officers, of course, will judge the advisability of releasing any information from their own states.

Table of Contents

- I. Summary of Information
- II. Epidemic and Case Reports
- III. Progress Reports
- IV. Influenza Virus Center Information
 - V. Reports of Influenza-Associated Deaths
- VI. Summary Tables Cases and Outbreaks
- VII. Proposed Voluntary Vaccine Allocation System. News Release from the Surgeon General

I. Summary of Information

Included in this Report is a daily record of patients seen with influenza-like illnesses at Charity Hospital, in New Orleans. It is hoped that eventually, by comparison with previous years' figures, some estimation of current morbidity due to influenza may be obtained. Additional enumerations from other areas will be added as they become available.

Evidence of infection with Asian strain has been reported from Oregon, with a camp outbreak of febrile respiratory disease also noted in the same area. An Oregon Boy Scout, who attended the Jamboree at Valley Forge, yielded this strain, also.

Asian strain influenza virus has been isolated from cases in an upstate New York labor camp and from foreign exchange students in New York City. One death has occurred in this latter group.

Probably the first report this season of influenza virus isolated from a fatal case of complicating staphylococcal pneumonia has come from California.

Evidence of influenza infection due to an A-prime virus has been obtained from a sporadic case in New Orleans, Louisiana.

Seed virus and fluid antigen of an egg-animal-line of influenza virus has been distributed by the Influenza Virus Center at Montgomery, Alabama. It is expected that hemagglutination-inhibition tests will be facilitated by this material.

Asian strain virus has been isolated from influenza patients in Colombia, South America. A/Denver/57 related strains have been implicated in several influenza cases which occurred in Saskatchewan, Canada.

As has been frequently observed recently, Asian strain influenza virus is widely "seeded" over the country. The mildness of the illness, coupled with the infrequency of diagnostic specimens, probably has resulted in only a very small percentage of cases being identified. What holds the virus in abeyance at this season is far from clear.

II. Epidemic and Case Reports

12-A. OREGON

(Reported by Drs. S. B. Osgood and M. A. Holmes, Oregon State Board of Health.)

Serologic evidence of infection with Asian strain influenza virus has been obtained in two sporadic cases occurring in military personnel. These men were on leave in Oregon from their California bases. Both C-F and H-I tests were performed on paired sera. 12-B. OREGON, Delake

(Reported by Drs. S. B. Osgood and M. A. Holmes, Oregon State Board of Health.)

An outbreak of influenza-like illness occurred at a girls' camp near Delake, Oregon. The camp was in session between July 26 and August 16. One hundred thirty-eight campers and 23 staff members were present. The campers, all from the Pacific Northwest, were girls between 8-13 years old.

On July 29 a Portland, Oregon, camper became ill 24 hours after arrival at the camp and 48 hours after exposure to her brother, who had just returned from the Valley Forge Boy Scout Jamboree. On August first the next campers became ill. By the peak day, August 6, approximately 10 staff members and 40 campers had been affected. The outbreak subsided rapidly after the peak day. The total attack rate was about 33%.

The illness was characterized by sudden onset, after an apparent incubation period of about 48 hours. Fever and conjunctivitis (palpebral injection) were noted in 100% of the cases. Headache, myalgia, backache, dry cough, and malaise were prominent. Gastrointestinal symptoms were infrequent. Duration of illness averaged 3 days, with a range of 2 to 7 days.

Blood specimens, but not throat washings, will be available for diagnostic studies.

12-C. MICHIGAN, Coldwater

(Reported by Dr. G. H. Agate, Michigan Department of Health, and Dr. Gordon Brown, University of Michigan.)

Soon after meeting a friend just arrived from Mexico City with a febrile respiratory disease, this Michigan woman developed typical influenza symptoms. Asian strain influenza virus was subsequently isolated from her throat washings obtained during the acute illness.

12-D. LOUISIANA, Livingston Parish

(Reported by Dr. J. D. Martin, Louisiana Department of Health.)

The school involved in this outbreak is only a few miles from Tangipahoa Parish (see CDC Influenza Report No. 10-A, 11-G) where influenza due to Asian strain virus has occurred.

School enrollment here is about 450. On August 13, 143 children were absent primarily because of febrile respiratory disease. About 125 were absent on August 14. Appropriate specimens are presently under laboratory study. As in Tangipahoa Parish, the lower socioeconomic groups are most heavily affected. Crowding and poor sanitary facilities are marked in this group. 12-E. LOUISIANA, New Orleans

(Reported by Dr. J. D. Martin, Louisiana Department of Health, and Dr. W. J. Mogabgab, Tulane University School of Medicine.)

A 26-year-old negro male was admitted to Charity Hospital on July 5 and discharged on July 9, with a clinical diagnosis of influenza. Acute and convalescent blood specimens were collected on July 6 and August 1. The hemagglutination-inhibition titer against A/Denver/57 rose from less than 1:4 to 1:64 in the convalescent specimen.

12-L. LOUISIANA, Baton Rouge

(Reported by Dr. J. D. Martin, Louisiana Department of Health.)

A sharp outbreak of influenza-like illness developed on August 14 in a Baton Rouge orphanage. On August 15, 61 children were reported ill and throat washings and blood specimens were obtained for laboratory study.

12-M. MICHIGAN, Bay County

(Reported by Dr. G. H. Agate, Michigan Department of Health, and Dr. G. Brown, University of Michigan.)

Migrant workers, travelling by bus from Mexico, were ill with febrile respiratory disease upon arrival in Eay County. Throat washings collected from acute cases on July 31 revealed Asian strain influenza virus.

(This report not noted in current summary charts.)

Report from Charity Hospital, New Orleans

Total Number of Patients Seen in the Admitting Room and Number with Influenza-like Illness

(Data provided by Charity Hospital administration through Dr. J. D. Martin, Louisiana State Department of Health)

Date	No. With influenza-like illness	No. seen	% with influenza-like illness
Aug. 4, 1957	25	304	8.0%
Aug. 5	35	642	5.5%
Aug. 6	33	500	6.0%
Aug. 7	34	470	7.2%
Aug. 8	33	398	8.3%
Aug. 12	89	710	12.5%

The above tabulation, when interpreted within the necessary limitations, may over the course of weeks give an estimation of influenza morbidity in New Orleans.* Patients visiting the Charity Hospital admitting room are usually of a low socioeconomic group, a population segment found most sus-

*Similar figures from other city hospitals might be equally valuable.

ceptible to epidemic influenza. Consequently, they provide a very sensitive index of influenza in an area. What relationship these figures may have to over-all community influenza morbidity is not easily determined. As absentee reports from schools and industry are available for comparison, this problem may be partially illuminated.

III. Progress Reports

12-F. NEW YORK, Cayuga County (See CDC Influenza Report No. 10-F)

- 5 -

(Reported by Dr. J. L. Freitag, New York Department of Health.)

Asian strain influenza virus has been isolated from throat washings obtained at the larger of the two previously reported labor camps.

12-G. NEW YORK, New York City (See CDC Influenza Report No. 11-B)

(Reported by Dr. H. Fuerst, New York City Health Department.)

Among the exchange students who arrived in New York City in early August were about 44 Turkish students, who had traveled on the Orient Express from Turkey to Western Europe. Cases of influenza-like illness broke out in this group in Belgrade and still more when they reached Vienna. Here an Austrian contingent boarded the train. By the time the students reached Rotterdam, 15 of the Turkish students were ill. The Turkish group was refused ship passage because of the illness. They therefore traveled to Amsterdam by bus and flew to New York, arriving August 7. Nine were ill on arrival and 3 were hospitalized. Asian strain virus has been isolated from two of these cases.

On August 12, more than 400 exchange students arrived in New York harbor on the Arosa Sky, with 50 active influenza cases. The Austrian contingent was the first to develop symptoms during the voyage. The students all traveled tourist class. About 740 other persons were on board the ship, including 200 first-class passengers and 200 crew members. Only one case occurred in each of these last groups.

The death of an 18-year-old exchange student is noted in section V of this Report.

The students will soon travel to many different sections of the United States to take up residence in private homes.

12-H. Boy Scout Jamboree (See CDC Influenza Report Nos. 3-D, 4-H, 5-F, 6-E, 7-C, 8-C, 11-F)

A report from Dr. M. E. Rindge, Connecticut Department of Health, states that serologic evidence of infection with Asian strain influenza virus has been obtained from a Boy Scout who became ill upon return from the Jamboree at Valley Forge, Pennsylvania. Sera collected on July 23 and August 1 were tested for hemagglutination-inhibition activity. A 16-fold increase was detected for Asian strain; an 8-fold increase for PR8; and a 4-fold increase for A/Denver/57.

Dr. D. S. Fleming, Minnesota Department of Health, reports that of 1423 Boy Scouts from the state who attended the Jamboree, about 206 were ill with influenza. About 66 secondary cases have occurred. Of 234 scouts queried by questionnaire, 34 had been ill, with 22 illnesses after return home.

Drs. S. B. Osgood and M. A. Holmes, Oregon State Health Department, have reported isolation of Asian strain influenza virus from a Boy Scout, ill on July 30, who returned home from the Jamboree on July 27. About 350 Boy Scouts returned to Portland on this latter date and, within three days, 3 scouts reported febrile respiratory disease, as did 4 family contacts.

12-I. CALIFORNIA, Monterey (See CDC Influenza Report No. 1-H, 2-F)

(Reported by Dr. P. K. Condit, California Department of Public Health.)

Among the more than 4000 influenza cases reported in late June and early July at Fort Ord, one death occurred and is noted in section V of this Report. Furthermore, 9 cases of lobar pneumonia were noted as complications in this group.

12-J. CALIFORNIA, Fresno

(Reported by Dr. R. M. Moldenhauer and Dr. J. Culver, California Department of Public Health.)

Asian strain influenza virus has now been isolated from cases of influenza that occurred in the Fresno County jail population in mid-July.

IV. Influenza Virus Center Information

Lyophilized seed virus and 15 ml. of fluid antigen of an egg-animalline of A/Japan 305/57 has been sent to all Influenza Collaborating Laboratories. As previously noted, this line of virus is more sensitive to specific antibody and should increase the usefulness and sensitivity of the hemagglutination-inhibition test when testing for Asian strain antibody. More complete details of this virus-line and necessary serum treatment are presented in CDC Influenza Report No. 8, section IV.

V. Reports of Influenza-Associated Deaths

(Reported by Drs. R. Moldenhauer and J. Culver, California Department of Public Health.)

Cal. 8 During the course of the influenza outbreak at a military

installation (CDC Influenza Report No. 1-H, 2-F) a young adult male developed an influenza-like illness and expired within 24 hours of onset. X-ray examination of the chest shortly before death showed signs of a left-sided bacterial pneumonia. Post-mortem examination revealed bilateral consolidation with Micrococcus pyogenes var. aureus cultured from lung tissue. Asian strain influenza virus has been isolated from bronchial mucosa. These specimens were obtained at autopsy and frozen prior to virus isolation.

N.Y..l An 18-year-old boy developed symptoms typical of influenza soon after he arrived with a group of foreign exchange students (CDC Influenza Report No. 11-B) in New York City. He was hospitalized the same day but expired within 24 hours. Post-mortem examination revealed a hemorrhagic pneumonitis. A specimen of lung tissue was frozen for virus isolation.

Additional Notes

Dr. A. Saenz, Pan American Sanitary Bureau, reported that Asian strain virus has been isolated from throat washings received at the WHO Influenza Center, Montgomery, Alabama, from Colombia, South America.

지 나 옷 가슴 것

Notification has been received at the WHO Influenza Center from Dr. F. P. Nagler, that the Laboratory of Hygiene in Ottawa, Canada, has recently isolated several strains of influenza virus closely related to A/Denver/57 strain. The illnesses occurred during July in Regina, Saskatchewan.

VII. Proposed Volumtary Vaccine Allocation System.

News Release from the Surgeon General, August 16, 1957.

The Public Health Service has proposed a voluntary interstate allocation system for the distribution of influenza vascine to the six licensed influenza vaccine manufacturers, Surgeon General Leroy E. Burney announced today.

The purpose of the system would be to assure "an equitable availability of vaccine supplies throughout all parts of the country," Dr. Burney said. He added that several of the companies already have informally indicated a willingness to go along with the plan so long as the vaccine remains in short supply.

The proposed voluntary system would be patterned along the following lines:

1. Each State would be entitled to shipments of a percentage of vaccine produced by each manufacturer equal to the percentage which that State's population bears to the total population of the United States;

2. Out of each week's vaccine production, manufacturers would establish for each State a quota of vaccine equal to the State's percentage entitlement;

3. Vaccine manufacturers would limit their sale of vaccine in each State to the quota established for that State;

4. Manufacturers would report weekly to the Public Health Service the net amount of vaccine released for sale that week and, on a State by State basis, the amount shipped into each State in terms of amounts shipped to public agencies and to other purchasers;

5. The Public Health Service would maintain data from the States regarding the adequacies of available vaccine supplies under the system and would recommend modifications in the quotas if it appeared that supplies were exceeding demand in some States.

Dr. Burney said that under this system the Public Health Service would not contemplate any allocation between public agency purchasers and commercial sales.

VII. Summary Tables - Cases and Outbreaks

TABLE I

Confirmed Outbreaks and Cases of Influenza Due to Asian Strains, United States June

I,		
	4773.2	
i	1957	
1	Š	
	<u></u>	
l		
ŀ	t 16,	
	m	
	جد	
	ġ	
ł	2	
	3	
	A	
	August	
	Н	
	ക	

	CDC Influenza Report Number	1-4		Ċ	H-	: 17	1-F 6-Note	6÷A	1-B 2-G	1 +B		2-4 4-F, 9-C
	nosis by Serology	Yes			Yes	Tes	Tes	Yes		Yes		Tes
	Lab. Diagnosis by Virus Isolation Serolog		Y⊕ (6-21-57)	Yes	Yes		Yes	Yes	Yes .		Yes	Yes
1	Deaths			•		••••••••••••••••••••••••••••••••••••••	0 0	0		00	0	0
	III .oN	8004	3159 753	78	t,000+	224 4	36	т,	Attack rates 18-45%	527+ 103+ 300+	case	Case es
)	Population at Risk	c.9500	c.4500 c.6600	130	~•	391 24	23	с.	5	~	Single	Single case 2 cases
	Type of Population	Ships from Orient	Naval Training Station recruits Station personnel	Naval vessel crew	Fort Ord Army Base Army personnel	High school girls and adult leaders	Boys camp, 15-17 year olds	Pakistani ship from Newport, R. I.	Crews of several naval vessels	Military personnel Military dependents Civilians	Military man from Far East	Hospital orderly Young females
	Location	CALIFORNIA San Francisco	San Diego	San Dîego	Monterey	Davis	San Mateo Co.	VIRGINIA Norfolk	RHODE ISLAND Newport	HAWALI	OHIO Cleveland	Cleveland
	Dates of Outbreaks	May 20 June 18	Mid-June	June 5-11	Late June	June 17-25	June 20-25	June 19-23	Early June	June	Nid-June	June 12-16

~
(mint
·CJ
- 35
w
- A
2
<u> </u>
- Net
••••
<u> </u>
- M
0
~
rn -
100 M
-
• •
e
Æ
-
m
-
and the second second

	CDC Influenza Report Number		1-K 2-B	3-A	-4-7 -4-7 -4-7	ዋ ም ምሌ	0 m 0 w 0	ц С Щ С Ц Ц	G .	8-B 10-G	10-B	1	10-A	H-LI
) } a=wi	nosis by Serology	Yes	Yes					Yes	Yes		Kana Kana Kana Kana Kana Kana Kana Kana			
.27	Lab. Diagnosis by Virus Isolation Serolog	Yes	Yes	Yes	Yes	Yes	Yes			Yes	Tes	Yes	Yes	Yes
	Deaths		00	0	0	0	2 (S.) 0 0		0	ېن د	0		د مر الم	0
	ILI .oN	200+	<u>ଜ</u> ମ	case	8	c•1000	33 60 speci- mens	c•320	S	کر	8		1,000+	c cases
	Population at Risk	1688 -	હ્ય	Single	211	53,000	~• ~•	5000	2 cases	2	99 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	.09	c.60,000	Sporadic
	Type of Population	College students and adult leaders	High School stu- dents and exposed residents	Traveller from the Philippines	Isolated encampment	International Boy Scout Jamboree	TXAS Corpus Christi Various cities Sporadic cases	Military transport from Orient	University faculty member and wife	Airline crewmen (from Chile)	Migrant workers, adults	Girl's camp	Entire population	Charity Hospítal OPD patients
	Location	ICWA Grinnell	UTAH Salt Lake City dents and exposed r	KENTUCKY Louisville	Morris Fork	PENNSYLVANIA Valley Forge	TEXAS Corpus Christi Various cities	WASHINGTON Seattle	NEBRASKA Omaha	FLORIDA Miami	MICHIGAN Calhoun Co.	LOUISIANA Grant Parish	Tangipahoa Parish	New Orleans
	Dates of Outbreaks	June 26 July 2	July 1-5	July 5	ft-tt vtut	July 11-18	Early July	July 17	Late June	July 28, Aug. 1	July 29	Mid-July	July 31- Aug. 8	Early Aug.

			1							
	CDC Influenža Report Number	A-11	11-6	10 -F 12 -F	11-B	-0- 	12-0	12-A	9-A 12-J	
رخم •	nosis by Serology	Sue La seconda de la seconda d La seconda de la seconda de			- 2			Yes		
	Lab. Diagnosis by Virus Isolation Serolog	Yes	Yes	Yes	Yes		Yes		Yes	
	Deaths	0	0	00	0		0			
	ILI .0N	Case	cases	c,75 70	9		case	c cases	17+	
	Population at Risk	Single	Sporadić cases	806 0TT	111		Single case	2 Sporadic cases	5	
	Type of Population	Army camp	2 Air Force Bases	Migrant workers, families, 2 camps		Airine and ship	Contact with case from Mexico	Military personnel	County jail	
	Location	NEW JERSEY Burlington Co.	TEXAS Bexar County	NEW YORK Cayuga Co.	New York City		MICHIGAN Coldwater	OREGON	CALLFORNIA Fresno	
	Dates of Outbreaks	July	July	July 20- Aug. 1	Aug. 7		Early Aug.	Late July	July 14	

TABLE I (Continued)

Ц
TABLE
• • • •

Unconfirmed Influenza-like Illness, Outbreaks - United States June 1-August 16, 1957

CDC Influenza Report Number	ImI	5-D 3-2 3-2 3-2 3-2 3-2 3-2 3-2 3-2 3-2 3-2		3-F	9-F	7-B	ቸ 6	9-6	9 - K	H-01
Obtained Blood	Yes	Tes	Yes	Yes Yes	Yes	Yes	Yes	Yes	Yes	Yes
Specimens Throat Washings	Yes	Yes		Yes Yes	Yes	Yes			Yes	Yes
Deaths	T	0 	0	00	0	0	0	0. 	° 0	0
LLI .ON	38 187	2511+	c.100	* 500+	ġ.	300-500	130	120	c.100	c•75
Population at Risk	75 ?	40,000	800	; 60	~•	C••	200 200	500	ç.	c•950
Type of Population	Mare Island Naval Yard - Marines Naval vessel crew	Camp Pendleton Marine recruits	Three summer chil- dren's camps	City Jail Teenapers	Migrant farm workers	Air Force Base personnel	Children's camps	Air Force Re- servists	Mental Hospital	Fishery workers, adult males
Location	CALIFORNIA Solano Co.**	Oceanside ^{**}	Fresno, Sonoma, Los Angeles Cos.	Los Angeles** Santa Clara**	Monterey and Sonoma Cos.	Marin County	Santa Clara and Calaveras Cos.	Butte County	Sonoma County	LOUISIANA Plaquemine Parish
Dates of Outbreaks	May 29 June 7 June 16	June 22 Early July	Mid-July	July 8-12 July 8	Mid-July	July	Late July- Early Aug.	Late July Early Aug.	Late July- Early Aug.	Aug. 1-6

**Identified as Type A influenza by C-F test.

TABLE II (Continued)

**Post-mortem examination performed.

TABLE III

Outbreaks of Febrile Respiratory Disease - Etiology Other Than Influenza or No Specimens Objerinable June 1-August 15, 1957

			CDC Influenza	Report Number	1		W_L			>		Ì		i jue	and the standard state of the s	12-E		
,	viatned .			Blood	Yes			>						Tes, In-	fluenza	A ADY Ime		
•	Specimens Obvained		Throat	Washings	Tes	Negative for Influenza	and the second sec)	6		2 3 .		an a	r I V		Yes		
			÷	Deaths	Q	1 •		•	τ-	-				y,	ļ	С		ŕ
	No. with	Influenza-	like III-	nesses	2004		06E•3				-	~		- -				
			Population	at Risk	¢.	and the second	c.2540			-						oporaule vase		and the second sec
			Type of	Population	Townspeople	 A second sec second second sec	15 Children's	Summer Camps		a superior of the second se					Ð		ь.,,	
2		3		Location	MISSOURI	Columbia	CALIFORNI A	San Mateo, Santa	Cruz, Sonoma,	Tuolumne, Plumas,	Fresno, San	Diego, Los Ange-	Setnunon set	I.OUTST AND	New Owleane	DIDGT TO MONT	- J #-	
	. 1		Date of	Outbreaks	Early July		Late June	through	M1d-July					July 5 -	July o			

TABLE IV

I

I

Reported Influenza-like Illness Among Returning Delegates from Grinnell (Iowa) Conference Through August 16, 1957

Omitted from this report. No additions since Report No. 11.

Reported Outbreaks of Influenza-like Illness Among Boy Scouts Returning from the Jamboree Through August 16, 1957 (See CDC Influenza Progress Reports 3-D, 4-H, 5-F, 6-E, 7-C, 7-D, 8-C, 11-F, 12-H)

والمتعادين والمحاولة			a ann an an ann ann ann an ann an ann a			
					Lab. Confirmation	
Date of	Final Destination	No.	Illness En	Illness After	Asian Strain	
Report	of Group	ш	Route Home	Arrival	Influenza	Reported From
	Southern California and					
July 23	Hawaii, except Los Angeles	27	Yes	8	Yes	Louisiana
July 23	CONNECTICUT	Ч	1	Yes	Yes	Connecticut
, 1	San Francisco via					Boston, Mass. and New
July 23	NEW ENGLAND	464	Yes	1		x
July 23	SOUTH CAROLINA	4	1	Yes		Marion County, S. C.
	LOUISIANA					
July 24	Tangipahoa Parish	5	Yes	1.		Roanoke, Va.
	TEXAS					
July 24	Austin	1	1	Yes	Yes	Texas
July 30	El Faso	24/	Yes	Yes		New Mexico
	CALIFORNIA					Yellowstone, Wyo. and
July 25	San Francisco	2004	Yes	1		•
	IddISSISSIM					
July 25	Jackson	~	1	Yes		Mississippi
	ALABAMA					
July 29	Jackson	4 -5	Yes	Yes		Alabama
July 29	Thomasville	5	Yes	Yes	Yes	Alabama
	MISSOURI					
July 31	Osceola	7001	8	Yes		Missouri
July 30	NEW MEXICO		Yes	Yes		New Mexico
	VIRGINIA	,				
Aug. o	Koanoke	77		Yes	**************************************	Virginia
Aug. 13	MINNESOTA	÷	Yes	Yes		Minnesota
Aug. 14	OREGON	m		Yes	Yes	Oregon

TABLE V

TABLE VI

Reported Instances of Influenza Associated Deaths, United States June 1, 1957 through August 16, 1957

				ų.			 The second s second second se second second s	
State	e de la companya de	1					Contributory Factors	
and No.	Locale of Death	Age	Sex	Date of Onset	Date of Death	Diagnosis of Influenga	and/or Reported Cause of Death	CDC Influenza Report No.
Cal. 1	San Diego	8 <u>5</u>	H	July 7	July 16	Clinical (CF Test 1:64)	Bronchopneumoni a**	6
Cal. 2	San Diego	11	W	July 17	July 21	Clinical	Coronary occlusion	6
Cal. 5	Davis	57	fizi,	June 29	July 4	Clinical	Acute Toxic Myocarditis**	9-1-0
Cal.	Mare Island	20	2	June 10	June 13	Clinical	Bilateral Lobar Pneu- monia with Consolida- tion (etiol. M. pyo- genes var. aureus)**	6
cal. 7	San Diego	ŤĒ	fz,	~	July 15	Clinical	"Fulminating Influenzal Pneumonia" (Hemolytic Streptococci also cultures)**	9
	Tangipahoa Parish	N	N M N		July 26	Clinical (Family out- break)	DOA- No further details yet	1
Cal. 8	Monterey	Young adult	æ	ç.	Early July	Virus isolation	Bilateral lobar Pneu- monia with Consolida- tion (etiol. M. pyo- genes var. aureus)**	
N. Y. 1	New York City	18	М	Aug. 13	Aug. 14	Clinical	Hemorrhagic pneumonitis**	12

**Post-mortem examination performed.

