



Models of toxicity of diacetyl and alternative diones

David M. Brass*, Scott M. Palmer

Division of Pulmonary, Allergy, and Critical Care Medicine, Duke University Medical Center, Durham, NC, USA



ARTICLE INFO

Article history:

Received 14 November 2016

Received in revised form 23 January 2017

Accepted 17 February 2017

Available online 21 February 2017

Keywords:

Diacetyl

2,3-Pantanedione

2,3-Hexanedione

Popcorn lung

Occupational lung disease

Bronchiolitis obliterans

ABSTRACT

Diacetyl (DA; 2,3-butanedione), with the chemical formula (CH_3CO)₂ is a volatile organic compound with a deep yellow color and a strong buttery flavor and aroma. These properties have made DA a particularly useful and common food flavoring ingredient. However, because of this increased occupational use, workers can be exposed to high vapor concentrations in the workplace. Despite being listed by the USFDA to be 'generally regarded as safe' (GRAS), multiple lines of evidence suggest that exposure to high concentrations of DA vapor causes long-term impairments in lung function with lung function testing indicating evidence of either restrictive or obstructive airway narrowing in affected individuals. A growing number of pre-clinical studies have now addressed the short and long-term toxicity associated with DA exposure providing further insight into the toxicity of DA and related diones. This review summarizes these observations.

© 2017 Elsevier B.V. All rights reserved.

1. Introduction

The ability to appreciate flavor is largely dependent on aroma (Mozell et al., 1969). The study of aroma and aroma perception is in its infancy, but is of increasing interest to science as well as the perfume and food industries (Silva Teixeira et al., 2016). Aromas arise largely from the volatilization of chemicals that are then perceived by olfactory receptors in the nares (Buck and Axel, 1991) and are a point of emphasis in marketing, highlighting their economic importance. Because the scale of food production requires enormous amounts of raw materials, workers may now be exposed occupationally to high concentrations of volatile food flavoring ingredients. Examples of such food flavoring ingredients include isoamyl acetate (banana), benzaldehyde (bitter almond or cherry), cinnamaldehyde (cinnamon), limonene (orange), methyl salicylate (wintergreen) and 2-,3-butanedione (butter). Because of the retrospective nature of hazard regulations in the US, the effects of occupational exposure to such food flavoring ingredients on workers is often not appreciated until after many years of use and occupational exposures. Such is the case with 2,3-butanedione.

2,3-butanedione (diacetyl; DA) is a naturally occurring α -diketone that is produced during fermentation. In its pure form, DA is a yellowish liquid with a strong buttery aroma. DA has a relatively high vapor pressure that renders small quantities particularly pungent (Starek-Swiechowicz and Starek, 2014). DA is present in

varying concentrations in a variety of foodstuffs including wine, dairy products, roasted coffee and beer (Shibamoto, 2014). Because DA imparts a buttery aroma and flavor it has been used in a variety of applications including the manufacture of microwave popcorn. Finally, DA is among those food additives classified by the United States Food and Drug Administration as 'generally regarded as safe' (GRAS).

The earliest mention of diacetyl we could find in the literature is from 1911 (Harden and Norris, 1911) in which it was used as a fluorescent indicator of the Voges and Proskauer reaction in which the presence of acetoin, an intermediate step in DA synthesis, may be detected in bacterial broth culture. In 1946, the likely chemical synthesis pathway for DA through the dehydrogenation of pyruvic acid was described in *Nature* (Suomalainen and Jannes, 1946) and was followed over 20 years later by a more mechanistic report also published in *Nature* in 1968 (Suomalainen and Ronkainen, 1968). In 1933, it was noted that "butter cultures with satisfactory character contained considerable quantities of acetyl methyl carbinol and diacetyl." (Michaelian et al., 1933). Oddly, in 1951 a patent was issued that described the use of DA in the manufacture of synthetic latex rubber (Messer and Reynolds, 1951). This manufacturing process was out-competed by developments in the petrochemical sector, but during a later oil crisis in the late '70's interest in large scale laboratory production of DA was revived briefly (Gupta et al., 1978). In 1969 the oral and intraperitoneal dose determined to cause 50% mortality (LD50) in both male and female rats was determined (Colley et al., 1969). A key finding of that study was that "The no-effect level (90 mg/kg/day) is equivalent to about 500 times the estimated daily intake by

* Corresponding author.

E-mail address: david.brass@duke.edu (D.M. Brass).

man (0.17 mg/kg)." that may, in part, account for the GRAS designation. The 1970's saw advances in understanding of DA chemistry and furthered the use of DA in laboratory tests for urea in biological samples. In 1979, Bjaldanes and Chew (Bjeldanes and Chew, 1979) recognizing that DA is present in a wide variety of foods reported on the mutagenic effects of DA. In 1979 it was recognized that DA could alter the binding of proteins to their known receptors and it was speculated that this could be due to direct protein damage (Rome and Miller, 1980). The 1980's and early 1990's saw the isolation and characterization of diacetyl reductase that converts DA to acetoin, while analytic methods for the detection of DA and its derivatives improved. In 1996, Sengupta et al. (Sengupta et al., 1996) identified a DA-specific odorant receptor in *C. elegans* that is used to affect chemotaxis towards or away from food sources (Langa et al., 2014). Notably, Linda Buck who first cloned this family of receptors in 1991 (Buck and Axel, 1991) was awarded the Nobel Prize in Physiology or Medicine for this fundamental observation. Interestingly, the DA-specific receptor does not sense the closely related 2-,3-pentanedione (PD) that is now being used as a popcorn flavoring alternative to DA. A confirmation of the same DA-specific odorant receptor in mammalian systems was reported in 1997 by Zhang et al. (Zhang et al., 1997). In 1998, Eriksson et al. (Eriksson et al., 1998) reported that DA can modify arginines, essentially confirming the report by Rome and Miller (Rome and Miller, 1980) and suggesting a mechanism by which DA may be toxic to cells and tissues. In a key observation in 2002, Kreiss et al. (Kreiss et al., 2002) noted the development of respiratory symptoms including spirometric evidence of airway obstruction as well as reduced forced vital capacity (FVC) measurements, used to evaluate restrictive and obstructive lung disease, in otherwise healthy popcorn factory workers. These changes were independent of smoking history. The aggregate of these symptoms were suggestive of bronchiolitis obliterans (BO). BO is a rare lung disease that is most commonly associated with the aftermath of lung or bone marrow transplantation. Therefore, this observation published in the New England Journal of Medicine was of considerable note. This report was followed by a number of others characterizing the full spectrum of the effects of DA on the human respiratory tract including cough, shortness of breath and wheezing, and biopsies showed histological and morphological changes in the lung that are consistent with bronchiolitis obliterans (Akpinar-Elci et al., 2004; Kanwal et al., 2006; van Rooy et al., 2007; Kreiss et al., 2012; Kreiss, 2014).

Because of these observations, alternatives to DA are now being used as food flavoring ingredients. These include 2-,3-pentanedione (PD) and possibly 2-,3-hexandione (HD). There are a number of excellent reviews on the subject of DA toxicity that include descriptions of how DA is synthesized in biological systems and its concentrations in different foods (Shibamoto, 2014), while others touch on chemical properties and summarize clinical reports of occupational exposures (Starek-Swiechowicz and Starek, 2014). Still others discuss physico-chemical properties of DA as a way of understanding its toxic effects on cells and tissues (Kovacic and Cooksy, 2010). The purpose of the present review is to synthesize what is known about the relative toxicities of DA, PD and HD with an emphasis on their hypothesized modes of action in *in vitro* cell culture and *in vivo* animal models.

2. *In vitro* toxicity

2.1. The epithelium

DA, with the chemical formula $(\text{CH}_3\text{CO})_2$, is an α -diketone, meaning that in this four carbon molecule the carbons double bonded to oxygens are directly adjacent. In this configuration the sharing of electrons between the adjacent carbonyl groups is

thought to render it particularly reactive as a nucleophile (Wondrak et al., 2002). PD and HD are also α -diketones and may thus be expected to have similar modes of action against similar substrates. It is thought that because of these chemical features α -diketones including DA can affect protein structure and function as a consequence (Miller and Gerrard, 2005). There are four representative *in vitro* studies that address the possible nature of the cellular injury that DA may cause (Zaccone et al., 2015; Fedan et al., 2006; More et al., 2012a,b; Kelly et al., 2014). For example, it has been shown that exposure to 10 mM DA increased epithelial layer permeability as determined by decreased measures of transepithelial resistance (Fedan et al., 2006). This observation has been supported by another study in which exposure of differentiated airway epithelial cells in air liquid interface cultures to DA for six hours at concentrations of 60 ppm and above caused a complete loss of transepithelial resistance (Zaccone et al., 2015). In yet another study, it was shown that DA forms guanosine adducts (More et al., 2012a,b) that the authors state contributes to DNA uncoiling and ultimately to cell death although the mechanism by which this occurs isn't clear. Finally, this laboratory has previously shown that exposure of NCI-H292 cells in liquid culture or primary human airway epithelial cells in air liquid interface culture induces robust shedding of the epidermal growth factor (EGFR) receptor ligand amphiregulin (Areg) (Kelly et al., 2014). In that study it was shown that inhibition of tumor necrosis factor- α converting enzyme (TACE) effectively abrogated Areg shedding, thereby providing a plausible mechanism through which the epithelium could be affecting the mesenchymal response to DA exposure *in vivo*, thereby leading to development of fibroproliferative airway lesions. Thus, *in vitro* studies have shown a loss of barrier function, formation of DNA adducts, protein modification and shedding of the EGFR ligand Areg. These observations are consistent with epithelial injury, which *in vivo* studies support after exposure to higher concentrations of DA.

2.2. Extrapulmonary toxicity *in vivo* and *in vitro*

The first reference we could find for the industrial use of DA as a food flavoring ingredient comes from a 1964 publication (Jenner et al., 1964) in which its large scale use provided the rationale for determining its LD50 in rats, which was identified to be between 1310 and 1920 mg/kg. In 1969 Colley et al. (Colley et al., 1969) determined this dose to be somewhat higher, reporting the oral LD50 for males at 3.4 g/kg and for females at 3 g/kg. The intraperitoneal LD50 for males was reported at 0.4 g/kg for males and 0.64 g/kg for females. In that follow-up report at a high dose of 540 mg/Kg/day in chronic studies the authors describe significant hematological alterations including increased hemoglobin concentrations, increased numbers of circulating neutrophils and lymphocytes, curiously only observed in females at the highest tested dose while circulating monocytes were increased at the highest dose only in males. At the highest dose tested the absolute weight of the brain, heart, liver, spleen, kidneys, adrenal glands and thyroid as well as terminal body weight were all significantly altered in male rats. All organs except the kidneys weighed significantly less than at baseline, while the mass of the adrenal glands increased significantly. Similar observations were made in the females with the difference being that the heart and spleen masses were unchanged. At the next highest dose tested (90 mg/Kg/day) no changes in organ weights were observed.

Methylglyoxal is closely related to DA in chemical structure and reactivity and has been hypothesized to play a role in development of Alzheimer's by virtue of its ability to cross the blood-brain barrier and to form adducts with amyloid- β peptides that increase their propensity to form aggregates that go on to become plaques in the brain (More et al., 2012a,b). To begin to understand the

neuronal toxicity of DA, the authors showed that DA altered amyloid- β peptides and potentiated their growth inhibitory effects on cells in culture. Carbonyl scavengers attenuated this effect. The study also reports that DA effectively crosses an in vitro model of the blood brain barrier and that the enzyme that inactivated methylglyoxal is ineffective against DA. In another study in human liver cells it was determined that DA is highly mutagenic (Whittaker et al., 2008) as measured by the L5178Y mouse lymphoma assay (Lloyd and Kidd, 2012) supporting the observation made by More et al. (More et al., 2012a,b) in epithelial cells that DA causes guanosine adduct formation.

Prior to 2002, such toxicological studies were justified as screens for the identification of potential mutagens. After 2002 all reports of the toxicology of DA refer back to the Kreiss finding that DA exposure causes BO in popcorn factory workers (Kreiss et al., 2002).

3. In vivo pulmonary toxicity

3.1. Rats

The USFDA has designated that diacetyl may be 'generally recognized as safe' (GRAS). According to federal regulations "the use of a food substance may be GRAS either through scientific procedures or, for a substance used in food before 1958, through experience based on common use in food.' It is likely that consumption of DA in the vanishing quantities required to add a buttery flavoring to foods is actually quite safe. However, the GRAS designation does not account for the possibility that workers involved in preparing foods with diacetyl may be exposed to high concentrations of the food flavoring ingredients that do not involve direct and intentional consumption. For example, in the Kreiss study (Kreiss et al., 2002) both the degree of airway obstruction and the proportion of workers with irregular spirometry correlated with increasing cumulative exposure to DA. These observations led directly to the first study of the toxicity of DA in rats (Hubbs et al., 2002). Given the findings of the Kreiss study, the study by Hubbs et al. (Hubbs et al., 2002) is very important in terms of how we consider the toxicity of food flavoring compounds and the GRAS designation more generally. In that study, Ann Hubbs and co-authors examined the hypothesis that inhalation of artificial butter flavoring containing DA causes airway epithelial injury (Hubbs et al., 2002). That careful study showed that inhalation of a combination of vapors released from a proprietary mixture of commercially available artificial butter flavoring that included DA at 285 ppm or greater for 6 h resulted in histopathologic abnormalities in the nasal epithelium at all concentrations of DA vapor and airway epithelial necrosis at all but the lowest DA concentration one day after exposure (Hubbs et al., 2002). In a follow-up study, the authors showed effectively that the component of artificial butter flavoring that was most closely associated with the observed epithelial injury was diacetyl itself (Hubbs et al., 2008). In that study, rats were exposed to DA vapor of up to 365 ppm in a time weighted average for 6 h and the animals were sacrificed 18–20 h later. DA exposure at levels well short of the maximum at 365 ppm caused significant damage to the nasal epithelium including dose dependent necrosuppurative rhinitis. In the lungs, epithelial injury was observed in only two of six animals exposed to the highest concentration of DA. That epithelial injury was characterized by SEM to show flattening of the airway epithelial cells accompanied by loss of cilia and the formation of gaps in the epithelial layer. However, it is worthwhile to note that the epithelium of four of the six rats in this group was grossly unaffected. These observations introduce a consistent theme in understanding the effects of DA and alternative diones on the lung,

namely that there appears to be a threshold exposure required for epithelial injury to occur.

In the first report to show a definitive connection between DA exposure and airway lesion development in rats, Palmer et al. (Palmer et al., 2011) showed that intratracheal instillation of 125 mg/kg of DA in rats causes profound injury to the airway epithelium, altered lung function, airway neutrophilia and lesions characteristic of BO. In that study it was shown that at one and three days after DA exposure, the airway epithelium was ablated in many airways and fully repaired by day 7. In that report it was also shown that mRNA expression of club cell secretory protein (CC10; CCSP), the definitive marker for airway secretory cells is significantly reduced at day 7 post-exposure. This observation is supported by immunofluorescent staining of histological sections in which prominently reduced CCSP immunoreactivity was shown. These observations support the notion that there is a rapid burst of epithelial repair after DA-induced injury and further suggest that the composition of the repaired epithelium is fundamentally altered at this stage in disease development. A follow up study published by Kelly et al. (Kelly et al., 2014) and discussed earlier showed that DA vapor exposure induces robust shedding of the epidermal growth factor receptor (EGFR) ligand amphiregulin (Areg). In that study it was shown that DA vapor exposure of immortalized epithelioid NCI-H292 cells as well as normal human bronchial epithelial cells grown at air liquid interface culture shed significantly increased quantities of Areg into the culture media in a TNF- α converting enzyme (TACE)-dependent manner. Areg is of considerable interest because it is known to participate in epithelial regeneration after injury in other organ systems including the small intestine (Shao and Sheng, 2010). These observations taken together provide a plausible connection between exposure to DA vapor, epithelial injury and repair and airway lesion development that may be EGFR dependent.

At roughly the same time as the Palmer report in PLoS One (Palmer et al., 2011), two separate reports characterized the dosimetry of inhaled DA in rats and compared this with theoretical deposition fractions in humans using computational modeling (Morris and Hubbs 2009; Gloede et al., 2011). These reports showed that a larger fraction of inhaled DA vapor would reach the smaller airways of a human lung than in a rat lung because of the convoluted complexity of rat nares.

Altered lung function is a key feature of BO that is recapitulated by day 7 in rats exposed to DA via intratracheal instillation (Palmer et al., 2011). Similarly, early denudation of the airways of epithelial cells is also now recognized as an established feature of DA exposure. Previous reports have shown that removing the epithelium increases the contractile response to bronchoprovocative agents such as acetylcholine (Hay et al., 1986; Spina, 1998). Therefore, investigators began to address whether DA exposure causes acute changes in lung function. For example, in 2013 Zaccone, et al. (2013) exposed rats to 100, 200, 300 or 360 ppm DA for 6 h and the animals were sacrificed for analysis 18 h later. Counter-intuitively, DA inhalation by itself had little to no effect on airway reactivity 18 h after DA exposure. Contrary to a previous report in which it was shown that guinea pig tracheas had increased contractility in response to DA exposure (Fedan et al., 2006), it was observed in denuded rat tracheas that exposure to DA increased contraction slightly followed by an overall net increase in relaxation (Zaccone et al., 2013). Overall, DA exposure decreased airway contractility and had no net effect on dynamic compliance in the lungs.

Another study that focused on how DA alters lung physiology observed that diacetyl inhalation dose dependently increases sensory innervation that can project into airway epithelium (Goravanahally et al., 2014). Substance P is important in the development of neurogenic inflammation and contributes to

coughing and wheezing (Pernow, 1985; Barnes, 1990; Nieber et al., 1992). That study reported on the significant airway epithelial injury observed after 356 ppm DA vapor exposure and further showed prominent substance P staining in neurons that innervate the airway epithelium. Similarly, this study shows increased innervation in the tracheal epithelium.

In yet another study, rats and mice were administered radiolabeled DA via intratracheal instillation to evaluate the ability of DA to move from the lung into the systemic circulation and to subsequently bind to cellular macromolecules. In this study it was shown that DA forms protein adducts with hemoglobin and albumin (Fennell et al., 2015). The authors suggest that such adducts could be used as markers of exposure.

Finally, in a report comparing the pulmonary effects of DA with some alternative diones, it was shown again that only the higher concentration vapor exposures caused airway epithelial injury and BO-like lesions (Morgan et al., 2016). Rats were exposed to 0, 100, 150 or 200 ppm of DA for six hours per day, five days per week for two weeks + two days or for two weeks followed by two weeks of recovery before evaluation. At 100 ppm DA, the exposed rats did not lose any weight, which is a consistent feature of vapor concentrations that cause epithelial injury and airway lesions. At 150 and 200 ppm the rats lost significant amounts of weight and had epithelial injury and airway lesions further supporting a general observation that there is a threshold exposure above which these vapors are highly toxic. Additionally, rats exposed to 200 ppm of DA developed parenchymal lesions consistent with pulmonary interstitial fibrosis. Rats exposed to 150 ppm did not have altered lung function after the end of the two weeks' exposure, but when evaluated two weeks after the end of the exposure had developed increased resistance and decreased compliance. All vapor concentrations of DA induced inflammation in the airways, the parenchyma and in the nares at the end of the exposure. This was largely resolved in the airways by the end of the two weeks' recovery period, but inflammation persisted through the recovery period in the parenchyma and the nares.

3.2. Mice

A central theme in the rat studies described above is that below a certain threshold of exposure the epithelium remains grossly unaffected by DA exposure. Associated with this is the observation that only above those exposure levels, in which the epithelium is injured, do the animals develop airway lesions consistent with BO. That mice are different from rats is almost a mantra in animal models of disease. Despite this, we do not have a clear understanding of what those differences are that lead to differences in responses to similar exposures. In this review we have separated out the discussion of rats and mice because of a fundamental difference in their responses to exposure to DA and alternative diones. The studies to be described below feature prominent epithelial injury and repair, but the lesions that develop are not consistent with BO.

The first report of mouse exposures to DA comes from Morgan et al. in 2008 (Morgan et al., 2008) in which it was reported that C57BL/6 mice were exposed to either: 0, 200 or 400 ppm DA vapor for six hours per day for five consecutive days; 0, 100, 200 or 400 ppm DA vapor for one hour per day for two or four weeks; to 0 or 1200 ppm DA vapor for 15 min per exposure twice daily for four weeks; to 0, 25, 50 or 100 ppm DA vapor for six hours per day, five days per week for 12 weeks; or finally to 400 mg/kg DA delivered by intratracheal instillation. This represents a truly comprehensive investigation of the effects of DA inhalation on the respiratory system in which the experimental design was to model relevant workplace exposures in the microwave popcorn industry. In this study it was determined that significant nasal and bronchial

epithelial injury, peribronchial lymphocytic inflammation and fibrohistiocytic lesions in the terminal bronchioles occur after the higher exposure concentrations. The conclusion of the study for the purposes of this review may be that despite significant epithelial injury and inflammation, C57BL/6 mice simply do not develop BO-like lesions after DA exposure. Because human subjects exposed to DA have altered lung function and constrictive and obstructive airway lesions and because rats develop similar lesions to humans, we must consider mice to be a less than ideal model of occupational DA exposure.

In another study of DA exposure in mice by Fennell et al. (Fennell et al., 2015) it was shown that, as with rats, less than 1% of orally administered DA entered the bloodstream, but that when it did a significant portion of that radiolabel was bound to hemoglobin and albumin. Also, as with the rats exposed to DA, protein-DA adducts were observed in both the bound hemoglobin and the albumin. Most recently Hubbs et al. (Hubbs et al., 2016) have shown in mice that DA injury of the respiratory epithelium is characterized by accumulation of ubiquitin, involved in protein degradation, and sequestosome-1, a marker of autophagy, as indicators of protein damage, which has been shown to be highly toxic to cells (Goldberg, 2003).

4. Pentanedione

4.1. Rats

In 2012, Hubbs et al. (Hubbs et al., 2012) reported on experiments in which rats were exposed to 120, 240, 320 or 360 ppm of PD for 6 h and evaluated for up to 20 h later. As with rats exposed to DA, described above, there was significant dose and time dependent injury to the nasal, trachea and mainstem bronchus epithelium in rats exposed to PD. In that study it was also shown that nasal epithelial cells that were olfactory marker protein (OMP) positive indicating that they are nasal epithelial neuroendocrine cells were also positive for activated caspase 3 by immunofluorescence, suggesting a mechanism by which there could be changes in epithelial innervation after PD exposure. Notably, this study reports alterations in cytokine and growth factor mRNA expression in diverse regions of the brain after PD exposure.

In 2015, Morgan et al. (Morgan et al., 2015) exposed male Wistar Han rats to 200 ppm PD or air for 6 h per day, 5 days per week for 2 weeks and performed laser capture microdissection on both exposed and affected airways and exposed and unaffected airways. RNA from the microdissected airways was isolated for microarray analysis. This study identified over 3800 differentially expressed genes in fibrotic bronchi. In pathway analysis, the authors identify that genes associated with vascular development were the most highly enriched followed by genes associated with extracellular matrix remodeling, internal cytoskeletal organization and cell adhesion. Perhaps not surprisingly, TGF- β family proteins in the pathway analysis, with TGF- β 2 being the most significantly differentially expressed member of the family. The authors point out that this is consistent with previous reports that TGF- β 2 is the primary family member expressed in airways after development (Tschumperlin et al., 2003; Balzar et al., 2005; Thompson et al., 2006)

4.2. Mice

In another comprehensive report, Morgan et al. (Morgan et al., 2012) show that both male and female mice exposed to 0, 50, 100 or 200 ppm PD vapor for six hours per day five days a week for up to two weeks develop a wide range of epithelial lesions in the nares at the two highest vapor concentrations. Similar observations were

made in the larynx and in the mainstem bronchus. Interestingly, in that report only male mice exposed to the 200 ppm PD vapor for the full two-week period had any change in the lavageable cell differential in the lung with ~50% of the cells being neutrophils.

5. Hexanedione

5.1. Rats

The first report in the literature of a rat HD exposure was in 1984, well before it was appreciated that DA and possibly alternative diones could cause BO. In that study by Iwasaki and Tsuruta (Iwasaki and Tsuruta, 1984) the goal was to determine the neurotoxicity of hexane indirectly. In that study, rats were exposed to 2-,3-hexanedione, 2-,4-hexanedione and 2-,5-hexanedione by gavage with each study animal receiving 4 mL of a 0.5% solution and sacrificed up to 24 h later. Concentrations of each form of HD were measured both in the blood and in neuronal tissue. The authors made the observation that the 2-,3- form of HD had the lowest retention time in neuronal tissue of any of the three forms.

More recently, as part of a comparative study between the effects of DA, PD and HD on rats, Morgan et al. (Morgan et al., 2012) exposed male rats to 0, 100, 150 or 200 ppm HD vapor and demonstrated that HD vapor exposure induced airway lesions in exposed rats only at 200 ppm and only in 2/12 rats in the exposed group. There was significant mortality in the DA and PD groups but none in the HD exposed group. Consistent with this the HD exposed rats did not lose weight in contrast with the rats exposed to either DA or PD. HD had no effect on measured lung function parameters including resistance and compliance, and caused considerably less epithelial injury.

5.2. Mice

To the best of our knowledge there are no studies that examine the effects of inhaled HD vapor on the mouse respiratory tract.

6. Conclusions

Two central themes have emerged from the body of literature we have reviewed. First, there appears to be a threshold of exposure, below which the nasal and airway epithelium appear essentially unchanged after exposure. This observation holds true for both rats and mice, currently the only study animals we are aware of that have been exposed to DA or alternative diones experimentally. Notably, while current OSHA guidelines require additional labeling stating that DA can cause damage to respiratory tract and lungs if inhaled, as recently as April 1 of 2016, the GRAS status of DA was reaffirmed. These observations, taken together, strongly suggest that the GRAS definition must be expanded to include other likely routes of exposure. While high concentration occupational exposures are encountered by a significantly smaller proportion of the general population, the effects of such exposures may be catastrophic. Therefore, a new metric that accounts for the likely route of exposure in industrial settings must be established.

The second theme that has emerged is that, while mice experience nasal and airway epithelial injury to the same general degree that rats do, they do not develop BO-like lesions. It is currently not known what the molecular differences between the responses of rats and mice to DA might be. Understanding this difference may have therapeutic value.

In summary, DA vapor causes toxic injury to and necrosis of the nasal, tracheal and bronchial epithelium. This has been demonstrated in the literature repeatedly since the initial observation in 2002 that DA exposure is associated with development of BO in popcorn factory workers. Recently, e-cigarette manufacturers have

begun using DA as a flavoring ingredient in e-cigarettes with marketing campaigns that highlight the 'fun' of products such as fruit and candy-flavored 'vape juice' (Farsalinos et al., 2015; Allen et al., 2016). These products are now ostensibly marketed to adults, but with flavor names such as 'Fruity Fun Cereal' and 'Bubblegum' it is hard to see how these would not be appealing to younger consumers. This raises the possibility that exposure to DA or related diones used in e-cigarettes could also induce pulmonary epithelial injury and increase the risk for development of BO in a cohort of young users of e-cigarettes. Given the rapid growth of e-cigarette usage, further research into the toxic effects of DA and related diones is greatly needed to protect individuals from the potential consequences of these exposures.

References

- Akpınar-Elci, M., Travis, W.D., Lynch, D.A., Kreiss, K., 2004. Bronchiolitis obliterans syndrome in popcorn production plant workers? *Eur. Respir. J.* 24 (2), 298–302.
- Allen, J.G., Flanigan, S.S., LeBlanc, M., Vallarino, J., MacNaughton, P., Stewart, J.H., Christiani, D.C., 2016. Flavoring chemicals in E-Cigarettes: diacetyl, 2,3-Pentanedione, and acetoin in a sample of 51 products, including fruit-, candy-, and cocktail-Flavored E-Cigarettes. *Environ. Health Perspect.* 124 (6), 733–739.
- Balzar, S., Chu, H.W., Silkoff, P., Cundall, M., Trudeau, J.B., Strand, M., Wenzel, S., 2005. Increased TGF-beta2 in severe asthma with eosinophilia? *J. Allergy Clin. Immunol.* 115 (1), 110–117.
- Barnes, P.J., 1990. Neurogenic inflammation in airways and its modulation. *Arch. Int. Pharmacodyn. Ther.* 303, 67–82.
- Bjeldanes, L.F., Chew, H., 1979. Mutagenicity of 1,2-dicarbonyl compounds: maltol, kojic acid, diacetyl and related substances. *Mutat. Res.* 67 (4), 367–371.
- Buck, L., Axel, R., 1991. A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* 65 (1), 175–187.
- Colley, J., Gaunt, I.F., Lansdown, A.B., Grasso, P., Gangolli, S.D., 1969. Acute and short-term toxicity of diacetyl in rats? *Food Cosmet. Toxicol.* 7 (6), 571–580.
- Eriksson, O., Fontaine, E., Bernardi, P., 1998. Chemical modification of arginines by 2,3-butanedione and phenylglyoxal causes closure of the mitochondrial permeability transition pore? *J. Biol. Chem.* 273 (20), 12669–12674.
- Farsalinos, K.E., Kistler, K.A., Gillman, G., Voudris, V., 2015. Evaluation of electronic cigarette liquids and aerosol for the presence of selected inhalation toxins? *Nicotine Tob. Res.* 17 (2), 168–174.
- Fedan, J.S., Dowdy, J.A., Fedan, K.B., Hubbs, A.F., 2006. Popcorn worker's lung: in vitro exposure to diacetyl an ingredient in microwave popcorn butter flavoring, increases reactivity to methacholine. *Toxicol. Appl. Pharmacol.* 215 (1), 17–22.
- Fennell, T.R., Morgan, D.L., Watson, S.L., Dhungana, S., Waidyanatha, S., 2015. Systemic uptake, albumin and hemoglobin binding of [(14)C]2,3-butanedione administered by intratracheal instillation in male Harlan Sprague Dawley rats and oropharyngeal aspiration in male B6C3F1/N mice. *Chem. Biol. Interact.* 227, 112–119.
- Gloege, E., Cichocki, J.A., Baldino, J.B., Morris, J.B., 2011. A validated hybrid computational fluid dynamics-physiologically based pharmacokinetic model for respiratory tract vapor absorption in the human and rat and its application to inhalation dosimetry of diacetyl? *Toxicol. Sci.* 123 (1), 231–246.
- Goldberg, A.L., 2003. Protein degradation and protection against misfolded or damaged proteins. *Nature* 426 (6968), 895–899.
- Goravanahally, M.P., Hubbs, A.F., Fedan, J.S., Kashon, M.L., Battelli, L.A., Mercer, R.R., Goldsmith, W.T., Jackson, M.C., Cumpston, A., Frazer, D.G., Dey, R.D., 2014. Diacetyl increases sensory innervation and substance P production in rat trachea? *Toxicol. Pathol.* 42 (3), 582–590.
- Gupta, K.G., Yadav, N.K., Dhawan, S., 1978. Laboratory-scale production of acetoin plus diacetyl by Enterobacter cloacae ATCC 27613? *Biotechnol. Bioeng.* 20 (12), 1895–1901.
- Harden, A., Norris, D., 1911. The diacetyl reaction for proteins. *J. Physiol.* 42 (4), 332–336.
- Hay, D.W., Farmer, S.G., Raeburn, D., Robinson, V.A., Fleming, W.W., Fedan, J.S., 1986. Airway epithelium modulates the reactivity of guinea-pig respiratory smooth muscle? *Eur. J. Pharmacol.* 129 (1–2), 11–18.
- Hubbs, A.F., Battelli, L.A., Goldsmith, W.T., Porter, D.W., Frazer, D., Friend, S., Schwegler-Berry, D., Mercer, R.R., Reynolds, J.S., Grote, A., Castranova, V., Kullman, G., Fedan, J.S., Dowdy, J., Jones, W.G., 2002. Necrosis of nasal and airway epithelium in rats inhaling vapors of artificial butter flavoring? *Toxicol. Appl. Pharmacol.* 185 (2), 128–135.
- Hubbs, A.F., Goldsmith, W.T., Kashon, M.L., Frazer, D., Mercer, R.R., Battelli, L.A., Kullman, G.J., Schwegler-Berry, D., Friend, S., Castranova, V., 2008. Respiratory toxicologic pathology of inhaled diacetyl in sprague-dawley rats? *Toxicol. Pathol.* 36 (2), 330–344.
- Hubbs, A.F., Cumpston, A.M., Goldsmith, W.T., Battelli, L.A., Kashon, M.L., Jackson, M.C., Frazer, D.G., Fedan, J.S., Goravanahally, M.P., Castranova, V., Kreiss, K., Willard, P.A., Friend, S., Schwegler-Berry, D., Fluharty, K.L., Sriram, K., 2012. Respiratory and olfactory cytotoxicity of inhaled 2,3-pentanedione in Sprague-Dawley rats? *Am. J. Pathol.* 181 (3), 829–844.
- Hubbs, A.F., Fluharty, K.L., Edwards, R.J., Barnabei, J.L., Grantham, J.T., Palmer, S.M., Kelly, F., Sargent, L.M., Reynolds, S.H., Mercer, R.R., Goravanahally, M.P., Kashon,

M.L., Honaker, J.C., Jackson, M.C., Cumpston, A.M., Goldsmith, W.T., McKinney, W., Fedan, J.S., Battelli, L.A., Munro, T., Bucklew-Moyers, W., McKinstry, K., Schwegler-Berry, D., Friend, S., Knepp, A.K., Smith, S.L., Sriram, K., 2016. Accumulation of ubiquitin and sequestosome-1 implicate protein damage in diacetyl-induced cytotoxicity. *Am. J. Pathol.*

Iwasaki, K., Tsuruta, H., 1984. Molecular mechanism of hexane neuropathy: significant differences in pharmacokinetics between 2.3-, 2.4-, and 2.5-hexanedione. *Ind. Health* 22 (3), 177–187.

Jenner, P., Hagan, E., Taylor, J., Cook, E., Fitzhugh, O., 1964. Food flavorings and compounds of related structure I. Acute oral toxicity. *Fd Cosmet. Toxicol.* 2, 327–343.

Kanwal, R., Kullman, G., Piacitelli, C., Boylstein, R., Sahakian, N., Martin, S., Fedan, K., Kreiss, K., 2006. Evaluation of flavorings-related lung disease risk at six microwave popcorn plants? *J. Occup. Environ. Med.* 48 (2), 149–157.

Kelly, F.L., Sun, J., Fischer, B.M., Voynow, J.A., Kummarapurugu, A.B., Zhang, H.L., Nugent, J.L., Beasley, R.F., Martinu, T., Gwinn, W.M., Morgan, D.L., Palmer, S.M., 2014. Diacetyl induces amphiregulin shedding in pulmonary epithelial cells and in experimental bronchiolitis obliterans? *Am. J. Respir. Cell Mol. Biol.* 51 (4), 568–574.

Kovacic, P., Cooksy, A.L., 2010. Electron transfer as a potential cause of diacetyl toxicity in popcorn lung disease. *Rev. Environ. Contam. Toxicol.* 204, 133–148.

Kreiss, K., Gomaa, A., Kullman, G., Fedan, K., Simoes, E.J., Enright, P.L., 2002. Clinical bronchiolitis obliterans in workers at a microwave-popcorn plant? *N. Engl. J. Med.* 347 (5), 330–338.

Kreiss, K., Fedan, K.B., Nasrullah, M., Kim, T.J., Materna, B.L., Prudhomme, J.C., Enright, P.L., 2012. Longitudinal lung function declines among California flavoring manufacturing workers? *Am. J. Ind. Med.* 55 (8), 657–668.

Kreiss, K., 2014. Work-related spirometric restriction in flavoring manufacturing workers. *Am. J. Ind. Med.* 57 (2), 129–137.

Langs, S., Martin-Cabrejas, I., Montiel, R., Landete, J.M., Medina, M., Arques, J.L., 2014. Short communication: combined antimicrobial activity of reuterin and diacetyl against foodborne pathogens? *J. Dairy Sci.* 97 (10), 6116–6121.

Lloyd, M., Kidd, D., 2012. The mouse lymphoma assay. *Methods Mol. Biol.* 817, 35–54.

Messer, W.E., Reynolds, J.A. (1951). Diacetyl as shortstop in butadiene-1,3 emulsion polymerization, Google Patents.

Michaelian, M.B., Farmer, R.S., Hammer, B.W., 1933. The relationship of acetyl methyl carbinol and diacetyl to butter cultures. *Iowa State Agr. Exp. Sta. Research Bull.* 155.

Miller, A.G., Gerrard, J.A., 2005. Assessment of protein function following cross-linking by alpha-dicarbonyls. *Ann. N. Y. Acad. Sci.* 1043, 195–200.

More, S.S., Raza, A., Vince, R., 2012a. The butter flavorant diacetyl, forms a covalent adduct with 2-deoxyguanosine, uncoils DNA, and leads to cell death. *J. Agric. Food Chem.* 60 (12), 3311–3317.

More, S.S., Vartak, A.P., Vince, R., 2012b. The butter flavorant diacetyl, exacerbates beta-amyloid cytotoxicity. *Chem. Res. Toxicol.* 25 (10), 2083–2091.

Morgan, D.L., Flake, G.P., Kirby, P.J., Palmer, S.M., 2008. Respiratory toxicity of diacetyl in C57BL/6 mice? *Toxicol. Sci.* 103 (1), 169–180.

Morgan, D.L., Jokinen, M.P., Price, H.C., Gwinn, W.M., Palmer, S.M., Flake, G.P., 2012. Bronchial and bronchiolar fibrosis in rats exposed to 2,3-pentanedione vapors: implications for bronchiolitis obliterans in humans? *Toxicol. Pathol.* 40 (3), 448–465.

Morgan, D.L., Merrick, B.A., Gerrish, K.E., Stockton, P.S., Wang, Y., Foley, J.F., Gwinn, W.M., Kelly, F.L., Palmer, S.M., Ton, T.V., Flake, G.P., 2015. Gene expression in obliterative bronchiolitis-like lesions in 2,3-pentanedione-exposed rats. *PLoS One* 10 (2), e0118459.

Morgan, D.L., Jokinen, M.P., Johnson, C.L., Price, H.C., Gwinn, W.M., Bousquet, R.W., Flake, G.P., 2016. Chemical reactivity and respiratory toxicity of the alpha-Diketone flavoring agents: 2,3-Butanedione, 2,3-Pentanedione, and 2,3-Hexanedione? *Toxicol. Pathol.* 44 (5), 763–783.

Morris, J.B., Hubbs, A.F., 2009. Inhalation dosimetry of diacetyl and butyric acid: two components of butter flavoring vapors. *Toxicol. Sci.* 108 (1), 173–183.

Mozell, M.M., Smith, B.P., Smith, P.E., Sullivan Jr., R.L., Swender, P., 1969. Nasal chemoreception in flavor identification. *Arch. Otolaryngol.* 90 (3), 367–373.

Nieber, K., Baumgarten, C.R., Rathsack, R., Ferkert, J., Oehme, P., Kunkel, G., 1992. Substance P and beta-endorphin-like immunoreactivity in lavage fluids of subjects with and without allergic asthma. *J. Allergy Clin. Immunol.* 90 (4 Pt 1), 646–652.

Palmer, S.M., Flake, G.P., Kelly, F.L., Zhang, H.L., Nugent, J.L., Kirby, P.J., Foley, J.F., Gwinn, W.M., Morgan, D.L., 2011. Severe airway epithelial injury, aberrant repair and bronchiolitis obliterans develops after diacetyl instillation in rats. *PLoS One* 6 (3), e17644.

Pernow, B., 1985. Role of tachykinins in neurogenic inflammation. *J. Immunol.* 135 (Suppl. 2), 812s–815s.

Rome, L.H., Miller, J., 1980. Butanedione treatment reduces receptor binding of a lysosomal enzyme to cells and membranes. *Biochem. Biophys. Res. Commun.* 92 (3), 986–993.

Sengupta, P., Chou, J.H., Bargmann, C.I., 1996. odr-10 encodes a seven transmembrane domain olfactory receptor required for responses to the odorant diacetyl? *Cell* 84 (6), 899–909.

Shao, J., Sheng, H., 2010. Amphiregulin promotes intestinal epithelial regeneration: roles of intestinal subepithelial myofibroblasts. *Endocrinology* 151 (8), 3728–3737.

Shibamoto, T., 2014. Diacetyl: occurrence, analysis, and toxicity. *J. Agric. Food Chem.* 62 (18), 4048–4053.

Silva Teixeira, C.S., Cerqueira, N.M., Silva Ferreira, A.C., 2016. Unravelling the olfactory sense: from the gene to odor perception? *Chem. Senses* 41 (2), 105–121.

Spina, D., 1998. Epithelium smooth muscle regulation and interactions. *Am. J. Respir. Crit. Care Med.* 158 (5 Pt 3), S141–145.

Starek-Swiechowicz, B., Starek, A., 2014. Diacetyl exposure as a pneumotoxic factor: a review. *Roczn. Panstw. Zakl. Hig.* 65 (2), 87–92.

Suomalainen, H., Jannes, L., 1946. Fermentative formation of diacetyl. *Nature* 157, 336.

Suomalainen, H., Ronkainen, P., 1968. Mechanism of diacetyl formation in yeast fermentation. *Nature* 220 (5169), 792–793.

Thompson, H.G., Mih, J.D., Krasieva, T.B., Tromberg, B.J., George, S.C., 2006. Epithelial-derived TGF-beta2 modulates basal and wound-healing subepithelial matrix homeostasis? *Am. J. Physiol. Lung Cell. Mol. Physiol.* 291 (6), L1277–1285.

Tschumperlin, D.J., Shively, J.D., Kikuchi, T., Drazen, J.M., 2003. Mechanical stress triggers selective release of fibrotic mediators from bronchial epithelium? *Am. J. Respir. Cell Mol. Biol.* 28 (2), 142–149.

Whittaker, P., Clarke, J.J., San, R.H., Begley, T.H., Dunkel, V.C., 2008. Evaluation of the butter flavoring chemical diacetyl and a fluorocarbon paper additive for mutagenicity and toxicity using the mammalian cell gene mutation assay in L5178Y mouse lymphoma cells? *Food Chem. Toxicol.* 46 (8), 2928–2933.

Wondrak, G.T., Cervantes-Laurean, D., Roberts, M.J., Qasem, J.G., Kim, M., Jacobson, E. L., Jacobson, M.K., 2002. Identification of alpha-dicarbonyl scavengers for cellular protection against carbonyl stress? *Biochem. Pharmacol.* 63 (3), 361–373.

Zaccone, E.J., Thompson, J.A., Ponnoth, D.S., Cumpston, A.M., Goldsmith, W.T., Jackson, M.C., Kashon, M.L., Frazer, D.G., Hubbs, A.F., Shimko, M.J., Fedan, J.S., 2013. Popcorn flavoring effects on reactivity of rat airways in vivo and in vitro? *J. Toxicol. Environ. Health A* 76 (11), 669–689.

Zaccone, E.J., Goldsmith, W.T., Shimko, M.J., Wells, J.R., Schwegler-Berry, D., Willard, P.A., Case, S.L., Thompson, J.A., Fedan, J.S., 2015. Diacetyl and 2,3-pentanedione exposure of human cultured airway epithelial cells: ion transport effects and metabolism of butter flavoring agents. *Toxicol. Appl. Pharmacol.* 291 (December (3)), 542–549. doi:<http://dx.doi.org/10.1016/j.taap.2015.10.004>.

Zhang, Y., Chou, J.H., Bradley, J., Bargmann, C.I., Zinn, K., 1997. The *Caenorhabditis elegans* seven-transmembrane protein ODR-10 functions as an odorant receptor in mammalian cells? *Proc. Natl. Acad. Sci. U. S. A.* 94 (22), 12162–12167.

van Rooy, F.G., Rooyackers, J.M., Prokop, M., Houba, R., Smit, L.A., Heederik, D.J., 2007. Bronchiolitis obliterans syndrome in chemical workers producing diacetyl for food flavorings? *Am. J. Respir. Crit. Care Med.* 176 (5), 498–504.