

## Biological monitoring of workers exposed to engineered nanomaterials

P. Schulte<sup>a,\*</sup>, V. Leso<sup>b</sup>, M. Niang<sup>c</sup>, I. Iavicoli<sup>b</sup>

<sup>a</sup> National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, 1090 Tusculum Avenue, MS C-14, Cincinnati, OH 45226, USA

<sup>b</sup> Department of Public Health, University of Naples Federico II, Via S. Pansini 5, 80131 Naples, Italy

<sup>c</sup> University of Cincinnati, Cincinnati, OH, USA



### ARTICLE INFO

#### Keywords:

Hazard identification  
Exposure assessment  
Risk management

### ABSTRACT

As the number of nanomaterial workers increase there is need to consider whether biomonitoring of exposure should be used as a routine risk management tool. Currently, no biomonitoring of nanomaterials is mandated by authoritative or regulatory agencies. However, there is a growing knowledge base to support such biomonitoring, but further research is needed as are investigations of priorities for biomonitoring. That research should be focused on validation of biomarkers of exposure and effect. Some biomarkers of effect are generally non-specific. These biomarkers need further interpretation before they should be used. Overall biomonitoring of nanomaterial workers may be important to supplement risk assessment and risk management efforts.

### 1. Introduction

The utility and novelty of engineered matter at the nanoscale is driving a revolution in science and commerce (Roco, 1997; Stirling, 2018). Engineered nanomaterials (ENMs) offer the opportunity to make products lighter, stronger, more conductive, and generally “smarter” (Iavicoli et al., 2014a,b; Navya and Daima, 2016; Thiruvengadam et al., 2018). ENMs entered commerce in the early 2000s and found utility in many sectors and commercial products. Indeed, workers have been and are expected to be increasingly involved and exposed to ENMs in various applications as well as along their whole life-cycle, from research to end-life or recycling (Schulte et al., 2016).

Simultaneously, occupational health concerns have emerged regarding the possible impact of such nano-sized materials on the health of exposed employees. These concerns were based on previous evidence about the effects of ultrafine particulate matter derived from air pollution epidemiological studies, data extrapolated from investigations on the health consequences of diesel and welding fume exposures that may contain incidental nanoparticles (NPs), and preliminary toxicological findings obtained in vitro and in vivo models with ENMs (Iavicoli et al., 2011, 2012, 2016; Leso et al., 2017; Oberdörster and Yu, 1999; Peters et al., 2011; Stone et al., 2017).

Assessing the occupational risks derived from ENM exposure is a challenging task. This is because of difficulties in characterizing their potential hazard due to their huge physico-chemical variability, as well as in adequately assessing environmental exposure and individual response. In fact, to date, although there has been extensive research, no standardized environmental monitoring strategies to assess workplace

and breathing zone exposures, nor the physico-chemical ENM features to be measured as the most effective dose metric parameters are widely agreed upon (Brouwer et al., 2009; Brouwer, 2010; Pietroiusti et al., 2018). Similarly there are no instances of regular monitoring of nanomaterial workers for biomarkers of effect. This is in part likely because these markers are not specific and have not been fully validated as indicators of adverse effects of specific exposure.

Nonetheless, biological monitoring may contribute to the identification of potential hazards of ENMs and to the assessment of occupational exposure to such xenobiotics; therefore, supporting a more adequate assessment and management of risks (Schulte and Hauser, 2012). In this study, the current status of biological monitoring for ENM workers will be addressed.

### 2. Methods

In order to review the available scientific literature on biological monitoring, a framework matrix was employed (Fig. 1). PubMed, Scopus, and Web of Science databases were searched. Our search strategy was aimed to identify issues concerning the development and use of biomarkers that may be helpful to address the different steps along (Fig. 1) the continuum of occupational safety and health actions from hazard identification to risk management (x-axis) (Schulte and Hauser, 2012). Additionally, the literature related to various classes of biomarkers (y-axis) was assessed for each category and subcategory in the continuum. The objective of the search was to identify papers that raised issues in the use of biomarkers for a cell or cells in the matrix. These issues will be discussed under the categories of biomonitoring,

\* Corresponding author.

E-mail address: [PSchulte@cdc.gov](mailto:PSchulte@cdc.gov) (P. Schulte).

Occupational Safety and Health Actions									
	Hazard Identification		Exposure Assessment		Risk Assessment		Risk Management		
Biomarker Types	In vivo	In vitro	Biomonitoring		Exposure-Response Modeling	Epidemiological Studies	Control Effectiveness	Medical Surveillance	Prevention through Design
			Direct	Indirect					
Pathological									
Genotoxic									
Transcriptomic									
Proteomic									
Metabolomic									
Epigenetic									

Fig. 1. Framework for literature search strategy.

epidemiological research, and medical surveillance. However, as the major findings were from toxicological research on ENMs, these will be presented to establish a foundation for thinking about the utility of various types of biomarkers for other purposes in the continuum. There is a rich literature on the validation of biomarkers that can be consulted regarding their utility (Bonassi and Au, 2002; Schulte and Hauser, 2012; Schulte and Talaska, 1995).

### 3. Hazard identification

Since the late 1990s, there has been a growing knowledge-base about the potential adverse effects of ENMs. This seems a challenging issue due to the diverse universe of materials characterized by combinations of a large number of physico-chemical characteristics and other parameters that may include size, shape, chemical composition, solubility, crystalline structure, charge, surface characteristics, attached functionalized groups, agglomeration, and impurities. In humans, ENMs generally have not yet been reported to cause adverse health effects (Bergamaschi, 2012). This may be due to a number of factors: the precautions adopted by the scientific and commercial communities to control exposure levels, the guidance from authoritative agencies responsible for the development of nanotechnology, the small number of involved workers, and the limited time since first exposure, which is generally 15–20 years, considering the periods that such products were introduced in commerce (Fig. 2).

Considering that nanotechnology is an enabling technology diffused through many productive sectors, the mass of produced ENMs is generally small. In these complex industrial scenarios, ENM dose-response relationships may be underestimated by the confounding influence of other possible chemical coexposures. Nevertheless, hazard research in the last 15 years has shown that some types of ENMs appears to have the ability to result in adverse effects in animals. Table 1 shows an overview of the toxicological findings. Interestingly, toxicological evidence may be helpful in extrapolating information for biological monitoring.

#### 3.1. In vitro studies

In vitro studies provide information concerning molecular effects of

ENMs on cellular models, therefore suggesting possible modes of action in relation also to the intrinsic ENM physico-chemical features. This information may be important to define the hazardous properties of such nanoscale materials, a critical step in the identification of suitable biomarkers. Different ENMs, including silver (Ag)- (Müller et al., 2018; Murphy et al., 2016), titanium dioxide (TiO<sub>2</sub>)- (Kongseng et al., 2016; Patil et al., 2016; Renwick et al., 2001), zinc oxide (ZnO)- (Patil et al., 2016), silica (SiO<sub>2</sub>)- (Breznan et al., 2017), magnesium oxide (MgO)-NPs (Mahmoud et al., 2016), and carbon based-NMs (Chatterjee et al., 2017; Chortarea et al., 2017; Kim and Yu, 2014), demonstrated cytotoxic, inflammatory, and oxidative stress effects, as well as the reduction of specific cellular function in vitro models. The small ENM size, resulting in a high surface area to volume ratio, may enhance ENM biological reactivity and relative toxicity (Guichard et al., 2016; Kim and Ryu, 2013; Soares et al., 2016; Uboldi et al., 2016).

Additionally, for adequate hazard identification, the chemical nature and surface chemistry should be also considered as possible influencing factors, since differently composed ENMs, i.e., metal oxide-NPs vs multi-walled-carbon nanotubes (MWCNTs) (Xia et al., 2013), Ag-NPs vs graphene oxide nano-sheets (Ivask et al., 2015), and differently functionalized NMs (Chatterjee et al., 2017; Magdolenova et al., 2015), resulted in diverse cytotoxic effects that could be also dependent on cellular type specificity (Chatterjee et al., 2017). The crystalline structure of the nano-compounds could also affect the toxicity of some ENMs, as in the case of anatase and rutile TiO<sub>2</sub>-NPs (Sayes et al., 2006). Shape-dependent effects have been reported for several ENMs, including Au- and TiO<sub>2</sub>-NPs, and carbon based-ENMs (Allegrri et al., 2016; Chithrani et al., 2006; Hamilton et al., 2009; Park et al., 2003; Porter et al., 2013; Öner et al., 2017). ENMs with a high aspect ratio morphology have raised particular concern in the field of respiratory toxicology, because they may induce toxicity similar to other fibrous substances, e.g., asbestos (Donaldson et al., 2011). Overall, these data may support a different behavior of ENMs according to their primary features as well as also in relation to those properties secondarily acquired when coming in contact with different culture media (Ritz et al., 2015).




Fig. 2. Conceptual timeline for growth of nanomaterial products.

Table 1

General Overview of Toxicology Findings for ENMs to date.

- Verified the correlation of particle surface area with biological effects (Duffin et al., 2001; Schmid and Stoeger, 2016)
- Confirmed that particle size is generally an important factor in toxicity but other physico-chemical factors may play major roles (Driscoll, 1996; Oberdörster et al., 2005)
- Confirmed that nanoparticles could reach the alveoli and could enter the interstitium and blood stream (Kreyling et al., 2009; Oberdörster et al., 1992)
- Demonstrated that pulmonary exposure to carbon nanotubes (CNTs) causes alveolar interstitial fibrosis, which develops rapidly and is persistent. Fibrotic potency appears related to the physicochemical properties of the CNT (Shvedova et al., 2005)
- Demonstrated that pulmonary exposure to nanoparticles can cause cardiovascular effects (alteration of heart rate and blood pressure, and microvascular dysfunction) (Li et al., 2007; Nurkiewicz et al., 2008)
- Demonstrated that pulmonary exposure to some nanomaterials may be carcinogenic (Kuempel et al., 2017; Rittinghausen et al., 2014; Sargent et al., 2014)
- Demonstrated that multi-walled carbon nanotubes (MWCNTs) (Mitsui-7) are a promoter of lung cancer (Kuempel et al., 2017)
- Demonstrated that nanoparticles deposited in the lung can translocate to distal sites (Aragon et al., 2017; Erdely et al., 2009; Mercer et al., 2013)
- Demonstrated that insoluble nanoparticles do not appear to rapidly cross intact skin (Prow et al., 2011)
- Demonstrated that nanoparticle penetration through intact skin is not likely but exposure to sensitizers and irritants still a concern (Gwinn and Vallyathan, 2006; Prow et al., 2011)

### 3.2. In vivo studies

In vivo studies are essential to define the toxico-kinetic and dynamic behavior of ENMs and particularly to identify early adverse effects and possible target organ damage whose investigation, through biomarkers detected in accessible biological matrices, may be important for a suitable planning of biological monitoring programs. As previously mentioned, the biomolecular interactions that ENMs may experience when in contact with biological fluids, strictly dependent on ENM exposure route, can determine the formation of the so called “protein corona,” which may affect ENM dynamic profile, and therefore, biological monitoring findings (Monopoli et al., 2012).

#### 3.2.1. Preclinical alterations

As reported in cellular models, and also in human and animal experiments, ENMs elicited the activation of oxidative stress responses

that could be monitored through the reduction in plasma levels of anti-oxidant defense systems. This was detected in workers involved in the manufacture and/or application of ENMs in Taiwan, and also found after a six month follow up period (Liao et al., 2014; Liou et al., 2012, 2016). Additionally, the increased exhaled breath condensate (EBC) concentrations of lipid, nucleic, and protein oxidation markers were detected in workers exposed to Fe<sub>2</sub>O<sub>3</sub>- and Fe<sub>3</sub>O<sub>4</sub>-NPs (Pelclova et al., 2016a), MWCNTs (Lee et al., 2015), and TiO<sub>2</sub>-NPs (Pelclova et al., 2017a, 2017b) compared to controls. In such epidemiological investigations, a dose-response correspondence was demonstrated according to the ENM environmental monitoring concentrations (Pelclova et al., 2016a, b) as well as the increase in risk levels established through a control binding nano-tool risk assessment approach (Liao et al., 2014).

Comparably, in animal models, reductions in plasma anti-oxidant defense systems and increased levels of Reactive Oxygen Species (ROS) and malondialdehyde (MDA) could be determined with SiO<sub>2</sub> (Du et al., 2013; Liu and Sun, 2013), Ag (Genter et al., 2012; Martins et al., 2017), Ag-NPs in coexposure with TiO<sub>2</sub>-NPs (Martins et al., 2017), Fe<sub>2</sub>O<sub>3</sub>- NPs (Sundarraj et al., 2017), and multi-walled carbon nanotube (MWCNTs) exposures (Reddy et al., 2011).

Regarding inflammatory response, blood levels of fibrinogen and pro-inflammatory cytokines were increased in workers exposed to nano-sized carbon black and MWCNTs (Fatkhutdinova et al., 2016; Liou et al., 2012; Vlaanderen et al., 2017; Zhang et al., 2014) as well as to TiO<sub>2</sub>-NPs compared to controls (Zhao et al., 2018). Conversely, when comparing serum proteins associated with inflammatory responses in ENM exposed and not-exposed workers in research laboratories, significant increases in CD40, TNFR2, and CD62P serum concentrations were evident during the first shift of a working week and at the end of the week in exposed employees (Glass et al., 2017). These findings may suggest an anti-inflammatory and suppressive homeostatic cellular response in relation to an immune activation.

Enhanced levels of leukotrienes were detected in the EBCs of subjects occupationally exposed to TiO<sub>2</sub>- NPs (Pelclova et al., 2016c) and MWCNTs (Lee et al., 2015). Comparably, increased concentrations of 8-isoprostane in EBC (Liou et al., 2017) and enhanced fractional exhaled nitric oxide were reported in workers exposed to metal oxide ENMs (Wu et al., 2014), also when the analysis was stratified according to the type of NM exposure (SiO<sub>2</sub>, TiO<sub>2</sub>, indium tin oxide-NPs), while potential fibrotic and pro-inflammatory biomarkers were detected in the sputum of MWCNT-involved workers (Fatkhutdinova et al., 2016). Conversely,

in research laboratory workers handling ENMs, no significant alterations in exhaled nitric oxide could be detected (Glass et al., 2017).

Alterations in blood acute-phase proteins and cytokine concentrations were also detected in animal models through acute to sub-acute administrations of different types of metallic or metal oxide-NPs such as TiO<sub>2</sub>- (Park et al., 2009), CeO<sub>2</sub>- (Nalabotu et al., 2011; Srinivas et al., 2011), Fe<sub>3</sub>O<sub>4</sub>- (Chen et al., 2010; Park et al., 2010; Srinivas et al., 2012)-, Ag- (Holland et al., 2015, 2016), and SiO<sub>2</sub>-NPs (Downs et al., 2012; Du et al., 2013; Lu et al., 2011) as well as with carbon-based NPs (Erdely et al., 2009). Interestingly, such preclinical alterations, whatever the biological matrix employed, may function both as indicators of early effects before clinical manifestations may occur and as indirect markers of exposure although with a low specificity for ENMs (Aragon et al., 2017; Erdely et al., 2009; Shvedova et al., 2016).

### 3.2.2. Pathological alterations

Pathological alterations pertain to a greater understanding of the potential for adverse effects following exposure. In highly exposed nanoscale carbon black workers, alterations in pulmonary functional parameters were observed compared to controls. This suggests possible adverse respiratory alterations (Zhang et al., 2014), although other investigations failed to detect such changes in MWCNT (Lee et al., 2015; Liao et al., 2014; Vlaanderen et al., 2017) and TiO<sub>2</sub>-NP exposed employees (Pelclova et al., 2016c, 2017b) as well as in workers handling ENMs in research laboratories (Glass et al., 2017).

Interestingly, a significant dose-dependent increase in the pulmonary surfactant protein D serum levels, as a biomarker of lung damage, was detected in workers employed in a packaging workshop of a nano-TiO<sub>2</sub> manufacturing plant in eastern China (Zhao et al., 2018). In these workers, alterations in cardiovascular disease markers, i.e., VCAM-1, ICAM-1, LDL, and TC, were associated with occupational ENM exposure (Zhao et al., 2018). Haematological alterations were reported in animals after oral or intra-gastric exposure to ZnO-NPs (Park et al., 2014) and TiO<sub>2</sub>-NPs (Vasantharaja et al., 2015) as well as after dermal exposure to hydroxyapatite-NPs (Parayanthala Valappil et al., 2014), while could not be detected after a sub-acute inhalation of CeO-NPs (Gosens et al., 2016). Increased ALT concentrations and decreased albumin levels in serum as markers of hepatocyte injury, confirmed also by histopathological alterations, were reported in rats intra-tracheally instilled with CeO<sub>2</sub>-NPs (Nalabotu et al., 2011). The hepatotoxic potential of ENMs has been also documented in animals orally treated with ZnO- (Park et al., 2014), Ag-, and Au-NPs (Shrivastava et al., 2016). Significant nephrotoxic alterations principally induced by metallic NPs have been demonstrated through the assessment of the kidney injury molecule-1 (KIM-1) urinary level (Blum et al., 2015; Iavicoli et al., 2016) as well as by the BUN and creatinine serum biomarkers following an oral exposure to Cu- (Chen et al., 2006; Lei et al., 2008; Liao and Liu, 2012; Sarkar et al., 2011), Au- (Shrivastava et al., 2016), and mesoporous silica-NPs (Li et al., 2015). Indicators for the abovementioned alterations may function as potential biomarkers helpful to define target organs of ENM toxicity and, in turn, suggest potential pathological health conditions susceptible to be aggravated by ENM exposure.

### 3.2.3. Genotoxicity

Due to their small particle size, large surface area, and physico-chemical characteristics, NPs exhibit unpredictable genotoxic properties. Studies on workers failed to show genotoxic effects (Liao et al., 2014; Liou et al., 2012) while a general dose-dependent increase in DNA strand breaks and micronucleus (MN) frequency was found in human peripheral blood cells treated in vitro with metal or metal oxide-NPs (Colognato et al., 2008; Di Bucchianico et al., 2013; Flower et al., 2012; Ghosh et al., 2010, 2012, 2013; Kang et al., 2008, 2011; Paino et al., 2012; Soni et al., 2017; Tavares et al., 2014), carbon-based NPs (Cveticanin et al., 2010; Öner et al., 2017; Tavares et al., 2014), and dendrimers (Ziemba et al., 2012). Concerning in vivo results, a

significant increase in such parameters was detected in the peripheral blood cells of TiO<sub>2</sub>- (Song et al., 2012; Trouiller et al., 2009) and ZnO-NP treated animals (Patil et al., 2016), as compared to controls, and a dose-dependent increase in DNA fragmentation percentage was detected in lymphocytes of SiO<sub>2</sub> and Fe<sub>2</sub>O<sub>3</sub>-NP treated rats (Jiménez-Villarreal et al., 2017). However, positive results were not always confirmed and conflicting results were reported also for other metallic or metal oxide ENMs (Balasubramanyam et al., 2009; Chen et al., 2014; Cordelli et al., 2017; Downs et al., 2012; Lindberg et al., 2012; Sadiq et al., 2012; Singh et al., 2013a, b; Song et al., 2012; Tiwari et al., 2011). This may suggest a possible influencing role of different ENM physico-chemical features, routes of exposure, and administered doses.

### 3.3. “Omic techniques” for hazard identification

Traditional toxicology is rapidly evolving into a system-based approach, able to capture almost all the interactions between living systems and endogenous and/or exogenous xenobiotics (Balbo et al., 2017). “Omic” techniques may reveal methods helpful to assess a wide range of biological responses induced by ENM exposures, therefore, they may be promising tools for the development of novel biomarkers of exposure and early effect (Fadde, 2015; Schulte and Hauser, 2012). These techniques currently include genomics, transcriptomics, proteomics, and more recent approaches, such as metabolomics, and adductomics involved into gene expression and its consequences.

#### 3.3.1. Genomics

Genomic studies may provide the opportunity to detect injury at the molecular level and the signaling pathways involved in organ damage long before the clinical symptoms occur (Andersen and Krewski, 2009; Klaper et al., 2014). Several toxicogenomic studies have investigated the in vitro and in vivo response, principally pulmonary gene expression changes, to NMs using DNA microarray analysis (Aydin et al., 2017; Costa et al., 2018; Decan et al., 2016; Ellinger-Ziegelbauer and Pauluhn, 2009; Gao et al., 2012; Husain et al., 2013; Li et al., 2013, 2017; Pacurari et al., 2011; Snyder-Talkington et al., 2015). However, accessible and more easily applicable biological matrices, i.e., blood or peripheral blood cells, have been rarely employed to assess systemic alterations; determining these correlations in animal models is an important part of practical biomarker validation. The expression of genes involved in immune, inflammatory and oxidative stress responses was affected by Au-NP-oligonucleotide complexes in human peripheral blood mononuclear cells (Kim et al., 2012), by graphene oxide also functionalized with amino groups in T lymphocytes and mononuclear cell lines (Orecchioni et al., 2017). In in vivo experiments, blood expression changes in genes involved in inflammation, oxidative stress, growth factors, tissue remodeling, and endothelial function were obtained in circulating blood cells of animals treated with SW- or MWCNTs (Erdely et al., 2009).

#### 3.3.2. Transcriptomics

Transcriptomics aims at quantifying changes in gene expression through the enumeration of the number of mRNA copies (Costa and Fadde, 2016). A significant increase in the mRNA expression of KL-6, a marker for the diagnosis and monitoring of interstitial lung diseases, as well as its up-stream and down-stream genes, were detected in accessible biological matrices, i.e., whole blood and sputum samples collected from workers exposed to MWCNTs compared to controls (Fatkhutdinova et al., 2016; Shvedova et al., 2016). In vitro studies also reported mRNA expression changes in macrophages exposed to mesoporous silica-NPs and PEGylated mesoporous silica-NPs at doses that do not elicit acute cytotoxicity (Yazdimamaghani et al., 2017). Numerous blood mRNAs were significantly up- or down-regulated post-MWCNT inhalation in animals developing lung pathological changes (Dymacek et al., 2015; Snyder-Talkington et al., 2013, 2016). Therefore, changes in blood mRNA expression, may potentially serve as suitable

biomarkers for ENM-induced lung pathological changes, providing less invasive measurements, compared to tissue analysis, that can be taken at shorter intervals with lower costs (Snyder-Talkington et al., 2016).

### 3.3.3. Proteomics

Proteomic investigation may be helpful to identify candidate protein biomarkers for the evaluation of ENM early effects (Matysiak et al., 2016). In vitro results reported differently expressed proteome profiles related to cellular viability, oxidative stress, and heat response processes in human lung cells exposed to MWCNTs (Phuyal et al., 2018) as well as in the human monocytes treated with Au-, CuO-, and CdTe-NPs (Tarasova et al., 2017). On the other hand, in vivo studies demonstrated an affected expression of proteins associated with metabolism, oxidative stress, and immune responses following TiO<sub>2</sub>-NP inhalatory exposure (Maurer et al., 2016). More recently, the possible employment of proteomic analysis for the detection of protein carbonylation patterns as persistent, sensitive, and indirect biomarkers of ENM induced oxidative stress reaction has been explored (Driessen et al., 2015; Riebeling et al., 2016). Overall, the suitability of these proteomic changes as biological indicators of effect should be firstly confirmed by traditional biochemistry, thus supporting their effectiveness in detecting early NP toxicity.

### 3.3.4. Metabolomics

Metabolomics is the comprehensive analysis of all the metabolites of an organism or specified biological samples (Robertson et al., 2011). In vitro metabolomic data, demonstrated that CeO<sub>2</sub>, SiO<sub>2</sub>- and CuO-NPs could affect the lipidome profile, and compromise the Phase II conjugational capacity in exposed hepatic cells (Kitchin et al., 2017). Differently hydroxylated or carboxylated functionalized CNTs induced alterations in amino acid metabolisms in hepatic and bronchial cells (Chatterjee et al., 2017) while an antioxidant effect was observed for chitosan-coated or ceria supported-Au-NPs in human peripheral blood cells (Palomino-Schätzlein et al., 2017). Concerning the metabolic profiling of ENM effects, as a rapid in vivo screening for nano-toxicity biomarkers, analyses performed in serum and urine of rats treated with Cu- (Lei et al., 2008) and TiO<sub>2</sub>-NPs (Bu et al., 2010; Tang et al., 2010, 2011) provided evidence for the hepatotoxicity, nephrotoxicity and alterations in energy metabolism induced by these NPs. Other studies reported the ability of Ag- (Hadrup et al., 2012; Xie et al., 2018), Fe<sub>2</sub>O<sub>3</sub>- (Feng et al., 2010), SiO<sub>2</sub>-NPs (Lu et al., 2011; Parveen et al., 2012), carbon 14-labeled C60 fullerenes (Sumner et al., 2010) and CNTs (Lin et al., 2013) to affect the liver, energy, lipid, glucose, and amino acid metabolism as assessed by the alterations found in serum and urine metabolic pathways, while amorphous SiO<sub>2</sub>, zirconium dioxides, and barium sulphate-NPs failed to show a relevant impact on plasma metabolome patterns (Buesen et al., 2014). Although interesting, the different and not always well characterized ENMs employed, as well as the lack of standardized analytical techniques and procedures, make the results difficult to compare, and no definite conclusions to be extrapolated.

### 3.3.5. Adductomics

Adductomic approaches have been developed to comprehensively describe toxicological features in response to a genotoxic xenobiotic insult (Hemeryck et al., 2016; Villalta and Balbo, 2017). Specific classes of DNA adducts are those resulting from the reaction of ROS ( $\cdot$ OH) with DNA, i.e., the 8-hydroxy-deoxy-guanosine (8-OH-dG). In field studies, 8-OH-dG concentrations were measured in easily available biological matrices collected from exposed workers, although with variable results. While no significant alterations were detected in urine, and plasma samples from employees of 14 manufacturing plants in Taiwan (Liao et al., 2014; Liou et al., 2012), increased concentrations were more recently reported in urine and white blood cells of workers exposed to metal oxide ENMs, whatever the type of ENM was considered (Liou et al., 2016, 2017), and in EBCs obtained from TiO<sub>2</sub>-NP exposed

subjects (Pelcova et al., 2016b). This DNA damage has been also investigated in vitro models exposed to metal or metal oxide ENMs, which demonstrated significant increases (Ng et al., 2017) although with a cellular type specificity (Mahmoud et al., 2016), and in Ag-NP treated animals that showed increased urinary concentrations of this biomarker (Chuang et al., 2013). The field of DNA adduct research is a highly promising area due to their potential. This is primarily related to the fact that, for instance, adducts may be interpreted as indicators of internal doses, biologically effective dose, and early effect, as well as susceptibility. Though the need exists to be validated specifically for each purpose.

### 3.3.6. Epigenetics

Epigenetics refers to heritable, reversible changes in gene expression occurring without alterations in DNA sequence (Shyamasundar et al., 2015; Sierra et al., 2016). DNA methylation, histone tail, and microRNA modifications are considered useful epigenetic markers. Recent research demonstrated that CNTs (Brown et al., 2016; Chatterjee et al., 2017; Li et al., 2016; Öner et al., 2017; Tabish et al., 2017), Au- (Tabish et al., 2017), TiO<sub>2</sub>- (Bai et al., 2015; Patil et al., 2016), SiO<sub>2</sub>- (Gong et al., 2010, 2012; Mytych et al., 2017), ZnO- (Choudhury et al., 2017), and CuO-NPs (Lu et al., 2016) were able to induce changes of specific methylation patterns, including tumor suppressor-, inflammatory-, and DNA repair genes in vitro and in vivo models. Interestingly, a significantly lower global DNA methylation in white blood cells collected from workers exposed to nanoscale indium tin oxide in plants in Taiwan was recently reported by Liou et al. (2017), while Ghosh et al. (2017), although failing to detect significant differences in global methylation, found significant changes in gene-specific DNA methylation in MWCNT exposed workers compared to controls.

Histone conformational modifications may either facilitate or depress the access of transcriptional machinery to the promoter region of some genes, leading to gene silencing or activation, respectively. As concerns ENMs, a global hypoacetylation in human breast carcinoma cells causing transcriptional repression of anti-apoptotic genes, thereby promoting cellular death, was detected after cadmium telluride quantum dots exposure (CdTe-QDs) (Choi et al., 2008). Ag-NPs were reported to affect histone post-translational modifications thus inducing a reduction in hemoglobin levels in mouse erythroleukemia cells (Qian et al., 2015). Non-coding microRNAs have been investigated in an attempt to identify fine, regulator molecules in ENM induced toxicity since they can modulate gene expression through the interaction with other epigenetic processes (Zhao et al., 2016; Peschansky and Wahlestedt, 2014). The changes in the microRNA expression profiling induced by exposure to Fe<sub>2</sub>O<sub>3</sub>-NPs, CdTe-QDs, Au-, Ag-NPs and MWCNTs were demonstrated to globally affect the mRNA and protein output of human treated cells, subsequently affecting many key biological patterns (Eom et al., 2014; Li et al., 2011a, b; Ng et al., 2017). Micro-RNA expression changes, implicated in inflammation and immune reactions, were found in the lungs of mice intra-tracheally exposed to TiO<sub>2</sub>-NPs (Halappanavar et al., 2011). Changes in the blood levels of liver-specific miRNAs were identified in mice exposed to SiO<sub>2</sub>-NPs (Nagano et al., 2013). However, as the functional consequences of the above mentioned epigenetic alterations were not assessed, further characterization of miRNA responsive genes and their role in adverse effects need to be defined.

## 4. Exposure assessment

Exposure assessments primarily involve area and breathing zone monitoring, job-exposure matrix modeling, or exposure-banding (Kauppinen et al., 2014; Moretto, 2015; NIOSH, 2009, 2017). Although, in recent years, there has been a growth of strong literature on task-based environmental monitoring (Asbach et al., 2017; Brouwer, 2010; Brouwer et al., 2012; Eastlake et al., 2016; Methner et al., 2010a,b,

2012), some challenging issues, including determining the best measurement metrics, availability of appropriate instrumentation, and diverse sampling strategies, prevent an adequate assessment of environmental exposure levels. Therefore, human biomonitoring (HBM) may function as a complementary means for exposure assessment that can take into account the inter-individual variability in absorption, metabolism, and excretion; the individual workload; and recent versus past exposure (Manno et al., 2010).

Currently there are no examples of HBM as a part of routine assessment of workers exposed to ENMs and no regulatory requirements are available for HBM. Nonetheless, there is a growing body of toxicological research that illustrates the possibility to verify some functional alterations detected in experimental settings as biomonitoring indicators for workers. For example, some ENMs can be detected in biological matrices, including blood or plasma, urine, and feces. Following metal- or metal-oxide-based ENM exposure, both via the inhalation or the intratracheal exposure (Balasubramanian et al., 2013; He et al., 2010; Semmler-Behnke et al., 2008; Sundarraj et al., 2017; Sung et al., 2009; Takenaka et al., 2001, 2006; Yu et al., 2007; Zhu et al., 2009) and via human skin application (Gulson et al., 2010, 2012), the elemental metal content is retrievable in blood, although generally in very small amounts. A positive dose-response relationship between the inhaled Ag-ENM and the Ag blood content was demonstrated in mice (Sung et al., 2009), although a differential size dependent biodistribution was reported for gold (Au)-NPs, because smaller 7-nm sized Au-NPs produced a greater metal content in blood compared to their larger 20-nm counterparts (Balasubramanian et al., 2013).

The duration of exposure can affect the interpretations of biomonitoring data, in fact, significant Ag-ENM accumulation in blood was evident after 15 days of treatment, but not following a shorter 5 day period of exposure. This suggests that the body burden of the ENMs is influenced by homeostatic processes (or the exposure could also be influencing homeostatic activities) rather than by the extent of the exposure (Iavicoli et al., 2014a,b). Detectable metal concentrations were reported also in urine samples collected from animals treated with metal or metal oxide ENMs via the respiratory or the dermal route of exposure (Balasubramanian et al., 2013; Gulson et al., 2010, 2012; Sundarraj et al., 2017; Zhu et al., 2009), but there would be no relationship described for any effect. Because of macrophage-mediated clearance mechanisms, the measurement of the elemental metal content in feces may be useful to evaluate the recent/current respiratory exposure to metal-NPs (Balasubramanian et al., 2013; Chuang et al., 2013; Li et al., 2016; Sundarraj et al., 2017; Zhu et al., 2009). However, it is rather difficult to routinely employ feces as a suitable biological matrix for occupational biomonitoring.

Exhaled breath condensate (EBC) is a promising matrix for human biological monitoring investigation, as demonstrated by the increased Ti levels in pre- and post-shift EBC samples collected from workers exposed to TiO<sub>2</sub>-NPs in a pigment production plant compared to unexposed controls (Pelclova et al., 2015, 2016b). Interestingly the biological levels of such an indicator resulted positively correlated with the environmental concentrations in different production or research areas of the plant. However, not all ENMs are easily found in body fluids. Carbon nanotubes are one example. In fact, there is minimal transport of carbon nanotubes from the alveoli to the blood, and they tend to move out of the blood rapidly in to various organs (Erdely et al., 2009).

## 5. Risk assessment

Taking into account what was previously described for ENM hazard assessment and exposure evaluation, it is possible to perceive the difficulties in defining suitable risk assessment strategies. Biological monitoring data, including biomarkers of exposure and effect, can be used in risk assessment involving nanomaterial workers, but thus far, there are no examples where biomarkers have been used in risk assessments. In this scenario, an exposure-response modeling should be

pursued to fully exploit the potential of possible exposure indicators. In fact, when the dose-response relationship is defined, the biomarker of exposure does not only indicate the dose actually adsorbed but provides also a reasonably accurate quantitative estimate of the occupational risks at the group and/or individual level (Iavicoli et al., 2014a). Additionally, scientific efforts should be focused at employing biological monitoring data to develop occupational exposure limits for ENM workers. To do this, the relationship between biomarker levels and adverse effects must be clearly demonstrated in a study design that includes no-effect exposure levels.

The efforts involving categorical approaches to developing occupational exposure limits (OELs) are beginning to identify biomarkers to predict toxicity. For example, the Nanosolutions project tested 31 different engineered ENMs and, out of 8 million data points, identified 11 biomarkers that support the toxicity of these xenobiotics (Nanosolutions, 2018). The response of these markers at human exposure levels must be determined. The 11 biomarkers will then be applied in untested engineered ENMs. These and other approaches that used ENMs with known hazards and the biomarkers of effect related to them may ultimately be used in dose-response modeling.

Another area that involves biomarkers that may be useful in risk assessment and OEL development for ENMs is the use of “omic” technologies that may be useful to understand the interactions and specific pathways impacted by different ENMs. Such biomarkers should be carefully verified in accessible biological matrices under low-dose, long-term conditions of exposure to define their specificity for different ENM insults, their predictive value, and therefore their realistic applicability. Investigations of the effect of functional silica ENMs on a human lung carcinoma cell line showed that it is possible to identify an “omic” analog to the NOAEL involving a no observed transcriptomic effect level or NOTEL (Pisani et al., 2015). This is the level of transcriptomic effect below which cellular responses are not seen to occur and may be much lower than the NOAEL. It could be used as a point of departure in deriving reference values. The development of pathways depicting the sequence of events between exposure to a stressor, progressing through intermediate events, and culminating in an adverse outcome is a growing concept for use in risk assessment (OECD, 2017). This concept may be increasingly used by regulatory agencies and, consequently, there may be more pressure for biomonitoring related to it.

Another relevant contribution to possible ENM-tailored risk assessment processes can be derived from epidemiologic studies. The review of the “first wave” of epidemiological studies of NM workers showed that most were too limited to indicate risk, although some found increased biomarkers of oxidative stress and inflammation in exposed workers (Liou et al., 2015). However, exposure assessment was generally qualitative, study designs were generally cross-sectional, and various selection biases were possible. Therefore, suitable epidemiological research should be planned in order to verify environmental exposure to ENMs and their relationships with biological indicators of exposure and effect. However, no examples where biomarkers have been used in such process are currently available. Studies aimed to address the effectiveness of control technology in a nanomaterial workplace should be pursued not only as a primary way to test control effectiveness in workplaces but also as a useful mean to demonstrate the failure of prevention (Kreider and Halperin, 2011). It could be possible, in laboratory studies, to evaluate control designs and utilize a biomonitoring component of laboratory animals to supplement airborne exposure measures. Biological monitoring may be even more important, in consideration of the possible absorption of ENMs through the dermal route of exposure especially in conditions of compromised skin integrity due to pre-existing diseases or damaging coexposures (Brouwer et al., 2012; Gulson et al., 2010, 2012; Lares Filon et al., 2016; Osmond-McLeod et al., 2014).

## 6. Risk management

Biomonitoring is a risk management tool along with workplace exposure assessment and exposure controls. In this context, biological monitoring data could be used effectively to verify the efficacy of the exposure limit in protecting the health of the workers (Iavicoli et al., 2014a,b). At present, the utility of biomonitoring nanomaterial workers depends on the status of evidence about the hazard of particular ENMs that could be the target of biomonitoring. Biomonitoring should not be used as a substitute for controlling exposure or for taking precautionary control measures.

An additional application of biological monitoring in risk management is the medical surveillance of workers. This is the effort to assess asymptomatic workers for early indications of health problems. There are currently no mandated medical surveillance guidance for ENM workers except for ones involving baseline and period respiratory assessment (pulmonary function testing, x-ray) (NIOSH, 2013). No cellular or molecular biomonitoring is mandated. There is a sparse literature studying the effectiveness of medical surveillance for nanomaterial workers. Gulumian et al. (2016) conducted a review of structures of medical surveillance of nanomaterial workers and identified seven studies that met inclusion criteria. They concluded that there was very low quality of evidence that screening might detect adverse health effects associated with workplace exposures to ENMs. However, this study conflated epidemiological studies and medical surveillance and, as noted earlier, there has not been much time since first worker exposure and the possibility of identifying health effects in epidemiological studies in routine medical surveillance.

One further area of risk management that involves biomarkers of exposure and effect and shows great promise is to design out potential hazards in ENMs. This is known as prevention through design or “safety by design” (Geraci et al., 2015). The concept is illustrated by the identification of specific toxic endpoints that were observed after zebra fish were exposed to functionalized ENMs. This study allowed for the development of predictive models for the design of inherently safer ENMs. Similarly biomarkers found in large arrays in ENM studies may be the basis for material designs that will have reduced ENM hazards.

## 7. Research needs

For biological monitoring of nanomaterial workers to go forward, there is need for robust research to develop biological monitoring protocols (Iavicoli et al., 2014a,b). This should include the physicochemical characterization of materials to which a population is exposed and well established understanding of toxico-kinetic and dynamic characteristics. This research may result in the determination of whether biological monitoring should measure direct parent compounds or possible metabolites or decomposition products. It may explain also the possible role of biomolecular corona formation in affecting ENM biodeposition and, consequently, biomonitoring results.

There is an evident need to link biomarkers in experimental animals to biomarkers in humans. Candidate biomarkers must be validated with exposure and/or effects (Bergamaschi et al., 2015). Additionally, considering the extremely high doses frequently employed in experimental settings and the extremely low ENM retrieved fractions in biological fluids, potential biomarkers should be validated under low-doses and longer periods of treatment, through analytical techniques sensitive enough to address trace levels of biological markers. Moreover, the predictive significance of such biological monitoring alterations in terms of long-term effects, the influences exerted by ENM characteristics, modes and periods of exposure, and inter- and intra-individual variabilities should be assessed to define suitable biomarkers and accurate interpretation of their results.

Validation is a process—a sense of degree rather than an all-or-none-state (IPCS, 2001). Validation includes assessment not only of sensitivity, specificity, and predictive value, but also utility in studies of

workers on a routine basis. Ultimately validation and developmental research to identify biomarkers for biomonitoring of workers will require human studies; these may be field, chamber, or epidemiologic studies. Such studies require attention to ethical legal and social issues. Of importance are issues of privacy, confidentiality, and notification of test and study results (Schulte and Smith, 2011).

## 8. Conclusion

There are no authoritative mandates or requirements currently available for biomonitoring of ENM workers. However, there is a growing body of research that is setting the stage for the use of biomonitoring to supplement environment exposure assessments to achieve a suitable ENM risk assessment and management in occupational settings. Biomarkers may be particularly useful when exposure results in systemic effects. Biomonitoring requires the determination of the extent of the hazard of candidate nanomaterial and evidence that demonstrates biomarker presence in workers correlates with exposure. Biomarkers of effect may also be used to assess exposure and have similar developmental requirements as biomarkers of exposure with the additional need to account for non-specificity and homeostatic influences. If biomonitoring is to be used to assess early or potential health effects, there will be a need for prospective studies to support that use. Overall, there is a rich research base for biomonitoring. However, further work is required before biomonitoring workers can be routinely implemented.

## Acknowledgements

The authors would like to thank: Drs. Kai Savolainen and Glenn Talaska for comments on earlier drafts; Amanda Keenan and Amanda Stammer for processing; and Cheryl Hamilton for editing.

## Disclaimer

The findings and conclusions in this report are those of the author(s) and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.toxlet.2018.06.003>.

## References

- Allegrì, M., Bianchi, M.G., Chiu, M., Varet, J., Costa, A.L., Ortelli, S., Blosi, M., Bussolati, O., Poland, C.A., Bergamaschi, E., 2016. Shape-related toxicity of titanium dioxide nanofibres. *PLoS One* 11, e0151365. <http://dx.doi.org/10.1371/journal.pone.0151365>.
- Andersen, M.E., Krewski, D., 2009. Toxicity testing in the 21st century: bringing the vision to life. *Toxicol. Sci.* 107, 324–330. <http://dx.doi.org/10.1093/toxsci/kfn255>.
- Aragon, J., Navascues, N., Mendoza, G., Irueta, S., 2017. Laser-treated electrospun fibers loaded with nano-hydroxyapatite for bone tissue engineering. *Int. J. Pharm.* 525, 112–122. <http://dx.doi.org/10.1016/j.ijpharm.2017.04.022>.
- Asbach, C., Alexander, C., Clavaguera, S., Dahmann, D., Dozol, H., Faure, B., Fierz, M., Fontana, L., Iavicoli, L., Kaminski, H., MacCalman, L., Meyer-Plath, A., Simonow, B., van Tongeren, M., Todea, A.M., 2017. Review of measurement techniques and methods for assessing personal exposure to airborne nanomaterials in workplaces. *Sci. Total Environ.* 603–604, 793–806. <http://dx.doi.org/10.1016/j.scitotenv.2017.03.049>.
- Aydın, E., Türkez, H., Hacimüftüoğlu, F., Tatar, A., Geyiköğlu, F., 2017. Molecular genetic and biochemical responses in human airway epithelial cell cultures exposed to titanium nanoparticles in vitro. *J. Biomed. Mater. Res. Part A* 105, 2056–2064. <http://dx.doi.org/10.1002/jbm.a.35994>.
- Bai, W., Chen, Y., Gao, A., 2015. Cross talk between poly(ADP-ribose) polymerase 1 methylation and oxidative stress involved in the toxic effect of anatase titanium dioxide nanoparticles. *Int. J. Nanomed.* 10, 5561–5569. <http://dx.doi.org/10.2147/IJN.S88059>.
- Balasubramanyam, A., Sailaja, N., Mahboob, M., Rahman, M.F., Hussain, S.M., Grover, P., 2009. In vivo genotoxicity assessment of aluminum oxide nanomaterials in rat

- peripheral blood cells using the comet assay and micronucleus test. *Mutagenesis* 24, 245–251. <http://dx.doi.org/10.1093/mutage/geb003>.
- Balasubramanian, S.K., Poh, K.W., Ong, C.N., Kreyling, W.G., Ong, W.Y., Yu, L.E., 2013. The effect of primary particle size on biodistribution of inhaled gold nano-agglomerates. *Biomaterials* 34, 5439–5452. <http://dx.doi.org/10.1016/j.biomaterials.2013.03.080>.
- Balbo, S., Turesky, R.J., Villalta, P.W., 2017. DNA adductomics. *Chem. Res. Toxicol.* 27, 356–366. <http://dx.doi.org/10.1021/tx4004352>.
- Bergamaschi, E., 2012. Human biomonitoring of engineered nanoparticles: an appraisal of critical issues and potential biomarkers. *J. Nanomater.* 2012, 564121. <http://dx.doi.org/10.1155/2012/564121>. 12 pp.
- Bergamaschi, E., Poland, C., Guseva Canu, I., Prina-Mello, A., 2015. The role of biological monitoring in nano-safety. *Nanotoday* 10 (3), 274–277. <http://dx.doi.org/10.1016/j.nantod.2015.02.001>.
- Blum, J.L., Edwards, J.R., Prozialeck, W.C., Xiong, J.Q., Zelikoff, J.T., 2015. Effects of maternal exposure to cadmium oxide nanoparticles during pregnancy on maternal and offspring kidney injury markers using a murine model. *J. Toxicol. Environ. Health Part A* 78, 711–724. <http://dx.doi.org/10.1080/15287394.2015.1026622>.
- Bonassi, S., Au, W.W., 2002. Biomarker epidemiology studies for health risk prediction. *Mutat. Res.* 511, 73–86. [http://dx.doi.org/10.1016/S1383-5742\(02\)00003-0](http://dx.doi.org/10.1016/S1383-5742(02)00003-0).
- Breznan, D., Das, D.D., O'Brien, J.S., MacKinnon-Roy, C., Nimesh, S., Vuong, N.Q., Bernatchez, S., DeSilva, N., Hill, M., Kumarathasan, P., Vincent, R., 2017. Differential cytotoxic and inflammatory potency of amorphous silicon dioxide nanoparticles of similar size in multiple cell lines. *Nanotoxicology* 11, 223–235. <http://dx.doi.org/10.1080/17435390.2017.1287313>.
- Brouwer, D., 2010. Exposure to manufactured nanoparticles in different workplaces. *Toxicology* 269, 120–127. <http://dx.doi.org/10.1016/j.tox.2009.11.017>.
- Brouwer, D., van Duuren-Stuurman, B., Berges, M., Jankowska, E., Bard, D., Mark, D., 2009. From workplace air measurement results towards estimates of exposure? Development of a strategy to assess exposure to manufactured nano-objects. *J. Nanopart. Res.* 11, 1867–1881. <http://dx.doi.org/10.1007/s11051-009-9772-1>.
- Brouwer, D., Berges, M., Virji, M.A., Fransman, W., Bello, D., Hodson, L., Gabriel, S., Tielemans, E., 2012. Harmonization of measurement strategies for exposure to manufactured nano-objects. Report of a workshop. *Ann. Occup. Hyg.* 56, 1–9. <http://dx.doi.org/10.1093/annhyg/mer099>.
- Brown, T.A., Lee, J.W., Holian, A., Porter, V., Fredriksen, H., Kim, M., Cho, Y.H., 2016. Alterations in DNA methylation corresponding with lung inflammation and as a biomarker for disease development after MWCNT exposure. *Nanotoxicology* 10, 453–461. <http://dx.doi.org/10.3109/17435390.2015.1078852>.
- Bu, Q., Yan, G., Deng, P., Peng, F., Lin, H., Xu, Y., Cao, Z., Zhou, T., Xue, A., Wang, Y., Cen, X., Zhao, Y.L., 2010. NMR-based metabolomic study of the sub-acute toxicity of titanium dioxide nanoparticles in rats after oral administration. *Nanotechnology* 21, 125105. <http://dx.doi.org/10.1088/0957-4484/21/12/125105>.
- Buesen, R., Landsiedel, R., Sauer, U.G., Wohlleben, W., Groeters, S., Strauss, V., Kamp, H., van Ravenzwaay, B., 2014. Effects of SiO<sub>2</sub>, ZnO, and BaSO<sub>4</sub> nanomaterials with or without surface functionalization upon 28-day oral exposure to rats. *Arch. Toxicol.* 88, 1881–1906. <http://dx.doi.org/10.1007/s00204-014-1337-0>.
- Chatterjee, N., Yang, J., Yoon, D., Kim, S., Joo, S.W., Choi, J., 2017. Differential crosstalk between global DNA methylation and metabolomics associated with cell type specific stress response by pristine and functionalized MWCNT. *Biomaterials* 115, 167–180. <http://dx.doi.org/10.1016/j.biomaterials.2016.11.005>.
- Chen, Z., Meng, H., Xing, G., Chen, C., Zhao, Y., Jia, G., Wang, T., Yuan, H., Ye, C., Zhao, F., Chai, Z., Zhu, C., Fang, X., Ma, B., Wan, L., 2006. Acute toxicological effects of copper nanoparticles in vivo. *Toxicol. Lett.* 163, 109–120. <http://dx.doi.org/10.1016/j.toxlet.2005.10.003>.
- Chen, B.A., Jin, N., Wang, J., Ding, J., Gao, C., Cheng, J., Xia, G., Gao, F., Zhou, Y., Chen, Y., Zhou, G., Li, X., Zhang, Y., Tang, M., Wang, X., 2010. The effect of magnetic nanoparticles of Fe<sub>3</sub>O<sub>4</sub> on immune function in normal ICR mice. *Int. J. Nanomed.* 5, 593–599. <http://dx.doi.org/10.2147/IJN.S12162>.
- Chen, Z., Wang, Y., Ba, T., Li, Y., Pu, J., Chen, T., Song, Y., Gu, Y., Qian, Q., Yang, J., Jia, G., 2014. Genotoxic evaluation of titanium dioxide nanoparticles in vivo and in vitro. *Toxicol. Lett.* 226, 314–319. <http://dx.doi.org/10.1016/j.toxlet.2014.02.020>.
- Chithrani, B.D., Ghazani, A.A., Chan, W.C., 2006. Determining the size and shape dependence of gold nanoparticle uptake into mammalian cells. *Nano Lett.* 6, 662–668. <http://dx.doi.org/10.1021/nl052396o>.
- Choi, A.O., Brown, S.E., Szyf, M., Maysinger, D., 2008. Quantum dot-induced epigenetic and genotoxic changes in human breast cancer cells. *J. Mol. Med. (Berlin Ger.)* 86, 291–302. <http://dx.doi.org/10.1007/s00109-007-0274-2>.
- Chortarea, S., Barosova, H., Clift, M.J.D., Wick, P., Petri-Fink, A., Rothen-Rutishauser, B., 2017. Human asthmatic bronchial cells are more susceptible to subchronic repeated exposures of aerosolized carbon nanotubes at occupationally relevant doses than healthy cells. *ACS Nano* 11, 7615–7625. <http://dx.doi.org/10.1021/acsnano.7b01992>.
- Choudhury, S.R., Ordaz, J., Lo, C.L., Damayanti, N.P., Zhou, F., Irudayaraj, J., 2017. From the cover: zinc oxide nanoparticles-induced reactive oxygen species promotes multimodal cyto- and epigenetic toxicity. *Toxicol. Sci.* 156, 261–274. <http://dx.doi.org/10.1093/toxsci/kfw252>.
- Chuang, H.C., Hsiao, T.C., Wu, C.K., Chang, H.H., Lee, C.H., Chang, C.C., Cheng, T.J., Taiwan CardioPulmonary Research Group (T-CPR), 2013. Allergenicity and toxicology of inhaled silver nanoparticles in allergen-provocation mice models. *Int. J. Nanomed.* 8, 4495–4506. <http://dx.doi.org/10.2147/IJN.S52239>.
- Colognato, R., Bonelli, A., Ponti, J., Farina, M., Bergamaschi, E., Sabbioni, E., Migliore, L., 2008. Comparative genotoxicity of cobalt nanoparticles and ions on human peripheral leukocytes in vitro. *Mutagenesis* 23, 377–382. <http://dx.doi.org/10.1093/mutage/gen024>.
- Cordelli, E., Keller, J., Eleuteri, P., Villani, P., Ma-Hock, L., Schulz, M., Landsiedel, R., Pacchierotti, F., 2017. No genotoxicity in rat blood cells upon 3- or 6-month inhalation exposure to CeO<sub>2</sub> or BaSO<sub>4</sub> nanomaterials. *Mutagenesis* 32, 13–22. <http://dx.doi.org/10.1093/mutage/gew005>.
- Costa, P.M., Fadeel, B., 2016. Emerging systems biology approaches in nanotoxicology: towards a mechanism-based understanding of nanomaterial hazard and risk. *Toxicol. Appl. Pharmacol.* 299, 101–111. <http://dx.doi.org/10.1016/j.taap.2015.12.014>.
- Costa, P.M., Gosens, I., Williams, A., Farcal, L., Pantano, D., Brown, D.M., Stone, V., Cassee, F.R., Halappanavar, S., Fadeel, B., 2018. Transcriptional profiling reveals gene expression changes associated with inflammation and cell proliferation following short-term inhalation exposure to copper oxide nanoparticles. *J. Appl. Toxicol.* 38, 385–397. <http://dx.doi.org/10.1002/jat.3548>.
- Cveticanin, J., Joksic, G., Leskovic, A., Petrovic, S., Sobot, A.V., Neskovic, O., 2010. Using carbon nanotubes to induce micronuclei and double strand breaks of the DNA in human cells. *Nanotechnology* 21, 015102. <http://dx.doi.org/10.1088/0957-4484/21/1/015102>.
- Decan, N., Wu, D., Williams, A., Bernatchez, S., Johnston, M., Hill, M., Halappanavar, S., 2016. Characterization of in vitro genotoxic, cytotoxic and transcriptomic responses following exposures to amorphous silica of different sizes. *Mutat. Res. Genet. Toxicol. Environ. Mutagen.* 796, 8–22. <http://dx.doi.org/10.1016/j.mrgentox.2015.11.011>.
- Di Bucchianico, S., Fabbri, M.R., Misra, S.K., Valsami-Jones, E., Berhanu, D., Reip, P., Bergamaschi, E., Migliore, L., 2013. Multiple cytotoxic and genotoxic effects induced in vitro by differently shaped copper oxide nanomaterials. *Mutagenesis* 28, 287–299. <http://dx.doi.org/10.1093/mutage/get014>.
- Donaldson, K., Murphy, F., Schinwald, A., Duffin, R., Poland, C.A., 2011. Identifying the pulmonary hazard of high aspect ratio nanoparticles to enable their safety-by-design. *Nanomedicine (Lond. U.K.)* 6, 143–156. <http://dx.doi.org/10.2217/nmm.10.139>.
- Downs, T.R., Crosby, M.E., Hu, T., Kumar, S., Sullivan, A., Sarlo, K., Reeder, B., Lynch, M., Wagner, M., Mills, T., Pfuhrer, S., 2012. Silica nanoparticles administered at the maximum tolerated dose induce genotoxic effects through an inflammatory reaction while gold nanoparticles do not. *Mutat. Res.* 745, 38–50. <http://dx.doi.org/10.1016/j.mrgentox.2012.03.012>.
- Drissen, M.D., Mues, S., Vennemann, A., Hellack, B., Bannuscher, A., Vimalakanthan, V., Riebeling, C., Ossig, R., Wiemann, M., Schneckeburger, J., Kuhlbusch, T.A., Renard, B., Luch, A., Haase, A., 2015. Proteomic analysis of protein carbonylation: a useful tool to unravel nanoparticle toxicity mechanisms. *Part. Fibre Toxicol.* 12, 36. <http://dx.doi.org/10.1186/s12989-015-0108-2>.
- Driscoll, K.E., 1996. Role of inflammation in the development of rate lung tumors in response to chronic particle exposure. *Inhal. Toxicol.* 8, 139–153.
- Du, Z., Zhao, D., Jing, L., Cui, G., Jin, M., Li, Y., Liu, X., Liu, Y., Du, H., Guo, C., Zhou, X., Sun, Z., 2013. Cardiovascular toxicity of different sizes amorphous silica nanoparticles in rats after intratracheal instillation. *Cardiovasc. Toxicol.* 13, 194–207. <http://dx.doi.org/10.1007/s12012-013-9198-y>.
- Duffin, R., Tran, C., Clouter, A., Brown, D., MacNee, W., Stone, V., Donaldson, K., 2001. The importance of surface area and specific reactivity in the acute pulmonary inflammatory response to particles. *Ann. Occup. Hyg.* 46, 242–245. <http://dx.doi.org/10.1093/annhyg/mef684>.
- Dymacek, J., Snyder-Talkington, B.N., Porter, D.W., Mercer, R.R., Wolfarth, M.G., Castranova, V., Qian, Y., Guo, N.L., 2015. mRNA and miRNA regulatory networks reflective of multi-walled carbon nanotube-induced lung inflammatory and fibrotic pathologies in mice. *Toxicol. Sci.* 144, 51–64. <http://dx.doi.org/10.1093/toxsci/kfu262>.
- Eastlake, A.C., Beauchamp, C., Martinez, K.F., Dahm, M.M., Sparks, C., Hodson, L.L., Geraci, C.L., 2016. Refinement of the nanoparticle emission assessment technique into the nanomaterial exposure assessment technique (NEAT 2.0). *Occup. Environ. Hyg.* 13, 708–717. <http://dx.doi.org/10.1080/15459624.2016.1167278>.
- Ellinger-Ziegelbauer, H., Pauluhn, J., 2009. Pulmonary toxicity of multi-walled carbon nanotubes (Baytubes) relative to alpha-quartz following a single 6h inhalation exposure of rats and a 3 months post-exposure period. *Toxicology* 266, 16–29. <http://dx.doi.org/10.1016/j.tox.2009.10.007>.
- Eom, H.J., Chatterjee, N., Lee, J., Choi, J., 2014. Integrated mRNA and micro RNA profiling reveals epigenetic mechanism of differential sensitivity of Jurkat T cells to AGNs and Ag ions. *Toxicol. Lett.* 229, 311–318. <http://dx.doi.org/10.1016/j.toxlet.2014.05.019>.
- Erdely, A., Hulderman, T., Salmen, R., Liston, A., Zeidler-Erdely, P.C., Schwegler-Berry, D., Castranova, V., Koyama, S., Kim, Y.A., Endo, M., Simeonova, P.P., 2009. Cross-talk between lung and systemic circulation during carbon nanotube respiratory exposure. Potential biomarkers. *Nano Lett.* 9, 36–43. <http://dx.doi.org/10.1021/nl801828z>.
- Fadeel, B., 2015. Systems biology in nanosafety research. *Nanomedicine (Lond. U.K.)* 10, 1039–1041. <http://dx.doi.org/10.2217/nmm.15.17>.
- Fathkhatdinova, L.M., Khaliullin, T.O., Vasil'yeva, O.L., Zalyalov, R.R., Mustafin, I.G., Kisin, E.R., Birch, M.E., Yanamala, N., Shvedova, A.A., 2016. Fibrosis biomarkers in workers exposed to MWCNTs. *Toxicol. Appl. Pharmacol.* 299, 125–131. <http://dx.doi.org/10.1016/j.taap.2016.02.016>.
- Feng, J., Liu, H., Zhang, L., Bhakoo, K., Lu, L., 2010. An insight into the metabolic responses of ultra-small superparamagnetic particles of iron oxide using metabolomic analysis of biofluids. *Nanotechnology* 21, 395101. <http://dx.doi.org/10.1088/0957-4484/21/39/395101>.
- Flower, N.A., Brabu, B., Revathy, M., Gopalakrishnan, C., Raja, S.V., Murugan, S.S., Kumaravel, T.S., 2012. Characterization of synthesized silver nanoparticles and assessment of its genotoxicity potentials using the alkaline comet assay. *Mutat. Res.* 742, 61–65. <http://dx.doi.org/10.1016/j.mrgentox.2011.12.003>.
- Gao, G., Ze, Y., Li, B., Zhao, X., Zhang, T., Sheng, L., Hu, R., Gui, S., Sang, X., Sun, Q., Cheng, J., Cheng, Z., Wang, L., Tang, M., Hong, F., 2012. Ovarian dysfunction and gene-expressed characteristics of female mice caused by long-term exposure to titanium dioxide nanoparticles. *J. Hazard. Mater.* 243, 19–27. <http://dx.doi.org/10.1016/j.jhazmat.2012.03.012>.

- 1016/j.jhazmat.2012.08.049.
- Genter, M.B., Newman, N.C., Shertzer, H.G., Ali, S.F., Bolon, B., 2012. Distribution and systemic effects of intranasally administered 25 nm silver nanoparticles in adult mice. *Toxicol. Pathol.* 40, 1004–1013. <http://dx.doi.org/10.1177/0192623312444470>.
- Geraci, C., Heidel, D., Sayes, C., Hodson, L., Schulte, P., Eastlake, A., Brenner, S., 2015. Perspectives on the design of safer nanomaterials and manufacturing processes. *J. Nanopart. Res.* 17, 366. [10.1007%2Fs11051-015-3152-9](http://dx.doi.org/10.1007%2Fs11051-015-3152-9).
- Ghosh, M., Bandyopadhyay, M., Mukherjee, A., 2010. Genotoxicity of titanium dioxide (TiO<sub>2</sub>) nanoparticles at two trophic levels: plant and human lymphocytes. *Chemosphere* 81, 1253–1262. <http://dx.doi.org/10.1016/j.chemosphere.2010.09.022>.
- Ghosh, M.J.M., Sinha, S., Chakraborty, A., Mallick, S.K., Bandyopadhyay, M., Mukherjee, A., 2012. In vitro and in vivo genotoxicity of silver nanoparticles. *Mutat. Res.* 749, 60–69. <http://dx.doi.org/10.1016/j.mrgentox.2012.08.007>.
- Ghosh, M., Chakraborty, A., Mukherjee, A., 2013. Cytotoxic, genotoxic and the hemolytic effect of titanium dioxide (TiO<sub>2</sub>) nanoparticles on human erythrocyte and lymphocyte cells in vitro. *J. Appl. Toxicol.* 33, 1097–1110. <http://dx.doi.org/10.1002/jat.2863>.
- Ghosh, M., Öner, D., Poels, K., Tabish, A.M., Vlaanderen, J., Pronk, A., Kuijpers, E., Lan, Q., Vermeulen, R., Bekaert, B., Hoet, P.H., Godderis, L., 2017. Changes in DNA methylation induced by multi-walled carbon nanotube exposure in the workplace. *Nanotoxicology* 11, 1195–1210. <http://dx.doi.org/10.1080/17435390.2017.1406169>.
- Glass, D.C., Mazhar, M., Xiang, S., Dean, P., Simpson, P., Priestly, B., Plebanski, M., Abramson, M., Sim, M.R., Dennekamp, M., 2017. Immunological effects among workers who handle engineered nanoparticles. *Occup. Environ. Med.* 74, 868–876. <http://dx.doi.org/10.1136/oemed-2016-104111>.
- Gong, C., Tao, G., Yang, L., Liu, J., Liu, Q., Zhuang, Z., 2010. SiO<sub>2</sub> nanoparticles induce global genomic hypomethylation in HaCaT cells. *Biochem. Biophys. Res. Commun.* 397, 397–400. <http://dx.doi.org/10.1016/j.bbrc.2010.05.076>.
- Gong, C., Tao, G., Yang, L., Liu, J., Liu, Q., Zhuang, Z., 2012. Methylation of PARP-1 promoter involved in the regulation of nano-SiO<sub>2</sub>-induced decrease of PARP-1 mRNA expression. *Toxicol. Lett.* 209, 264–269. <http://dx.doi.org/10.1016/j.toxlet.2012.01.007>.
- Gosens, I., Cassee, F.R., Zanella, M., Manodori, L., Brunelli, A., Costa, A.L., Bokkers, B.G., de Jong, W.H., Brown, D., Hristozov, D., Stone, V., 2016. Organ burden and pulmonary toxicity of nano-sized copper (II) oxide particles after short-term inhalation exposure. *Nanotoxicology* 10, 1084–1095. <http://dx.doi.org/10.3109/17435390.2016.1172678>.
- Guichard, Y., Fontana, C., Chavinier, E., Terzetti, F., Gaté, L., Binet, S., Darne, C., 2016. Cytotoxic and genotoxic evaluation of different synthetic amorphous silica nanomaterials in the V79 cell line. *Toxicol. Ind. Health* 32, 1639–1650. <http://dx.doi.org/10.1177/0748233715572562>.
- Gulson, B., McCall, M., Korsch, M., Gomez, L., Casey, P., Oytam, Y., Taylor, A., McCulloch, M., Trotter, J., Kinsley, L., Greenoak, G., 2010. Small amounts of zinc from zinc oxide particles in sunscreens applied outdoors are absorbed through human skin. *Toxicol. Sci.* 118, 140–149. <http://dx.doi.org/10.1093/toxsci/kfq243>.
- Gulson, B., Wong, H., Korsch, M., Gomez, L., Casey, P., McCall, M., McCulloch, M., Trotter, J., Stauber, J., Greenoak, G., 2012. Comparison of dermal absorption of zinc from different sunscreen formulations and differing UV exposure based on stable isotope tracing. *Sci. Total Environ.* 420, 313–318. <http://dx.doi.org/10.1016/j.scitotenv.2011.12.046>.
- Gulmian, M., Verbeek, J., Andraos, C., Sanabria, N., de Jager, P., 2016. Systematic review of screening and surveillance programs to protect workers from nanomaterials. *PLoS One* 11, e0166071. <http://dx.doi.org/10.1371/journal.pone.0166071>.
- Gwinn, M.R., Vallyathan, V., 2006. Nanoparticles: health effects—pros and cons. *Environ. Health Perspect.* 114, 1818–1825. <http://dx.doi.org/10.1289/ehp.8871>.
- Hadrup, N., Lam, H.R., Loeschner, K., Mortensen, A., Larsen, E.H., Frandsen, H., 2012. Nanoparticulate silver increases uric acid and allantoin excretion in rats, as identified by metabolomics. *J. Appl. Toxicol.* 32, 929–933. <http://dx.doi.org/10.1002/jat.2779>.
- Halappanavar, S., Jackson, P., Williams, A., Jensen, K.A., Hougaard, K.S., Vogel, U., Yauk, C.L., Wallin, H., 2011. Pulmonary response to surface-coated nanotitanium dioxide particles includes induction of acute phase response genes, inflammatory cascades, and changes in microRNAs: a toxicogenomic study. *Environ. Mol. Mutagen.* 52, 425–439. <http://dx.doi.org/10.1002/em.20639>.
- Hamilton, R.F., Wu, N., Porter, D., Buford, M., Wolfarth, M., Holian, A., 2009. Particle length-dependent titanium dioxide nanomaterials toxicity and bioactivity. *Part. Fibre Toxicol.* 6, 35. <http://dx.doi.org/10.1186/1743-8977-6-35>.
- He, X., Zhang, H., Ma, Y., Bai, W., Zhang, Z., Lu, K., Ding, Y., Zhao, Y., Chai, Z., 2010. Lung deposition and extrapulmonary translocation of nano-ceria after intratracheal instillation. *Nanotechnology* 21, 285103. <http://dx.doi.org/10.1088/0957-4484/21/28/285103>.
- Hemeryck, L.Y., Moore, S.A., Vanhaecke, L., 2016. Mass spectrometric mapping of the DNA adductome as a means to study genotoxin exposure, metabolism, and effect. *Anal. Chem.* 88, 7436–7446. <http://dx.doi.org/10.1021/acs.analchem.6b00863>.
- Holland, N.A., Becak, D.P., Shannahan, J.H., Brown, J.M., Carratt, S.A., Winkle, L., Pinkerton, K.E., Wang, C.M., Munusamy, P., Baer, D.R., Sumner, S.J., Fennell, T.R., Lust, R.M., Wingard, C.J., 2015. Cardiac ischemia reperfusion injury following instillation of 20 nm citrate-capped nanosilver. *J. Nanomed. Nanotechnol.* [56](http://dx.doi.org/10.4172/2157-7439.S6-006). <http://dx.doi.org/10.4172/2157-7439.S6-006>.
- Holland, N.A., Thompson, L.C., Vidanapathirana, A.K., Urankar, R.N., Lust, R.M., Fennell, T.R., Wingard, C.J., 2016. Impact of pulmonary exposure to gold core silver nanoparticles of different size and capping agents on cardiovascular injury. *Part. Fibre Toxicol.* 13, 1. <http://dx.doi.org/10.1186/s12989-016-0159-z>.
- Husain, M., Saber, A.T., Guo, C., Jacobsen, N.R., Jensen, K.A., Yauk, C.L., Williams, A., Vogel, U., Wallin, H., Halappanavar, S., 2013. Pulmonary instillation of low doses of titanium dioxide nanoparticles in mice leads to particle retention and gene expression changes in the absence of inflammation. *Toxicol. Appl. Pharmacol.* 269, 250–262. <http://dx.doi.org/10.1016/j.taap.2013.03.018>.
- Iavicoli, L., Leso, V., Fontana, L., Bergamaschi, A., 2011. Toxicological effects of titanium dioxide nanoparticles: a review of in vitro mammalian studies. *Eur. Rev. Med. Pharmacol. Sci.* 15, 481–508. <http://dx.doi.org/10.1155/2012/964381>.
- Iavicoli, L., Leso, V., Bergamaschi, A., 2012. Toxicological effects of titanium dioxide nanoparticles: a review of in vivo studies. *J. Nanomater.* 36. <http://dx.doi.org/10.1155/2012/964381>. Article ID 964381.
- Iavicoli, L., Leso, V., Manno, M., Schulte, P.A., 2014a. Biomarkers of nanomaterial exposure and effect: current status. *J. Nanopart. Res.* 16, 2302. <http://dx.doi.org/10.1007/s11051-014-2302-9>.
- Iavicoli, L., Leso, V., Ricciardi, W., Hodson, L.L., Hoover, M.D., 2014b. Opportunities and challenges of nanotechnology in the green economy. *Environ. Health* 13, 78. <http://dx.doi.org/10.1186/1476-069X-13-78>.
- Iavicoli, L., Fontana, L., Nordberg, G., 2016. The effects of nanoparticles on the renal system. *Crit. Rev. Toxicol.* 46, 490–560. <http://dx.doi.org/10.1080/10408444.2016.1181047>.
- IPCS, 2001. Environmental Health Criteria 222. Biomarkers in Risk Assessment: Validity and Validation. International Programme on Chemical Safety (IPCS) (Accessed 30 January 2018). <http://www.inchem.org/documents/ehc/ehc/ehc222.htm>.
- Ivask, A., Voelcker, N.H., Seabrook, S.A., Hor, M., Kirby, J.K., Fenech, M., Davis, T.P., Ke, P.C., 2015. DNA melting and genotoxicity induced by silver nanoparticles and graphene. *Chem. Res. Toxicol.* 28, 1023–1035. <http://dx.doi.org/10.1021/acs.chemrestox.5b00052>.
- Jiménez-Villarreal, J., Rivas-Armendáriz, D.I., Arellano Pérez-Vertti, R.D., Olivás Calderón, E., García-Garza, R., Betancourt-Martínez, N.D., Serrano-Gallardo, L.B., Morán-Martínez, J., 2017. Relationship between lymphocyte DNA fragmentation and dose of iron oxide (Fe<sub>2</sub>O<sub>3</sub>) and silicon oxide (SiO<sub>2</sub>) nanoparticles. *Genet. Mol. Res.* 16, 1–12. <http://dx.doi.org/10.4238/gmr16019206>.
- Kang, S.J., Kim, B.M., Lee, Y.J., Chung, H.W., 2008. Titanium dioxide nanoparticles trigger p53-mediated damage response in peripheral blood lymphocytes. *Environ. Mol. Mutagen.* 49, 399–405. <http://dx.doi.org/10.1002/em.20399>.
- Kang, S.J., Lee, Y.J., Kim, B.M., Choi, Y.J., Chung, H.W., 2011. Cytotoxicity and genotoxicity of titanium dioxide nanoparticles in UVA-irradiated normal peripheral blood lymphocytes. *Drug Chem. Toxicol.* 34, 277–284. <http://dx.doi.org/10.3109/01480545.2010.546800>.
- Kauppinen, T., Uuksulainen, S., Saalo, A., Mäkinen, I., Pukkala, E., 2014. Use of the Finnish information system on occupational exposure (FINJEM) in epidemiologic, surveillance, and other applications. *Ann. Occup. Hyg.* 58, 380–396. <http://dx.doi.org/10.1093/annhyg/met074>.
- Kim, S., Ryu, D.Y., 2013. Silver nanoparticle-induced oxidative stress, genotoxicity and apoptosis in cultured cells and animal tissues. *J. Appl. Toxicol.* 33, 78–89. <http://dx.doi.org/10.1002/jat.2792>.
- Kim, J.S., Yu, I.J., 2014. Single-wall carbon nanotubes (SWCNT) induce cytotoxicity and apoptosis produced by reactive oxygen species (ROS) generation in phytohemagglutinin (PHA)-stimulated male human peripheral blood lymphocytes. *J. Toxicol. Environ. Health Part A* 77, 1141–1153. <http://dx.doi.org/10.1080/15287394.2014.917062>.
- Kim, E.Y., Schulz, R., Swantek, P., Kunstman, K., Malim, M.H., Wolinsky, S.M., 2012. Gold nanoparticle-mediated gene delivery induces widespread changes in the expression of innate immunity genes. *Gene Ther.* 19, 347–353. <http://dx.doi.org/10.1038/gt.2011.95>.
- Kitchin, K.T., Stirdivant, S., Robinette, B.L., Castellon, B.T., Liang, X., 2017. Metabolic effects of CeO<sub>2</sub>, SiO<sub>2</sub> and CuO metal oxide nanomaterials on HepG2 cells. *Part. Fibre Toxicol.* 14, 1–16. <http://dx.doi.org/10.1186/s12989-017-0230-4>.
- Klaper, R., Arndt, D., Bozich, J., Dominguez, G., 2014. Molecular interactions of nanomaterials and organisms: defining biomarkers for toxicity and high-throughput screening using traditional and next-generation sequencing approaches. *Analyst* 139, 882–895. <http://dx.doi.org/10.1039/c3an01644g>.
- Kongseng, S., Yoovathaworn, K., Wongprasert, K., Chunhabundit, R., Sukwong, P., Pissuwan, D., 2016. Cytotoxic and inflammatory responses of TiO<sub>2</sub> nanoparticles on human peripheral blood mononuclear cells. *J. Appl. Toxicol.* 36, 1364–1373. <http://dx.doi.org/10.1002/jat.3342>.
- Kreider, T., Halperin, W., 2011. Engineered nanomaterials: learning from the past, planning for the future. *J. Occup. Environ. Med.* 53, S108–S112. <http://dx.doi.org/10.1097/JOM.0b013e31821b146a>.
- Kreyling, W.G., Semmler-Behnke, M., Seitz, J., Scymczak, W., Wenk, A., Mayer, P., Takenaka, S., Oberdörster, G., 2009. Size dependence of the translocation of inhaled iridium and carbon nanoparticle aggregates from the lung of rats to the blood and secondary target organs. *Inhal. Toxicol.* 21, 55–60. <http://dx.doi.org/10.1080/08958370902942517>.
- Kuempel, E.D., Jaurand, M.C., Møller, P., Morimoto, Y., Kobayashi, N., Pinkerton, K.E., Sargent, L.M., Vermeulen, R.C., Fubini, B., Kane, A.B., 2017. Evaluating the mechanistic evidence and key data gaps in assessing the potential carcinogenicity of carbon nanotubes and nanofibers in humans. *Crit. Rev. Toxicol.* 47, 1–58. <http://dx.doi.org/10.1080/10408444.2016.1206061>.
- Laresse Filon, F., Crosera, M., Mauro, M., Baracchini, E., Bovenzi, M., Montini, T., Fornasiero, P., Adami, G., 2016. Palladium nanoparticles exposure: evaluation of permeation through damaged and intact human skin. *Environ. Pollut.* 214, 497–503. <http://dx.doi.org/10.1016/j.envpol.2016.04.077>.
- Lee, J.S., Choi, Y.C., Shin, J.H., Lee, J.H., Lee, Y., Park, S.Y., Baek, J.E., Park, J.D., Ahn, K., Yu, I.J., 2015. Health surveillance study of workers who manufacture multi-walled carbon nanotubes. *Nanotoxicology* 9, 802–811. <http://dx.doi.org/10.3109/17435390.2014.978404>.

- Lei, R., Wu, C., Yang, B., Ma, H., Shi, C., Wang, Q., Wang, Q., Yuan, Y., Liao, M., 2008. Integrated metabolomic analysis of the nano-sized copper particle-induced hepatotoxicity and nephrotoxicity in rats: a rapid in vivo screening method for nanotoxicity. *Toxicol. Appl. Pharmacol.* 232, 292–301. <http://dx.doi.org/10.1016/j.taap.2008.06.026>.
- Leso, V., Fontana, L., Mauriello, M.A., Iavicoli, I., 2017. Occupational risk assessment of engineered nanomaterials: limits, challenges and opportunities. *Curr. Nanosci.* 13, 55–78. <http://dx.doi.org/10.2174/157341371266616101714934>.
- Li, Z., Hulderman, T., Salmen, R., Chapman, R., Leonard, S.S., Young, S.H., Shvedova, A., Luster, M.I., Simeonova, P.P., 2007. Cardiovascular effects of pulmonary exposure to single-walled carbon nanotubes. *Environ. Health Perspect.* 115, 377–382.
- Li, S., Wang, H., Qi, Y., Tu, J., Bai, Y., Tian, T., Huang, N., Wang, Y., Xiong, F., Lu, Z., Xiao, Z., 2011a. Assessment of nanomaterial cytotoxicity with SOLiD sequencing-based microRNA expression profiling. *Biomaterials* 32, 9021–9030. <http://dx.doi.org/10.1016/j.biomaterials.2011.08.033>.
- Li, S., Wang, Y., Wang, H., Bai, Y., Liang, G., Wang, Y., Huang, N., Xiao, Z., 2011b. MicroRNAs as participants in cytotoxicity of CdTe quantum dots in NIH/3T3. *Biomaterials* 32, 3807–3814. <http://dx.doi.org/10.1016/j.biomaterials.2011.01.074>.
- Li, B., Ze, Y., Sun, Q., Zhang, T., Sang, X., Cui, Y., Wang, X., Gui, S., Tan, D., Zhu, M., Zhao, X., Sheng, L., Wang, L., Hong, F., Tang, M., 2013. Molecular mechanisms of nanosized titanium dioxide-induced pulmonary injury in mice. *PLoS One* 8, e55563. <http://dx.doi.org/10.1371/journal.pone.0055563>.
- Li, L., Liu, T., Fu, C., Tan, L., Meng, X., Liu, H., 2015. Biodistribution, excretion, and toxicity of mesoporous silica nanoparticles after oral administration depend on their shape. *Nanomedicine* 11, 1915–1924. <http://dx.doi.org/10.1016/j.nano.2015.07.004>.
- Li, J., Tian, M., Cui, L., Dwyer, J., Fullwood, N.J., Shen, H., Martin, F.L., 2016. Low-dose carbon-based nanoparticle-induced effects in A549 lung cells determined by biospectroscopy are associated with increases in genomic methylation. *Sci. Rep.* 6, 20207. <http://dx.doi.org/10.1038/srep20207>.
- Li, Y., Yan, J., Ding, W., Chen, Y., Pack, L.M., Chen, T., 2017. Genotoxicity and gene expression analyses of liver and lung tissues of mice treated with titanium dioxide nanoparticles. *Mutagenesis* 32, 33–46. <http://dx.doi.org/10.1093/mutage/gew065>.
- Liao, M., Liu, H., 2012. Gene expression profiling of nephrotoxicity from copper nanoparticles in rats after repeated oral administration. *Environ. Toxicol. Pharmacol.* 34, 67–80. <http://dx.doi.org/10.1016/j.etap.2011.05.014>.
- Liao, H.Y., Chung, Y.T., Lai, C.H., Wang, S.L., Chiang, H.C., Li, L.A., Tsou, T.C., Li, W.F., Lee, H.L., Wu, W.T., Lin, M.H., Hsu, J.H., Ho, J.J., Chen, C.J., Shih, T.S., Lin, C.C., Liou, S.H., 2014. Six-month follow-up study of health markers of nanomaterials among workers handling engineered nanomaterials. *Nanotoxicology* 8, 100–110. <http://dx.doi.org/10.3109/17435390.2013.858793>.
- Lin, B., Zhang, H., Lin, Z., Fang, Y., Tian, L., Yang, H., Yan, J., Liu, H., Zhang, W., Xi, Z., 2013. Studies of single-walled carbon nanotubes-induced hepatotoxicity by NMR-based metabolomics of rat blood plasma and liver extracts. *Nanoscale Res. Lett.* 8, 1–11. <http://dx.doi.org/10.1186/1556-276X-8-236>.
- Lindberg, H.K., Falck, G.C., Catalan, J., Koivisto, A.J., Suhonen, S., Jarventaus, H., Rossi, E.M., Nykassenoja, H., Peltonen, Y., Moreno, C., Alenius, H., Tuomi, T., Savolainen, K.M., Norppa, H., 2012. Genotoxicity of inhaled nanosized TiO<sub>2</sub> in mice. *Mutat. Res.* 745, 58–64. <http://dx.doi.org/10.1016/j.mrgentox.2011.10.011>.
- Liou, S.H., Tsou, T.C., Wang, A.L., Li, L.A., Chiang, H.C., Li, W.F., Lin, P.P., Lai, C.H., Lee, H.L., Lin, M.H., Hsu, J.H., Chen, C.R., Shih, T.S., Liao, A.Y., Chung, Y.T., 2012. Epidemiological study of health hazards among workers handling engineered nanomaterials. *J. Nanopart. Res.* 14, 1–15. <http://dx.doi.org/10.1007/s11051-012-0878-5>.
- Liou, S.H., Tsai, C.S.J., Pelclova, D., Schubauer-Berigan, M.K., Schulte, P.A., 2015. Assessing the first wave of epidemiological studies of nanomaterial workers. *J. Nanopart. Res.* 17, 1–19. <http://dx.doi.org/10.1007/s11051-015-3219-7>.
- Liou, S.H., Chen, Y.C., Liao, H.Y., Wang, C.J., Chen, J.S., Lee, H.L., 2016. Increased levels of oxidative stress biomarkers in metal oxides nanomaterial-handling workers. *Biomarkers* 21, 600–606. <http://dx.doi.org/10.3109/1354750X.2016.1160432>.
- Liou, S.H., Wu, W.T., Liao, H.Y., Chen, C.Y., Tsai, C.Y., Jung, W.T., Lee, H.L., 2017. Global DNA methylation and oxidative stress biomarkers in workers exposed to metal oxide nanoparticles. *J. Hazard. Mater.* 331, 329–335. <http://dx.doi.org/10.1016/j.jhazmat.2017.02.042>.
- Liu, X., Sun, J., 2013. Time-course effects of intravenously administered silica nanoparticles on blood coagulation and endothelial function in rats. *J. Nanosci. Nanotechnol.* 13, 222–228. <http://dx.doi.org/10.1166/jnn.2013.6910>.
- Lu, X., Tian, Y., Zhao, Q., Jin, T., Xiao, S., Fan, X., 2011. Integrated metabolomics analysis of the size-response relationship of silica nanoparticles-induced toxicity in mice. *Nanotechnology* 22, 055101. <http://dx.doi.org/10.1088/0957-4484/22/5/055101>.
- Lu, X., Miousse, I.R., Pirela, S.V., Moore, J.K., Melnyk, S., Koturbash, I., Demokritou, P., 2016. In vivo epigenetic effects induced by engineered nanomaterials: a case study of copper oxide and laser printer-emitted engineered nanoparticles. *Nanotoxicology* 10, 629–639. <http://dx.doi.org/10.3109/17435390.2015.1108473>.
- Magdolenova, Z., Drlickova, M., Henjum, K., Rundén-Pran, E., Tulinska, J., Bilanovicova, D., Pojana, G., Kazimirova, A., Barancokova, M., Kuricova, M., Liskova, A., Staruchova, M., Ciampor, F., Vavra, I., Lorenzo, Y., Collins, A., Rinna, A., Fjellsbø, L., Volkovova, K., Marcomini, A., Amiry-Moghaddam, M., Dusinska, M., 2015. Coating-dependent induction of cytotoxicity and genotoxicity of iron oxide nanoparticles. *Nanotoxicology* 9, 44–56. <http://dx.doi.org/10.3109/17435390.2013.847505>.
- Mahmoud, A., Ezgi, Ö., Merve, A., Özhan, G., 2016. In vitro toxicological assessment of magnesium oxide nanoparticle exposure in several mammalian cell types. *Int. J. Toxicol.* 35, 429–437. <http://dx.doi.org/10.1177/1091581816648624>.
- Manno, M., Viau, C., in collaboration with, Cocker, J., Colosio, C., Lowry, L., Muttii, A., Nordberg, M., Wang, S., 2010. Biomonitoring for occupational health risk assessment (BOHRA). *Toxicol. Lett.* 192, 3–16. <http://dx.doi.org/10.1016/j.toxlet.2009.05.001>.
- Martins Jr, A.D.C., Azevedo, L.F., de Souza Rocha, C.C., Carneiro, M.F.H., Venancio, V.P., de Almeida, M.R., Antunes, L.M.G., de Carvalho Hott, R., Rodrigues, J.L., Ogunjimi, A.T., Adeyemi, J.A., Barbosa Jr, F., 2017. Evaluation of distribution, redox parameters, and genotoxicity in Wistar rats co-exposed to silver and titanium dioxide nanoparticles. *J. Toxicol. Environ. Health Part A* 80, 1156–1165. <http://dx.doi.org/10.1080/15287394.2017.1357376>.
- Matysiak, M., Kapka-Skrzypczak, L., Brzóska, K., Gutleb, A.C., Kruszewski, M., 2016. Proteomic approach to nanotoxicity. *J. Proteom.* 137, 35–44. <http://dx.doi.org/10.1016/j.jpropt.2015.10.025>.
- Maurer, M.M., Donohoe, G.C., Maleki, H., Yi, J., McBride, C., Nurkiewicz, T.R., Valentine, S.J., 2016. Comparative plasma proteomic studies of pulmonary TiO<sub>2</sub> nanoparticle exposure in rats using liquid chromatography tandem mass spectrometry. *J. Proteom.* 130, 85–93. <http://dx.doi.org/10.1016/j.jpropt.2015.09.010>.
- Mercer, R.R., Scabillon, J.F., Hubbs, A.F., Wang, L., Battelli, L.A., McKinney, W., Castranova, V., Porter, D.W., 2013. Extrapulmonary transport of MWNT following inhalation exposure. *Part. Fibre Toxicol.* 10, 38. <http://dx.doi.org/10.1186/1743-8977-10-38>.
- Methner, M., Hodson, L., Geraci, C., 2010a. Nanoparticle emission assessment technique (NEAT) for the identification and measurement of potential inhalation exposure to engineered nanomaterials—part. *J. Occup. Environ. Hyg.* 7, 127–132. <http://dx.doi.org/10.1080/10545962093476355>.
- Methner, M., Hodson, L., Dames, A., Geraci, C., 2010b. Nanoparticle Emission Assessment Technique (NEAT) for the identification and measurement of potential inhalation exposure to engineered nanomaterials—part b: results from 12 field studies. *J. Occup. Environ. Hyg.* 7, 163–176. <http://dx.doi.org/10.1080/10545962093508066>.
- Methner, M., Beaucham, C., Crawford, C., Hodson, L., Geraci, C., 2012. Field application of the Nanoparticle Emission Assessment Technique (NEAT): task-based air monitoring during the processing of engineered nanomaterials (ENM) at four facilities. *J. Occup. Environ. Hyg.* 9, 543–555. <http://dx.doi.org/10.1080/105459624.2012.699388>.
- Monopoli, M.P., Aberg, C., Salvati, A., Dawson, K.A., 2012. Biomolecular coronas provide the biological identity of nanosized materials. *Nat. Nanotechnol.* 7, 779–786. <http://dx.doi.org/10.1038/nnano.2012.207>.
- Moretto, A., 2015. Exposure assessment for chemical and physical agents. *Handb. Clin. Neurol.* 131, 47–59. <http://dx.doi.org/10.1016/B978-0-444-62627-1.00004-4>.
- Müller, L., Steiner, S.K., Rodriguez-Lorenzo, L., Petri-Fink, A., Rothen-Rutishauser, B., Latzin, P., 2018. Exposure to silver nanoparticles affects viability and function of natural killer cells, mostly via the release of ions. *Cell Biol. Toxicol.* 34, 167–176. <http://dx.doi.org/10.1007/s10565-017-9403-z>.
- Murphy, A., Casey, A., Byrne, G., Chambers, G., Howe, O., 2016. Silver nanoparticles induce pro-inflammatory gene expression and inflammasome activation in human monocytes. *J. Appl. Toxicol.* 36, 1311–1320. <http://dx.doi.org/10.1002/jat.3315>.
- Mytych, J., Zebrowski, J., Lewinska, A., Wnuk, M., 2017. Prolonged effects of silver nanoparticles on p53/p21 pathway-mediated proliferation, DNA damage response, and methylation parameters in HT22 hippocampal neuronal cells. *Mol. Neurobiol.* 54, 1285–1300. <http://dx.doi.org/10.1007/s12035-016-9688-6>.
- Nagano, T., Higashisaka, K., Kunieda, A., Iwahara, Y., Tanaka, K., Nagano, K., Abe, Y., Kamada, H., Tsunoda, S., Nabeshi, H., Yoshikawa, T., Yoshioka, Y., Tsutsumi, Y., 2013. Liver-specific microRNAs as biomarkers of nanomaterial-induced liver damage. *Nanotechnology* 24, 405102. <http://dx.doi.org/10.1088/0957-4484/24/40/405102>.
- Nalabatu, S.K., Kolli, M.B., Triest, W.E., Ma, J.Y., Manne, N.D., Katta, A., Addagarla, H.S., Rice, K.M., Blough, E.R., 2011. Intratracheal instillation of cerium oxide nanoparticles induces hepatic toxicity in male Sprague-Dawley rats. *Int. J. Nanomed.* 6, 2327–2335. <http://dx.doi.org/10.2147/IJN.S25119>.
- Nanosolutions, 2018. <http://nanosolutionsfp7.com/nanosolutions-explained/> (Accessed 30 January 2018).
- Navya, P.N., Daima, H.K., 2016. Rational engineering of physicochemical properties of nanomaterials for biomedical applications with nanotoxicological perspectives. *Nano Converg.* 3, 1. <http://dx.doi.org/10.1186/s40580-016-0064-z>.
- Ng, C.T., Yong, L.Q., Hande, M.P., Ong, C.N., Yu, L.E., Bay, B.H., Baeg, G.H., 2017. Zinc oxide nanoparticles exhibit cytotoxicity and genotoxicity through oxidative stress responses in human lung fibroblasts and *Drosophila melanogaster*. *Int. J. Nanomed.* 12, 1621–1637. <http://dx.doi.org/10.2147/IJN.S124403>.
- NIOSH, 2009. Approaches to Safe Nanotechnology: Managing the Health and Safety Concerns Associated With Engineered Nanomaterials. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, Atlanta (Accessed 30 January 2018). <https://www.cdc.gov/niosh/docs/2009-125/pdfs/2009-125.pdf>.
- NIOSH, 2013. Occupational Exposure to Carbon Nanotubes and Nanofibers. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, Cincinnati, OH. In: <https://www.cdc.gov/niosh/docs/2013-145/pdfs/2013-145.pdf>.
- NIOSH, 2017. The NIOSH Occupational Exposure Banding Process: Guidance for the Evaluation of Chemical Hazards. External Review Draft. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, Atlanta (Accessed 31 January 2018). [https://www.cdc.gov/niosh/docket/review/docket290/pdfs/clean-cib-niosh-oebprocess-guidancefortheevaluationofchemicalhazards\\_3.8.17.pdf](https://www.cdc.gov/niosh/docket/review/docket290/pdfs/clean-cib-niosh-oebprocess-guidancefortheevaluationofchemicalhazards_3.8.17.pdf).
- Nurkiewicz, T.R., Porter, D.W., Hubbs, A.F., Cumpston, J.L., Chen, B.T., Frazer, D.G., Castranova, V., 2008. Nanoparticle inhalation augments particle-dependent systemic microvascular dysfunction. *Part. Fibre Toxicol.* 5, 1. <http://dx.doi.org/10.1186/1743-8977-5-1>.
- Oberdorster, G., Yu, C.P., 1999. Lung dosimetry—considerations for noninhalation studies. *Exp. Lung Res.* 25, 1–6. <http://dx.doi.org/10.1080/0190214992070385>.
- Oberdorster, G., Ferin, J., Gelein, R., Soderholm, S.C., Finkelstein, J., 1992. Role of the

- alveolar macrophage in lung injury: studies with ultrafine particles. *Environ. Health Perspect.* 97, 193–199. <http://dx.doi.org/10.2307/3431353>.
- Oberdörster, G., Maynard, A., Donaldson, K., Castranova, V., Fitzpatrick, J., Ausman, K., Carter, J., Karn, B., Kreyling, W., Lai, D., Olin, S., Monteiro-Riviere, N., Warheit, D., Yang, H., I.L.S.I. Research Foundation/Risk Science Institute Nanomaterial Toxicity Screening Working Group, 2005. Principles for characterizing the potential human health effects from exposure to nanomaterials: elements of a screening strategy. *Part. Fibre Toxicol.* 2, 8. <http://dx.doi.org/10.1186/1743-8977-2-8>.
- OECD, 2017. Revised Guidance Document on Developing and Assessing Adverse Outcome Pathways. Series on Testing and Assessment, No. 184. 2013. ENM/JM/MONO, pp. 6. (Accessed 30 January 2018). [http://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=env/jm/mono\(2013\)6&doclanguage=en](http://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=env/jm/mono(2013)6&doclanguage=en).
- Öner, D., Moisse, M., Ghosh, M., Duca, R.C., Poels, K., Luyts, K., Putzeys, E., Cokic, S.M., Van Landuyt, K., Vanoirbeek, J., Lambrechts, D., Godderis, L., Hoet, P.H., 2017. Epigenetic effects of carbon nanotubes in human monocytic cells. *Mutagenesis* 32, 181–191. <http://dx.doi.org/10.1093/mutage/gew053>.
- Orecchioni, M., Bedognetti, D., Newman, L., Fuoco, C., Spada, F., Hendrickx, W., Marincola, F.M., Sgarrella, F., Rodrigues, A.F., Ménard-Moyon, C., Cesareni, G., Kostarelos, K., Bianco, A., Delogu, L.G., 2017. Single-cell mass cytometry and transcriptome profiling reveal the impact of graphene on human immune cells. *Nat. Commun.* 8, 1109. <http://dx.doi.org/10.1038/s41467-017-01015-3>.
- Osmond-McLeod, M.J., Oytam, Y., Kirby, J.K., Gomez-Fernandez, L., Baxter, B., McCall, M.J., 2014. Dermal absorption and short-term biological impact in hairless mice from sunscreens containing zinc oxide nano- or larger particles. *Nanotoxicology* 8 (Suppl. 1), 72–84. <http://dx.doi.org/10.3109/17435390.2013.855832>.
- Pacurari, M., Qian, Y., Porter, D.W., Wolfarth, M., Wan, Y., Luo, D., Ding, M., Castranova, V., Guo, N.L., 2011. Multi-walled carbon nanotube-induced gene expression in the mouse lung: association with lung pathology. *Toxicol. Appl. Pharmacol.* 255, 18–31. <http://dx.doi.org/10.1016/j.taap.2011.05.012>.
- Paino, I.M., Marangoni, V.S., de Oliveira Rde, C., Antunes, L.M., Zucolotto, V., 2012. Cytotoxicity and genotoxicity of gold nanoparticles in human hepatocellular carcinoma and peripheral blood mononuclear cells. *Toxicol. Lett.* 215, 119–125. <http://dx.doi.org/10.1016/j.toxlet.2012.09.025>.
- Palomino-Schätzlein, M., García, H., Gutiérrez-Carcedo, P., Pineda-Lucena, A., Herance, J.R., 2017. Assessment of gold nanoparticles on human peripheral blood cells by metabolic profiling with 1H-NMR spectroscopy, a novel translational approach on a patient-specific basis. *PLoS One* 12, e0182985. <http://dx.doi.org/10.1371/journal.pone.0182985>.
- Parayanthala Valappil, M., Santhakumar, S., Arumugam, S., 2014. Determination of oxidative stress related toxicity on repeated dermal exposure of hydroxyapatite nanoparticles in rats. *Int. J. Biomater.* 2014, 476942. <http://dx.doi.org/10.1155/2014/476942>.
- Park, K.H., Chhowalla, M., Iqbal, Z., Sesti, F., 2003. Single-walled carbon nanotubes are a new class of ion channel blockers. *J. Biol. Chem.* 278, 50212–50216.
- Park, E.J., Yoon, J., Choi, K., Yi, J., Park, K., 2009. Induction of chronic inflammation in mice treated with titanium dioxide nanoparticles by intratracheal instillation. *Toxicology* 260, 37–46. <http://dx.doi.org/10.1016/j.tox.2009.03.005>.
- Park, E.J., Kim, H., Kim, Y., Yi, J., Choi, K., Park, K., 2010. Inflammatory responses may be induced by a single intratracheal instillation of iron nanoparticles in mice. *Toxicology* 275, 65–71. <http://dx.doi.org/10.1016/j.tox.2010.06.002>.
- Park, H.S., Kim, S.J., Lee, T.J., Kim, G.Y., Meang, E., Hong, J.S., Kim, S.H., Koh, S.B., Hong, S.G., Sun, Y.S., Kang, J.S., Kim, Y.R., Kim, M.K., Jeong, J., Lee, J.K., Son, W.C., Park, J.H., 2014. A 90-day study of sub-chronic oral toxicity of 20 nm positively charged zinc oxide nanoparticles in Sprague Dawley rats. *Int. J. Nanomed.* 9, 93–107. <http://dx.doi.org/10.2147/IJN.S57927>.
- Parveen, A., Rizvi, S.H., Gupta, A., Singh, R., Ahmad, I., Mahdi, F., Mahdi, A.A., 2012. NMR-based metabolomics study of sub-acute hepatotoxicity induced by silica nanoparticles in rats after intranasal exposure. *Cell. Mol. Biol. (Noisy-le-grand)* 58, 196–203. <http://dx.doi.org/10.1170/T941>.
- Patil, N.A., Gade, W.N., Deobagkar, D.D., 2016. Epigenetic modulation upon exposure of lung fibroblasts to TiO<sub>2</sub> and ZnO nanoparticles: alterations in DNA methylation. *Int. J. Nanomed.* 11, 4509–4519.
- Pelclova, D., Barosova, H., Kukutschova, J., Zdimal, V., Navratil, T., Fenclova, Z., Vlkova, S., Schwarz, J., Zikova, N., Kacer, P., Komarc, M., Belacek, J., Zakharov, S., 2015. Raman microspectroscopy of exhaled breath condensate and urine in workers exposed to fine and nano TiO<sub>2</sub> particles: a cross-sectional study. *J. Breath Res.* 9, 036008. <http://dx.doi.org/10.1088/1752-7155/9/3/036008>.
- Pelclova, D., Zdimal, V., Kacer, P., Fenclova, Z., Vlkova, S., Syslova, K., Navratil, T., Schwarz, J., Zikova, N., Barosova, H., Turci, F., Komarc, M., Pelcl, T., Belacek, J., Kukutschova, J., Zakharov, S., 2016a. Oxidative stress markers are elevated in exhaled breath condensate of workers exposed to nanoparticles during iron oxide pigment production. *J. Breath Res.* 10, 016004. <http://dx.doi.org/10.1088/1752-7155/10/1/016004>.
- Pelclova, D., Zdimal, V., Fenclova, Z., Vlkova, S., Turci, F., Corazzari, I., Kacer, P., Schwarz, J., Zikova, N., Makes, O., Syslova, K., Komarc, M., Belacek, J., Navratil, T., Machajova, M., Zakharov, S., 2016b. Markers of oxidative damage of nucleic acids and proteins among workers exposed to TiO<sub>2</sub> (nano) particles. *Occup. Environ. Med.* 73, 110–118. <http://dx.doi.org/10.1136/oemed-2015-103161>.
- Pelclova, D., Zdimal, V., Kacer, P., Fenclova, Z., Vlkova, S., Komarc, M., Navratil, T., Schwarz, J., Zikova, N., Makes, O., Syslova, K., Belacek, J., Zakharov, S., 2016c. Leukotrienes in exhaled breath condensate and fractional exhaled nitric oxide in workers exposed to TiO<sub>2</sub> nanoparticles. *J. Breath Res.* 10, 036004. <http://dx.doi.org/10.1088/1752-7155/10/3/036004>.
- Pelclova, D., Zdimal, V., Kacer, P., Zikova, N., Komarc, M., Fenclova, Z., Vlkova, S., Schwarz, J., Makeš, O., Syslova, K., Navratil, T., Turci, F., Corazzari, I., Zakharov, S., Bello, D., 2017a. Markers of lipid oxidative damage in the exhaled breath condensate of nano TiO<sub>2</sub> production workers. *Nanotoxicology* 11, 52–63. <http://dx.doi.org/10.1080/17435390.2016.1262921>.
- Pelclova, D., Zdimal, V., Kacer, P., Komarc, M., Fenclova, Z., Vlkova, S., Zikova, N., Schwarz, J., Makes, O., Navratil, T., Zakharov, S., Bello, D., 2017b. Markers of lipid oxidative damage among office workers exposed intermittently to air pollutants including nanoTiO<sub>2</sub> particles. *Rev. Environ. Health* 32, 193–200. <http://dx.doi.org/10.1515/reveh-2016-0030>.
- Peschansky, V.J., Wahlestedt, C., 2014. Non-coding RNAs as direct and indirect modulators of epigenetic regulation. *Epigenetics* 9, 3–12. <http://dx.doi.org/10.4161/epi.27473>.
- Peters, A., Rückerl, R., Cyrys, J., 2011. Lessons from air pollution epidemiology for studies of engineered nanomaterials. *J. Occup. Environ. Med.* 53, S8–S13. <http://dx.doi.org/10.1097/JOM.0b013e31821ad5c0>.
- Phuyal, S., Kasem, M., Knittelfelder, O., Sharma, A., Fonseca, D.M., Vebrat, V., Shaposhnikov, S., Slupphaug, G., Skaug, V., Zienoldiny, S., 2018. Characterization of the proteome and lipidome profiles of human lung cells after low dose and chronic exposure to multiwalled carbon nanotubes. *Nanotoxicology* 12, 138–152. <http://dx.doi.org/10.1080/17435390.2018.1425500>.
- Pietroiusti, A., Stockmann-Juvala, H., Lucaroni, F., Savolainen, K., 2018. Nanomaterial exposure, toxicity, and impact on human health. *Wiley Interdiscip. Rev.: Nanomed. Nanobiotechnol.* e1513. <http://dx.doi.org/10.1002/wnan.1513>.
- Pisani, C., Gaillard, J.C., Nouvel, V., Odorico, M., Armengaud, J., Prat, O., 2015. High-throughput, quantitative assessment of the effects of low-dose silica nanoparticles on lung cells: grasping complex toxicity with a great depth of field. *BMC Genomics* 16, 315. <http://dx.doi.org/10.1186/s12864-015-1521-5>.
- Porter, D.W., Wu, N., Hubbs, A.F., Mercer, R.R., Funk, K., Meng, F., Li, J., Wolfarth, M.G., Battelli, L., Friend, S., Andrew, M., Hamilton, R.Jr., Sriram, K., Yang, F., Castranova, V., Holian, A., 2013. Differential mouse pulmonary dose and time course responses to titanium dioxide nanospheres and nanobelts. *Toxicol. Sci.* 131, 179–193. <http://dx.doi.org/10.1093/toxsci/kfs261>.
- Prow, T.W., Grice, J.E., Lin, L.L., Faye, R., Butler, M., Becker, W., Wurm, E.M., Yoong, C., Robertson, T.A., Soyer, H.P., Roberts, M.S., 2011. Nanoparticles and microparticles for skin drug delivery. *Adv. Drug Deliv. Rev.* 63, 470–491. <http://dx.doi.org/10.1016/j.addr.2011.01.012>.
- Qian, Y., Zhang, J., Hu, Q., Xu, M., Chen, Y., Hu, G., Zhao, M., Liu, S., 2015. Silver nanoparticle-induced hemoglobin decrease involves alteration of histone 3 methylation status. *Biomaterials* 70, 12–22. <http://dx.doi.org/10.1016/j.biomaterials.2015.08.015>.
- Reddy, A.R., Rao, M.V., Krishna, D.R., Himabindu, V., Reddy, Y.N., 2011. Evaluation of oxidative stress and anti-oxidant status in rat serum following exposure of carbon nanotubes. *Regul. Toxicol. Pharmacol.* 59, 251–257. <http://dx.doi.org/10.1016/j.yrtph.2010.10.007>.
- Renwick, L.C., Donaldson, K., Clouter, A., 2001. Impairment of alveolar macrophage phagocytosis by ultrafine particles. *Toxicol. Appl. Pharmacol.* 172, 119–127. <http://dx.doi.org/10.1006/taap.2001.9128>.
- Riebeling, C., Wiemann, M., Schnekenburger, J., Kuhlbusch, T.A., Wohlleben, W., Luch, A., Haese, A., 2016. A redox proteomics approach to investigate the mode of action of nanomaterials. *Toxicol. Appl. Pharmacol.* 299, 24–29. <http://dx.doi.org/10.1016/j.taap.2016.01.019>.
- Rittinghausen, S., Hackbarth, A., Creutzenberg, O., Ernst, H., Heinrich, U., Leonhardt, A., Schaudien, D., 2014. The carcinogenic effect of various multi-walled carbon nanotubes (MWCNTs) after intraperitoneal injection in rats. *Part. Fibre Toxicol.* 11, 59. <http://dx.doi.org/10.1186/s12989-014-0059-z>.
- Ritz, S., Schöttler, S., Kotman, N., Baier, G., Musyanovych, A., Kuharev, J., Landfester, K., Schild, H., Jahn, O., Tenzer, S., Mailänder, V., 2015. Protein corona of nanoparticles: distinct proteins regulate the cellular uptake. *Biomacromolecules* 16, 1311–1321. <http://dx.doi.org/10.1021/acs.biomac.5b00108>.
- Robertson, D.G., Watkins, P.B., Reily, M.D., 2011. Metabolomics in toxicology: preclinical and clinical applications. *Toxicol. Sci.* 120, S146–S170. <http://dx.doi.org/10.1093/toxsci/kfq358>.
- Roco, M.C., 1997. Introduction, WTEC Workshop Report on R&D Status and Trends in Nanoparticles, Nanostructured Materials, and Nanodevices in the United States. World Technology Evaluation Center (WTEC), Baltimore (Accessed 30 January 2018). <http://www.wtec.org/loyola/pdf/nanousws.pdf>.
- Sadiq, R., Bhalli, J.A., Yan, J., Woodruff, R.S., Pearce, M.G., Li, Y., Mustafa, T., Watanabe, F., Pack, L.M., Biris, A.S., Khan, Q.M., Chen, T., 2012. Genotoxicity of TiO<sub>2</sub> anatase nanoparticles in B6C3F1 male mice evaluated using pig-a and flow cytometric micronucleus assays. *Mutat. Res.* 745, 65–72. <http://dx.doi.org/10.1016/j.mrgentox.2012.02.002>.
- Sargent, L.M., Porter, D.W., Staska, L.M., Hubbs, A.F., Lowry, D.T., Battelli, L., Siegrist, K.J., Kashon, M.L., Mercer, R.R., Bauer, A.K., Chen, B.T., Salisbury, J.L., Frazer, D., McKinney, W., Andrew, M., Tsuruoka, S., Endo, M., Fluharty, K.L., Castranova, V., Reynolds, S.H., 2014. Promotion of lung adenocarcinoma following inhalation exposure to multi-walled carbon nanotubes. *Part. Fibre Toxicol.* 11 (3). <http://dx.doi.org/10.1186/1743-8977-11-3>.
- Sarkar, A., Das, J., Manna, P., Sil, P.C., 2011. Nano-copper induces oxidative stress and apoptosis in kidney via both extrinsic and intrinsic pathways. *Toxicology* 290, 208–217. <http://dx.doi.org/10.1016/j.tox.2011.09.086>.
- Sayes, C.M., Wahj, R., Kurian, P.A., Liu, Y., West, J.L., Ausman, K.D., Warheit, D.B., Colvin, V.L., 2006. Correlating nanoscale titania structure with toxicity: a cytotoxicity and inflammatory response study with human dermal fibroblasts and human lung epithelial cells. *Toxicol. Sci.* 92, 174–185. <http://dx.doi.org/10.1093/toxsci/kfj197>.
- Schmid, O., Stoeger, T., 2016. Surface area is the biologically most effective metric for acute nanoparticle toxicity of the lung. *J. Aerosol. Sci.* 99, 133–143. <http://dx.doi.org/10.1016/j.jaerosci.2015.12.006>.

- Schulte, P.A., Hauser, J.E., 2012. The use of biomarkers in occupational health research, practice, and policy. *Toxicol. Lett.* 213, 91–99. <http://dx.doi.org/10.1016/j.toxlet.2011.03.027>.
- Schulte, P.A., Smith, A., et al., 2011. Ethical issues in molecular epidemiologic research. In: Rothman (Ed.), *Molecular Epidemiology: Principles and Practices*. International Agency for Research on Cancer, Lyon, pp. 9–21.
- Schulte, P.A., Talaska, G., 1995. Validity criteria for the use of biological markers of exposure to chemical agents in environmental epidemiology. *Toxicology* 101, 73–88. [http://dx.doi.org/10.1016/0300-483X\(95\)03020-G](http://dx.doi.org/10.1016/0300-483X(95)03020-G).
- Schulte, P.A., Iavicoli, I., Rantanen, J.H., Dahmann, D., Iavicoli, S., Pipke, R., Guseva Canu, I., Bocconi, F., Ricci, M., Polci, M.L., Sabbioni, E., Pietrousti, A., Mantovani, E., 2016. Assessing the protection of the nanomaterial workforce. *Nanotoxicology* 10, 1013–1019. <http://dx.doi.org/10.3109/17435390.2015.1132347>.
- Semmler-Behnke, M., Kreyling, W.G., Lipka, J., Fertsch, S., Wenk, A., Takenaka, S., Schmid, G., Brandau, W., 2008. Biodistribution of 1.4- and 18-nm gold particles in rats. *Small* 4, 2108–2111. <http://dx.doi.org/10.1002/sml.200800922>.
- Shrivastava, R., Kushwaha, P., Bhutia, Y.C., Flora, S., 2016. Oxidative stress following exposure to silver and gold nanoparticles in mice. *Toxicol. Ind. Health* 32, 1391–1404. <http://dx.doi.org/10.1177/0748233714562623>.
- Shvedova, A.A., Kisin, E.R., Mercer, R., Murray, A.R., Johnson, V.J., Potapovich, A.I., Tyurina, Y.Y., Gorelik, O., Arepalli, S., Schwegler-Berry, D., Hubbs, A.F., Antonini, J., Evans, D.E., Ku, B.K., Ramsey, D., Maynard, A., Kagan, V.E., Castranova, V., Baron, P., 2005. Unusual inflammatory and fibrogenic pulmonary responses to single-walled carbon nanotubes in mice. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 289, L698–L708. <http://dx.doi.org/10.1152/ajplung.00084.2005>.
- Shvedova, A.A., Yanamala, N., Kisin, E.R., Khailullin, T.O., Birch, M.E., Fatkhutdinova, L.M., 2016. Integrated analysis of dysregulated ncRNA and mRNA expression profiles in humans exposed to carbon nanotubes. *PLoS One* 11, e0150628. <http://dx.doi.org/10.1371/journal.pone.0150628>.
- Shyamasundar, S., Ng, C.T., Yung, L.Y., Dheen, S.T., Bay, B.H., 2015. Epigenetic mechanisms in nanomaterial-induced toxicity. *Epigenomics* 7, 395–411. <http://dx.doi.org/10.2217/epi.15.3>.
- Sierra, M.I., Valdés, A., Fernández, A.F., Torrecillas, R., Fraga, M.F., 2016. The effect of exposure to nanoparticles and nanomaterials on the mammalian epigenome. *Int. J. Nanomed.* 11, 6297–6306. <http://dx.doi.org/10.2147/IJN.S120104>.
- Singh, S.P., Kumari, M., Kumari, S.I., Rahman, M.F., Mahboob, M., Grover, P., 2013a. Toxicity assessment of manganese oxide micro and nanoparticles in Wistar rats after 28 days of repeated oral exposure. *J. Appl. Toxicol.* 33, 1165–1179. <http://dx.doi.org/10.1002/jat.2887>.
- Singh, S.P., Rahman, M.F., Murty, U.S., Mahboob, M., Grover, P., 2013b. Comparative study of genotoxicity and tissue distribution of nano and micron sized iron oxide in rats after acute oral treatment. *Toxicol. Appl. Pharmacol.* 266, 56–66. <http://dx.doi.org/10.1016/j.taap.2012.10.016>.
- Snyder-Talkington, B.N., Dymacek, J., Porter, D.W., Wolfarth, M.G., Mercer, R.R., Pacurari, M., Denvir, J., Castranova, V., Qian, Y., Guo, N.L., 2013. System-based identification of toxicity pathways associated with multi-walled carbon nanotube-induced pathological responses. *Toxicol. Appl. Pharmacol.* 272, 47–89. <http://dx.doi.org/10.1016/j.taap.2013.06.026>.
- Snyder-Talkington, B.N., Dong, C., Zhao, X., Dymacek, J., Porter, D.W., Wolfarth, M.G., Castranova, V., Qian, Y., Guo, N.L., 2015. Multi-walled carbon nanotube-induced gene expression in vitro: concordance with in vivo studies. *Toxicology* 328, 66–74. <http://dx.doi.org/10.1016/j.tox.2014.12.012>.
- Snyder-Talkington, B.N., Dong, C., Sargent, L.M., Porter, D.W., Staska, L.M., Hubbs, A.F., Raese, R., McKinney, W., Chen, B.T., Battelli, L., Lowry, D.T., Reynolds, S.H., Castranova, V., Qian, Y., Guo, N.L., 2016. mRNAs and miRNAs in whole blood associated with lung hyperplasia, fibrosis, and bronchiolo-alveolar adenoma and adenocarcinoma after multi-walled carbon nanotube inhalation exposure in mice. *J. Appl. Toxicol.* 236, 161–174. <http://dx.doi.org/10.1002/jat.3157>.
- Soares, T., Ribeiro, D., Proença, C., Chisté, R.C., Fernandes, E., Freitas, M., 2016. Size-dependent cytotoxicity of silver nanoparticles in human neutrophils assessed by multiple analytical approaches. *Life Sci.* 145, 247–254. <http://dx.doi.org/10.1016/j.lfs.2015.12.046>.
- Song, M.F., Li, Y.S., Kasai, H., Kawai, K., 2012. Metal nanoparticle-induced micronuclei and oxidative DNA damage in mice. *J. Clin. Biochem. Nutr.* 50, 211–216. <http://dx.doi.org/10.3164/jcbn.11-70>.
- Soni, D., Gandhi, D., Tarale, P., Bafana, A., Pandey, R.A., Sivanesan, S., 2017. Oxidative stress and genotoxicity of zinc oxide nanoparticles to pseudomonas species, human promyelocytic leukemia (HL-60), and blood cells. *Biol. Trace Elem. Res.* 178, 218–227. <http://dx.doi.org/10.1007/s12011-016-0921-y>.
- Srinivas, A., Rao, P.J., Selvam, G., Murthy, P.B., Reddy, P.N., 2011. Acute inhalation toxicity of cerium oxide nanoparticles in rats. *Toxicol. Lett.* 205, 105–115. <http://dx.doi.org/10.1016/j.toxlet.2011.05.1027>.
- Srinivas, A., Rao, P.J., Selvam, G., Goparaju, A., Murthy, P.B., Reddy, P.N., 2012. Oxidative stress and inflammatory responses of rat following acute inhalation exposure to iron oxide nanoparticles. *Hum. Exp. Toxicol.* 31, 1113–1131. <http://dx.doi.org/10.1177/0960327112446515>.
- Stirling, P.A., 2018. *The Nanotechnology Revolution: A Global Bibliographic Perspective*. CRC Press, Taylor Francis Group.
- Stone, V., Miller, M.R., Clift, M.J.D., Elder, A., Mills, N.L., Møller, P., Schins, R.P.F., Vogel, U., Kreyling, W.G., Alstrup, Jensen, K., Kuhlbusch, T.A.J., Schwarze, P.E., Hoet, P., Pietrousti, A., De Vizcaya-Ruiz, A., Baeza-Squiban, A., Teixeira, J.P., Tran, C.L., Cassee, F.R., 2017. Nanomaterials versus ambient ultrafine particles: an opportunity to exchange toxicology knowledge. *Environ. Health Perspect.* 125, 106002. <http://dx.doi.org/10.1289/EHP424>.
- Sumner, S.C., Fennell, T.R., Snyder, R.W., Taylor, G.F., Lewin, A.H., 2010. Distribution of carbon-14 labeled C60 (14C)C60 in the pregnant and in the lactating dam and the effect of C60 exposure on the biochemical profile of urine. *J. Appl. Toxicol.* 30, 354–360. <http://dx.doi.org/10.1002/jat.1503>.
- Sundarraj, K., Raghunath, A., Panneerselvam, L., Perumal, E., 2017. Iron oxide nanoparticles modulate heat shock proteins and organ specific markers expression in mice male accessory organs. *Toxicol. Appl. Pharmacol.* 317, 12–24. <http://dx.doi.org/10.1016/j.taap.2017.01.002>.
- Sung, J.H., Ji, J.H., Park, J.D., Yoon, J.U., Kim, D.S., Jeon, K.S., Song, M.Y., Jeong, J., Han, B.S., Han, J.H., Chung, Y.H., Chang, H.K., Lee, J.H., Cho, M.H., Kelman, B.J., Yu, I.J., 2009. Subchronic inhalation toxicity of silver nanoparticles. *Toxicol. Sci.* 108, 452–461. <http://dx.doi.org/10.1093/toxsci/kfn246>.
- Tabish, A.M., Poels, K., Byun, H.M., Luyts, K., Baccarelli, A.A., Martens, J., Kerkhofs, S., Seys, S., Hoet, P., Godderis, L., 2017. Changes in DNA methylation in mouse lungs after a single intra-tracheal administration of nanomaterials. *PLoS One* 12, e0169886. <http://dx.doi.org/10.1371/journal.pone.0169886>.
- Takenaka, S., Karg, E., Roth, C., Schulz, H., Ziesenis, A., Heinzmann, U., Schramel, P., Heyder, J., 2001. Pulmonary and systemic distribution of inhaled ultrafine silver particles in rats. *Environ. Health Perspect.* 109, 547–551. <http://dx.doi.org/10.2307/3454667>.
- Takenaka, S., Karg, E., Kreyling, W.G., Lentner, B., Möller, W., Behnke-Semmler, M., Jennen, L., Walch, A., Michalke, B., Schramel, P., Heyder, J., Schulz, H., 2006. Distribution pattern of inhaled ultrafine gold particles in the rat lung. *Inhal. Toxicol.* 18, 733–740. <http://dx.doi.org/10.1080/08958370600748281>.
- Tang, M., Zhang, T., Xue, Y., Wang, S., Huang, M., Yang, Y., Lu, M., Lei, H., Kong, L., Yuepu, P., 2010. Dose dependent in vivo metabolic characteristics of titanium dioxide nanoparticles. *J. Nanosci. Nanotechnol.* 10, 8575–8583. <http://dx.doi.org/10.1166/jnn.2010.2482>.
- Tang, M., Zhang, T., Xue, Y., Wang, S., Huang, M., Yang, Y., Lu, M., Lei, H., Kong, L., Wang, Y., Pu, Y., 2011. Metabonomic studies of biochemical changes in the serum of rats by intratracheally instilled TiO<sub>2</sub> nanoparticles. *J. Nanosci. Nanotechnol.* 11, 3065–3074. <http://dx.doi.org/10.1166/jnn.2011.3604>.
- Tarasova, N.K., Gallud, A., Ytterberg, A.J., Chernobrovkin, A., Aranzas, J.R., Astruc, D., Antipov, A., Fedutik, Y., Fadeel, B., Zubarev, R.A., 2017. Cytotoxic and proinflammatory effects of metal-based nanoparticles on THP-1 monocytes characterized by combined proteomics approaches. *J. Proteome Res.* 16, 689–697. <http://dx.doi.org/10.1021/acs.jproteome.6b00747>.
- Tavares, A., Louro, H., Antunes, S., Quarre, S., Simar, S., De Temmerman, P.J., Verleysen, E., Mast, J., Jensen, K.A., Norrpa, H., Nesslany, F., Silva, M.J., 2014. Genotoxicity evaluation of nanosized titanium dioxide, synthetic amorphous silica and multi-walled carbon nanotubes in human lymphocytes. *Toxicol. In Vitro* 28, 60–69. <http://dx.doi.org/10.1016/j.tiv.2013.06.009>.
- Thiruvengadam, M., Rajakumar, G., Chung, I.M., 2018. Nanotechnology: current uses and future applications in the food industry. *Biotechnology* 8, 74. <http://dx.doi.org/10.1007/s13205-018-1104-7>.
- Tiwari, D.K., Jin, T., Behari, J., 2011. Dose-dependent in vivo toxicity assessment of silver nanoparticle in Wistar rats. *Toxicol. Mech. Methods* 21, 13–24. <http://dx.doi.org/10.3109/15376516.2010.529184>.
- Trouiller, B., Reliene, R., Westbrook, A., Solaimani, P., Schiestl, R.H., 2009. Titanium dioxide nanoparticles induce DNA damage and genetic instability in vivo in mice. *Cancer Res.* 69, 8784–8789. <http://dx.doi.org/10.1158/0008-5472.CAN-09-2496>.
- Uboldi, C., Urbán, P., Gilliland, D., Bajak, E., Valsami-Jones, E., Ponti, J., Rossi, F., 2016. Role of the crystalline form of titanium dioxide nanoparticles: rutile, and not anatase, induces toxic effects in Balb/3T3 mouse fibroblasts. *Toxicol. In Vitro* 31, 137–145. <http://dx.doi.org/10.1016/j.tiv.2015.11.005>.
- Vasantharaja, D., Ramalingam, V., Aadinaath, Reddy, G.G., 2015. Titanium dioxide nanoparticles induced alteration in haematological indices of adult male wistar rats. *J. Acad. Ind. Res.* 3, 632–635.
- Villalta, P.W., Balbo, S., 2017. The future of DNA adductomic analysis. *Int. J. Mol. Sci.* 18, 1870. <http://dx.doi.org/10.3390/ijms18091870>.
- Vlaenderen, J., Pronk, A., Rothman, N., Hildesheim, A., Silverman, D., Hosgood, H.D., Spaan, S., Kuijpers, E., Godderis, L., Hoet, P., Lan, Q., Vermeulen, R., 2017. A cross-sectional study of changes in markers of immunological effects and lung health due to exposure to multi-walled carbon nanotubes. *Nanotoxicology* 11, 395–404. <http://dx.doi.org/10.1080/17435390.2017.1308031>.
- Wu, W.T., Liao, H.Y., Chung, Y.T., Li, W.F., Tsou, T.C., Li, L.A., Lin, M.H., Ho, J.J., Wu, T.N., Liou, S.H., 2014. Effect of nanoparticles exposure on fractional exhaled nitric oxide (FENO) in workers exposed to nanomaterials. *Int. J. Mol. Sci.* 15, 878–894. <http://dx.doi.org/10.3390/ijms15010878>.
- Xia, T., Hamilton, R.F., Bonner, J.C., Crandall, E.D., Elder, A., Fazlollahi, F., Girtsman, T.A., Kim, K., Mitra, S., Ntim, S.A., Orr, G., Tagmount, M., Taylor, A.J., Telesca, D., Tolic, A., Vulpe, C.D., Walker, A.J., Wang, X., Witzmann, F.A., Wu, N., Xie, Y., Zink, J.I., Nel, A., Holian, A., 2013. Interlaboratory evaluation of in vitro cytotoxicity and inflammatory responses to engineered nanomaterials: the NIEHS nano GO consortium. *Environ. Health Perspect.* 121, 683–690. <http://dx.doi.org/10.1289/ehp.1306561>.
- Xie, J., Dong, W., Liu, R., Wang, Y., Li, Y., 2018. Research on the hepatotoxicity mechanism of citrate-modified silver nanoparticles based on metabolomics and proteomics. *Nanotoxicology* 12, 18–31. <http://dx.doi.org/10.1080/17435390.2017.1415389>.
- Yazdimaghani, M., Moos, P.J., Ghandehari, H., 2017. Global gene expression analysis of macrophage response induced by nonporous and porous silica nanoparticles. *Nanomedicine* 14, 533–545. <http://dx.doi.org/10.1016/j.nano.2017.11.021>.
- Yu, L.E., Yung, L.Y.L., Ong, C.N., Tan, Y.L., Balasubramanian, K.S., Hartono, D., Shui, G.H., Wenk, M.R., Ong, W.Y., 2007. Translocation and effects of gold nanoparticles after inhalation exposure in rats. *Nanotoxicology* 1, 235–242. <http://dx.doi.org/10.1080/17435390701763108>.
- Zhang, R., Dai, Y., Zhang, X., Niu, Y., Meng, T., Li, Y., Duan, H., Bin, P., Ye, M., Jia, X.,

- Shen, M., Yu, S., Yang, X., Gao, W., Zheng, Y., 2014. Reduced pulmonary function and increased pro-inflammatory cytokines in nanoscale carbon black-exposed workers. Part. Fibre Toxicol. 11, 73. <http://dx.doi.org/10.1186/s12989-014-0073-1>.
- Zhao, Y., Sun, H., Wang, H., 2016. Long noncoding RNAs in DNA methylation: new players stepping into the old game. Cell Biosci. 6, 45. <http://dx.doi.org/10.1186/s13578-016-0109-3>.
- Zhao, L., Zhu, Y., Chen, Z., Xu, H., Zhou, J., Tang, S., Xu, Z., Kong, F., Li, X., Zhang, Y., Li, X., Zhang, J., Jia, G., 2018. Cardiopulmonary effects induced by occupational exposure to titanium dioxide nanoparticles. Nanotoxicology 11, 1–16. <http://dx.doi.org/10.1080/17435390.2018.1425502>.
- Zhu, M.T., Feng, W.Y., Wang, Y., Wang, B., Wang, M., Ouyang, H., Zhao, Y.L., Chai, Z.F., 2009. Particokinetics and extrapulmonary translocation of intratracheally instilled ferric oxide nanoparticles in rats and the potential health risk assessment. Toxicol. Sci. 107, 342–351. <http://dx.doi.org/10.1093/toxsci/kfn245>.
- Ziemba, B., Matuszko, G., Appelhans, D., Voit, B., Bryszewska, M., Klajnert, B., 2012. Genotoxicity of poly(propylene imine) dendrimers. Biopolymers 97, 642–648. <http://dx.doi.org/10.1002/bip.22056>.