



Preliminary assessment of exposure to persistent organic pollutants among pregnant women in Puerto Rico

Deborah J. Watkins^{a,*}, Carmen M. Vélez-Vega^b, Zaira Rosario^b, José F. Cordero^c, Akram N. Alshawabkeh^d, John D. Meeker^a

^a Department of Environmental Health Sciences, University of Michigan School of Public Health, Ann Arbor, MI, USA

^b University of Puerto Rico Graduate School of Public Health, Medical Sciences Campus, San Juan, PR, USA

^c Department of Epidemiology and Biostatistics, University of Georgia, Athens, GA, USA

^d College of Engineering, Northeastern University, Boston, MA, USA

ARTICLE INFO

Keywords:

Puerto Rico
Persistent organic pollutants
Polychlorinated biphenyls (PCBs)
Organochlorine pesticides (OCPs)
Per- and polyfluoroalkyl substances (PFASs)
Polybrominated diphenyl ethers (PBDEs)

ABSTRACT

Background: Puerto Rico has over 200 hazardous waste sites, as well as higher rates of several adverse health outcomes compared to the mainland US. In response to concerns of potential links between environmental contaminant exposure and preterm birth, the Puerto Rico Testsite for Exploring Contamination Threats (PROTECT) birth cohort was formed. Biomarkers of exposure for several contaminants have been found to be elevated in this cohort compared to women of child-bearing age in the National Health and Nutrition Examination Survey (NHANES). However, exposure to persistent organic pollutants (POPs) has not been evaluated.

Methods: In this preliminary analysis, we measured four classes of POPs, including perfluoroalkyl substances (PFASs), polybrominated diphenyl ethers (PBDEs), polychlorinated biphenyls (PCBs), and several persistent pesticides in serum samples collected at 16–20 weeks gestation from the first 48 women enrolled in PROTECT. We performed descriptive analyses for each analyte, assessed correlations between analytes using Spearman correlations, and compared serum levels in PROTECT women to levels in NHANES women aged 18–40.

Results: Several PFASs were detected in 96–100% of samples, with moderate to strong correlations between most PFASs (range $r = 0.44$ – 0.88). BDE47, BDE153, PCB 138–158, PCB153 and p,p'-dichlorodiphenyldichloroethene (p,p'-DDE) were detected in the majority of samples, with strong correlations between PCBs and p,p'-DDE (range $r = 0.59$ – 0.74). The median concentration for each analyte was lower than, sometimes by a factor of 4 (e.g. BDE47, p,p'-DDE), the median concentration reported in NHANES women aged 18–40.

Conclusion: Although participants in the ongoing PROTECT birth cohort have higher exposure to many environmental contaminants, this preliminary study suggests that they have lower exposure to several POPs, specifically PCBs, OCPs, PFASs, and PBDEs.

1. Introduction

Since the 1950s, the island of Puerto Rico has undergone rapid industrialization, resulting in a legacy of toxic spills, chemical waste, air pollution, and contamination of the surrounding environment (Gioda et al., 2007; Hunter and Arbona, 1995; Padilla et al., 2011). Currently, Puerto Rico has over 200 hazardous waste sites, including 18 active Superfund sites (EPA, 2018). In addition, Puerto Rico has higher rates of several adverse health outcomes, such as preterm birth (Martin et al., 2018), childhood asthma (CDC, 2017b), autism (Cordero, 2013), and obesity (Garza et al., 2011; Otero-Gonzalez and Garcia-Fragoso, 2008;

Rivera-Soto et al., 2010), compared to the mainland US. In response to concerns of potential links between environmental contaminant exposure and preterm birth, the Puerto Rico Testsite for Exploring Contamination Threats (PROTECT) birth cohort was formed as part of a NIEHS Superfund Research Program. Biomarkers of exposure for several contaminants have been found to be elevated among pregnant women in this cohort, including several phthalates (Cantonwine et al., 2014), phenols (Meeker et al., 2013), certain polycyclic aromatic hydrocarbons (Cathey et al., 2018), and certain metals, compared to female participants in the National Health and Nutrition Examination Survey (NHANES) aged 18–40. Notably, we recently reported that the

* Corresponding author. Department of Environmental Health Sciences, University of Michigan School of Public Health, 6611C SPH I, 1415 Washington Heights, Ann Arbor, MI, 48109, USA.

E-mail address: debjwat@umich.edu (D.J. Watkins).

<https://doi.org/10.1016/j.ijheh.2019.02.001>

Received 25 October 2018; Received in revised form 23 January 2019; Accepted 1 February 2019

1438-4639/© 2019 Elsevier GmbH. All rights reserved.

Abbreviations

AAA	Autoridad de Acueductos y Alcantarillados
β-HCCH	hexachlorocyclohexane
DDT	dichlorodiphenyltrichloroethane
DDE	dichlorodiphenyldichloroethene
Et-FOSAA	2-(N-ethyl-perfluorooctane sulfonamide) acetate
GC-IDHRMS	gas chromatography isotope dilution high resolution mass spectrometry
γ-HCCH	γ-hexachlorobenzene
HCB	hexachlorobenzene
LOD	limit of detection
Me-FOSAA	2-(N-methyl-perfluorooctane sulfonamide) acetate
NHANES	National Health and Nutrition Examination Survey
OCP	organochlorine pesticide

PBDE	polybrominated diphenyl ether
PCB	polychlorinated biphenyl
PFAS	per- and polyfluoroalkyl substance
PFDeA	perfluorodecanoate
PFHxS	perfluorohexane sulfonate
PFNA	perfluorononanoate
PFOA	perfluorooctanoic acid
PFOS	perfluorooctane sulfonate
FOSA	perfluorooctane sulfonamide
POP	persistent organic pollutant
PROTECT	Puerto Rico Testsite for Exploring Contamination Threats
SPE-HPLC-MS/MS	solid phase extraction, isotope dilution-high performance liquid chromatography tandem mass spectrometry

antibacterial compounds triclosan and triclocarban were significantly higher in PROTECT women compared to NHANES (Ashrap et al., 2018). However, exposure to persistent organic pollutants (POPs), specifically polychlorinated biphenyls (PCBs), organochlorine pesticides (OCPs), per- and polyfluoroalkyl substances (PFASs), and polybrominated diphenyl ethers (PBDEs), has not previously been evaluated within the PROTECT cohort.

POPs are toxic chemicals that resist degradation, and as a result, can be transported far from their point of release in the environment. POPs also bioaccumulate in wildlife and in humans, with wide-ranging adverse health effects (EPA, 2017d; Fry and Power 2017). In response to concerns for environmental and human health, over 150 countries have signed the Stockholm Convention on Persistent Organic Pollutants since 2001, indicating agreement to reduce or eliminate the production, use, or release of specific POPs (Secretariat of the Stockholm Convention Clearing House, 2008; UNEP, 2009), including the four classes of compounds discussed here (PCBs, OCPs, PFASs, and PBDEs) and many others. Importantly, the US signed the original Convention, but has not yet ratified the agreement.

PCBs were used as coolants, lubricants, and hydraulic fluids in heat transfer and electrical equipment, as well as plasticizers and dyes, and exposure to PCBs has been associated with cancer, as well as effects on immune function, birth outcomes, endocrine disruption, and neurodevelopment (EPA, 2017b). In 1979, production of PCBs was banned as a result of environmental and human health concerns, but PCB exposure continues to be widespread due to their persistence in the environment and presence in contaminated food (CDC, 2017a; Xue et al., 2014).

OCPs include dichlorodiphenyltrichloroethane (DDT), lindane, dieldrin, hexachlorobenzene, chlordane, mirex, and others, many of which were used to control a range of insects. Human exposure to OCPs has been associated with a variety of adverse health effects, including cancer, neurotoxicity, and reproductive toxicity (Mrema et al., 2013). Again, due to environmental and health concerns they have been largely restricted or banned in the US. As a result, exposure to OCPs has been greatly reduced, but their environment persistence means that exposure to specific OC pesticides continues (CDC, 2017a). In addition, DDT is still widely used in malaria-endemic areas (Mrema et al., 2013).

PFASs are man-made chemicals that are oil and water resistant and highly stable. Because of these properties, PFASs have been used in numerous consumer products, including stain repellants, textiles, cookware, food packaging, cleaning products, and fire-fighting foams (EPA, 2018). Exposure to longer chain PFASs, such as perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS), has been associated with increased risk of cancer, low birth weight, and immune and endocrine system disruption (Bach et al., 2015; DeWitt et al., 2018; Grandjean et al., 2017; Rappazzo et al., 2017; Steenland et al., 2010). Although longer chain PFASs have been phased out of use in the US, use of shorter chain replacement PFASs has increased despite limited

information on their toxicity (Birnbaum and Grandjean, 2015; Brendel et al., 2018). Due to their persistence in the environment, PFASs have also become emerging drinking water contaminants in several communities in the US and worldwide (Banzhaf et al., 2017; Essumang et al., 2017; Guelfo and Adamson, 2018; Hu et al., 2016).

Finally, PBDEs are a class of chemicals that have been widely used as flame retardants in a range of consumer products, including polyurethane foam furniture, textiles, and electronics. In human and experimental studies, PBDE exposure has been associated with cancer, neurotoxicity, reproductive toxicity, and thyroid hormone disruption (EPA, 2017a). PBDEs were phased out of use in the US between 2004 and 2013, but exposure continues due to continued use of older PBDE-containing products, importation of PBDE-containing products from other countries, and persistence of these chemicals in the environment (EPA, 2017c).

To our knowledge, no biomonitoring data exists on POP exposure in Puerto Rico. The goal of this preliminary analysis was to characterize exposure to the above chemicals among pregnant women in the PROTECT cohort, as identification of elevated exposures in this population is crucial to our overall research objectives. Specifically, understanding how POPs may contribute to the overall mixture of environmental contaminant exposures will aid our investigation of exposures, preterm birth, and child development.

2. Methods

2.1. Study population

The PROTECT project, which began in 2011, is an ongoing prospective birth cohort study in Northern Puerto Rico designed to investigate relationships between environmental exposures and adverse birth outcomes. The present analysis was conducted on 48 of the first PROTECT participants with sufficient serum sample volume availability, for whom recruitment and inclusion criteria have been previously described (Cantonwine et al., 2014; Meeker et al., 2013). Briefly, women were recruited early in pregnancy (< 20 weeks gestation) in 2011–2013 from prenatal clinics and hospitals in Northern Puerto Rico, and were eligible to participate if they were between the ages of 18 and 40 years, had a singleton pregnancy, did not use oral contraceptives within three months before pregnancy or in vitro fertilization to get pregnant, and were free of known medical and obstetric complications. At the first study visit (16–20 weeks gestation), women provided a blood sample, urine sample, and demographic information via nurse-administered questionnaires. Although additional biological samples and questionnaire data were collected at subsequent prenatal study visits as previously described (Aker et al., 2016; Johns et al., 2015), because the chemicals of interest are biologically persistent they were measured only in samples collected at the first study visit for this

preliminary analysis. The study was approved by the ethics committees of the University of Puerto Rico, Northeastern University, the University of Michigan School of Public Health, and the University of Georgia. All participants provided full informed consent prior to enrollment.

2.2. Measurement of persistent organic pollutants

Serum collected from participants at the first prenatal study visit was separated into aliquots, frozen and stored at -80°C , and then shipped on dry ice to laboratories at the Centers for Disease Control and Prevention (CDC) for analysis. Perfluoroalkyl substances (PFASs) were measured using on-line solid phase extraction coupled to isotope dilution-high performance liquid chromatography tandem mass spectrometry (SPE-HPLC-MS/MS) as previously described (Kato et al., 2011). Measured PFASs comprised linear and branched perfluorooctanoate isomers (n-PFOA and Sb-PFOA, respectively), linear and branched isomers of perfluorooctane sulfonate (n-PFOS and Sm-PFOS, respectively), perfluorooctane sulfonamide (FOSA), perfluorohexane sulfonate (PFHxS), perfluorononanoate (PFNA), perfluorodecanoate (PFDeA), 2-(N-ethyl-perfluorooctane sulfonamide) acetate (Et-FOSAA), and 2-(N-methyl-perfluorooctane sulfonamide) acetate (Me-FOSAA). Total PFOS was calculated by summing n-PFOS and Sm-PFOS isomers, while only n-PFOA was used in analyses as Sb-PFOA was detected in only one sample. Polybrominated diphenyl ethers (PBDEs), polychlorinated biphenyls (PCBs), and persistent pesticides were measured using liquid/liquid extraction and gas chromatography isotope dilution high resolution mass spectrometry (GC-IDHRMS) as previously described (Jones et al., 2012). Eleven PBDE congeners were measured (BDE-17, 28, 47, 66, 85, 99, 100, 153, 154, 183, 209) as well as hexabromobiphenyl (PBB-153). Measured PCB congeners comprised PCB-28, 66, 74, 99, 105, 114, 118, 138–158, 146, 153, 156, 157, 167, 170, 178, 180, 183, 187, 189, 194, 196–203, 199, 206, and 209. Persistent pesticides included hexachlorobenzene (HCB), β -hexachlorocyclohexane (β -HCCH), γ -hexachlorobenzene (γ -HCCH, or lindane), oxychlorodane, trans-nonachlor, mirex, p,p'-dichlorodiphenyltrichloroethane (p,p'-DDT), o,p'-DDT, and p,p'-dichlorodiphenyldichloroethene (DDE, a DDT metabolite). Total cholesterol and total triglycerides were measured using commercially available test kits (Roche Diagnostics Corp.; Indianapolis, IN) and a Hitachi 912 chemistry analyzer (Hitachi; Tokyo, Japan). All values below their respective limit of detection (LOD) were replaced with the $\text{LOD}/\sqrt{2}$. Total PBDE, PCB, and pesticide concentrations were adjusted for serum total lipid concentration and reported in ng/g lipid weight.

Table 1

Distributions of persistent organic pollutant concentrations in serum samples collected from 2011 to 2013 from a subset of PROTECT participants (n = 48) and NHANES women aged 18–40 years.

Percentiles												
Analyte	LOD ^a	% > LOD	Mean	Min	25 th	50 th	75 th	95 th	NHANES 50 th women 18–40	NHANES 95th women 18–40	NHANES n	NHANES cycle
PFASs (μg/L)												
n-PFOA	0.1	100	1.16	0.20	0.60	0.90	1.13	3.10	1.10	2.90	375	2013–2014
n-PFOS	0.1	100	1.66	0.20	0.78	1.50	2.33	3.30	2.30	9.00	375	2013–2014
PFDeA	0.1	64.6	0.17	0.07	0.07	0.10	0.20	0.60	0.20	0.70	760	2009–2010
PFHxS	0.1	95.8	0.34	0.07	0.20	0.20	0.30	0.90	0.70	3.25	760	2009–2010
PFNA	0.1	97.9	0.42	0.07	0.20	0.30	0.50	1.00	0.70	2.03	760	2009–2010
Sm-PFOS	0.1	97.9	0.61	0.07	0.30	0.55	0.93	1.20	0.60	2.00	375	2013–2014
PBDEs (ng/g lipid)												
BDE153	2.3	70.8	5.55	1.27	2.08	3.87	6.57	12.8	4.90	36.6	363	2003–2004
BDE47	3.6	68.8	7.50	1.56	3.26	4.66	7.94	21.8	22.8	127	361	2003–2004
PCBs (ng/g lipid)												
PCB138-158	2.3	56.3	2.84	1.34	1.84	2.33	3.31	5.96	6.67	27.5	343	2003–2004
PCB153	2.3	77.1	3.64	1.41	2.23	3.15	3.82	9.12	8.23	31.7	342	2003–2004
Pesticides (ng/g lipid)												
p,p'-DDE	11.5	95.8	32.3	8.49	21.1	30.9	40.6	55.5	135	850	345	2003–2004

^a Median LOD; BDE153, PCB138–158, PCB153 range = 1.4–3.3 ng/g lipid; BDE47 range = 2.1–5.2 ng/g lipid; p,p'-DDE range = 6.8–17 ng/g lipid.

2.3. Statistical analysis

Descriptive analyses were performed for each analyte, including determination of the percent of serum samples with concentrations below the LOD for each chemical, means, and distribution percentiles, as well as Spearman correlations between chemicals detected in at least 50% of samples.

3. Results

The LODs, means, and distributions of all analytes detected in greater than 50% of serum samples are shown in Table 1. Several PFASs were highly detected, including the linear PFOA isomer, linear and branched PFOS isomers, PFNA, and PFHxS (96–100% above LOD). BDE47 and BDE153 were the only PBDEs detected in the majority of samples, while PCB 138–158 and PCB153 were the only commonly detected PCBs. Of the persistent pesticides that were measured, only p,p'-DDE was detected in the majority of samples. LODs and the number and percent of samples above the LOD for all other analytes are presented in Table 2.

The median concentration for each analyte was lower than, sometimes by a factor of 4, the median concentration reported in women aged 18–40 years participating in the National Health and Nutrition Examination Survey (NHANES) (Table 1). For example, the median BDE47 level in PROTECT was 4.7 ng/g lipid, while the median among women of child-bearing age in NHANES was 22.8 ng/g lipid. With the exception of PFOA, this trend was also evident, or even more pronounced, at the 95th percentile for many analytes. Specifically, the 95th percentile for p,p'-DDE among PROTECT women was 56 ng/g lipid but more than an order of magnitude higher in NHANES at 850 ng/g lipid. However, PBDEs, PCBs, and the pesticides measured in the current study have not been reported in NHANES since 2003–2004, 7–10 years before our samples were collected. On the other hand, NHANES PFAS levels were measured in samples collected during the same time frame as PROTECT samples.

Serum concentrations of analytes within each chemical group were significantly correlated (Table 2). For example, Spearman correlation coefficients for PFASs ranged from 0.44 to 0.88 ($p < 0.05$ for all), and PCB138/158 and PCB153 were strongly correlated ($r = 0.74$, $p < 0.05$). p,p'-DDE was correlated with several analytes from other chemical groups, with significant correlation coefficients ranging from 0.29 to 0.38 for PFASs, 0.59 to 0.72 for PCBs, and 0.41 for BDE47. In addition, there were significant weak correlations between certain PCBs and PFASs: $r = 0.29$ for PFDeA and PCB138/158; $r = 0.33$ for PFHxS

Table 2

Spearman correlations between chemicals (n = 48).

	BDE47	BDE153	PFOA	PFOS	PFHxS	PFNA	PFDeA	SmPFOS	PCB138/158	PCB153	DDE
BDE47	1	0.35*	0.13	0	0.01	0.08	−0.03	0.08	0.14	0.18	0.41*
BDE153		1	−0.09	0.13	0.08	−0.07	−0.12	0.1	0.18	0.2	0.21
PFOA			1	0.44*	0.55**	0.71**	0.63**	0.57**	0.08	0.06	0.15
PFOS				1	0.75**	0.75**	0.6**	0.88**	0.06	0.21	0.22
PFHxS					1	0.69**	0.51**	0.82**	0.19	0.33*	0.38*
PFNA						1	0.85**	0.7**	0.14	0.26	0.29*
PFDeA							1	0.56**	0.29*	0.28	0.23
SmPFOS								1	0.1	0.26	0.36*
PCB138/158									1	0.74*	0.59*
PCB153										1	0.72*
DDE											1

*p < 0.05, **p < 0.001.

and PCB153.

4. Discussion

In this preliminary analysis, we characterized distributions of four classes of persistent organic pollutants among pregnant women in an ongoing birth cohort in Puerto Rico. Although this cohort has elevated exposure to several environmental contaminants, we found that serum concentrations of PCBs, OCPs, PFASs, and PBDEs were lower compared to levels among female NHANES participants aged 18–40 years. However, many of the chemicals discussed here have not been measured in NHANES since 2004 to 2010, and population levels have been shown to be declining (Olsen et al., 2012; Parry et al., 2018; Sjödin et al., 2004; Zota et al., 2013). As a result, we were not able to make direct comparisons to current concentrations among women in the US, with the exception of PFAS. Regardless, our findings suggest that our continuing research efforts should focus on contaminants known to be elevated in this population.

In comparison with other birth cohorts in the US, geometric mean concentrations of BDE47 and p,p'-DDE were higher among women from the Center for Health Assessment of Mothers and Children of Salinas (CHAMACOS) cohort in 2011 (post pregnancy), compared to median concentrations among pregnant PROTECT women, while concentrations of other analytes were similar (Sjödin et al., 2018). Pregnant women participating in the Health Outcomes and Measures of the Environment (HOME) Study in Cincinnati, OH from 2003 to 2006 had higher levels of several PFASs, BDE47, and BDE153 compared to PROTECT women (Braun et al., 2016; Vuong et al., 2015), although PROTECT samples were collected 5–10 years later. These comparisons are consistent with the reported NHANES comparisons and suggest that PROTECT women are not highly exposed to a number of POPs. In addition, the observed correlations of analytes within the same chemical group are as expected based on similar sources of exposure and metabolism within groups, and are consistent with previous reports (Bramwell et al., 2016; Gladen et al., 2003; Yorita Christensen and White, 2011).

Given the numerous hazardous waste sites in northern Puerto Rico, it is unclear why PROTECT participants, and possibly the region's general population, have relatively low serum levels of several POPs, suggesting lower exposure. Interestingly, PCB levels were very low, despite reports of extensive PCB contamination in other regions of Puerto Rico (Alegría et al., 2016; Kumar et al., 2016). This suggests that PCB contamination may be variable around the island, and potentially lower in the areas where PROTECT women live in northern Puerto Rico. Given the tropical climate, it is also surprising that OCP levels were mostly below detection, suggesting that usage of other pesticide classes is likely. Another explanation for low serum PCB and OCP concentrations in this cohort is the relatively young age of PROTECT participants, with a mean age of 27, as many were not yet born when these chemicals were phased out of use in the 1970s. Regarding PFASs and PBDEs, it is

possible that consumer products utilizing these chemicals were not widely distributed in Puerto Rico, despite their apparent ubiquity.

This exploratory study has several limitations, most notably the small sample size. As a result, our ability to detect statistically significant associations was limited, and we may have missed individuals or areas with higher exposure. With the exception of PFASs, the limited volume of serum available for analyses led to relatively high limits of detection for many compounds, which may have affected our findings. In addition, blood samples were collected prior to the 2017 hurricane season, during which Hurricanes Irma and Maria likely spread existing environmental contamination on the Island, potentially leading to higher post-hurricane exposure.

5. Conclusion

Although participants in the ongoing PROTECT birth cohort have higher exposure to many environmental contaminants, this preliminary study suggests that they may have lower exposure to several PCBs, OCPs, PFASs, and PBDEs. Therefore, our continuing work in with this population will likely prioritize and focus on contaminants to which they have demonstrated increased exposure.

Acknowledgements

This work is supported by the National Institute of Environmental Health Sciences grants P42ES017198, P50ES026049, UG3OD023251, and UH3OD023251, and U.S. Environmental Protection Agency (EPA) grant R836155. We would like to thank Dr. Antonia Calafat and Dr. Andreas Sjödin at the Centers for Disease Control and Prevention (CDC) for the measurement of persistent organic pollutants in serum samples. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIEHS, EPA, or CDC.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijheh.2019.02.001>.

References

- US Environmental Protection Agency (EPA), 2018. Basic information on PFAS. Available: <https://www.epa.gov/pfas/basic-information-pfas>, Accessed date: 21 May 2018.
- Aker, A.M., Watkins, D.J., Johns, L.E., Ferguson, K.K., Soldin, O.P., Anzalota Del Toro, L.V., et al., 2016. Phenols and parabens in relation to reproductive and thyroid hormones in pregnant women. *Environ. Res.* 151, 30–37.
- Alegría, H., Martínez-Colon, M., Birgul, A., Brooks, G., Hanson, L., Kurt-Karakus, P., 2016. Historical sediment record and levels of PCBs in sediments and mangroves of Jobos Bay, Puerto Rico. *Sci. Total Environ.* 573, 1003–1009.
- Ashrap, P., Watkins, D.J., Calafat, A.M., Ye, X., Rosario, Z., Brown, P., et al., 2018. Elevated concentrations of urinary triclocarban, phenol and paraben among pregnant women in northern Puerto Rico: predictors and trends. *Environ. Int.* 121, 990–1002.
- Bach, C.C., Bech, B.H., Brix, N., Nohr, E.A., Bonde, J.P., Henriksen, T.B., 2015. Perfluoroalkyl and polyfluoroalkyl substances and human fetal growth: a systematic

- review. *Crit. Rev. Toxicol.* 45, 53–67.
- Banzhaf, S., Filipovic, M., Lewis, J., Sparrenbom, C.J., Barthel, R., 2017. A review of contamination of surface-, ground-, and drinking water in Sweden by perfluoroalkyl and polyfluoroalkyl substances (PFASs). *Ambio* 46, 335–346.
- Birnbaum, L.S., Grandjean, P., 2015. Alternatives to PFASs: perspectives on the science. *Environ. Health Perspect.* 123, A104–A105.
- Bramwell, L., Glinianaia, S.V., Rankin, J., Rose, M., Fernandes, A., Harrad, S., et al., 2016. Associations between human exposure to polybrominated diphenyl ether flame retardants via diet and indoor dust, and internal dose: a systematic review. *Environ. Int.* 92–93, 680–694.
- Braun, J.M., Chen, A., Romano, M.E., Calafat, A.M., Webster, G.M., Yoltan, K., et al., 2016. Prenatal perfluoroalkyl substance exposure and child adiposity at 8 years of age: the HOME study. *Obesity* 24, 231–237.
- Brendel, S., Fetter, E., Staude, C., Vierke, L., Biegel-Engler, A., 2018. Short-chain perfluoroalkyl acids: environmental concerns and a regulatory strategy under REACH. *Environ. Sci. Eur.* 30, 9.
- Cantonwine, D.E., Cordero, J.F., Rivera-Gonzalez, L.O., Anzalota Del Toro, L.V., Ferguson, K.K., Mukherjee, B., et al., 2014. Urinary phthalate metabolite concentrations among pregnant women in northern Puerto Rico: distribution, temporal variability, and predictors. *Environ. Int.* 62, 1–11.
- Cathey, A., Ferguson, K.K., McElrath, T.F., Cantonwine, D.E., Pace, G., Alshawabkeh, A., et al., 2018. Distribution and predictors of urinary polycyclic aromatic hydrocarbon metabolites in two pregnancy cohort studies. *Environ. Pollut.* 232, 556–562.
- Centers for Disease Control and Prevention (CDC), 2017a. Fourth National Report on Human Exposure to Environmental Chemicals. Updated tables, January 2017. Atlanta, GA.
- Centers for Disease Control and Prevention (CDC), 2017b. National Health Interview survey data, Table 4-1. Current asthma prevalence percentages by age. Available: <https://www.cdc.gov/asthma/nhis/2015/table4-1.htm>.
- Cordero, J., 2013. Prevalence of autism in Puerto Rico: results of the 2011 survey. In: *Proceedings of the Interagency Autism Coordinating Committee*. U.S. Department of Health & Human Services January 29, 2013.
- DeWitt, J.C., Blossom, S.J., Schaidler, L.A., 2018. Exposure to per-fluoroalkyl and poly-fluoroalkyl substances leads to immunotoxicity: epidemiological and toxicological evidence. *J. Expo. Sci. Environ. Epidemiol.* <https://doi.org/10.1038/s41370-018-0097-y>.
- US Environmental Protection Agency (EPA), 2017a. Technical fact sheet - polybrominated diphenyl ethers (PBDEs). Available: https://www.epa.gov/sites/production/files/2014-03/documents/ffrofactsheet_contaminant_perchlorate_january2014_final_0.pdf, Accessed date: 21 May 2018.
- US Environmental Protection Agency (EPA), 2017b. Learn about polychlorinated biphenyls. Available: <https://www.epa.gov/pbcs/learn-about-polychlorinated-biphenyls-pbcs>, Accessed date: 11 April 2018.
- US Environmental Protection Agency (EPA), 2017c. Assessing and managing chemicals under TSCA - polybrominated diphenyl ethers (PBDEs). Available: <https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/polybrominated-diphenyl-ethers-pbdes>, Accessed date: 21 May 2018.
- US Environmental Protection Agency (EPA), 2017d. Persistent organic pollutants: a global issue, a global response. Available: <https://www.epa.gov/international-cooperation/persistent-organic-pollutants-global-issue-global-response>, Accessed date: 10 April 2018.
- US Environmental Protection Agency (EPA), 2018. National priorities list (NPL) sites - by state. Available: <https://www.epa.gov/superfund/national-priorities-list-npl-sites-state#PR>, Accessed date: 9 March 2018.
- Essumang, D.K., Eshun, A., Hogarth, J.N., Bentum, J.K., Adjei, J.K., Negishi, J., et al., 2017. Perfluoroalkyl acids (PFAAs) in the Pra and Kakum river basins and associated tap water in Ghana. *Sci. Total Environ.* 579, 729–735.
- Fry, K., Power, M.C., 2017. Persistent organic pollutants and mortality in the United States, NHANES 1999–2011. *Environ. Health* 16, 105.
- Garza, J.R., Perez, E.A., Prelip, M., McCarthy, W.J., Feldman, J.M., Canino, G., et al., 2011. Occurrence and correlates of overweight and obesity among island Puerto Rican youth. *Ethn. Dis.* 21, 163–169.
- Gioda, A., Perez, U., Rosa, Z., Jimenez-Velez, B.D., 2007. Particulate matter (pm10 and pm2.5) from different areas of Puerto Rico. *Fresenius Environ. Bull.* 861–868.
- Gladen, B.C., Doucet, J., Hansen, L.G., 2003. Assessing human polychlorinated biphenyl contamination for epidemiologic studies: lessons from patterns of congener concentrations in Canadians in 1992. *Environ. Health Perspect.* 111, 437–443.
- Grandjean, P., Heilmann, C., Weihe, P., Nielsen, F., Mogensen, U.B., Timmermann, A., et al., 2017. Estimated exposures to perfluorinated compounds in infancy predict attenuated vaccine antibody concentrations at age 5-years. *J. Immunol.* 14, 188–195.
- Guelfo, J.L., Adamson, D.T., 2018. Evaluation of a national data set for insights into sources, composition, and concentrations of per- and polyfluoroalkyl substances (PFASs) in U.S. drinking water. *Environ. Pollut.* 236, 505–513.
- Hu, X.C., Andrews, D.Q., Lindstrom, A.B., Bruton, T.A., Schaidler, L.A., Grandjean, P., et al., 2016. Detection of poly- and perfluoroalkyl substances (PFASs) in U.S. drinking water linked to industrial sites, military fire training areas, and wastewater treatment plants. *Environ. Sci. Technol. Lett.* 3, 344–350.
- Hunter, J.M., Arbona, S.I., 1995. Paradise lost: an introduction to the geography of water pollution in Puerto Rico. *Soc. Sci. Med.* 40, 1331–1355.
- Johns, L.E., Ferguson, K.K., Soldin, O.P., Cantonwine, D.E., Rivera-Gonzalez, L.O., Del Toro, L.V., et al., 2015. Urinary phthalate metabolites in relation to maternal serum thyroid and sex hormone levels during pregnancy: a longitudinal analysis. *Reprod. Biol. Endocrinol.* 13, 4.
- Jones, R., Edenfield, E., Anderson, S., Zhang, Y., Sjodin, A., 2012. Semi-automated extraction and cleanup method for measuring persistent organic pollutants in human serum. *Organohalogen Compd.* 74, 97–98.
- Kato, K., Basden, B.J., Needham, L.L., Calafat, A.M., 2011. Improved selectivity for the analysis of maternal serum and cord serum for polyfluoroalkyl chemicals. *J. Chromatogr. A* 1218, 2133–2137.
- Kumar, N., Ramirez-Ortiz, D., Solo-Gabriele, H.M., Treaster, J.B., Carrasquillo, O., Toborek, M., et al., 2016. Environmental PCBs in guanica bay, Puerto Rico: implications for community health. *Environ. Sci. Pollut. Res. Int.* 23, 2003–2013.
- Martin, J.A., Hamilton, B.E., Osterman, M.J.K., Driscoll, A.K., Drake, P., 2018. Births: Final Data for 2016. National Vital Statistics Reports, Hyattsville, MD.
- Meeker, J.D., Cantonwine, D.E., Rivera-Gonzalez, L.O., Ferguson, K.K., Mukherjee, B., Calafat, A.M., et al., 2013. Distribution, variability, and predictors of urinary concentrations of phenols and parabens among pregnant women in Puerto Rico. *Environ. Sci. Technol.* 47, 3439–3447.
- Mrema, E.J., Rubino, F.M., Brambilla, G., Moretto, A., Tsatsakis, A.M., Colosio, C., 2013. Persistent organochlorinated pesticides and mechanisms of their toxicity. *Toxicology* 307, 74–88.
- Olsen, G.W., Lange, C.C., Ellefson, M.E., Mair, D.C., Church, T.R., Goldberg, C.L., et al., 2012. Temporal trends of perfluoroalkyl concentrations in American Red Cross adult blood donors, 2000–2010. *Environ. Sci. Technol.* 46, 6330–6338.
- Otero-Gonzalez, M., Garcia-Fragoso, L., 2008. Prevalence of overweight and obesity in a group of children between the ages of 2 to 12 years old in Puerto Rico. *Puert. Rico Health Sci. J.* 27, 159–161.
- Padilla, I., Irizarry, C., Steele, K., 2011. Historical contamination of groundwater resources in the north coast karst aquifers of Puerto Rico. *Revista Dimension* 3, 7–12.
- Parry, E., Zota, A.R., Park, J.S., Woodruff, T.J., 2018. Polybrominated diphenyl ethers (PBDEs) and hydroxylated PBDE metabolites (OH-PBDEs): a six-year temporal trend in northern California pregnant women. *Chemosphere* 195, 777–783.
- Rappazzo, K.M., Coffman, E., Hines, E.P., 2017. Exposure to perfluorinated alkyl substances and health outcomes in children: a systematic review of the epidemiologic literature. *Int. J. Environ. Res. Publ. Health* 14.
- Rivera-Soto, W.T., Rodriguez-Figueroa, L., Calderon, G., 2010. Prevalence of childhood obesity in a representative sample of elementary school children in Puerto Rico by socio-demographic characteristics. 2008. *Puert. Rico Health Sci. J.* 29, 357–363.
- Secretariat of the Stockholm Convention Clearing House, 2008. Stockholm convention clearing house. Available: <http://chm.pops.int/>, Accessed date: 10 April 2018.
- Sjodin, A., Jones, R.S., Focant, J.F., Lapeza, C., Wang, R.Y., McGahee 3rd, E.E., et al., 2004. Retrospective time-trend study of polybrominated diphenyl ether and polybrominated and polychlorinated biphenyl levels in human serum from the United States. *Environ. Health Perspect.* 112, 654–658.
- Sjodin, A., Jones, R.S., Gunier, R.B., Wong, L.Y., Holland, N., Eskenazi, B., et al., 2018. Polybrominated diphenyl ethers, polychlorinated biphenyls, and 2,2-bis(4-chlorophenyl)-1,1-dichloroethane in 7- and 9-year-old children and their mothers in the Center for the Health Assessment of Mothers and Children of Salinas cohort. *Environ. Sci. Technol.* 52, 2287–2294.
- Steenland, K., Fletcher, T., Savitz, D.A., 2010. Epidemiologic evidence on the health effects of perfluorooctanoic acid (pfoa). *Environ. Health Perspect.* 118, 1100–1108.
- United Nations Environment Programme (UNEP), 2009. Stockholm convention on persistent organic pollutants (POPs) as amended in 2009. Available: <http://chm.pops.int/Home/tabid/2121/Default.aspx>, Accessed date: 10 April 2018.
- Vuong, A.M., Webster, G.M., Romano, M.E., Braun, J.M., Zoeller, R.T., Hoofnagle, A.N., et al., 2015. Maternal polybrominated diphenyl ether (PBDE) exposure and thyroid hormones in maternal and cord sera: the HOME study, Cincinnati, USA. *Environ. Health Perspect.* 123, 1079–1085.
- Xue, J., Liu, S.V., Zartarian, V.G., Geller, A.M., Schultz, B.D., 2014. Analysis of NHANES measured blood PCBs in the general US population and application of sheds model to identify key exposure factors. *J. Expo. Sci. Environ. Epidemiol.* 24, 615–621.
- Yorita Christensen, K.L., White, P., 2011. A methodological approach to assessing the health impact of environmental chemical mixtures: PCBs and hypertension in the National Health and Nutrition Examination Survey. *Int. J. Environ. Res. Publ. Health* 8, 4220–4237.
- Zota, A.R., Linderholm, L., Park, J.S., Petreas, M., Guo, T., Privalsky, M.L., et al., 2013. Temporal comparison of PBDEs, OH-PBDEs, PCBs, and OH-PCBs in the serum of second trimester pregnant women recruited from San Francisco General Hospital, California. *Environ. Sci. Technol.* 47, 11776–11784.