





ORIGINAL ARTICLE

Biological and environmental exposure monitoring of volatile organic compounds among nail technicians in the Greater Boston area

Diana M. Ceballos^{1,2}  | Jessica Craig¹  | Xianqiang Fu³  | Chunrong Jia³  |
David Chambers⁴ | MyDzung T. Chu² | Alai T. Fernandez⁴ | Victoria Fruh¹ |
Zoe E. Petropoulos¹ | Joseph G. Allen² | Jose Vallarino² | Lydia Thornburg⁴ |
Thomas F. Webster¹

¹Department of Environmental Health, Boston University School of Public Health, Boston, Massachusetts

²Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts

³School of Public Health, University of Memphis, Memphis, Tennessee

⁴Volatile Organic Compounds Laboratory, Tobacco and Volatiles Branch, Centers for Disease Control and Prevention, Atlanta, Georgia

Correspondence

Diana M. Ceballos, Department of Environmental Health, Boston University School of Public Health, 715 Albany Street, Talbot 4th Floor West, Boston, MA 02118. Email: ceballos@bu.edu

Funding information

This work was supported by funds contributed by Harvard NIEHS Research Center for Environmental Health pilot grant (NIH P30ES000002), NIH/NIEHS 2R25ES023635-04, T32ES014562, and T32ES007069; NIOSH T42 OH008416, NIH, Harvard Catalyst NIH UL1 TR001102, Harvard Hoffman Program in Chemicals and Health, and Harvard JPB Environmental Health Fellowship. We also acknowledge pilot funding from Boston University School of Public Health. Dr. Webster was supported in part by NIH/NIEHS R01ES028800, R01ES016099, and USEPA grant 83564201.

Abstract

Nail technicians are exposed to volatile organic compounds (VOCs) from nail products, but no studies have previously measured VOC biomarkers for these workers. This study of 10 nail technicians aimed to identify VOCs in nail salons and explore relationships between air concentrations and biomarkers. Personal and area air samples were collected using thermal desorption tubes during a work shift and analyzed using gas chromatography/mass spectrometry (GC/MS) for 71 VOCs. Whole blood samples were collected pre-shift and post-shift, and analyzed using GC/MS for 43 VOCs. Ventilation rates were determined using continuous CO₂ measurements. Predominant air VOC levels were ethyl methacrylate (median 240 µg/m³), methyl methacrylate (median 205 µg/m³), toluene (median 100 µg/m³), and ethyl acetate (median 639 µg/m³). Blood levels were significantly higher post-shift than pre-shift for toluene (median pre-shift 0.158 µg/L and post-shift 0.360 µg/L) and ethyl acetate (median pre-shift <0.158 µg/L and post-shift 0.510 µg/L); methacrylates were not measured in blood because of their instability. Based on VOCs measured in these seven nail salons, we estimated that emissions from Greater Boston area nail salons may contribute to ambient VOCs. Ventilation rates did not always meet the ASHRAE guideline for nail salons. There is a need for changes in nail product formulation and better ventilation to reduce VOC occupational exposures.

KEYWORDS

biomarker, indoor air, nail salon, toluene, ventilation, volatile organic compound

1 | INTRODUCTION

Nail services is an 8.5 billion-dollar business in the United States, with an estimated workforce of 380 000 nail technicians.¹ Nail

salons are largely owned and staffed by female (94%), Vietnamese (51%), and young workers (43% age <40).¹ Most nail salons are small businesses, which usually provide limited knowledge and resources for health and safety, employing fewer than 10 nail technicians.²

This article has been contributed to by US Government employees and their work is in the public domain in the USA.

Limited English language skills in some nail technicians create additional occupational health and safety barriers.³

Nail technicians are exposed to a wide variety of chemicals during their work shift, while using products for doing manicures, pedicures, nail art, and artificial nails. Products common to the trade include those used for nail sculpting, as well as nail polishes and polish removers, which contain volatile organic compounds (VOCs) such as toluene, acetone, isopropyl acetate, ethyl acetate, and butyl acetate.^{4,5} Although most of these chemicals are listed as ingredients in safety data sheets (SDS or MSDS), some are not, even if present in significant quantities.⁶ Exposure to nail product-related VOCs may occur through inhalation and dermal contact as technicians prepare, apply, or remove these products,⁷ with the highest levels of exposure occurring during acrylic nail sculpting and gel nail polish procedures.^{8,9} There is also potential for dermal absorption of nail product VOCs by direct contact and permeation through gloves, and to a lesser extent from air.¹⁰

Exposure assessment of nail technicians has been limited by a number of occupational constraints requiring control and characterization including numerous product formulations, limited accessibility to worker populations, and work environments that are often small, confined, and have client visibility. Most research has involved assessing ventilation^{11,12} and educating nail technicians.^{13,14} These studies have found that nail salons with limited ventilation often have the highest levels of total volatile organic compound (TVOC, the sum of VOCs detected with a real-time instrument).^{15,16} Nail salons do not always provide appropriate personal protective equipment (PPE), and management and employees have limited knowledge of appropriate health and safety precautions.^{14,15} Only a few studies have assessed worker exposure to VOCs, most finding strong odors and air levels well below occupational exposure limits (OELs) for individual VOCs or mixture of VOCs.^{5,9,16-20} There has been one study measuring nail technicians' urinary VOC biomarkers.²¹ In that study, urinary acetone and hippuric acid were found to be weakly correlated with acetone airborne concentrations (correlation coefficients of 0.49 and 0.45, respectively) and not significantly higher than those measured in a control group.²¹ The authors concluded that these acetone biomarkers were not ideal for assessing exposures of nail technicians.

A few studies have examined possible respiratory, dermal, neurological, and reproductive health outcomes in nail technicians.²²⁻²⁶ Negative health risks identified are the same as those typically associated with VOC exposure.²⁷⁻²⁹ Because most of these epidemiological studies relied on self-reported measures of exposure such as job title or number of nail procedures, negative health outcomes cannot be confidently related to the actual exposures. This ambiguity could be abated through biomonitoring, a direct exposure measurement method that quantifies exposures resulting from multiple exposure routes.³⁰

We performed an in-depth assessment of nail technicians' exposure to VOCs. The objectives of this pilot study were to (a) characterize air concentrations of specific VOCs in salons and the breathing zone of nail technicians; (b) examine the relationship between personal VOC exposure and blood biomarker concentrations;

Practical Implications

- Nail technicians' elevated blood levels of toluene and ethyl acetate suggest occupational exposure to volatile organic compounds from nail products.
- Ventilation was found to be a key variable in the air levels of volatile organic compounds found in the salons.
- Changes in nail product formulation and better ventilation in nail salons would reduce volatile organic compounds occupational exposures.

(c) understand occupational contribution to exposures during the work shift; and (d) test the appropriateness of the exposure assessment methods used in this study for this worker population.

2 | METHODS

2.1 | Study design

We assessed VOC exposure of 10 nail technicians over one workday during the period of November 2016 to June 2017. Nail technicians were recruited from seven nail salons in the Greater Boston area, Massachusetts, United States, with the support of the Massachusetts Healthy Cosmetology Community (HCC). Nail salons included some located in the City of Boston and others in the Boston metropolitan area. Inclusion criteria for participation were (a) female, (b) non-smoker (smoking can be a source of VOCs), (c) >18 years of age, and (d) working full-time (at least 35 hours a week) in a nail salon. Vietnamese language services were provided when needed. Once a participant confirmed interest in participating, we coordinated a first visit to the nail salon with both the participant and the salon owner. Potential participants read and signed a consent form. We scheduled sampling toward the end of a participant's workweek when workload typically increases to capture highest exposure levels and following biomonitoring guidelines.³¹ After collecting pre-shift biological (blood and breath) samples and setting up other sampling equipment, we performed a ventilation check and observed work practices. At the end of the shift, we collected post-shift biological samples and administered a survey. Study protocols were approved by the Institutional Review Boards of the Harvard T.H. Chan School of Public Health and Boston University School of Public Health.

2.2 | Survey instrument

We administered a short questionnaire to all participants to document occupational factors—work practices, workload, and controls used such as local exhaust ventilation (LEV) and PPE—and non-occupational factors—demographics, use of nail polish, and health-related behaviors—that may have contributed to VOC exposure.

2.3 | Nail salon observations and ventilation check

During a walk-through on the day of sampling, we documented processes and controls, including nail procedures performed, nail products used, room ventilation, and PPE used. Temperature, relative humidity, number of salon occupants, and CO₂ level indoors were continuously recorded at 1-minute intervals during the work shift using a TSI Q-track indoor air quality monitor 7575, which was bump calibrated before the sampling day with a 400 ppm CO₂ gas standard. CO₂ level outdoors were also measured for estimating ventilation rates. Any existing general and LEV were checked using a Gastec Smoke Tube Kit (SKC Int.). Smoke generated with smoke tubes was effectively drawn into an LEV system and away from the nail technician's breathing zone when LEVs were operating adequately.

2.4 | Nail products safety data sheet review

We used Safety Data Sheets (SDS) to identify VOC ingredients in nail polish brands and other nail products found in the salons. Nail polish included two main products: standard defined as polish that lasts 1-2 weeks and does not require ultraviolet light to cure, and gel defined as polish that lasts 2-4 weeks and usually requires ultraviolet light to cure. Both nail polish products included a base, color, and top coat. Gel products also used a primer and a nail bonding agent. Other products included artificial nail sculpting ingredients, nail glues, nail cleaners, and nail polish removers. SDSs were obtained from online resources or directly from the nail product manufacturer as these were not found in the nail salons.

2.5 | Personal and area air VOC sampling and analysis

Volatile organic compound air monitoring used passive thermal desorption (TD) tube-type samplers during the work shift, following manufacturer instructions. The TD tubes were standard industrial tubes (89 mm L × 6.4 mm o.d. × 5 mm i.d) packed with 200 mg of Tenax TA (Catalog No. C1-AXXX-5003; Markes International). For personal air sampling, a tube was fixed at the breast pocket of each nail technician using a proprietary pen clip with the sampling end turned up; sampling time averaged 493 minutes with a standard deviation (SD) of 86 minutes capturing their work shift. For area sampling, at least one tube per salon was fixed on a stand with the sampling end facing down in an area used during the day by the participant; sampling time averaged 531 minutes with a SD of 110 minutes. In two of the seven salons, we placed replicate TD tubes side-by-side as part of quality control. These standard TD tubes have validated sampling rates ranging from 0.25 to 0.68 mL/min depending on the compound in occupational settings.³²

After sample collection, tube samples were sealed with storage caps and stored in sealed aluminum containers. Samples were shipped with field blanks to the Environmental Analytical Laboratory at the University of Memphis within 2 days of collection. The samples were analyzed within 5 days upon receipt as previously

described by Jia and Fu.³³ In brief, a tube was thermally desorbed at 260°C on an automated TD system (ULTRA 2+UNITY 2; Markes International), and the desorbed VOCs were then separated and analyzed on a gas chromatography/mass spectrometry (GC/MS) system (Agilent 7890A/5975C; Agilent). Data analysis was performed for 71 target VOCs, selected from EPA TO-15 Method and common compounds previously found in nail salons. Method detection limits (MDLs) were derived from the analytical limit of detection (LOD) of the instrument for 8-hour sampling (Table S1). Samples below the MDL were imputed as $MDL/\sqrt{2}$.³⁴

2.6 | Sampling and analysis of VOC biomarkers

Two gray top vacutainer tubes of whole blood (7 mL) were collected from each participant at the work site, one at the beginning and one immediately at the end of the work shift. Contaminant-free tubes were provided by the Centers for Disease Control and Prevention (CDC) VOC Laboratory. Trained nurses from the Beth Israel Deacons Medical Center performed venipuncture following the universal precautions for working with blood and blood products.³⁵ The tubes of whole blood were sent overnight (5°C) to the CDC VOC Laboratory. Blood was analyzed within 5 days following headspace solid-phase microextraction with benchtop GC-MS CDC Method VOC54, as previously described.^{36,37} MDLs were provided by the laboratory (Table S2). Concentrations below the MDL were imputed as $MDL/\sqrt{2}$.³⁴ Of the 43 blood VOCs tested, eight were detected in blood; of the 71 air VOCs that were screened, 22 were detected in air using TD tubes.

Sampled exhaled breath pre-shift and post-shift using 1-L Tedlar bags were also measured at CDC. VOCs were analyzed using headspace solid-phase microextraction (SPME) GC/MS method for the analytes detected in blood, as previously described by Sampson et al.³⁸ However, semiquantative breath data (not shown) were below lowest calibration standard levels. For predominant blood VOCs detected, toluene and ethyl acetate, the lowest standard concentrations for exhaled breath measurements were 3.65 and 62.4 µg/m³ (1.06 and 17.21 ppb_v), respectively.

2.7 | Data analysis

The workload was defined as the number of nail procedures a technician performed on the day of sampling. Data on workload were asked to technicians and verified by observations during the day. We estimated ventilation rates, or air exchanges per hour (ACHs), using CO₂ measurements and number of occupants in the space at steady state, and using a steady-state box model³⁹⁻⁴²; results were compared to ventilation guidelines for nail salons by the American Society of Heating, Refrigerating and Air-Conditioning Engineering (ASHRAE).⁴⁰ Details are shown in Supporting Information 7.

Air VOC samples were not blank corrected because monitored compounds were non-detectable in field blanks. If there was more than one area air sample per salon during the same visit (ie, replicate samples or two technicians sampled in the same salon), measurements were averaged. Salon 7 was visited twice for different

TABLE 1 Nail salon characteristics (n = 7)

Nail salon ID	Area m ²	Number manicure stations	Number pedicure stations	General exhaust ventilation	Measured CFM per person (m ³ /min-person)	ACH	LEV	Typical services	Days per week open
1	55	3	2	No	7.0 (2.1)	0.32	Yes	Acrylic nails, natural care, hair (once a week without chemical treatments)	7
2	92	7	5	Yes	38 (11)	0.88	No	Acrylic nails, natural care, waxing	6
3	279	12	9	Yes	15 (4.5)	0.42	No	Acrylic nails, natural care (rarely done), waxing	6
4	124	10	8	Yes	ND	ND	No	Natural care, waxing	7
5	45	3	1	Yes	6.4 (1.9)	0.21	Yes	Acrylic nails, natural care (rarely done)	5
6	57	5	4	Yes	88 (26)	3.8	No	Acrylic nails, natural care, waxing	7
7	217	6	6	Yes	27 (8.1)	0.32	No	Acrylic nails, natural care, waxing	6

Note: Data missing due to instrument logging failure. Natural care refers to using standard nail polish lines.

Abbreviations: ACH, air exchanges per hour of ventilation (#/h); CFM, cubic feet per minute of ventilation (ft³/min, equivalent to 0.3 m³/min); LEV, local exhaust ventilation; ND, not determined.

TABLE 2 Demographic and employment characteristics of nail technician participants (n = 10)

Participant characteristics	No. of respondents	Mean (SD)
Demographics		
Current age, y	10	45.0 (10.8)
Country of origin (survey language)		
United States (English)	4	
Vietnam (Vietnamese)	6	
Employment		
Occupation at current salon		
Length, y		8.4 (6.4)
Nail technician	8	
Owner (also nail technician)	2	
Employment in nail industry		
No other job as nail technician	10	
Full-time, y	10	10 (7.7)
Part-time, y	8	10 (12.4)
Work schedule		
Weekly, h		37.9 (6.1)
Day of sampling, h		8.73 (1.4)
Nail procedures on day of sampling, by type		
Manicure, count	8	2.1 (1.3)
Pedicure, count	8	1.5 (0.6)
Types of manicure/pedicure reported, count		
Acrylic (ie, new artificial nails)		6.0 (3.0)
Refill (ie, redo of artificial nails)		1.0 (0.0)
Gel nail polish (ie, UV light needed for curing)		4.0 (0.0)

participants, and the samples were treated as independent samples considering the large temporal variation.

We compared medians and ranges of blood concentrations with those from the 2013-2014 US non-smoking female general population.⁴³ We compared personal air and blood concentrations to the lowest OELs and biological exposure indices (BEIs).^{31,44,45} For correlations between variables, we used Spearman correlation coefficients; for contrasts between paired variables (eg, pre- and post-shift blood VOCs) or to compare post-shift blood and NHANES blood levels, we used Wilcoxon signed-rank tests. Statistical analyses used AIHA IHstats V229, JMP 13.2.0, and SPSS 17.0.2, and statistical significance level, α , was set at 0.05.

3 | RESULTS

3.1 | Characteristics of nail salons and nail technicians

Table 1 describes the seven nail salons we studied. The salons were small (45 m²)-to-medium size (279 m²) and opened 5-7 days per

week. Salons had up to nine manicure and 12 pedicure stations, although not all stations were occupied in some salons on the day of our visit. All but one of the salons performed acrylic nail enhancements. Salons kept temperatures throughout the work shift between 18–25°C and 32%–53% relative humidity. Six salons had general exhaust ventilation while one relied on natural ventilation. The salon estimated general ventilation rates of 6.4–88 cfm/person and did not always meet the ASHRAE guideline for acceptable indoor air quality in nail salons of 25 cfm/person (equivalent to 7.5 m³/min-person).³⁹ Two salons used LEV systems (extraction hoods to filter contaminants before recirculating the air indoors) over the manicure stations, which we verified were in good operating condition.

Table 2 summarizes demographic and employment characteristics of the ten nail technicians in our study. Six participants were Vietnamese, and four were White non-Hispanic. The participants averaged 9 years in the profession. Participants worked an average of 9 hours on the day of our visit and usually worked 39 hours per week. The workload for technicians ranged from three to eight clients on the sampling day. A few nail technicians used limited PPE during their work shift, including gloves (thin latex and nitrile), uniforms, safety glasses, or respirators (disposable N95 with charcoal cartridges). Seven of the ten nail technicians reported washing their hands before eating and before leaving work. Additional information on the use of PPE and personal behaviors (eg, handwashing) is in Tables S3 and S4, but was not further analyzed due to the small sample size.

3.2 | Nail products safety data sheet review

Nail polish was the most common nail product used in salons (typical nail salon carrying 4–5 different nail polish brands, while some carrying as many as 10). Of the 22 brands we reviewed, standard nail polish products reported higher SDS-disclosed percent by weight (% WT) content than gel products for ethyl acetate (<60% WT for standard vs <50% WT for gels) and toluene (<30% WT for standard vs 0% WT for gels). However, gel products had higher percent by weight content of acrylates (<100% WT for gels vs <30% WT for standard). Acrylates listed in nail polish SDSs included hydroxyethyl methacrylate, isobornyl acrylate, tetrahydrofurfuryl methacrylate, and ethyl methacrylate. The nail bonding agent listed 100% ethyl acetate.

Nail products used for sculpting artificial nails, commonly known as acrylic nails or enhancements, included liquid, powder, or pre-mixed powder, composed of <100% acrylates. Acrylates listed in acrylic nail products' SDSs listed different polymerization fragments including ethyl and methyl methacrylate, butyl methacrylate, and ethyl 2-cyanoacrylate. Other nail products used during nail procedures that contained VOCs in their ingredients included nail glue or superglue (<100% acrylates) and nail cleaners (ie, 100% isopropyl alcohol). Further, nail polish removers used in the salons were all 100% acetone. Nail polish removers that do not contain acetone were not found in the salons studied.

3.3 | VOCs in personal and indoor air

Of the 71 monitored VOCs analyzed in air, 22 were detected in the nail salons studied (Table S1). The most abundant and frequently measured VOCs in air were toluene, ethyl acetate, ethyl methacrylate, methyl methacrylate, and d-limonene (Table 3). Other VOCs detected in more than 50% of the samples included benzene, ethylbenzene, all the xylenes, styrene, p-isopropyltoluene (p-cymene), and carbon tetrachloride. Personal VOC air concentrations were significantly higher than area air ($P < 0.001$) for ethyl acetate, m,p-xylenes (not o-xylene), methyl methacrylate, and d-limonene. Conversely, personal VOC air concentrations were significantly lower than area air ($P < 0.001$) for benzene and ethyl methacrylate. All detected personal VOC levels were well below the lowest OELs, although OELs are not available for ethyl acrylate and p-isopropyltoluene (Table 3).

Ventilation (ACH) was significantly correlated with a reduction in toluene personal air levels (Spearman $r = -0.76$, $P = 0.017$). However, ACH was significantly associated with an increase in methyl methacrylate salon air concentrations (Spearman $r = 0.71$, $P = 0.034$); although unexpected, perhaps a higher airflow in the salon increases evaporation of the acrylic monomer when products were opened. Other VOCs in personal air, salon air, and blood were not significantly correlated with ventilation.

Workload (ie, number of procedures during the work shift) was only significantly associated with ethyl acetate area air concentrations (Spearman $r = 0.84$, $P = 0.002$), similarly to that described by Quach et al.¹³

3.4 | Blood VOCs

Of the 43 VOCs analyzed in blood, only two had detection rates above 50% in either pre- or post-shift samples from nail salon participants (Table S2). VOCs detected in air that were not quantified in blood and breath samples were methyl and ethyl methacrylate because they are not stable in blood, and d-limonene and p-isopropyltoluene are not included in the blood VOC method because they have low toxicity.^{46,47} Hazardous VOCs such as toluene and ethyl acetate were detected at >80% in post-shift blood samples. Median concentrations of these two compounds increased significantly from pre-shift to post-shift ($P < 0.02$, Figure 1). Nevertheless, there was variability among the nail technician blood levels. Table 4 and Figure 1 show that post-shift median blood concentrations of toluene (0.36 µg/L, maximum of 0.79 µg/L) and ethyl acetate (0.51 µg/L, maximum of 1 µg/L) significantly exceeded the levels estimated in the general female non-smoking US population for toluene (0.056 µg/L, 95th percentile of 0.225 µg/L) and ethyl acetate (<0.158 µg/L, 95th percentile of 0.158 µg/L), with $P < 0.002$.

Pre-shift blood concentrations and personal air exposures displayed significant associations for toluene (Spearman's $r = 0.68$, $P = 0.042$), but ethyl acetate was not significant (Spearman's $r = 0.27$, $P = 0.48$). Blood post-shift concentrations and personal air exposures displayed positive but insignificant associations: toluene $r = 0.53$, $P = 0.12$; ethyl acetate $r = 0.27$, $P = 0.46$.

TABLE 3 Nail technicians full-shift personal and area air sampling results measured with thermal desorption tubes

Analyte ^a	Nail technicians personal Air concentrations (n = 10)				Nail salon area Air concentrations (n = 8)			ACGIH
	MDL μg/m ³	%> MDL	Median μg/m ³	Range μg/m ³	%> MDL	Median μg/m ³	Range μg/m ³	TLV TWA μg/m ³
BETX								
Benzene	0.16	90	0.46	<MDL-3.4	100	2.4 ^c	1.6-4.7	1600
Toluene	0.26	100	39	4.8-85	100	21	2.2-105	75 300
Ethyl benzene	0.16	80	0.32	<MDL-2.1	100	0.35	0.21-0.91	434 000
o-Xylene	0.23	60	0.33	<MDL-2.6	88	0.50	<MDL-0.74	434 000
m,p-xylene	0.35	90	0.79 ^b	<MDL-7.8	100	0.43	0.52-1.8	434 000
Acrylates								
Ethyl methacrylate	0.28	70	24	<MDL-6900	75	240 ^c	<MDL-5430	NA
Methyl methacrylate	0.37	80	190	<MDL-2100	88	205	<MDL-2580	410 000
Other VOCs documented in nail products								
Ethyl acetate	0.28	100	760 ^b	140-2800	100	639	142-2630	1 440 000
Styrene	0.17	50	0.17	<MDL-0.50	88	0.39	<MDL-0.86	85 141
VOC likely as a fragrance in cleaning products, lotions, makeup, or waxes								
p-isopropyltoluene	0.16	80	1.2	0.12-3.3	90	0.52	<MDL-2.52	NA
d-limonene	0.53	100	9.3 ^b	5.3-180	100	6.4	4.2-130	110 000 ^d
Legacy VOC								
Carbon tetrachloride	0.14	70	0.76	0.10-1.5	90	0.29	0.1-1.0	1078 ^d
Total measured VOCs								
Sum of 71 VOCs		100	2750	677-7910	100	1970	584-6280	

Note: Sum of 71 VOCs = sum of all analytes identified and quantified by the analysis of the thermal desorption tubes.

Abbreviations: ACGIH TLV TWA, threshold limit values, American Conference of Governmental Industrial Hygienists⁴⁶ for an 8-h time-weighted average; BETX, benzene, toluene, ethylbenzene, and the xylenes; MDL, method detection limit.

^aAnalytes listed are those with at least 50% detected samples from Table S1.

^bPersonal air significantly higher than area air.

^cArea air significantly higher than personal air.

^dGerman limit is reported for d-limonene as TLV is not established; OSHA PEL was reported for carbon tetrachloride as is lower than the TLV.⁴⁵

4 | DISCUSSION

4.1 | Personal and area air VOC levels compared to previous studies

This pilot study documents the presence in nail salon air of several VOCs typically found in nail products: toluene, ethyl acetate, and ethyl and methyl methacrylates. Levels were similar to ranges previously documented in nail salons in the United States (Table 5). The levels of benzene, ethylbenzene, and the xylenes in our study were very low (non-detect to low μg/m³) and comparable to ambient levels.²⁸

While personal air concentration exceeded area air concentrations for four of the most frequently detected VOCs, the reverse was true for two VOCs, while there was no significant difference for the other six VOCs. Additional information (eg, technician use of specific products) and a larger data set may be needed to understand such patterns.

4.2 | Biomarkers of nail product-related VOCs

Most of the VOCs detected in the salon air were below detection limits of the blood analysis, but of the VOCs detected in the air and blood samples, toluene and ethyl acetate predominated. Post-shift blood concentrations of these two compounds increased significantly during the shift, suggesting that nail salon work made an important contribution to nail technician's exposures. Median post-shift blood sample concentrations exceeded those found in the female, non-smoking US population.

Half-lives for many stable VOCs such as toluene and ethyl acetate can extend up to 10 hours if the VOC has permeated into less polar tissues such as adipose,⁴⁸⁻⁵¹ making them able to be measured throughout a work shift. However, we found venipuncture to be challenging in this worker population (ie, we failed to draw blood once after many attempts). Our attempt to use less invasive exhaled breath to measure VOCs was not successful at the low exposure levels (all non-detects). Development of a more sensitive breath




FIGURE 1 Toluene and ethyl acetate blood levels measured before and after the work shift (2016-2017) and compared to the US general population (NHANES 2013-2014 for female non-smokers). *Post-shift blood levels significantly higher from pre-shift or NHANES. †All NHANES ethyl acetate and all but two pre-shift ethyl acetate samples were <MDL. Each color in the figure corresponds to the same participant

analysis method with pre-concentration or analysis of intact VOCs in urine may be worth pursuing.

4.3 | Exposure sources and factors

Pre-shift blood levels may partly reflect work on previous days or non-work activities. Toluene was the only VOC measured in pre-shift blood at high detection rates (>89%). Toluene is ubiquitous,⁵² and air salon concentrations were relatively low (median personal air of 39 µg/m³). Further, pre-shift blood samples were collected in the salon approximately 20-45 minutes after the technician arrived before nail services were provided, which may have given time for toluene blood level to equilibrate with the salon environment.⁵³

To control for other VOC sources such as tobacco smoke and personal use of nail salon products, we recruited only non-smokers and asked participants to refrain from painting their nails during the week before the start of the sampling day. Only one participant reported alcohol consumption on the day before sampling, and the laboratory did not report that ethanol interfered in the exhaled breath analysis. Nevertheless, there may be the potential for contributions from other VOC sources outside of work.

Of the four VOCs with the highest median concentrations in personal air—ethyl acetate highest median followed by methyl methacrylate, ethyl methacrylate, and then toluene—ethyl acetate and toluene were also detected at high rates in post-shift blood.

Methacrylates were not examined in blood because of their high instability in blood. Ethyl acetate, toluene, and acrylates were also identified as ingredients of nail polish and artificial nail products using SDS information. This information suggests that nail products are a source of exposure for nail technicians, similarly to that confirmed by Zhong et al.¹⁶

Although d-limonene and p-isopropyltoluene occurred in the air of the nail salons, we did not identify these compounds as ingredients in the SDSs we reviewed. It is possible that they may be classified as a fragrance, or are present in other products used in the salons that we did not review, such as lotions, scrubs, waxes, and cleaning products. Both substances are irritating agents for both eyes and skin, and d-limonene may cause dermatitis.^{46,47} Carbon tetrachloride was also found in most air samples but in very low levels typical of background indoor levels; this chemical is a legacy air contaminant that was previously used in industrial cleaning, dry cleaning, and aerosols.⁵⁴

Besides environmental and occupational sources, other factors may affect VOC exposure of nail technicians such as ventilation, workload, and use of PPE. We estimated ventilation rates to be 0.2-3.8 ACH (6.4-88 cfm per person), comparable to those in seven nail salons in the Los Angeles area⁵⁵ (ACH from 0.2 to 5.7) and 17 nail salons in Michigan¹⁶ (ACH from 0.7 to 45.2) that also used CO₂ measurements in their studies. As reported above, we found an association between ventilation and toluene personal air concentration and workload and area air concentration for ethyl acetate; however,

TABLE 4 VOC concentrations in blood collected from nail technicians before and after the work shift and compared to the US general population

Analyte ^b	Nail technicians (n = 9 ^a)				Nail technicians (n = 10)				US general population					BEI µg/L
									2013-2014 NHANES female non-smokers ⁴³					
	Pre-shift				Post-shift									
	Blood concentrations µg/L				Blood concentrations µg/L				Blood concentrations µg/L					
MDL µg/L	%> MDL	Median	Range	%> MDL	Median	Range	%> MDL	Median	Range	n	MDL	Median	Range	95th percentile
Ethyl acetate	0.16	22	<0.16	<0.16-0.77	80	0.51 ^c	<0.16-1.5	1160	0.16	<0.16	<0.16-7.1	<0.158	NA	
Toluene	0.025	89	0.16	<0.025-0.23	100	0.36 ^c	0.028-0.79	1147	0.025	0.056	<0.025-3.79	0.225	20	

Abbreviations: BEI, Biological Exposure Index ACGIH.⁴⁶; MDL, method detection limit; N, number of samples.

^aOne pre-shift sample was not collected because of challenges drawing blood during phlebotomy.

^bAnalytes listed are those with >50% detect in at either pre- or post-shift samples from Table S2.

^cPost-shift blood levels significantly higher from pre-shift blood levels.

more associations were not possible due to low statistical power or chance. Due to our small sample size, we could not adequately analyze the effect of PPE.

4.4 | Significant contribution to ambient VOCs from nail salons

To understand the potential impact that the measured levels of VOCs in nail salons may have on annual emissions to the outdoor air, we estimated emissions per salon (calculation details described in Supporting Information 5). If we make a series of assumptions: (a) The seven nail salons studied were representative of a typical nail salon in the City of Boston, (b) City of Boston has 200 salons (data from the Boston Public Health Commission), and (c) salons operated all year, emissions from nail salons in the City of Boston (benzene 0.37, toluene 2.97, metric tons/y) would be comparable or higher than the emissions reported for all industrial facilities in Suffolk County in 2016 (benzene 0.58, toluene 1.2, metric tons/y).⁵⁶ In contrast, there were no emissions for these VOCs reported to the toxic release inventory for the City of Boston in 2016.⁵⁶ Nail salon emissions would be higher for the Greater Boston area but an accurate count of nail salons for the area was not available. Although individual salons are not required to report emissions, the nail salon industry collectively may be a non-negligible VOC emission source. Our rough estimates indicate that nail salon outdoor emissions would be important to explore in future studies because in the United States, urban air VOCs are increasingly derived from consumer products rather than industrial sites or transport-related emissions.⁵⁷

4.5 | Nail salon VOCs and nail technician's health

Salon air concentrations were below lowest OELs, and nail technician blood levels were less than BEIs (Tables 3 and 4). These measured levels have not been identified to cause acute and chronic health effects.

Health effects typical of long-term toluene exposures in the ppmV range of a month, some of which are commonly reported by nail technicians, include the following: respiratory and eye irritation; irritated, dry, and cracked skin; nervous system dysfunction; headaches; dizziness; fatigue; and developmental impairment.^{45,58,59} While many nail polish brands are eliminating toluene,⁶⁰ it may continue to be found as a residue in nail products.

Ethyl acetate is a major ingredient in many nail polishes. Chronic ethyl acetate exposure in the ppmV range on the order of a month has been associated with eye, skin, nose, and throat irritation, as well as nervous system depression, edema of the liver and kidney, corneal clouding, and even anemia.^{61,62} There have been no studies of developmental toxicity to the unborn child,⁶³ which would be important for the high percentage of nail technicians in reproductive age.

Likely sources of methyl and ethyl methacrylate are nail polishes (especially gel products) and acrylic liquid or powder used for sculpting nails. Methyl methacrylate has been banned in 32 US States, but not Massachusetts.⁶⁴ The Food and Drug Administration (FDA) has

TABLE 5 Range of VOC concentrations detected in nail salons in this study compared to other studies

Country	Reference	Personal or area air (n)	# VOCs	Sampling method	VOC air concentrations range ^a				
					Toluene μg/m ³	Ethyl methacrylate μg/m ³	Methyl methacrylate μg/m ³	Ethyl acetate μg/m ³	d-limonene μg/m ³
US	This study	Personal (10)	72	Passive	4.8-85	ND-6900	ND-2100	140-2800	5.3-180
		Area (8)	72		2.2-105	ND-5430	ND-2580	142-2630	4.2-130
	Zhong 2019 ¹⁶	Personal (34)	12	Passive	ND-650	ND-1920	ND-36 000	170-9650	ND-300
		Area (34)			ND-380	ND-1350	ND-34 200	84-6900	ND-200
	Quach 2018 ¹³	Personal	3	Passive	ND-325 ^b	-	ND-41 000 ^b	-	-
		Personal	3	Passive	188-640	-	82-27 880	-	-
	Alaves 2013 ¹⁹	Area (12)	7	Active	38-1168	-	ND-16 810	180-7200	-
		Personal (167)	3	Passive	75-3766	-	-	72-19 800	-
	Marty 2007 ⁷⁷	Area (3)	6	Active	38-226	-	492-5330	72-540	-
		Personal	2	Active	-	187-145 000	615-21 320	-	-
	Spencer 1997 ¹¹	Personal, LEV (8)	2	Active	-	1870-7013	ND	-	-
		Personal, no LEV (10)	3		-	32 725-65 450	ND	-	-
	Almaguer 1998 ¹⁷	Personal (1)	4	Active	-	-	ND	-	-
		Area (6)	4		-	ND-32 725	ND	-	-
	Almaguer 1992 ⁷⁸	Personal (2)	2	Active	-	11 000-12 000	-	-	-
		Personal (17)	4	Active	377-7532	ND-79 475	-	-	-
	Froines 1986 ⁸	Area (≤25)	3	Active and passive	-	11 220-43 010	8610-27 880	-	-
		Area	70	Active	75-1168	421-14 960	82-328	36-4284	-
Korea	Park 2014 ⁸⁰	Area		Passive	ND-1 849 000	ND-285 000	ND-2 500 000	720-608 000	-
	Yang 2010 ²¹	Area (≤25)	13	Active	490-5300	-	-	756-9000	-
Greece	Tsigonia 2010 ¹⁸	Area	3	Active	-	-	41-492	-	-
		Area	3	Active	3.0-67	-	-	ND-44	ND-227

Abbreviations: n, number of samples when available in the publication; ND, not detected; VOCs, volatile organic compounds.

^aRange was chosen as the best comparison metric because of the different central tendencies reported across studies.

^bApproximate range deduced from statistics given in the publication.

recommended since the late 1970s that methyl methacrylate not be used in pure liquid form because of adverse skin reactions (sensitizing),^{65,66} but it is not listed as a prohibited or restricted substance.⁶⁶ Methyl and ethyl acrylates may occur as residues in products that only disclose the presence of acrylated oligomer or copolymers. The wide variety of acrylate formulations in nail products makes measuring and controlling exposure challenging. We found methyl methacrylate in salon air and listed as an ingredient in some nail products (ie, acrylic powder SDS). The lowest air levels measured for both methacrylates were in salons that did not provide artificial nail procedures.

Potential health concerns for ethyl and methyl methacrylate include irritation of the nose and throat, coughing, shortness of breath, nervous system effects, drowsiness, light-headedness, dizzy spells, trembling of hands (suggested neurotoxicity), and skin sensitization or dermatitis.^{45,67,68} Of particular concern was the listing of polyurethanes as an ingredient in one of the nail polish brands found in the salons studied. Polyurethanes have been well documented as potent sensitizers in the autobody industry.⁶⁹ This was particularly concerning, as the nail polish containing polyurethanes was labeled as non-allergenic.

4.6 | Strengths and limitations

Major strengths of our study include the measurement of the occupational contribution to VOC blood levels for nail technicians and the use of TD tubes. The latter proved to be an ideal sampling technique because they are easy to use, small, inconspicuous, had a 100% participation rate (ie, all technicians in the study agreed to wear TD tubes as part of the study) in the client-oriented salon environment, and follow well-defined sampling and analysis guidelines.^{32,70-72} TD tubes may serve for future studies to collect breath samples as it has been previously described.⁷³⁻⁷⁵ We characterized exposure to more VOCs than previous studies at nail salons. Nevertheless, we were not able to measure all VOCs typically found in nail salons, as our methods did not include acetone, formaldehyde, ethanol, butyl acetate, isopropyl alcohol, and other acrylates beyond ethyl and methyl methacrylates. An important limitation of our study was the small sample size, preventing conclusions regarding many exposure factors such as PPE. Further, data from only seven nail salons may not be representative of all nail salons in the Greater Boston area. However, the demographics of nail technicians we were able to recruit (60% Vietnamese and 40% Caucasian) were similar to the national statistics in the nail salon industry (56% Vietnamese and 36% Caucasian).¹

Since October 2018, nail salons in the City of Boston need to mechanically exhaust air to the outside—instead of relying on air being passively exhausted through windows, doors, and cracks in the building like the salons in our study.⁷⁶ Even if the ventilation changes result in improved air quality indoors, nail salons would need to effectively filter exhausted air from VOCs to reduce current emissions to the ambient air. Thus, this study will serve as the basis for designing future studies to assess VOC exposure of nail technicians, the

impact of safer alternatives in nail products, better controls including improved ventilation, and improved training and outreach to this worker population.

5 | CONCLUSIONS

Nail technicians are exposed to a variety of VOCs throughout their work life. Higher concentrations of some VOCs in the post-shift blood of nail technicians compared to those measured pre-shift or in the general US population indicate the importance of occupational exposure. Ethyl acetate and toluene found in air and blood were also listed as ingredients in nail products in the investigated salons. TD tubes were effective and easy to use for measuring a wide variety of VOC concentrations in indoor and personal air in nail salons. While blood was an effective VOC biomarker for investigating toluene and ethyl acetate exposures, venipuncture was challenging in this work environment, given that a whole tube of blood is needed for current analytical methods. While full-shift VOC air concentrations were well below established OELs (not available for ethyl acrylate and p-isopropyltoluene), they could still contribute to chronic health effects such as respiratory and skin irritation, headaches, dizziness, dermatitis, and neurocognitive and reproductive toxicity. Reducing VOC exposure among nail technicians may require change in nail product formulation, better ventilation, and more consistent and effective use of personal protective equipment.

ACKNOWLEDGEMENTS

The authors are extremely grateful to the participating nail salons and nail technicians without whom this work would not have been possible. The authors express sincere acknowledgements to the Massachusetts Healthy Cosmetology Committee for their support and technical guidance during the protocol preparation and recruitment. Special thanks to Nan Pham and Lori Ducharme for their industry expertise and dedicated assistance during recruiting. Thanks to Nan Pham, Tammy Nguyen, Vy Nguyen, Anna Young, Audrey Nathanson, Wendy Hori, and Zhao Tong for their assistance in the study. Thanks to the mentoring from John Spengler, Robert Herrick, Russ Hauser, David Christiani, Roberta White, and Petros Koutrakis.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

ORCID

Diana M. Ceballos  <https://orcid.org/0000-0002-8789-2528>

Jessica Craig  <https://orcid.org/0000-0002-9565-5557>

Xianqiang Fu  <https://orcid.org/0000-0001-8789-4737>

Chunrong Jia  <https://orcid.org/0000-0001-6541-6478>

REFERENCES

1. NAILS. 2017–2018 NAILS big book statistics. In: *Nails Magazine*. Vol 2018: Michelle Mullen; 2018.
2. Lentz TJ, Sieber WK, Jones JH, Piacitelli GM, Catlett LR. Surveillance of safety and health programs and needs in small U.S. businesses. *Appl Occup Environ Hyg*. 2001;16(11):1016–1021.
3. Cunningham TR, Guerin RJ, Keller BM, Flynn MA, Salgado C, Hudson D. Differences in safety training among smaller and larger construction firms with non-native workers: Evidence of overlapping vulnerabilities. *Saf Sci*. 2018;103:62–69.
4. Gjolstad M, Thorud S, Molander P. Occupational exposure to airborne solvents during nail sculpturing. *J Environ Monit*. 2006;8(5):537–542.
5. Quach T, Gunier R, Tran A, et al. Characterizing workplace exposures in Vietnamese women working in California nail salons. *Am J Public Health*. 2011;101(Suppl 1):S271–S276.
6. Kopelovich L, Perez AL, Jacobs N, Mendelsohn E, Keenan JJ. Screening-level human health risk assessment of toluene and dibutyl phthalate in nail lacquers. *Food Chem Toxicol*. 2015;81:46–53.
7. Shakibaei N. Reducing workers' exposures to chemicals and dust in nail salons using local exhaust ventilation systems: environmental and occupational health sciences, University of Washington; 2014.
8. Froines JR, Garabrant DH. Quantitative evaluation of manicurists exposure to methyl, ethyl and isobutyl methacrylate during production of synthetic fingernails. *Appl Ind Hyg*. 1986;1(2):70–74.
9. Quach T, Von Behren J, Nelson DO, et al. Evaluating an owner-to-worker training intervention in California nail salons using personal air monitoring. *Am J Ind Med*. 2018;61(10):831–841.
10. Chao KP, Lee PH, Wu MJ. Organic solvents permeation through protective nitrile gloves. *J Hazard Mater*. 2003;99(2):191–201.
11. Spencer AB, Estill CF, McCammon JB, Mickelsen RL, Johnston OE. Control of ethyl methacrylate exposures during the application of artificial fingernails. *Am Ind Hyg Assoc J*. 1997;58(3):214–218.
12. Local Hazardous Waste Management Program (LHWMP). Nail salon ventilation 2018; 2018. <http://www.hazwastehelp.org/health/nail-salons-Ventilation.aspx>. Accessed October 24, 2018.
13. Quach T, Von Behren J, Tsoh J, et al. Improving the knowledge and behavior of workplace chemical exposures in Vietnamese-American nail salon workers: a randomized controlled trial. *Int Arch Occup Environ Health*. 2018;91(8):1041–1050.
14. Goldin LJ, Ansher L, Berlin A, et al. Indoor air quality survey of nail salons in Boston. *J Immigr Minor Health*. 2014;16(3):508–514.
15. Pavilonis B, Roelofs C, Blair C. Assessing indoor air quality in New York City nail salons. *J Occup Environ Hyg*. 2018;15(5):422–429.
16. Zhong L, Batterman S, Milando CW. VOC sources and exposures in nail salons: a pilot study in Michigan, USA. *Int Arch Occup Environ Health*. 2019;92(1):141–153.
17. Almaguer D, Malkin R. *The Grand Experience Salon*. Chicago, IL: National Institute for Occupational Safety and Health; 1998.
18. Tsigonia A, Lagoudi A, Chandrinou S, Linos A, Evlogias N, Alexopoulos EC. Indoor air in beauty salons and occupational health exposure of cosmetologists to chemical substances. *Int J Environ Res Public Health*. 2010;7(1):314–324.
19. Alaves VM, Sleeth DK, Thiese MS, Larson RR. Characterization of indoor air contaminants in a randomly selected set of commercial nail salons in Salt Lake County, Utah, USA. *Int J Environ Health Res*. 2013;23(5):419–433.
20. Garcia E, Sharma S, Pierce M, et al. Evaluating a county-based healthy nail salon recognition program. *Am J Ind Med*. 2015;58(2):193–202.
21. Yang JH, Kim JY, Eom A, Hyoung HK, Koh S, Sciences EH. Relationships between airborne exposure and urinary metabolites of nail technicians. *Toxicol Environ Health Sci*. 2010;2(3):175–181.
22. Roelofs C, Azaroff LS, Holcroft C, Nguyen H, Doan T. Results from a community-based occupational health survey of Vietnamese-American nail salon workers. *J Immigr Minor Health*. 2008;10(4):353–361.
23. Quach T, Von Behren J, Goldberg D, Layefsky M, Reynolds P. Adverse birth outcomes and maternal complications in licensed cosmetologists and manicurists in California. *Int Arch Occup Environ Health*. 2015;88(7):823–833.
24. Harris-Roberts J, Bowen J, Sumner J, et al. Work-related symptoms in nail salon technicians. *Occup Med (Lond)*. 2011;61(5):335–340.
25. Peretz J, Gallicchio L, Miller S, Greene T, Zacur H, Flaws JA. Infertility among cosmetologists. *Reprod Toxicol*. 2009;28(3):359–364.
26. Gallicchio L, Miller SR, Greene T, Zacur H, Flaws JA. Adverse health outcomes among cosmetologists and noncosmetologists in the Reproductive Outcomes of Salon Employees (ROSE) study. *J Toxicol Environ Health A*. 2011;74(1):52–61.
27. Jones AP. Indoor air quality and health. *Atmos Environ*. 1999;33(28):4535–4564.
28. Bolden AL, Kwiatkowski CF, Colborn T. New look at BTEX: are ambient levels a problem? *Environ Sci Technol*. 2015;49(9):5261–5276.
29. Woodall GM, Smith RL. The air toxics health effects database (ATHED). *Toxicol Appl Pharmacol*. 2008;233(1):20–24.
30. McIntosh DL, Spengler JD. *Environmental Health Criteria* 214. Geneva, Switzerland: World Health Organization; 2000.
31. ACGIH. American Conference of Governmental Industrial Hygienist 2018 TLVs® and BEIs®. 2018.
32. ASTM International. *Standard practice for choosing sorbents, sampling parameters and thermal desorption analytical conditions for monitoring volatile organic chemicals in air*. West Conshohocken, PA: ASTM International
33. Jia C, Fu X. Diffusive uptake rates of volatile organic compounds on standard ATD tubes for environmental and workplace applications. *Environments*. 2017;4(4):87.
34. Hornung RW, Reed LD. Estimation of average concentration in the presence of nondetectable values. *Appl Occup Environ Hyg*. 1990;5(1):46–51.
35. OSHA. OSHA factsheet: OSHA's bloodborne pathogens standard.
36. Blount B, Kobelski R, McElprang D, et al. Quantification of 31 volatile organic compounds in whole blood using solid-phase microextraction and gas chromatography-mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2006;832(2):292–301.
37. Chambers DM, Blount BC, McElprang DO, Waterhouse MG, Morrow JC. Picogram measurement of volatile n-alkanes (n-hexane through n-dodecane) in blood using solid-phase microextraction to assess nonoccupational petroleum-based fuel exposure. *Anal Chem*. 2008;80(12):4666–4674.
38. Sampson MM, Chambers DM, Pazo DY, Moliere F, Blount BC, Watson CH. Simultaneous analysis of 22 volatile organic compounds in cigarette smoke using gas sampling bags for high-throughput solid-phase microextraction. *Anal Chem*. 2014;86(14):7088–7095.
39. Tucker W. *ASHRAE Standard 62: Ventilation for Acceptable Indoor Air Quality*. In: Agency EP, ed. Washington, DC: USEPA; 1992.
40. ASHRAE. Aircraft, design conditions, ventilation. 2011 (ASHRAE Handbook, HVAC applications).
41. Siconolfi SMCLSF. Carbon dioxide and water vapor production at rest and during exercise: a report on data collection for the crew and thermal systems division, NASA Technical Paper no 3500. Vol 1. Washington, DC: United States. National Aeronautics and Space Administration; 1994.
42. Persily AK. *Evaluating building IAQ and ventilation with indoor carbon dioxide*. Atlanta, GA: American Society of Heating, Refrigerating and Air-Conditioning Engineers, Inc.; 1997.
43. CDC. Fourth report on human exposure to environmental chemicals, updated tables, (March 2018). Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; October 24, 2018.

44. OSHA. Annotated OSHA Z-2 table. 2018; <https://www.osha.gov/dsg/annotated-pels/tablez-2.html>. Accessed October 24, 2018.
45. NIOSH. *NIOSH Pocket Guide to Chemical Hazards*. Ann Arbor, MI: Centers for Disease Control and Prevention; 2007.
46. NIH. Toxic summary for D-limonene. <https://toxnet.nlm.nih.gov/cgi-bin/sis/search/a?dbs+hsdb:@term+@DOCNO+1809>. Accessed October 24, 2018.
47. NIH. Substance p-cymene. <https://chem.nlm.nih.gov/chemidplus/rn/99-87-6>. Accessed October 24, 2018.
48. USHHS. Toxicological profile for toluene.
49. Nise G, Attewell R, Skerfving S, Orbaek P. Elimination of toluene from venous blood and adipose tissue after occupational exposure. *Br J Ind Med*. 1989;46(6):407-411.
50. Pellizzari ED, Wallace LA, Gordon SM. Elimination kinetics of volatile organics in humans using breath measurements. *J Expo Anal Environ Epidemiol*. 1992;2(3):341-355.
51. Brugnone F, Perbellini L, Apostoli P, Locatelli M, Mariotto P. Decline of blood and alveolar toluene concentration following two accidental human poisonings. *Int Arch Occup Environ Health*. 1983;53(2):157-165.
52. McDonald BC, de Gouw JA, Gilman JB, et al. Volatile chemical products emerging as largest petrochemical source of urban organic emissions. *Science*. 2018;359(6377):760-764.
53. Ashley DL, Prah JD. Time dependence of blood concentrations during and after exposure to a mixture of volatile organic compounds. *Arch Environ Health*. 1997;52(1):26-33.
54. ATSDR. Carbon tetrachloride poisoning. https://www.atsdr.cdc.gov/cssem/carbon_tetrachloride/docs/Carbon_Tetra_Patient_Ed_Sheet-508.pdf. Accessed December 19, 2018.
55. Nguyen C. *Indoor Air Quality of Nail Salons in the Greater Los Angeles Area: Assessment of Chemical and Particulate Matter Exposures and Ventilation*. Los Angeles, CA: University of California, Los Angeles; 2016: Open Access via eScholarship <http://escholarship.org/uc/item/0rv805bd>
56. USEPA. Toxics release inventory (TRI) program. 2018. <https://www.epa.gov/toxics-release-inventory-tri-program>. Accessed October 24, 2018.
57. Lewis AC. The changing face of urban air pollution. *Science*. 2018;359(6377):744-745.
58. NIH. Toxic summary for toluene. <https://toxnet.nlm.nih.gov/cgi-bin/sis/search/a?dbs+hsdb:@term+@DOCNO+131>. Accessed October 24, 2018.
59. Grandjean P, Landrigan PJ. Neurobehavioural effects of developmental toxicity. *Lancet Neurol*. 2014;13(3):330-338.
60. Young AS, Allen JG, Kim U-J, et al. Phthalate and organophosphate plasticizers in nail polish: evaluation of labels and ingredients. *Environ Sci Technol*. 2018;52(21):12841-12850.
61. Lewis R. *Sax's Dangerous Properties of Industrial Materials*. Hoboken, NJ: Wiley-Interscience, Wiley & Sons, Inc.; 2004.
62. NIH. Toxic summary for ethyl acetate. <https://toxnet.nlm.nih.gov/cgi-bin/sis/search2/f?./temp/~cGq4fg:3>. Accessed October 24, 2018.
63. Grandjean P, Landrigan PJ. Developmental neurotoxicity of industrial chemicals. *Lancet*. 2006;368(9553):2167-2178.
64. Safe Cosmetics. Acrylates. <http://www.safecosmetics.org/get-the-facts/chemicals-of-concern/2978/>. Accessed October 24, 2018.
65. USFDA. Nail care products. <https://www.fda.gov/Cosmetics/ProductsIngredients/Products/ucm127068.htm>. Accessed October 24, 2018.
66. USFDA. Prohibited & restricted ingredients. <https://www.fda.gov/cosmetics/guidanceregulation/lawsregulations/ucm127406.htm>. Accessed October 24, 2018.
67. Romita P, Foti C, Masciopinto L, et al. Allergic contact dermatitis to acrylates. *J Biol Regul Homeost Agents*. 2017;31(2):529-534.
68. Gatica-Ortega ME, Pastor-Nieto MA, Mercader-Garcia P, Silvestre-Salvador JF. Allergic contact dermatitis caused by (meth)acrylates in long-lasting nail polish - are we facing a new epidemic in the beauty industry? *Contact Dermatitis*. 2017;77(6):360-366.
69. Reeb-Whitaker C, Anderson NJ, Bonauto DK. Prevention guidance for isocyanate-induced asthma using occupational surveillance data. *J Occup Environ Hyg*. 2013;10(11):597-608.
70. ISO. *Indoor, Ambient and Workplace Air – Sampling and Analysis of Volatile Organic Compounds by Sorbent Tube/Thermal Desorption/Capillary Gas Chromatography – Part 2: Diffusive Sampling*. Geneva, Switzerland: ISO 16017-2:2003; 2003:35.
71. USEPA. *Compendium Method TO-17 Determination of Volatile Organic Compounds in Ambient Air Using Active Sampling Onto Sorbent Tubes*. Cincinnati, OH: U.S. Environmental Protection Agency; 1999.
72. USEPA. EPA Method 325b, Volatile Organic Compounds from Fugitive and Area Sources: Sampler Preparation and Analysis. Washington, DC: U.S. Environmental Protection Agency; 2015.
73. Reynolds JC. Volatile organic compound mixtures in human breath: Monitoring and data interpretation. Workshop on New Directions and Advances in Biological and Chemical Exposure Assessment for Epidemiologic and Risk Characterization; 2008; University of California, Los Angeles.
74. Batterman S, Godwin C, Jia CR. Design and evaluation of a new breath monitoring system for volatile organic compounds. Paper presented at: International Council of Chemical Association (ICCA) Biomonitoring Workshop2006; Minneapolis, MN, USA.
75. Markers International. Application note 013. The Bio-VOC – A low-cost, simple device for biological monitoring of VOCs in breath. 2015.
76. Boston Public Health Commission. Ventilation requirements: nail salons ventilation requirements. <http://www.bphc.org/whatwedo/healthy-homes-environment/safe-shops/nail-salons/Pages/Ventilation-Requirements.aspx>. Accessed January 5, 2019.
77. Marty A. *Ethyl Methacrylate and Methyl Methacrylate Exposure among Fingernail Sculptors*. Tampa, FL: University of South Florida; 2007: Open Access via eScholarship <https://scholarcommons.usf.edu/etd/2278/>.
78. Almaguer D, Malkin R. *Haute Nails*. Norman, OK: National Institute for Occupational Safety and Health; 1992.
79. Hiipakka D, Samimi B. Exposure of acrylic fingernail sculptors to organic vapors and methacrylate dusts. *A Ind Hyg Assoc J*. 1987;48(3):230-237.
80. Park SA, Gwak S, Choi S. Assessment of occupational symptoms and chemical exposures for nail salon technicians in Daegu City. Korea. *J Prev Med Public Health*. 2014;47(3):169-176.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Ceballos DM, Craig J, Fu X, et al. Biological and environmental exposure monitoring of volatile organic compounds among nail technicians in the Greater Boston area. *Indoor Air*. 2019;29:539–550. <https://doi.org/10.1111/ina.12564>