



Contents lists available at ScienceDirect

Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv

Short Communication


Urinary phthalate metabolite concentrations in relation to levels of circulating matrix metalloproteinases in pregnant women

Leah D. Bedrosian^a, Kelly K. Ferguson^b, David E. Cantonwine^c, Thomas F. McElrath^c, John D. Meeker^{a,*}^a Department of Environmental Health Sciences, University of Michigan School of Public Health, Ann Arbor, MI, USA^b Epidemiology Branch, National Institute of Environmental Health Sciences, Research Triangle Park, North Carolina, USA^c Division of Maternal Fetal Medicine, Department of Obstetrics and Gynecology, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

HIGHLIGHTS

- Exposure to phthalates is common, including among pregnant women, and may be associated with adverse pregnancy outcomes.
- MMPs are enzymes that participate in extracellular matrix degradation that may play a role in adverse pregnancy outcomes.
- Experimental studies suggest phthalate exposure may upregulate MMP, but human data are lacking.
- In 134 pregnant women we found that urinary phthalate metabolites were associated with higher circulating MMP levels.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 10 August 2017

Received in revised form 22 September 2017

Accepted 22 September 2017

Available online 28 September 2017

Editor: Jay Gan

Keywords:

Biomarker

Enzyme

Environment

Epidemiology

Exposure

Pregnancy

ABSTRACT

Phthalate exposure has been shown to be associated with adverse pregnancy outcomes. However, human studies informing relevant mechanistic pathways are lacking. Experimental studies have reported that matrix metalloproteinases (MMPs), which are responsible for extracellular protein degradation, may be upregulated in response to phthalate exposure. In this exploratory study we measured urinary phthalate metabolite concentrations, plasma MMP levels, and relevant covariates among 134 pregnant women. There were statistically significant or suggestive positive relationships between several phthalates, particularly between metabolites of di-(2-ethylhexyl) phthalate, with MMP-1 and MMP-9 levels. Further research is needed to confirm these results and how they may inform the mechanisms involved between phthalate exposure and adverse pregnancy outcomes.

© 2017 Elsevier B.V. All rights reserved.

* Corresponding author at: Department of Environmental Health Sciences, University of Michigan School of Public Health, 1835 SPH I, 1415 Washington Heights, Ann Arbor, MI 48109, United States.

E-mail address: meekerj@umich.edu (J.D. Meeker).

1. Introduction

Phthalates are a family of chemicals that are widely present in the environment due to their use in industry as well as in personal care and consumer products. High molecular weight phthalates such as di-(2-ethylhexyl) phthalate (DEHP) are widely used as plasticizers with exposure occurring through food packaging, bottled water, children's

Table 1
Distribution of urinary phthalate metabolite and plasma MMP concentrations (N = 134).

Biomarker	Geometric mean	Percentiles				
		25th	50th	75th	95th	Maximum
<i>Urine (ng/ml)</i>						
MEHP	7.52	3.05	7.12	15.1	83.4	888
MEHHP	21.6	6.91	19.8	42.9	390	2100
MEOHP	12.9	4.25	12.1	25.9	178	1200
MECPP	32.6	10.8	29.6	72.4	439	4290
ΣDEHP	26.8	10.6	24.9	56.6	387	2140
MBZP	4.75	1.93	4.36	12.4	42.7	255
MBP	12.2	6.10	13.6	31.4	66.6	217
MIBP	5.60	2.31	6.62	12.3	29.8	158
MEP	105	31.4	80.2	308	2180	5500
MCPP	1.65	0.642	1.53	3.50	14.2	116
<i>Plasma (ng/ml)</i>						
MMP-1	0.59	0.21	0.59	0.98	2.62	7.52
MMP-2	50.7	36.0	54.8	75.8	123	182
MMP-9	51.0	36.4	51.2	78.5	122	211

toys, medical devices, and other products (Marie et al., 2015). Low molecular weight phthalates such as diethyl phthalate (DEP) or dibutyl phthalate (DBP) are commonly found in lotions, perfumes and deodorants, posing a primary concern for women (Marie et al., 2015).

Matrix metalloproteinases (MMPs) are enzymes that participate in extracellular matrix degradation. MMPs and their inhibitors rely on a delicate balance whose disruption can result in a range of adverse pregnancy outcomes and other health effects (Geng et al., 2016). Several animal and in vitro studies have shown that phthalates induce increased MMPs in a range of cell types. Specifically, DEHP (and its metabolite MEHP) upregulated both MMP-2 and MMP-9, while DBP upregulated MMP-9 (Kim et al., 2015; Scarano et al., 2009; Yao et al., 2012; Zhang et al., 2016; Zhu et al., 2010). Increased circulating levels of MMPs have been reported in human studies of preeclampsia (Eleuterio et al., 2015), gestational hypertension (Ab Hamid et al., 2012), pregnancy loss (Nissi et al., 2013), and preterm birth (Kramer et al., 2010). Increased exposure to phthalates have also been associated with these same adverse health measures in recent epidemiologic studies (Cantonwine et al., 2016; Ferguson et al., 2014; Messerlian et al., 2016; Werner et al., 2015). However, to our knowledge no human studies to date have assessed the relationship between phthalate exposure and circulating MMP levels. The present exploratory study aimed to investigate these relationships to determine the potential for MMPs to help inform mechanism(s) of action between phthalates and adverse pregnancy outcomes.

2. Methods

Women who planned to deliver at Brigham and Women's Hospital in Boston between the years 2006 to 2008 were recruited into a

prospective cohort study as described previously (Ferguson et al., 2014). Written informed consent was obtained from the participants and the institutional review boards of Brigham and Women's Hospital and the University of Michigan approved this study. During the study visit, the women filled out demographic questionnaires and supplied urine and blood samples for biomarker analysis. During their following three study visits, further biological samples were collected. Specimens were stored at -80 degrees Celsius until analysis. For this exploratory study, samples of urine and plasma from 134 women collected at the third study visit (median = 26 weeks gestation; range = 23–29 weeks) were analyzed for phthalate metabolites and MMPs, respectively.

Nine phthalate metabolites were measured in each urine sample by NSF International using a modified version of the LC-MS/MS method developed at the Centers for Disease Control and Prevention (Lewis et al., 2013). To account for urine dilution, specific gravity (SG) was measured using a handheld refractometer.

Maternal plasma samples were analyzed for matrix metalloproteinases by the University of Michigan Cancer Center Immunology Core. Five MMPs (1, 2, 7, 9 and 10) were measured using EMD Millipore's MILLIPLEX Multiplex assay kit using the Luminex xMAP platform. Only MMP-1, 2 and 9 were included further in our data analysis since MMP-7 and MMP-10 were not detected in any of the samples. Coefficients of variation for replicate standards at varying concentrations were below 8% for MMP-1, 11% for MMP-2 and 7% for MMP-9.

All statistical analysis was performed using R version 3.3.2. Phthalate metabolite and MMP concentrations were all right-skewed and log-transformed for statistical analysis. All phthalate metabolites were analyzed separately, in addition to a summed value (ΣDEHP) for multiple metabolites of DEHP (MEHP, MEHHP, MEOHP, MECPP) based on molecular weight. Demographic information was collected for covariates of interest, including maternal age, race/ethnicity, education level, health insurance provider and body mass index. Differences in MMP or phthalate levels between categorical covariates were tested using *t*-tests or one-way ANOVA, and Pearson correlations were calculated for continuous covariates. Next, a crude analysis of the association between phthalates and MMPs was conducted using simple linear regression. Final adjusted models were constructed by including covariates either associated with both phthalates and MMPs or changed effect estimates by 10% or more. The same covariates were included in all models for consistency. To further examine potential non-linear effects, phthalate metabolite concentrations were divided into tertiles and regressed on MMP levels while adjusting for the same covariates.

3. Results

The majority of women in the study were white and college-educated as described previously (Ferguson et al., 2014). There was a suggestive difference in MMP-1 levels by insurance provider ($p = 0.16$). We also found a significant difference between BMI categories in MMP-2

Table 2
Adjusted percent change (95% Confidence Intervals) in MMP level associated with an interquartile range (IQR) increase in phthalate metabolite concentrations. N = 134.

	MMP-1				MMP-2				MMP-9		
	% Δ	95% CI	p-Value		% Δ	95% CI	p-Value		% Δ	95% CI	p-Value
MEHP	18.10	−3.92, 44.9	0.11		1.94	−11.2, 17.0	0.79		10.43	−4.84, 28.3	0.19
MEHHP	15.11	−8.23, 44.4	0.22		2.78	−11.5, 19.6	0.71		17.45	−0.16, 38.4	0.05
MEOHP	23.77	−1.08, 54.6	0.06		5.57	−8.97, 22.7	0.47		14.52	−2.50, 34.7	0.10
MECPP	25.89	0.19, 58.2	0.05		8.32	−6.97, 26.4	0.30		13.60	−3.92, 34.1	0.13
ΣDEHP	23.50	−0.33, 52.8	0.05		6.04	−8.19, 22.3	0.42		14.92	−1.66, 34.3	0.08
MBZP	−1.29	−24.9, 29.5	0.92		18.22	−1.11, 41.1	0.07		−2.75	−20.2, 18.4	0.78
MBP	36.52	−1.63, 89.5	0.06		14.38	−8.02, 42.5	0.22		−0.65	−21.9, 26.4	0.96
MIBP	28.08	−6.63, 76.0	0.12		16.44	−5.69, 43.8	0.16		−12.96	−30.9, 9.63	0.24
MEP	−1.81	−21.9, 23.1	0.87		13.64	−2.26, 31.8	0.10		−4.46	−19.0, 12.6	0.59
MCP	25.09	−0.84, 58.1	0.06		9.96	−5.92, 28.5	0.23		−0.17	−15.9, 18.5	0.98

Adjusted for BMI, insurance, specific gravity, age.




Fig. 1. Adjusted regression coefficients for multiplicative change MMP-1 level in relation to tertiles of urinary Σ DEHP concentration. p-Value for trend = 0.008. Adjusted for BMI, insurance, specific gravity, age.

levels ($p = 0.03$) and a suggestive difference in MMP-9 levels ($p = 0.15$) by BMI category. In Table 1, geometric means as well as 25th, 75th, 95th and 100th percentiles of phthalate and MMP levels are presented. Crude results (not presented) were similar to those in the adjusted models (Table 2). We observed a significant positive association between MMP-1 and MECPP as well as suggestive associations for MEHP, MEOHP, Σ DEHP, MBP, MIBP and MCP. For MMP-2, we found suggestive positive associations with MBZP and MEP. For MMP-9, we observed suggestive associations between MEHP, MEOHP, MECPP and Σ DEHP. When tertiles of summed DEHP metabolites were regressed on MMP levels, results were consistent with the models using continuous variables and some strong increasing trends were observed. For example, the second and third tertiles of Σ DEHP were associated with increases in MMP-1 of 42% and 87%, respectively, compared to the lowest Σ DEHP tertile (Fig. 1).

4. Discussion

The results of this exploratory study suggest that urinary phthalate metabolites are associated with increased circulating MMP levels in pregnant women. We observed the strongest relationships between DEHP metabolites and MMP-1 and MMP-9 levels. We found suggestive positive associations between DBP and DiBP metabolites and MMP-1 as well as between BzBP and DEP metabolites and MMP-2. To our knowledge this is the first human study to test these associations.

Our findings in a human population are supported by several experimental studies that have shown a positive association between phthalate levels and MMP expression in various cell types. In an in vitro study by Kim et al. (2015), endometrial cells treated with DEHP were found to have increased MMP-9 levels. Another in vitro study of breast cancer cells by Zhang et al. (2016) also reported that DEHP was associated with the overexpression of MMP-2 and MMP-9. A third in vitro study by Zhu et al. (2010) reported similar trends in human neuroblastoma cells. Among other phthalates, an in vivo study by Scarano et al. (2009) found that rats treated with DBP had higher MMP-9 activity in the prostate, whereas we found that the DBP metabolites MBP and MCP were associated with MMP-1, but not associated with MMP-2 or MMP-9. To our knowledge, there are no studies that have tested the effect of phthalates on MMP-1. Future research is needed to obtain a comprehensive understanding of the effects of phthalates on MMP expression and activity.

Despite the present study's exploratory nature and relatively small sample size, our findings may help inform the existing body of research concerning environmental as they relate to various adverse pregnancy outcomes influenced by inflammatory pathways. MMPs are involved in significant degradation and remodeling of tissues. The role of MMPs

prior to parturition are less defined, though MMP-8 levels in amniotic fluid are predictive of intra-amniotic inflammation/infection and early spontaneous delivery (Chaemsaitong et al., 2017; Kim et al., 2016). MMP-1, 2 and 9 have been shown to increase with increasing cervical dilation and play an important role in uterine postpartum involution in animals (Manase et al., 2006; Winkler et al., 1999), while MMP-9 facilitates fetal membrane rupture and placental separation (Tsatsas et al., 1999).

Since MMPs are a vital part of the reproductive process, there is a need to assess which chemicals may impact their balance that could play a role in adverse pregnancy outcomes such as preeclampsia and preterm birth (Geng et al., 2016). Phthalates have been found to be associated with adverse pregnancy outcomes, but knowledge on relevant mechanisms is lacking. The present study observed a positive association between metabolites of DEHP, and potentially other phthalates, and MMP levels. However, as an exploratory study our sample size and statistical power were limited and future full-scale studies should address the potential for MMP levels to mediate associations between phthalates or other environmental exposures and adverse pregnancy outcomes.

Acknowledgments

Funding was provided by the National Institute of Environmental Health Sciences, National Institutes of Health (R01ES018872, P42ES017198, P50ES026049, and U2CES026553). The authors have no competing interests to declare.

References

- Ab Hamid, J., Mohtarrudin, N., Osman, M., Andi Asri, A.A., Wan Hassan, W.H., Aziz, R., 2012. Matrix metalloproteinase-9 and tissue inhibitors of metalloproteinases 1 and 2 as potential biomarkers for gestational hypertension. *Singap. Med. J.* 53, 681–683.
- Cantonwine, D.E., Meeker, J.D., Ferguson, K.K., Mukherjee, B., Hauser, R., McElrath, T.F., 2016. Urinary concentrations of bisphenol A and phthalate metabolites measured during pregnancy and risk of preeclampsia. *Environ. Health Perspect.* 124, 1651–1655.
- Chaemsaitong, P., Romero, R., Docheva, N., Chaiyasit, N., Bhatti, G., Pacora, P., et al., 2017. Comparison of rapid MMP-8 and interleukin-6 point-of-care tests to identify intra-amniotic inflammation/infection and impending preterm delivery in patients with preterm labor and intact membranes. *J. Matern. Fetal Neonatal Med.* 1–17.
- Eleuterio, N.M., Palei, A.C., Rangel Machado, J.S., Tanus-Santos, J.E., Cavalli, R.C., Sandrim, V.C., 2015. Positive correlations between circulating adiponectin and MMP2 in preeclampsia pregnant. *Pregnancy Hypertens* 5, 205–208.
- Ferguson, K.K., McElrath, T.F., Meeker, J.D., 2014. Environmental phthalate exposure and preterm birth. *JAMA Pediatr.* 168, 61–67.
- Geng, J., Huang, C., Jiang, S., 2016. Roles and regulation of the matrix metalloproteinase system in parturition. *Mol. Reprod. Dev.* 83, 276–286.
- Kim, S.H., Cho, S., Ihm, H.J., Oh, Y.S., Heo, S.H., Chun, S., et al., 2015. Possible role of phthalate in the pathogenesis of endometriosis: in vitro, animal, and human data. *J. Clin. Endocrinol. Metab.* 100, E1502–E1511.

- Kim, S.M., Romero, R., Lee, J., Chaemsaitong, P., Lee, M.W., Chaivayit, N., et al., 2016. About one-half of early spontaneous preterm deliveries can be identified by a rapid matrix metalloproteinase-8 (MMP-8) bedside test at the time of mid-trimester genetic amniocentesis. *J. Matern. Fetal Neonatal Med.* 29, 2414–2422.
- Kramer, M.S., Kahn, S.R., Platt, R.W., Genest, J., Chen, M.F., Goulet, L., et al., 2010. Mid-trimester maternal plasma cytokines and CRP as predictors of spontaneous preterm birth. *Cytokine* 49, 10–14.
- Lewis, R.C., Meeker, J.D., Peterson, K.E., Lee, J.M., Pace, G.G., Cantoral, A., et al., 2013. Predictors of urinary bisphenol A and phthalate metabolite concentrations in Mexican children. *Chemosphere* 93, 2390–2398.
- Manase, K., Endo, T., Chida, M., Nagasawa, K., Honnma, H., Yamazaki, K., et al., 2006. Coordinated elevation of membrane type 1-matrix metalloproteinase and matrix metalloproteinase-2 expression in rat uterus during postpartum involution. *Reprod. Biol. Endocrinol.* 4, 32.
- Marie, C., Vendittelli, F., Sauviant-Rochat, M.P., 2015. Obstetrical outcomes and biomarkers to assess exposure to phthalates: a review. *Environ. Int.* 83, 116–136.
- Messerlian, C., Wylie, B.J., Minguez-Alarcon, L., Williams, P.L., Ford, J.B., Souter, I.C., et al., 2016. Urinary concentrations of phthalate metabolites and pregnancy loss among women conceiving with medically assisted reproduction. *Epidemiology* 27, 879–888.
- Nissi, R., Talvensaari-Mattila, A., Kotila, V., Niinimäki, M., Jarvela, I., Turpeenniemi-Hujanen, T., 2013. Circulating matrix metalloproteinase MMP-9 and MMP-2/TIMP-2 complex are associated with spontaneous early pregnancy failure. *Reprod. Biol. Endocrinol.* 11, 2.
- Scarano, W.R., Toledo, F.C., Guerra, M.T., de Campos, S.G., Junior, L.A., Felisbino, S.L., et al., 2009. Long-term effects of developmental exposure to di-n-butyl-phthalate (DBP) on rat prostate: proliferative and inflammatory disorders and a possible role of androgens. *Toxicology* 262, 215–223.
- Tsatas, D., Baker, M.S., Rice, G.E., 1999. Differential expression of proteases in human gestational tissues before, during and after spontaneous-onset labour at term. *J. Reprod. Fertil.* 116, 43–49.
- Werner, E.F., Braun, J.M., Yolton, K., Khoury, J.C., Lanphear, B.P., 2015. The association between maternal urinary phthalate concentrations and blood pressure in pregnancy: the HOME study. *Environ. Health* 14, 75.
- Winkler, M., Oberpichler, A., Tschesche, H., Ruck, P., Fischer, D.C., Rath, W., 1999. Collagenolysis in the lower uterine segment during parturition at term: correlations with stage of cervical dilatation and duration of labor. *Am. J. Obstet. Gynecol.* 181, 153–158.
- Yao, P.L., Lin, Y.C., Richburg, J.H., 2012. Mono-(2-ethylhexyl) phthalate (MEHP) promotes invasion and migration of human testicular embryonal carcinoma cells. *Biol. Reprod.* 86 (160) (161–110).
- Zhang, S., Ma, J., Fu, Z., Zhang, Z., Cao, J., Huang, L., et al., 2016. Promotion of breast cancer cells MDA-MB-231 invasion by di(2-ethylhexyl)phthalate through matrix metalloproteinase-2/-9 overexpression. *Environ. Sci. Pollut. Res. Int.* 23, 9742–9749.
- Zhu, H., Zheng, J., Xiao, X., Zheng, S., Dong, K., Liu, J., et al., 2010. Environmental endocrine disruptors promote invasion and metastasis of SK-N-SH human neuroblastoma cells. *Oncol. Rep.* 23, 129–139.