


Pregnancy Nutrition Surveillance



2009 Report

This report summarizes selected data on maternal health and nutrition indicators received from states, U.S. territories, and Indian Tribal Organizations that contributed to the Centers for Disease Control and Prevention's (CDC) Pregnancy Nutrition Surveillance System.

This report was developed by CDC's Division of Nutrition, Physical Activity, and Obesity in the National Center for Chronic Disease Prevention and Health Promotion.

Suggested Citation

Reinold C, Dalenius K, Brindley P, Smith B, Grummer-Strawn L. *Pregnancy Nutrition Surveillance 2009 Report*. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2011.

Acknowledgements

We gratefully acknowledge and thank all contributors to the national Pregnancy Nutrition Surveillance System. The efforts of state, territorial, and Indian Tribal Organization surveillance coordinators; informatics staff; and local clinic staff to collect data and use nutrition surveillance systems make the Pregnancy Nutrition Surveillance System possible.

This report is available online at www.cdc.gov/pednss/pdfs/PNSS_2009.pdf.

For more information, contact

Division of Nutrition, Physical Activity, and Obesity
National Center for Chronic Disease Prevention and Health Promotion
Centers for Disease Control and Prevention

4770 Buford Highway NE, Mail Stop K-25
Atlanta, GA 30341-3717
800-CDC-INFO (232-4636)
TTY: 888-232-6348

Pregnancy Nutrition Surveillance

The Pregnancy Nutrition Surveillance System (PNSS) is a public health surveillance system that monitors the prevalence of nutrition problems, behavioral risk factors, and birth outcomes among low-income women who are enrolled in public health programs in states or U.S. territories or through Indian Tribal Organizations (ITOs).

The goal of the PNSS is to collect, analyze, and disseminate surveillance data to guide public health policy and action. The collected data are used to set public health priorities and to plan, implement, and evaluate nutrition programs for pregnant women. This report summarizes PNSS data from 2009 and highlights data trends from 2000 through 2009.

The PNSS collects demographic data about maternal age, race and ethnicity, education level, household income, migrant status, and participation in food and medical assistance programs. Information about a mother's height; weight before, during, and after pregnancy; hemoglobin and hematocrit levels; parity; medical care during pregnancy; and enrollment in the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) also is collected.

In addition, the PNSS collects information about patterns of maternal smoking and alcohol consumption 3 months before and during pregnancy. Data collected about infants include date of birth, birthweight, and breastfeeding status.

Data are collected at the clinic level, aggregated at the state level, and then submitted to the Centers for Disease Control and Prevention (CDC) for analysis. The PNSS generates surveillance reports for each contributor and for the nation, based on results from participating contributors. A *contributor* is defined as a state, U.S. territory, or ITO.


In 2009, the PNSS included 31 states, the District of Columbia, 5 ITOs, and 1 U.S. territory (Figure 1). These contributors provided more than 1.3 million records to the system, nearly double the number of records collected in 2000. This gain is largely due to the increased number of contributors to the PNSS. Fluctuations in the number of contributors and records can affect trends. The number of PNSS contributors shown differs slightly from year to year because some contributors did not provide data every year during the 10-year period from 2000 through 2009 (Table 1). The WIC program has consistently been the primary source of PNSS data, contributing nearly 100% of the records in 2009.

Demographic Characteristics

Income

More than 60% of women who participated in the 2009 PNSS and were eligible for WIC had gross incomes equal to or less than 100% of the U.S. poverty level. To be eligible for the WIC program, an applicant's gross income must be equal to or less than 185% of the 2009 U.S. Department of Health and Human Services' poverty guidelines.

Figure 1. Contributors* to the 2009 Pregnancy Nutrition Surveillance System



* In addition to states shown, includes the District of Columbia, Puerto Rico, and the following Indian Tribal Organizations: the Cheyenne River Sioux Tribe (SD), the Inter Tribal Council of Arizona, the Northern Arapaho Tribe (WY), the Rosebud Sioux Tribe (SD), and the Standing Rock Sioux Tribe (ND).

Table 1. Contributors to the Pregnancy Nutrition Surveillance System, 2000–2009*

Contributor	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Alabama										
Arizona										
American Samoa										
Cheyenne River Sioux Tribe (SD)										
Chickasaw Nation (OK)										
Connecticut										
District of Columbia										
Florida										
Georgia										
Hawaii										
Idaho										
Illinois										
Indiana										
Inter Tribal Council (AZ)										
Iowa										
Kansas										
Massachusetts										
Michigan										
Minnesota										
Missouri										
Montana										
Nebraska										
New Hampshire										
New Jersey										
New Mexico										
New York										
Northern Arapaho Tribe (WY)										
North Carolina										
North Dakota										
Ohio										
Oregon										
Puerto Rico										
Rhode Island										
Rosebud Sioux Tribe (SD)										
South Carolina										
Standing Rock Sioux Tribe (ND)										
Three Affiliated Tribes (ND)										
Utah										
Vermont										
Virginia										
West Virginia										
Wichita Caddo Tribe (OK)										
Wisconsin										
Number of Contributors	26	29	27	29	31	31	32	34	38	38
Total Records Submitted (x 1,000)	751	739	727	769	856	857	1,143	1,253	1,308	1,304

* Shaded blocks indicate years that data were contributed.

Race and Ethnicity

Of the women in the 2009 PNSS, 41.6% were non-Hispanic white, 24.1% were non-Hispanic black, 28.3% were Hispanic or Latina, 3.2% were Asian or Other Pacific Islander, 1.1% were American Indian or Alaska Native, and 1.2% were of multiple races. In the 2009 PNSS, a smaller proportion of women was non-Hispanic white (41.6%) compared with the proportion of women in the 2000 PNSS (50.0%), and more women were Hispanic or Latina in 2009 (28.4%) than in 2000 (19.8%).

Age

In the 2009 PNSS, 59.4% of women were aged 20–29 years, 20.4% were aged 30–39 years, and 1.5% were aged 40 years or older. The proportion of teenaged mothers declined from 24.2% in 2000 to 18.4% in 2009. The proportion of pregnant women aged 30 years or older increased from 17.3% in 2000 to 21.9% in 2009.

Education

In the 2009 PNSS, 30.9% of women had less than a high school education, and this proportion has changed little during the past decade. Among all women in the United States who gave birth in 2007 and for whom education data were available, 23.1% had less than a high school education.¹

Maternal Health Indicators

Prepregnancy Weight Status

Prepregnancy weight is a determinant of infant birthweight. For example, studies suggest an association between being underweight before pregnancy and giving birth to an infant with low birthweight.² Obese women are at increased risk of preeclampsia, gestational diabetes, cesarean delivery, and failure to initiate breastfeeding.^{2,3}

In 2009, the Institute of Medicine (IOM) released updated guidelines for categorizing body mass index (BMI) values for women as well as appropriate weight gain during pregnancy. Before this update, it had been nearly 20 years since the IOM had issued guidelines and, in this time, more research had been conducted on the effects of weight gain in pregnancy

on the health of both mother and baby. Significant changes also have occurred in the population of U.S. women having babies. American women are a diverse group, they are having more twin and triplet pregnancies, and they tend to be older when they become pregnant. In addition, a greater percentage of them are entering pregnancy overweight or obese, and many are gaining too much weight during pregnancy. Many of these changes make both mothers and their babies vulnerable to additional health risks.⁴

Prepregnancy BMI was calculated for each woman in the 2009 PNSS and was based on self-reported weight and measured height. Women were classified by the IOM's new 2009 BMI categories.⁴ *Underweight* is defined as a BMI less than 18.5 (1990 IOM definition was a BMI less than 19.8). *Normal weight* is defined as a BMI of 18.5–24.9 (1990 IOM definition was a BMI of 19.8–26.0). *Overweight* is defined as a BMI of 25.0–29.9 (1990 IOM definition was a BMI of 26.1–29.0), and *obese* is defined as a BMI greater than or equal to 30.0 (1990 IOM definition was a BMI greater than 29.0). These same adult BMI categories apply to teenagers who are pregnant.

In the 2009 PNSS, 4.5% of women were underweight, 42.5% were normal weight, 25.9% were overweight, and 27.0% were obese. The overall proportion of women in the 2009 PNSS who were overweight or obese was 52.9%. Proportions varied by contributor, from 44.9% (Puerto Rico) to 69.5% (Inter Tribal Council of Arizona) (Table 2).

To ensure consistency in its trend analysis, CDC applied the BMI cutoffs recommended in the 2009 IOM report to historical data. Based on the new standards, the prevalence of prepregnancy underweight is lower, the prevalence of overweight is substantially higher, and the prevalences of normal weight and obesity are lower.

From 2000 through 2009, the prevalence of women who were overweight or obese before pregnancy increased from 46.9% to 52.9%, but the percentage of women who were underweight before they became pregnant decreased from 6.2% to 4.5% (Figure 2).

Table 2. State-specific prevalence of selected indicators of maternal health, 2009 Pregnancy Nutrition Surveillance System*

Contributors	Under-weight [†]	Over-weight [†]	< Ideal wt gain [‡]	> Ideal wt gain [‡]	3rd trim anemia [§]	Post p anemia	1st trim med	No med	1st trim WIC	Smoking					House postp	LBW [¶]	HBW ^{**}	Pre-term ^{††}	Ever BF ^{‡‡}
										3 mos b/f preg	3rd trim	House prenatal	House postp	3 mos b/f preg					
Alabama	5.7	57.0	*	*	41.3	38.3	*	41.6	*	*	*	*	*	10.1	4.6	14.4	*		
Arizona	4.7	51.7	20.0	51.4	23.7	20.7	*	35.7	6.8	3.1	5.6	5.7	7.1	6.8	10.7	67.0	*		
Cheyenne River Sioux Tribe (SD)	2.2	62.6	*	*	*	*	*	34.7	51.7	*	39.2	*	*	*	*	*	*		
Connecticut	3.8	52.9	*	*	30.9	53.0	85.1	4.3	24.1	24.6	2.9	9.2	*	7.0	7.3	7.5	68.9		
District of Columbia	3.8	56.9	27.3	43.5	49.0	41.8	85.3	4.7	24.1	6.4	4.2	12.6	9.7	9.9	5.5	15.7	54.3		
Florida	4.8	50.8	22.2	49.0	40.6	36.6	*	33.6	16.0	19.3	*	*	*	8.6	5.9	12.5	73.0		
Georgia	4.1	55.2	22.6	48.9	36.4	26.5	81.5	3.6	41.2	17.2	9.5	*	*	*	*	10.9	55.6		
Hawaii	4.9	48.0	18.9	53.1	31.1	20.8	76.0	5.5	23.7	*	*	*	*	7.5	6.6	5.0	86.2		
Idaho	3.7	52.9	17.7	49.2	22.2	18.7	*	34.4	25.4	13.0	18.6	13.2	6.4	7.3	10.0	84.1			
Illinois	4.0	54.5	23.5	47.7	32.5	34.2	*	38.4	19.3	10.8	17.9	13.8	6.2	7.7	7.6	66.5			
Indiana	5.0	54.1	18.0	54.5	37.2	27.1	*	37.9	29.0	21.1	37.4	32.4	8.4	6.3	14.3	63.3			
Inter Tribal Council (AZ)	1.7	69.5	19.4	54.8	28.8	28.0	74.2	4.0	27.4	12.0	1.5	12.0	9.8	6.1	9.1	8.5	67.7		
Iowa	2.6	58.2	16.7	52.1	30.1	22.9	*	44.1	39.8	23.1	20.8	8.8	7.6	8.5	10.2	*	*		
Kansas	4.6	54.3	19.9	51.1	29.5	22.7	75.4	5.7	35.7	31.5	17.8	18.8	15.9	7.7	6.5	8.8	70.3		
Massachusetts	4.2	49.0	19.6	51.8	34.7	29.9	73.7	11.3	35.5	21.9	10.6	16.9	12.4	7.9	7.9	9.1	75.1		
Michigan	4.2	54.2	21.8	49.4	36.8	27.6	86.8	1.0	32.5	35.8	19.3	17.0	6.5	8.4	7.1	11.5	57.2		
Minnesota	3.6	54.2	29.2	41.8	28.5	19.3	*	37.1	28.3	10.8	*	*	*	6.7	9.0	9.6	74.2		
Montana	4.3	51.8	19.9	51.8	14.6	22.4	79.2	8.4	27.4	36.2	6.6	20.4	17.9	6.5	7.7	10.0	77.4		
Nebraska	3.7	55.2	20.5	47.5	27.2	26.4	87.5	2.7	28.2	32.7	15.7	*	*	7.3	7.4	10.7	71.2		
New Hampshire	4.8	51.7	17.4	54.8	29.4	23.0	90.5	1.4	33.9	43.2	25.4	31.8	28.1	7.3	10.0	7.1	72.4		
New Jersey	3.9	52.4	20.2	45.4	36.6	49.9	*	25.2	7.5	4.0	7.2	6.9	7.2	6.6	6.6	13.1	61.8		
New Mexico	4.7	52.6	19.7	45.7	25.2	18.8	*	36.4	21.8	4.8	7.9	9.5	8.3	4.7	4.7	12.1	73.8		
New York	4.6	50.1	22.2	41.7	37.0	32.3	85.4	1.8	29.1	16.2	8.1	7.3	*	8.0	6.8	10.5	72.6		
North Carolina	4.9	55.1	22.5	46.2	35.3	32.2	78.7	0.7	27.2	24.1	14.5	*	*	9.1	6.6	14.1	62.5		
North Dakota	2.8	56.5	15.8	52.6	27.2	21.3	*	43.0	46.6	25.5	18.0	7.6	8.2	8.7	8.7	12.4	*		
Northern Arapaho Tribe (WY)	4.0	63.4	*	*	*	*	*	36.4	*	*	*	*	*	*	*	*	*		
Ohio	5.1	53.3	20.8	52.2	31.9	25.6	*	33.3	38.1	21.5	37.7	34.8	8.7	6.2	6.2	10.7	48.6		
Oregon	3.3	55.8	17.2	49.6	28.1	20.0	*	39.6	*	*	*	*	*	6.1	9.8	8.1	91.2		
Puerto Rico	8.0	44.9	25.5	40.3	41.1	31.1	90.7	0.4	*	*	*	*	*	15.1	1.9	16.6	55.8		
Rhode Island	4.5	50.7	19.7	49.9	32.0	33.2	86.4	5.4	43.5	20.4	11.0	17.4	14.4	7.7	8.0	14.0	62.3		
Rosebud Sioux Tribe (SD)	3.4	56.0	27.5	46.1	*	42.6	76.4	10.0	31.8	42.4	16.4	27.5	16.6	3.8	8.5	11.0	64.8		
South Carolina	5.0	54.8	17.3	51.4	35.7	32.8	*	34.9	27.2	14.1	13.3	10.3	9.8	5.3	*	*	*		
Standing Rock Sioux Tribe (ND)	3.0	57.7	*	*	*	*	*	29.6	52.5	*	*	*	*	*	*	*	*		
Utah	4.0	50.6	19.4	48.7	20.5	11.7	82.1	7.7	20.1	3.3	1.6	5.5	3.3	7.1	6.1	10.8	91.5		
Vermont	4.2	52.1	19.5	48.8	22.0	23.2	79.0	0.4	34.1	43.8	29.3	*	*	7.5	10.6	9.9	77.7		
Virginia	4.8	53.6	18.7	49.0	36.3	25.8	74.9	2.5	31.9	29.2	11.0	22.6	44.6	9.1	6.6	10.6	61.2		
West Virginia	7.0	51.3	15.3	52.5	28.3	21.2	85.4	5.3	49.2	51.1	33.7	28.5	19.2	9.0	6.3	12.1	46.3		
Wisconsin	3.7	55.1	21.6	50.9	31.9	25.0	85.6	2.8	34.1	36.1	19.7	20.4	6.9	7.8	7.8	9.7	69.0		
National PNSS	4.5	52.9	21.2	48.2	33.9	28.8	82.2	3.1	34.1	22.9	13.0	17.1	16.8	8.2	6.7	11.1	66.7		

Abbreviations: Wt = weight; Trim = trimester; Post p = postpartum; Med = entry to medical care; WIC = Special Supplemental Nutrition Program for Women, Infants, and Children; Mos = months; b/f preg = before pregnant; House = household; LBW = low birthweight; HBW = high birthweight; Ever BF = ever breastfed.
 * Percentages are not calculated if <100 records were available for analysis after exclusion of missing or misclassified data.
 † Prepregnancy weight status: Underweight = body mass index (BMI) <18.5; Overweight = BMI ≥25.0 (includes overweight and obese). Source: *Weight Gain During Pregnancy: Reexamining the Guidelines*. National Academies Press; 2009.
 ‡ Recommended pregnancy weight gain = 28–40 lb if underweight, 25–35 lb if normal weight, 15–25 lb if obese. Source: *Weight Gain During Pregnancy: Reexamining the Guidelines*. National Academies Press; 2009.
 § Adjusted for altitude and smoking. Source: *MMWR* 1998;47(RR-3):1–36.
 ¶ Defined as birthweight <2,500 grams.
 ** Defined as birthweight >4,000 grams.
 †† Defined as gestational period <37 weeks.
 ‡‡ Among infants born during the reporting period. *Healthy People 2010* objective 16-19: Increase the proportion of mothers who breastfeed their babies during the early postpartum period to 75%. Source: *Healthy People 2010*. Volume II. U.S. Government Printing Office; 2000.

Maternal Weight Gain

IOM recommendations for maternal weight gain are unchanged for underweight, normal weight, and overweight women. The IOM recommends a pregnancy weight gain of 28–40 pounds for underweight women, 25–35 pounds for women of normal weight, and 15–25 pounds for overweight women. For obese women, IOM now recommends a pregnancy weight gain of 11–20 pounds (prior recommendation was at least 15 pounds). Recommended weight gain for short women and for individual racial and ethnic groups are the same as those for the whole population.⁴

Women who gain less than the IOM's recommendation during pregnancy are at increased risk of giving birth to an infant with low birthweight. Women who gain more than the IOM's recommended weight gain during pregnancy are at increased risk of giving birth to an infant with high birthweight, which can cause difficulty with delivery.⁴

Additionally, women who gain excess weight during pregnancy may have more difficulty returning to their prepregnancy weight.⁴ In the 2009


PNSS, 21.2% of women gained less weight than recommended during pregnancy, 30.6% gained the recommended amount of weight, and 48.2% gained more weight than recommended. The proportion of women in the 2009 PNSS who gained more than the recommended amount of weight during pregnancy varied by contributor, from 40.3% (Puerto Rico) to 54.8% (New Hampshire and the Inter Tribal Council of Arizona) (Table 2).

Whether the women in the 2009 PNSS gained an ideal amount of weight during pregnancy varied significantly by the women's prepregnancy BMI (Figure 3). Women who were overweight or obese before pregnancy were more likely to exceed the IOM's recommended maximum weight gain for their body size. Excess weight gain during pregnancy and failure to lose weight after pregnancy are important and identifiable predictors of long-term obesity.⁵

From 2000 through 2009, the prevalence of women who gained too much weight during pregnancy remained constant (48.3% and 48.2%), and the prevalence of women who gained less than the

IOM's recommended weight gain during pregnancy dropped slightly, from 22.4% to 21.2%.

Figure 2. Trends in prevalence* of prepregnancy overweight and underweight†



* State, U.S. territory, and Indian Tribal Organization contributors to the PNSS change slightly each year. Differences in prevalence trends should be interpreted cautiously.

† Overweight = body mass index (BMI) ≥ 25.0 (includes overweight and obese); underweight = BMI < 18.5 .

Anemia

Iron deficiency is common among women of reproductive age, including during pregnancy.⁶ Because pregnant women require higher amounts of iron, iron supplementation during pregnancy is often recommended. Pregnant women may not consume an adequate amount of iron if they do not take iron supplements during pregnancy, especially during the first trimester.⁶

Iron-deficiency anemia* during the first two trimesters of pregnancy has been associated with inadequate gestational weight gain, a two-fold risk of preterm delivery, and a three-fold risk of giving birth to an infant with low birthweight.⁷ Iron-deficiency anemia during the third trimester of pregnancy is associated with a two-fold risk of an inadequate weight gain for gestation.⁷

In the 2009 PNSS, 7.7% of women had anemia when they enrolled in the WIC program during their first trimester, 12.2% had anemia when they enrolled in the program during their second trimester, and 33.9% had anemia when they enrolled during their third trimester. Women who enrolled in the WIC program during their third trimester were more likely to start prenatal care late in pregnancy and may not be representative of all low-income women in their third trimester of pregnancy.

Healthy People 2010 objective 19-13 proposed reducing the prevalence of third trimester anemia among low-income pregnant females to no more than 20%.⁸

In the 2009 PNSS, the prevalence of anemia in the third trimester of pregnancy was highest for black mothers (49.5%). The prevalences for other racial and ethnic groups were all above the *Healthy People 2010* target of 20% (27.2% for whites, 29.9% for Hispanics or Latinas, 30.7% for American Indians or Alaska Natives, and 29.6% for Asians and Native Hawaiians or Other Pacific Islanders).


From 2000 through 2009, the overall prevalence of anemia during the third trimester of pregnancy rose slightly, from 29.7% to 33.9%. In the 2009 PNSS, only Montana met the *Healthy People 2010* objective for the prevalence of anemia during the third trimester of pregnancy.

Women at risk of anemia at 4–6 weeks postpartum should be screened.⁸ In the 2009 PNSS, 28.8% of postpartum women were anemic at 6 or more weeks after delivery. Racial and ethnic disparities were similar to those described previously for pregnant women, with anemia rates ranging from 22.1% among white mothers to 46.0% among black mothers.

Interpregnancy Interval

Interpregnancy interval is the time between the end of one pregnancy and the last menstrual period before the next pregnancy. Women with an interpregnancy interval of less than 6 months are at higher risk of experiencing maternal mortality and morbidity, giving birth to infants with low birthweight or infants who are small for gestational age, and giving birth preterm than are women who conceive after an interpregnancy interval of 18–23 months.^{9,10} In addition, a shorter interpregnancy interval means less time for repletion of nutrient

Figure 3. Maternal weight gain* by prepregnancy body mass index[†]



* The Institute of Medicine recommends a weight gain of 28–40 lb if underweight prior to pregnancy, 25–35 lbs if normal weight, 15–25 lb if overweight, and 11–20 lb if obese. Source: *Weight Gain During Pregnancy: Reexamining the Guidelines*. National Academies Press; 2009.

[†] Underweight = body mass index (BMI) <18.5, normal weight = BMI 18.5–24.9, overweight = BMI 25.0–29.9, obese = BMI ≥30.0.

* Anemia: Low hemoglobin (Hb) or low hematocrit (Hct). CDC defines anemia during pregnancy as follows: first trimester, Hb <11.0 g/dl or Hct <33%; second trimester, Hb <10.5 g/dl or Hct <32%; and third trimester, Hb <11.0 g/dl or Hct <33%.⁶

stores.¹¹ In the 2009 PNSS, 12.0% of all women had an interpregnancy interval of less than 6 months, and 26.7% had an interval of 6 months to less than 18 months.

Maternal Behavioral Indicators

Prenatal Care

The quality, quantity, and timing of prenatal care influence pregnancy outcomes, and inadequate prenatal care increases a woman's risk of a poor pregnancy outcome.¹² *Healthy People 2010* objective 16-6a proposed increasing the proportion of pregnant women who receive prenatal care during the first trimester to 90%. According to national health statistics in 2006, 68.3% of women sought prenatal care during their first trimester, and 8.2% of women received no prenatal care during pregnancy.¹³

In the 2009 PNSS, 82.2% of women began prenatal care during their first trimester—an increase from 73.9% reported in 2000—and 3.1% received no prenatal care. Mothers aged 19 years or younger were less likely than older mothers to obtain prenatal care during the first trimester. By age group, 60.4% of teenagers younger than age 15 years, 74.0% of those aged 15–17 years, and 79.5% of those 18–19 years obtained prenatal care during the first trimester.

White women (86.3%) were more likely to obtain prenatal care during the first trimester of pregnancy than blacks (79.1%), Hispanics (79.5%), American Indians or Alaska Natives (78.3%), or Asians and Native Hawaiians or Other Pacific Islanders (80.0%).

WIC Enrollment

Several studies have concluded that participation in the WIC program is associated with improved birthweights and fewer preterm deliveries.^{12,14–16} One study concluded that participation in the WIC program during pregnancy resulted in fewer deliveries of infants who are small for gestational age; the study also found that having a healthier infant was linked to longer enrollment in the program.¹⁵

In the 2009 PNSS, 34.1% of women enrolled in the WIC program during their first trimester, 34.7%

during their second trimester, 17.5% during their third trimester, and 13.7% after giving birth. The proportion of women who enrolled during their first trimester increased from 26.8% in 2000. Percentages varied among contributors, from 20.1% (Utah) to 49.2% (West Virginia) (Table 2).

Maternal Smoking

Infants born to mothers who smoke cigarettes during pregnancy have a 40.0% higher rate of mortality than infants born to nonsmoking mothers.¹⁷ Small gestational size, rather than preterm birth, is the main mechanism through which smoking causes excess infant mortality.¹⁷ The percentage of preterm births is not significantly different for smoking mothers when compared with nonsmoking mothers, but the percentages of low birthweight and full-term low birthweight are significantly higher for infants born to smoking mothers (Figure 4).

Maternal smoking also increases the risk of sudden infant death syndrome (SIDS)¹⁸ and spontaneous abortion,¹⁹ and it has long-term negative effects on the growth, development, behavior, and cognition of the infant.^{20,21} *Healthy People 2010* objective 16-17c proposed increasing the rate of abstinence from smoking during the last month of pregnancy to 98%.

In the 2009 PNSS, 22.9% of women reported smoking during the 3 months before they became pregnant, and 13.0% reported smoking during the last 3 months of their pregnancy. The proportions of women who smoke before pregnancy or during the last trimester of pregnancy have consistently declined since 2000, when 31.3% of women reported that they smoked before becoming pregnant and 21.4% reported smoking during their last trimester.

In the 2009 PNSS, the prevalence of smoking during the last 3 months of pregnancy was highest among white women (22.8%), women aged 20–29 years (14.1%), and women with less than a high school education (15.6%). The proportion of women in the 2009 PNSS who smoked during their last trimester of pregnancy varied widely by contributor, from 1.5% (Inter Tribal Council of Arizona) to 33.7% (West Virginia) (Table 2).

The PNSS also monitors the proportion of women who quit smoking. Of the 22.9% of women who reported smoking during the 3 months before they became pregnant, 43.1% reported that they quit smoking by the time they enrolled in WIC, and 36.7% of these women abstained from smoking during the last 3 months of their pregnancies.

Household Smoking

Studies of the association between passive smoking during pregnancy and birth outcomes have found a small effect of passive smoking on infant birthweight.²² However, the effects of household smoking on infants' health are better documented.^{22,23} Exposure to passive smoke during the first 2 years of a child's life is associated with a higher incidence of SIDS, respiratory infection, ear infections, and more severe asthma.^{22,23}

In the 2009 PNSS, 17.1% of women reported that, during their pregnancy, someone other than themselves smoked in their household, and 16.8% reported that someone else in the household smoked during the weeks following the birth of the infant.

Birth Outcomes

Low Birthweight

Low birthweight (<2,500 grams) is associated with neonatal and postneonatal mortality.²⁴ Infants with low birthweight who survive are at increased risk of health problems that range from neurodevelopmental handicaps to conditions of the lower respiratory tract.²⁵


Healthy People 2010 objective 16-10a proposed reducing the incidence of low birthweight among U.S. infants to 5%.⁹ In the 2009 PNSS, 8.2% of infants had low birthweights; this proportion is the same as the proportion of U.S. infants born with low birthweights in 2009.²⁶ The incidence of low birthweight varied by contributor, from 3.8% (Rosebud Sioux Tribe) to 10.1% (Alabama) (Table 2).

The overall incidence of low birthweight in 2009 (8.2%) was slightly higher than the incidence in 2000 (7.7%). By racial and ethnic group, the incidence was higher for black infants (11.9%) than for white (7.2%), Hispanic (7.0%), American Indian or Alaska Native (7.4%), or Asian and Native Hawaiian or Other Pacific Islander (7.4%) infants.

High Birthweight

A high birthweight (>4,000 grams) is associated with an increased risk of birth injuries such as shoulder dystocia.²⁷ In the 2009 PNSS, 6.7% of infants had high birthweights, compared with 8.3% in 2000. The prevalence of high birthweight varied by contributor, from 1.9% (Puerto Rico) to 10.6% (Vermont) (Table 2). American Indian or Alaska Native infants had the highest rate (9.4%), followed by white (7.9%), Hispanic (6.8%), Asian and Native Hawaiian or Other Pacific Islander (4.9%), and black (4.2%) infants.

Figure 4. Birth outcomes by mother's smoking status*



* A smoker is defined as a woman who uses any tobacco in the last 3 months of pregnancy.

[†] Defined as birthweight <2,500 grams.

[‡] Defined as gestational period <37 weeks.

[§] Defined as born at or after 37 weeks gestation with birthweight <2,500 grams.

Preterm Births

Preterm birth refers to infants who are born before 37 weeks gestation. It is associated with increased risk of newborn health complications; long-term disabilities such as mental retardation, cerebral palsy, lung and gastrointestinal problems, vision and hearing loss; and death.²⁸ Factors related to preterm birth include iron-deficiency anemia during pregnancy, low gestational weight gain, low income, race and ethnicity, young age of the mother, smoking, and low educational attainment.¹¹

Healthy People 2010 objective 16-11a proposed reducing preterm births to 7.6%. In the 2009 PNSS, 11.1% of infants were born prematurely, up very slightly from 10.9% in 2000. The prevalence of preterm births differed by race, with black infants having a higher prevalence (13.5%) than American Indian or Alaska Native (11.2%), Hispanic (11.1%), white (10.0%), or Asian and Native Hawaiian or Other Pacific Islander (8.8%) infants.

Full-Term Low Birthweight

The PNSS monitors the proportion of full-term, low-birthweight infants and uses this information as an indicator to diagnose intrauterine growth retardation or fetal growth restriction.¹² An infant is considered to be full term with low birthweight if the infant is born at or after 37 weeks gestation but weighs less than 2,500 grams. Poor nutrition during pregnancy is cited as one of the causes of full-term, low-birthweight infants.¹¹ An infant's size at birth is important because fetal growth restriction contributes to the risk of respiratory distress, hypoglycemia, and other health problems in infants.¹¹

In the 2009 PNSS, 3.5% of infants who were born at full term had low birthweight, which is slightly lower than the 3.8% reported in 2000. Black infants (5.1%) had a higher prevalence of being full term with low birthweight than infants from other racial or ethnic groups.

Infant Feeding Practices

Breastmilk, which is nutritionally superior to any other milk supply, provides infants with many benefits, including a reduction in acute otitis media, nonspecific gastroenteritis, severe lower respiratory tract infection, asthma, obesity, type 1 and type 2 diabetes, childhood leukemia, SIDS, and necrotizing enterocolitis.²⁹

The proportion of breastfed infants in the PNSS has steadily increased in recent years. In 2009, 66.7% of infants were breastfed, compared with 52.9% in 2000. *Healthy People 2010* objective 16-19 proposed increasing the proportion of infants ever breastfed to 75%.¹⁰ Only seven states in the 2009 PNSS met this objective: Hawaii, Idaho, Massachusetts, Montana, Oregon, Utah, and Vermont. The percentage of infants ever breastfed varied widely among contributors, from 46.3% (West Virginia) to 91.5% (Utah) (Table 2).


Breastfeeding initiation varied by race and ethnicity. In the 2009 PNSS, initiation rates ranged from 56.8% among black mothers and 64.4% among white mothers to 77.3% among Hispanic mothers. Among all mothers who gave birth in the United States in 2007, 59.7% of black women and 77.7% of white women initiated breastfeeding, while 80.6% of Hispanic women reported initiation.²⁹

Maternal Health Progress Review

Advances in several indicators were observed in the PNSS population from 2000 through 2009 (Figure 5).[†] The prevalence of initiation of breastfeeding increased from 52.9% in 2000 to 66.7% in 2009, a relative increase of 26%. The 2009 rate was the highest ever reported. In addition, since 2000, the proportion of women who enroll in the WIC program during their first trimester increased 27%, and the proportion of women who gain inadequate weight during pregnancy decreased 5%.

[†] For trend data comparing 2000 with 2009 in Figure 5 and throughout this publication, state, U.S. territory, and ITO PNSS contributors change slightly each year. Thus, the differences in prevalence must be interpreted cautiously.

Figure 5. Changes* in maternal and infant health status, 2000–2009



* State, U.S. territory, and Indian Tribal Organization contributors to the PNSS change slightly each year. Differences in prevalence trends should be interpreted cautiously.

[†] WIC = Special Supplemental Food Program for Women, Infants, and Children.

[‡] Prepregnancy overweight and obesity = body mass index >25.0. Source: *Weight Gain During Pregnancy: Reexamining the Guidelines*. National Academies Press; 2009.

The prevalence of women who smoked during the last trimester of pregnancy has dropped 39% during the past 10 years. The prevalence of women who smoked during the last trimester of pregnancy was 13.0% in 2009, the lowest prevalence ever reported in the PNSS.

The 2009 PNSS report also indicated areas of concern, such as the 6% increase in low birthweight. The prevalence of iron-deficiency anemia during pregnancy also remains high.

Overweight is a major public health problem that has steadily increased in the United States, and this problem is mirrored in the PNSS population. The proportion of women who are overweight or obese before pregnancy has increased 13% since 2000, and nearly half of women now gain excess weight during pregnancy.

Recommendations

The PNSS data indicate that national and state public health programs are needed to support the following activities:

- Implement strategies to continue to reduce the prevalence of tobacco use among pregnant women and women of reproductive age.
- Promote and support breastfeeding through effective programs, medical care systems, work sites, and communities.
- Prevent preterm delivery and low birthweight by providing preconception nutrition, including iron supplementation.⁷
- Conduct outreach activities to promote early awareness of pregnancy status and encourage early entry into comprehensive prenatal care, including the WIC program.
- Provide information to prenatal participants, especially women who are overweight or obese before pregnancy, about the importance of appropriate weight gain during pregnancy and the health risks of excess weight gain and postpartum weight retention.

References

1. Martin JA, Hamilton BE, Sutton PD, Ventura SJ, Mathews MS, Kirmeyer S, et al. Births: final data for 2007. *National Vital Statistics Reports* 2010;58(24):1–86. Available at www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_24.pdf.
2. Doherty DA, Magaan EF, Francis J, Morrison JC, Newnham JP. Pre-pregnancy body mass index and pregnancy outcomes. *International Journal of Gynecology & Obstetrics* 2006;95(3):242–247.
3. Li R, Jewells S, Grummer-Strawn L. Maternal obesity and breast-feeding practices. *American Journal of Clinical Nutrition* 2003;77(4):931–936.
4. Rasmussen KM, Yaktine AL, eds.; Committee to Reexamine IOM Pregnancy Weight Guidelines; Institute of Medicine; National Research Council. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington, DC: National Academies Press; 2009.
5. Rooney BL, Schauburger CW. Excess pregnancy weight gain and long-term obesity: one decade later. *Obstetrics & Gynecology* 2002;100(2):245–252.
6. Centers for Disease Control and Prevention. Recommendations to prevent and control iron deficiency in the United States. *MMWR* 1998;47(RR-3):1–36.
7. Scholl, TO. Iron status during pregnancy: setting the stage for mother and infant. *American Journal of Clinical Nutrition* 2005;81(5):1218S–1222S.
8. U.S. Department of Health and Human Services. *Healthy People 2010*. Volume II. Washington, DC: U.S. Government Printing Office; 2000. Available at www.healthypeople.gov.
9. Zhu BP, Rolfs RT, Nangle BE, Horan JM. Effect of the interval between pregnancies on perinatal outcomes. *New England Journal of Medicine* 1999;340(8):589–594.
10. Conde-Agudelo A, Beliza'n JM. Maternal morbidity and mortality associated with interpregnancy interval: cross sectional study. *BMJ* 2000;321(7271):1255–1259.
11. Institute of Medicine. *WIC Nutrition Risk Criteria: A Scientific Assessment*. Washington, DC: National Academies Press; 1996.
12. Devaney B, Bilheimer L, Schore J. Medicaid costs and birth outcomes: the effects of prenatal WIC participation and the use of prenatal care. *Journal of Policy Analysis and Management* 1992;11(4): 573–592.
13. Centers for Disease Control and Prevention. *Health, United States, 2009: With Special Feature on Medical Technology*. Hyattsville, MD: U.S. Department of Health and Human Services; 2010.
14. Abrams B. Preventing low birthweight: does WIC work? A review of evaluations of the Special Supplemental Food Program for Women, Infants, and Children. *Annals of the New York Academy of Sciences* 1993;678:306–315.

15. Ahluwalia I, Hogan VK, Grummer-Strawn L, Colville WR, Peterson A. The effect of WIC participation on small-for-gestational-age births: Michigan, 1992. *American Journal of Public Health* 1998;88(9):1374–1377.
16. Bitler MP, Currie J. Does WIC work? The effects of WIC on pregnancy and birth outcomes. *Journal of Policy Analysis and Management* 2005;24(1):73–91.
17. Salihu HM, Aliyu MH, Pierre-Louis BJ, Alexander GR. Levels of excess infant deaths attributable to maternal smoking during pregnancy in the United States. *Maternal and Child Health Journal* 2003;7(4):219–227.
18. Anderson ME, Johnson DC, Batal HA. Sudden infant death syndrome and prenatal maternal smoking: rising attributed risk in the Back to Sleep era. *BMC Medicine* 2005;3:4.
19. U.S. Department of Health and Human Services. *Women and Smoking: A Report of the Surgeon General*. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2001.
20. Samet JM, Yoon SY, editors. *Women and the Tobacco Epidemic: Challenges for the 21st Century*. Canada: World Health Organization; 2001.
21. U.S. Department of Health and Human Services. *The Health Consequences of Smoking: A Report of the Surgeon General*. Washington, DC: U.S. Department of Health and Human Services; 2004.
22. U.S. Department of Health and Human Services. *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2006.
23. Hofhuis W, de Jongste JC, Merkus PJ. Adverse health effects of prenatal and postnatal tobacco smoke exposure on children. *Archives of Disease in Childhood* 2003;88(12):1086–1090.
24. Mathews TJ, MacDorman MF. Infant mortality statistics from the 2005 period linked birth/ infant death set. *National Vital Statistics Reports* 2008;57(2):1–32.
25. Goldenberg RL, Culhane JF. Low birth weight in the United States. *American Journal of Clinical Nutrition* 2007;85(2):584S–590S.
26. Hamilton BE, Martin JA, Ventura SJ. Births: preliminary data for 2009. *National Vital Statistics Reports* 2010;59(3):1–19. Available at www.cdc.gov/nchs/data/nvsr/nvsr59/nvsr59_03.pdf.
27. Jolly MC, Sebire NJ, Harris JP, Regan L, Robinson S. Risk factors for macrosomia and its clinical consequences: a study of 350,311 pregnancies. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 2003;11:9–14.
28. March of Dimes. Your Premature Baby. Available at www.marchofdimes.com/professionals/14332_1157.asp.
29. Ip S, Chung M, Raman G, Chew P, Magula N, DeVine D, et al. *Breastfeeding and Maternal and Infant Health Outcomes in Developed Countries*. Rockville, MD: Agency for Healthcare Research and Quality; 2007. Evidence Report/Technology Assessment No. 153. Available at www.ahrq.gov/downloads/pub/evidence/pdf/brfout/brfout.pdf.
30. Centers for Disease Control and Prevention. Breastfeeding Among U.S. Children Born 2000–2008, CDC National Immunization Survey. Available at www.cdc.gov/breastfeeding/data/NIS_data.

