



An algorithm for quantitatively estimating non-occupational pesticide exposure intensity for spouses in the Agricultural Health Study

Nicole C. Deziel^{1,2} · Laura E Beane Freeman² · Jane A. Hoppin³ · Kent Thomas⁴ · Catherine C. Lerro² · Rena R. Jones² · Cynthia J. Hines⁵ · Aaron Blair² · Barry I. Graubard² · Jay H. Lubin² · Dale P. Sandler⁶ · Honglei Chen^{6,7} · Gabriella Andreotti² · Michael C. Alavanja² · Melissa C. Friesen²

Received: 9 June 2018 / Accepted: 5 October 2018 / Published online: 30 October 2018
© Springer Nature America, Inc. 2018

Abstract

Residents of agricultural areas experience pesticide exposures from sources other than direct agricultural work. We developed a quantitative, active ingredient-specific algorithm for cumulative (adult, married lifetime) non-occupational pesticide exposure intensity for spouses of farmers who applied pesticides in the Agricultural Health Study (AHS). The algorithm addressed three exposure pathways: take-home, agricultural drift, and residential pesticide use. Pathway-specific equations combined (i) weights derived from previous meta-analyses of published pesticide exposure data and (ii) information from the questionnaire on frequency and duration of pesticide use by applicators, home proximity to treated fields, residential pesticide usage (e.g., termite treatments), and spouse's off-farm employment (proxy for time at home). The residential use equation also incorporated a published probability matrix that documented the likelihood active ingredients were used in home pest treatment products. We illustrate use of these equations by calculating exposure intensities for the insecticide chlorpyrifos and herbicide atrazine for 19,959 spouses. Non-zero estimates for ≥ 1 pathway were found for 78% and 77% of spouses for chlorpyrifos and atrazine, respectively. Variability in exposed spouses' intensity estimates was observed for both pesticides, with 75th to 25th percentile ratios ranging from 7.1 to 7.3 for take-home, 6.5 to 8.5 for drift, 2.4 to 2.8 for residential use, and 3.8 to 7.0 for the summed pathways. Take-home and drift estimates were highly correlated (≥ 0.98), but were not correlated with residential use (0.01–0.02). This algorithm represents an important advancement in quantifying non-occupational pesticide relative exposure differences and will facilitate improved etiologic analyses in the AHS spouses. The algorithm could be adapted to studies with similar information.

Electronic supplementary material The online version of this article (<https://doi.org/10.1038/s41370-018-0088-z>) contains supplementary material, which is available to authorized users.

✉ Nicole C. Deziel
nicole.deziel@yale.edu
Melissa C. Friesen
friesenmc@mail.nih.gov

- ¹ Yale School of Public Health, Yale University, New Haven, CT, USA
- ² Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA
- ³ Department of Biological Sciences, Center for Human Health and the Environment, North Carolina State University, Raleigh, NC, USA

Introduction

Pesticide exposure for people living in agricultural areas can occur via multiple pathways. These pathways may include occupational exposure from personal use of pesticides on

- ⁴ National Exposure Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, Durham, NC, USA
- ⁵ Division of Surveillance, Hazard Evaluation and Field Studies, National Institute for Occupational Safety and Health, Cincinnati, OH, USA
- ⁶ Department of Health and Human Services, Epidemiology Branch, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, Durham, NC, USA
- ⁷ Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI, USA

farms, take-home exposure from the transfer of pesticides from skin, clothes, and shoes of agricultural workers to the family home, agricultural drift exposure from living near fields or orchards where pesticides are applied, and residential use exposure from pesticide treatments of insects and weeds in and around the home [1]. Most epidemiologic studies of cancer risk from agricultural pesticide exposure have focused on a single pathway, i.e., the occupational pathway, which likely provides the largest contribution to total exposure. However, non-occupational pathways may be important contributors to pesticide exposure for residents in agricultural areas and need to be characterized to maximize power to detect associations. This may be particularly important for family members of occupational pesticide users, who may not experience direct occupational pesticide exposures, or may experience these exposures at a lower frequency, duration, or magnitude.

In epidemiologic studies, non-occupational pesticide exposure in agricultural areas has typically been characterized using surrogate exposure metrics from participant-reported information. These metrics usually account for only one exposure pathway and have limited information on specific chemicals or on the frequency and duration of exposure [2, 3]. Surrogates used to characterize non-occupational exposures include duration of living on a farm [4, 5], proximity to treated fields [6], treatment of residential pests [7], or laundering pesticide-laden clothing [4, 8]. Other studies have used geographic information system-based methods that incorporated crop maps or satellite imagery to calculate the density of crops near a residence to predict possible pesticide exposures [7, 9–11], although such data may only be intermittently available or may lack the spatial resolution needed to identify individual crop characteristics [12]. Some studies have also used biomarkers of exposure, which are aggregate measures of pesticide exposure from all pathways [13–16]. Due to the short half-lives of many pesticide biomarkers, they rarely represent the long-term exposure relevant for cancer or other chronic disease evaluations [17].

An alternative approach is to use a transparent decision rule-based framework that links subject-reported information to quantitative measures, which has been used in several studies to estimate historical occupational exposures [18, 19]. For example, an occupational pesticide exposure algorithm, previously developed for the prospective cohort of >52,000 licensed pesticide applicators in the Agricultural Health Study (AHS), combines subject-reported information on the frequency of mixing pesticides, application methods, and personal protective equipment usage with weighting factors derived from the literature, expert judgment, and field evaluation studies [20–22]. The occupational algorithm-based intensity estimates have

demonstrated good correlation with exposure monitoring data [23, 24]. These intensity estimates have been combined with frequency and duration of pesticide use to obtain cumulative exposure estimates for epidemiologic analyses.

Here, we used a decision rule-based approach to develop a deterministic algorithm that quantitatively estimates cumulative (i.e., over adult married life) exposure intensity for a specific active ingredient from three non-occupational pesticide exposure pathways for spouses of the AHS pesticide applicators. The “AHS spouses” comprise 32,345 spouses (99% female) of pesticide applicators who sought private, restricted-use pesticide licenses in Iowa and North Carolina in 1993–97. These spouses reported diverse levels of participation in farming activities, with ~50% not engaged in any pesticide application-related activities at enrollment [25, 26]. They provide a unique population in which to study the impact of pesticide exposure on chronic disease risks, including female-specific outcomes, at pesticide exposure levels anticipated to be at the high end of the non-occupationally exposed population [27, 28]. In addition, this group will allow follow-up on suggestive findings in the female AHS applicators ($n = 1563$) within a much larger population of exposed; e.g., the observation of a suggested increased risk of ovarian cancer among female applicators exposed to atrazine [29].

The non-occupational spouse pesticide algorithm described here provides quantitative estimates of cumulative pesticide exposure intensity for the take-home, agricultural drift, and residential pesticide use pathways for single-pesticide active ingredients. These pathways were selected based on evidence from the literature that higher pesticide concentrations in house dust were associated with farm work done by one or more home residents (take-home), shorter distances between homes and treated fields (drift), and greater home use of pesticides to treat pests in and around the home (residential use) [1]. Other exposure pathways, though plausible, were not consistently supported by the literature (e.g., dietary ingestion). The non-occupational spouse algorithm described here uses a similar approach to the above-mentioned applicator occupational pesticide algorithm [20, 21]; however, it directly incorporates frequency and duration of pesticide applications into pathway-specific equations to estimate cumulative exposure and represents multiple pathways rather than a single pathway. In this paper, we present the derivation of the non-occupational algorithm. To illustrate its application, we calculated pathway-specific intensity estimates separately for the insecticide chlorpyrifos and herbicide atrazine in a subset of AHS spouses. Chlorpyrifos and atrazine were selected because they were commonly measured in the published pesticide house dust studies and were frequently used in the AHS cohort [1, 30].




Fig. 1 Conceptual framework for non-occupational pesticide exposure algorithm

Methods: algorithm development and example calculations

Algorithm overview

To estimate spouse-specific, cumulative (adult, married life) non-occupational exposure intensity for a single-active ingredient (ai) ($E_{\text{non-occ,ai}}$), we developed active ingredient (ai)-specific equations for each of the take-home ($E_{\text{take-home,ai}}$), agricultural drift ($E_{\text{drift,ai}}$), and residential pesticide use ($E_{\text{res,ai}}$) pathways in units of intensity-weighted years. The overall framework is shown in Fig. 1 and each pathway is described in detail below. These pathway equations can be used singly or combined depending on the plausibility of their contributions for a given active ingredient. For example, the residential use pathway can be dropped for those active ingredients never licensed for residential use. If all pathways are deemed relevant and assumed to have equal weight, the pathway-specific estimates may be summed to obtain an overall cumulative pesticide exposure intensity across the three pathways (Eq. (1)).

$$E_{\text{non-occ,ai}} = E_{\text{take-home,ai}} + E_{\text{drift,ai}} + E_{\text{res,ai}} \quad (1)$$

Each of the equations were based on our findings from a qualitative literature review that identified factors predicting relative differences in pesticide exposure among households and individuals in agricultural areas [1]. We used pesticide concentrations in house dust as surrogates for non-occupational exposures, because house dust measurements integrate pesticide levels over months or years [31, 32] and exhibited sufficient variability and detection rates among agricultural households in published studies. We abstracted published dust concentrations across a range of pesticide active ingredients and used meta-regression models to quantify the average relative contributions of the pathways

to pesticide concentrations in house dust [30] to provide the pathway weights. We used individual-level information from responses to AHS questionnaires administered to the spouses and applicators to identify exposure differences among the AHS spouses. The questionnaires provided active ingredient-specific information, including duration and frequency of agricultural use of each active ingredient by the applicator, as well as information on household characteristics (e.g., distance between the home and nearest treated fields; home pest treatment practices), and the spouse's off-farm employment as a surrogate for estimating time spent at home (<http://aghealth.nih.gov/collaboration/questionnaires.html>).

For illustrative purposes, we present descriptive statistics in the Results section for the calculated pathway-specific exposure intensities for chlorpyrifos and atrazine focusing on pesticide use data collected at enrollment (1993–1997). We also required information related to cohabitation between spouses and partners, which was collected later in the study (2005–2010). We present these results for the subset of AHS spouses with complete information on cohabitation ($n = 19,064$ for chlorpyrifos, $n = 19,077$ for atrazine), which represents 60% of the total spouse cohort. Current efforts to reconstruct the cohabitation data for all spouses will facilitate future application of the algorithm to the full spouse cohort.

Duration metrics

Duration metrics obtained from AHS questionnaires were used in all pathways to estimate cumulative exposure over adult, married life up to time of AHS enrollment. The take-home and agricultural drift pathways incorporated the applicator's number of days (Days Applicator Applied_{ai}) per year and years of active ingredient use while living with the

spouse (Years Applicator Applied While Together_{ai}). We did not include the spouse's application days to avoid double-counting days when both spouse and applicator applied pesticides and to avoid overlap with the occupational pathway. Because the pathway-specific weights represent the average contribution aggregated over a long time span rather than the contribution per pesticide application day, we standardized Days Applicator Applied_{ai} by dividing by the median number of days per year of any pesticide application reported across all applicators with at least one day of use at enrollment (Median Application Days = 14.5), which reflected typical usage over the course of a substantial proportion of a farmer's working lifetime. Years Applicator Applied While Together_{ai} was derived from the applicator's reported start and stop years of active ingredient use as well as the start and stop years of cohabitation, assuming continuous cohabitation. The residential pesticide use exposure pathway incorporated the number of years the spouse and applicator cohabitated (Years Together).

To illustrate the calculation of the algorithm, we present a hypothetical female spouse who lived with her husband for 18 years at enrollment (Years Together). Her husband applied both chlorpyrifos and atrazine for 15.5 of the 18 years in which they lived together (Years Applicator Applied While Together_{chlorpyrifos} = Years Applicator Applied While Together_{atrazine} = 15.5 y). Her husband applied chlorpyrifos for "5–9 days in an average year" (Days Applicator Applied_{chlorpyrifos} = 7 d/y) and atrazine "10–19 days" (Days Applicator Applied_{atrazine} = 14.5 d/y).

Take-home exposure pathway

The take-home exposure pathway captured pesticide exposure from the transfer of pesticides into the home from the skin, clothes, and shoes of applicators. Evidence that take-home exposure influenced pesticide concentrations in house dust came primarily from studies that identified higher concentrations of pesticides in house dust in homes of farmers who applied pesticides more frequently or recently compared to homes of farmers who did not apply pesticides or applied them less frequently or recently [1]. Thus, for each active ingredient, we estimated the spouse's take-home exposure intensity ($E_{\text{take-home,ai}}$) as proportional to the spouse's average daily hours spent at the home divided by 24 (Hours per Day Spouse at Home/24 h), the Days Applicator Applied_{ai}, and the Years Applicator Applied While Together_{ai}, with weight $k_{\text{take-home}}$ (Eq. (2)). We derived $k_{\text{take-home}}$ from our previous meta-analysis of house dust pesticide concentrations that found that homes of farmers with more frequent or recent pesticide application had 2.3 (95% CI: 1.5–3.3) times higher house dust pesticide concentrations than those of farmers with less frequent or recent

pesticide use, based on 15 summary statistics reported in 5 studies [30]. Because the number of hours at home was not queried in the AHS questionnaires, Hours per Day Spouse at Home was based on the spouses' self-reported information about their longest-held job and employment-specific patterns of proportion of hours per day spent at home from the 2003 American Time Use Survey of women living in non-metropolitan areas (the earliest year of data available) [33]. Spouses who had a year-round off-farm job ("full time"), spouses who had a part-time job or who had a part-year full-time job ("part-time"), and spouses who worked neither part-time nor full-time outside the home ("no off-farm job") were assigned the survey's averages of 15.6, 17.8, and 21.0 h/day at home, respectively. While we had data available on other behaviors typically used to characterize "take-home exposure", i.e., laundering pesticide-laden clothing or whether the applicator removed shoes/clothing prior to entering the house, we did not include these behaviors as modifiers at this time because our previously conducted literature review did not find strong support that these behavioral factors modified spouse's exposure to pesticides [1].

$$E_{\text{take-home,ai}} = k_{\text{take-home}} \times [\text{Hours per Day Spouse at Home}/24 \text{ Hours per Day}] \times [\text{Days Applicator Applied}_{\text{ai}}/\text{Median Application Days}] \times \text{Years Applied While Together}_{\text{ai}} \quad (2)$$

Therefore, if the hypothetical spouse described above never had an off-farm job (Hours per Day Spouse at Home = 21 h), her cumulative take-home exposure for chlorpyrifos would be 15 intensity-weighted years (calculation provided below); for atrazine it would be 31 intensity-weighted years (not shown).

$$E_{\text{take-home,chlorpyrifos}} = (2.3 \times \frac{21}{24}) \times \frac{7}{14.5} \times 15.5 = 15 \text{ intensity-weighted years}$$

Agricultural drift

The agricultural drift pathway captured pesticide exposure from the airborne transport of pesticides from the site of pesticide application to the home (rather than direct transfer from applicator). Evidence that agricultural drift influenced house dust pesticide concentrations came predominantly from studies that examined these concentrations at varying distances of the home to treated fields [1]. Several studies also found that incorporating the quantity (mass) or density (mass per unit area) of active ingredients applied near the home yielded stronger associations with dust pesticide concentrations compared to proximity alone [11, 34, 35]. In

Table 1 Agricultural drift exposure: data-derived weighting factors based on the distance between residence and treated fields

Questionnaire response categories ^a	Distance (ft) ^b	Predicted GM concentration (µg/g) ^c	k_{drift} (95% CI) ^{d,e}
<100 yd	150	0.13	3.0 (1.7, 5.4)
100–199 yards	450	0.08	1.9 (1.4, 2.6)
200–299 yards	750	0.07	1.5 (1.2, 1.9)
>300 yards or more	1350	0.05	1.2 (1.1, 1.3)
>1/4 mile (Ref) ^f	1980	0.044	1.0
No pesticides applied on farm			1.0

^aQuestionnaire item: “How far is your home from the nearest field or orchard where pesticides are applied?” Response category units reflect those on the questionnaire (English units)

^bDistance was assigned the mid-point of the range of the self-reported category or 1.5 times the lower bound. Note that 100 ft = 30.5 m.

^cCalculated using the formula Predicted GM = distance^{-0.43} e^{0.15}, from Eq. (13) in ref. [30]

^dRelative increase in GM pesticide concentrations relative to the reference group (GM_{Distance}/GM_{Ref})

^eConfidence intervals calculated using formula Lower CL = exp(-0.21 * ln(distance/distance_{Ref})) and Upper CL = exp(-0.65 * ln(distance/distance_{Ref})) based on confidence limits on the LogDistance regression parameter from Table 4 in ref. [30]

^fReference category of 1980 ft was based on 1.5 times a quarter mile, the farthest distance category queried in a follow-up questionnaire (Phase 2) of the AHS

the AHS, however, spouses reported a categorical distance from the home to the nearest treated field (in English units), and we lacked information on amount applied, application rate, and formulation used with which to derive metrics of quantity or density of applied active ingredients. Additionally, the question about distance to treated fields was not linked to the applicator’s pesticide use, so we assumed the applicator used the active ingredient at the fields nearest to the home and we were unable to account for drift occurring from pesticide usage on other nearby farms.

We estimated the spouse’s exposure from agricultural drift ($E_{\text{drift,ai}}$) to be proportionally related to k_{drift} , the Days Applicator Applied_{ai}, and the Years Applicator Applied While Together_{ai} (Eq. (3)). We derived k_{drift} from our previous meta-regression analysis of the geometric mean (GM) dust pesticide concentrations at varying distances in ft (d) of the house from the fields for each distance response category (Table 1). For example, dust pesticide concentrations were predicted to be 3.0 times higher in homes < 100 yd (< 91 m) from treated fields, compared to homes >1/4 mile away (>402 m).

$$E_{\text{drift,ai}} = k_{\text{drift}} \times (\text{Days Applicator Applied}_{\text{ai}} / \text{Median Application Days}) \times \text{Years Applied While Together}_{\text{ai}} \quad (3)$$

For example, if our hypothetical spouse reported living “100–199 yards” (91 to 182 m) within treated fields, her k_{drift} value would be 1.9 (Table 1). Her cumulative agricultural drift estimate would be 14 intensity-weighted years for chlorpyrifos (calculation below) and 38 intensity-

weighted years for atrazine.

$$E_{\text{drift,chlorpyrifos}} = (1.9 \times \left[\frac{7}{14.5}\right] \times 15.5) = 14 \text{ intensity-weighted years}$$

Residential pesticide use

The residential pesticide use pathway captured pesticide exposure that occurred from treating the home or pets for insects and the lawn, yard, or garden (hereafter, “lawn”) for weeds. Evidence that residential pesticide use contributed to pesticide concentrations in house dust came primarily from findings of higher concentrations or detection rates in rural/agricultural households reporting treatment for specific home and lawn pests (e.g., weeds, fleas/ticks) compared to households that did not treat for those particular pests [1, 36, 37]. In these published studies, homeowners reported the type of pest treatment but did not provide specific active ingredients. Similarly, the AHS questionnaires queried spouses about whether household members or professionals performed several residential pest treatments (identified with subscript trt): termites, non-termite insects, flea/tick treatments applied in the home, flea/tick treatment directly to pets, and lawn weeds; these questions were not active ingredient-specific.

To obtain the likelihood of exposure to specific active ingredients for each treatment, we used a previously developed pesticide exposure probability matrix that synthesized market data on sales of commercial pest treatment products and active ingredients to estimate the probability

that an active ingredient was used for different pest treatment scenario across various time frames (1976, 1980, 1990, and 2000) [38] (<http://dceg.cancer.gov/tools/design/pesticide>). Using this pesticide probability matrix, we mapped the spouses' questionnaire responses about the types of pest treatments and who applied them (household member, professional applicator, or other/do not know) to one or more of the matrix's pest treatment scenarios as described in Supplemental Table S1. Calculated probabilities by active ingredient are provided in Supplemental Table S2. We averaged the probabilities from 1976, 1980, and 1990 to derive the probabilities for treatments reported at enrollment.

We estimated that the residential use contribution to house dust concentrations for each pesticide active ingredient, $E_{res,ai}$, was proportional to whether the treatment occurred ($Treated_{tr}$) and the Years Together, with weight k_{res} . The value for k_{res} was either 1.3 (95% CI: 1.1–1.6) or 1.5 (95% CI: 1.2–1.9) if the probability of use for the pest treatment scenario from Supplemental Table S2 was $\geq 1\%$ to $<20\%$ or $\geq 20\%$, respectively, based on our previous meta-regression analysis comparing treated versus untreated homes (88 statistics, 5 studies) [30]. Years Together was used because the frequency and duration of most of these treatments were not directly queried in the AHS questionnaire. The residential use contribution was summed across pest treatment types (Eq. (4)).

$$E_{res,ai} = \left(\begin{array}{l} k_{termite,ai} \times Treated_{termite} + k_{insect,ai} \times \\ Treated_{insect} + k_{fleas\ home,ai} \times Treated_{fleas\ home} + \\ k_{fleas\ pets,ai} \times Treated_{fleas\ pets} + k_{weeds,ai} \times \\ Treated_{weeds} \end{array} \right) \times \text{Years Together} \quad (4)$$

For example, our hypothetical spouse reported that her home was treated for termites ($Treated_{termite} = 1$), treated for non-termite insects by a professional ($Treated_{insect} = 1$) and that her lawn was treated for weeds by a household member ($Treated_{weeds} = 1$). She also reported that a flea/tick shampoo was used to treat a pet ($Treated_{fleas\ pets} = 1$), but that her home was never treated for fleas/ticks by a fumigant/fogger bomb product ($Treated_{fleas\ home} = 0$). The probability that chlorpyrifos was used in a termite treatment at enrollment was 0.24 (Supplemental Table S2), yielding a value of $k_{termite,chlorpyrifos}$ of 1.5. Similarly, based on probabilities of use, $k_{insect,chlorpyrifos}$, $k_{fleas\ home,chlorpyrifos}$, $k_{fleas\ pets,chlorpyrifos} = 1.3$; these values were zero for atrazine because it is not an insecticide. For weed treatment by a household member, the pesticide exposure matrix yielded a probability of zero for both chlorpyrifos and atrazine ($k_{weeds,chlorpyrifos} = k_{weeds,atrazine} = 0$). Chlorpyrifos is an insecticide and therefore not used on weeds and atrazine

is not commonly found in residential lawn products [38]; its residential use is generally confined to Florida and the Southeast [39]. Her cumulative residential pesticide use exposure estimate would be 74 and 0 intensity-weighted years for chlorpyrifos (calculation below) and atrazine, respectively.

$$E_{res,chlorpyrifos} = (1.5 \times 1 + 1.3 \times 1 + 1.3 \times 0 + 1.3 \times 1 + 0 \times 1) \times 18 = 74 \text{ intensity-weighted years}$$

Results

Prevalence of use and duration metrics

A substantial proportion of spouses experienced exposures through the take-home and drift pathways due to the applicator's use, with 44% and 76% of applicators reporting usage of chlorpyrifos and atrazine, respectively (Table 2). The duration of use among those applying was shorter and spanned a narrower range for chlorpyrifos with a median of 3.5 y (inter-quartile range [IQR]: 3.5–8.0) compared to 8.0 y for atrazine (IQR: 3.5–25.5). Median application frequency among users was 7.0 d/y for both active ingredients, with a narrower IQR reported for chlorpyrifos (2.5–7.0) compared to atrazine (2.5–14.5).

Take-home exposure pathway

Most spouses (73%) reported long-term, full-time off-farm jobs (and therefore were estimated to spend less time home), while 18% held part-time off-farm jobs, and 9% never held an off-farm job. The resulting median (IQR) cumulative exposure intensity estimates for the take-home pathway among those exposed was 2.5 (0.9–6.6) intensity-weighted years for chlorpyrifos and 5.8 (2.1–15) intensity-weighted years for atrazine, demonstrating a sevenfold ratio between the 75th and 25th percentiles for both active ingredients. The median intensity-weighted take-home estimate for exposed spouses was over two times higher for atrazine than for chlorpyrifos (5.8 vs. 2.5, respectively).

Agricultural drift

The majority of spouses reported living within 200 yards of treated fields, with 47% of spouses assigned a distance between a home and treated field (d) of 150 ft (46 m), and 20% assigned 450 ft (137 m) (Table 2). Only 4% lived in the farthest category (1950 ft or 594 m). The median (IQR) cumulative exposure intensity estimates for the agricultural drift pathway among those exposed were 4.1 (1.7–11) for chlorpyrifos and 8.0 (2.6–22) for atrazine (Table 2), demonstrating ratios for the 75th to 25th percentiles of 6.5

Table 2 Distributions of subject-specific pathway components and cumulative estimates of non-occupational exposure intensity for atrazine and chlorpyrifos for AHS spouses with complete information

Pathway Components	Chlorpyrifos (Insecticide) (<i>N</i> =19,064) ^a <i>N</i> (%)	Atrazine (Herbicide) (<i>N</i> =19,077) ^a <i>N</i> (%)
Applicator-husband applied active ingredient		
No	10,762 (56%)	4596 (24%)
Yes	8302 (44%)	14,481 (76%)
Take-home		
Hours per Day Spouse at Home		
Full-time off-farm job (15.6 h)	13,920 (73%)	13,923 (73%)
Part-time off-farm job (17.8 h)	3345 (18%)	3353 (18%)
No off-farm job (21.0 h)	1799 (9%)	1801 (9%)
Drift		
Distance (ft) between house and fields (d) [corresponding k_{drift} value] ^{b,c}		
150 [3]	8879 (47%)	8907 (47%)
450 [1.9]	3832 (20%)	3829 (20%)
750 [1.5]	1308 (7%)	1302 (7%)
1350 [1.2]	4216 (22%)	4215 (22%)
1950 [1.0]	829 (4%)	824 (4%)
Residential treatments		
Treated _{termintes} = Yes	2946 (15%)	2944 (15%)
Treated _{insects} = Yes	9814 (51%)	9843 (52%)
Treated _{fleas home} = Yes	629 (3%)	632 (3%)
Treated _{fleas pet} = Yes	7484 (39%)	7495 (39%)
Treated _{weeds} = Yes	6978 (37%)	6999 (37%)
Exposed via residential use pathway (>0 probability of active ingredient)	11,999 (63%)	881 (5%)
	Median (IQR), Exposed spouses	Median (IQR), Exposed spouses
Duration Metrics, exposed subjects^b		
Years Applicator Applied While Together _{ai} (y)	3.5 (3.5–8.0)	8.0 (3.5–25.5)
Days Applicator Applied _{ai} (d/y)	7.0 (2.5–7.0)	7.0 (2.5–14.5)
Pathway estimates, exposed subjects		
$E_{take-home,ai}$ (intensity-weighted y) ^d	2.5 (0.9–6.6)	5.8 (2.1–15)
$E_{drift,ai}$ ^e	4.1 (1.7–11)	8.0 (2.6–22)
$E_{res,ai}$ ^{f,g}	40 (27–75)	33 (19–45)
$E_{non-occ,ai}$ ^h	39 (19–73)	16 (5.4–38)

^aNumber of spouses with complete information for an active ingredient. See Methods section for inclusion criteria/how “complete information” defined

^bAll reported statistics based on those exposed to the active ingredient

^c k_{drift} values obtained from Table 1

^dEquation (2)

^eEquation (3)

^fTreatment and active ingredient-specific weights provided in Supplemental Table S1 and S2

^gEquation (4)

^hEquation (1)

Algorithm inputs with constant values: Median Application Days = 14.5 d/y, $k_{take-home} = 2.3$; $k_{termintes, chlorpyrifos} = 1.5$, $k_{insects, chlorpyrifos}$, $k_{fleas home, chlorpyrifos}$, $k_{fleas pets, chlorpyrifos} = 1.3$, $k_{weeds, chlorpyrifos} = 0$; $k_{termintes, atrazine} = 1.5$, $k_{insects, atrazine}$, $k_{fleas home, atrazine}$, $k_{fleas pets, atrazine} = 1.3$, $k_{weeds, atrazine} = 0$

and 8.5, respectively. The median intensity-weighted take-home estimate for exposed spouses was two times higher for atrazine than for chlorpyrifos (8.0 vs. 4.1).

Residential pesticide use

Residential pest treatments were common, with treatment of non-termite insects (52%) being the most prevalent, followed by fleas on pets (39%), weeds (37%), termites (15%), and fleas in the home (3%) (Table 2). Chlorpyrifos is an insecticide that was commonly used at enrollment. All three types of insect treatments had a $\geq 1\%$ probability of containing chlorpyrifos (Supplemental Table S2), and therefore all contributed to the estimated exposure estimate among spouses reporting the given treatment. The k_{termites} value was 1.5, k_{insects} was 1.3, $k_{\text{fleas home}}$ was 1.3, and $k_{\text{fleas pets}}$ was 1.3. As with all insecticides, the k_{weeds} for chlorpyrifos was 0 (Supplemental Table S2). Of the 19,064 spouses with complete information on chlorpyrifos, 11,999 (63%) were estimated to be exposed via the residential pesticide use pathway (i.e., $E_{\text{res}} > 0$). Among those exposed, the median (IQR) exposure intensity estimate was 40 (27–75), yielding a ratio of 75th to 25th percentiles of 2.8.

Atrazine is an herbicide, and thus only weed treatments had the potential to contribute to atrazine's residential pesticide use exposure intensity estimate. Because atrazine had low probability of use in any consumer products, only spouses reporting treatments by "professionals" or "both household members and professionals" or an "other" or "unknown" person had a $\geq 1\%$ probability and contributed to the residential pesticide use exposure intensity (Supplemental Table S2). Of the 6999 spouses reporting a weed treatment, only 881 were estimated to be exposed to atrazine via the residential pesticide use pathway (5% of the population with complete atrazine data) (Table 2). Among those exposed, the median (IQR) was 33 (19–45) intensity-weighted years, yielding a ratio of 75th to 25th percentiles of 2.4. The median and IQR intensity-weighted residential use estimate for exposed spouses were higher and broader for chlorpyrifos vs. atrazine (median 40 vs. 33, respectively; IQR 27–75 vs. 19–45, respectively) (Table 2).

Integration of the three pathways

Total non-occupational exposure intensity ($E_{\text{non-occ,ai}}$) was calculated by summing each pathway, assuming the resulting exposure from the three pathways is additive. While the proportion of spouses with non-zero estimates for total non-occupational exposure intensity were nearly identical (78% and 77% of the population for chlorpyrifos and atrazine, respectively), the prevalence and magnitude of exposure intensity for the specific pathways differed for these two active ingredients. Among those with $E_{\text{non-occ}} > 0$,

Table 3 Correlations among pathway-specific exposure intensity estimates for chlorpyrifos and atrazine

	Spearman correlation coefficients			
	Take-home	Agricultural drift	Residential use	Total non-occupational
Chlorpyrifos, spouses with $E_{\text{non-occ}} > 0$ ($n = 19,064$)				
Take-home	1.0	0.99*	0.02	0.31*
Agricultural drift		1	0.01	0.31*
Residential use			1	0.90*
Atrazine, spouses with $E_{\text{non-occ}} > 0$ ($n = 19,077$)				
Take-home	1	0.98*	0.02	0.96*
Agricultural drift		1	0.02	0.96*
Residential use			1	0.24*

* $p < 0.001$

the median (IQR) was 39 (19–73) for chlorpyrifos and 16 (5.4–38) for atrazine.

The Spearman correlations between pathways among spouses with total, non-occupational exposure > 0 are shown in Table 3. For chlorpyrifos, the take-home and drift exposure intensity estimates were almost perfectly positively correlated with each other ($r_{\text{Spearman}} = 0.99$), but each was not correlated with the residential use pathway ($r_{\text{Spearman}} = 0.02$ for take-home and $r_{\text{Spearman}} = 0.01$ for drift) (Table 3). Neither the take-home nor drift pathways were strongly correlated with the total, non-occupational exposure for chlorpyrifos ($r_{\text{Spearman}} = 0.31$ for each), whereas the residential pesticide use pathway was highly correlated with the total non-occupational exposure intensity ($r_{\text{Spearman}} = 0.90$). For atrazine, the take-home and drift exposure intensity estimates were also highly correlated ($r_{\text{Spearman}} = 0.98$) and were highly correlated with the total, non-occupational exposure intensity ($r_{\text{Spearman}} = 0.96$ for each) (Table 3). For atrazine, the residential pesticide use pathway was not correlated with either the take-home or drift pathways ($r_{\text{Spearman}} = 0.02$ for each), and was only weakly correlated with total non-occupational exposure intensity ($r_{\text{Spearman}} = 0.24$).

Discussion

We developed a deterministic algorithm based on subject-level information and quantitative synthesis of published exposure data to assign quantitative, cumulative married lifetime estimates of pesticide exposure intensity from three non-occupational exposure pathways. This algorithm fills an important gap that provides for better identification of

differences in cumulative pesticide exposure intensity among the AHS spouses, a population that presumably experiences higher non-occupational exposures than the general population because of their proximity to agricultural pesticide uses. Application of this algorithm to the AHS spouses demonstrated a wide distribution of cumulative exposure intensity estimates for two active ingredients, supporting use of the algorithm in future etiologic analyses, potentially advancing our understanding of non-occupational pesticide exposure on chronic health outcomes. Our systematic approach and the transparency of our assumptions will allow the algorithm to be revised as new information becomes available and will allow for sensitivity analyses that examine the impact of our assumptions on exposure–response relationships.

Development of the algorithm to refine exposure intensity estimates for AHS spouses beyond previously used simpler metrics, such as ever/never use of a specific pesticide ([26]; Lerro et al. 2015; [40, 41]) or whether the applicator applied a specific pesticide [41, 42], required several assumptions necessitated by the study design, the types of information collected in the AHS, and the availability of published data, which we describe below. We strove to align the information from the AHS questionnaire with the evidence in the literature. Although the development of this algorithm often involved assumptions or imperfect proxies, we expect that the algorithm's use will reduce exposure misclassification over previously used metrics and will improve our ability to evaluate risk in this population. Although exposure misclassification will remain a concern, it is expected to be non-differential, because the information from the spouses and applicators was largely obtained prior to diagnosis of diseases.

Pathway-specific, questionnaire-based exposure surrogates

The take-home pathway described here takes into account the over two times increase in house dust pesticide concentrations reported in the literature in homes of farmers applying pesticides and the AHS applicators' pesticide use patterns, with a modifier that represented the AHS spouses' time spent in or around the home, because increased time at home increases the likelihood or amount of contact with pesticides that enter the home from the applicator's activities. This time modifier was based on the spouse's longest-held off-farm job due to the lack of a full occupational history. It was also based on the 2003 American Time Use Survey of women in non-metropolitan areas, which we assumed was representative of the predominantly rural population and earlier years.

The agricultural drift pathway takes into account the observed exponential decrease in house dust pesticide

concentrations with increasing distance from treated fields and the applicators' pesticide use patterns. We assumed that self-reported distance to fields was relevant to all active ingredients used by the applicator, as pesticide-specific distance was not queried. In addition, we assumed that the nearest treated field was the applicator's field, because the distance question did not distinguish between the applicators' fields and their neighbors' fields. A more accurate measure of distance to fields could in future efforts use geocoded residential addresses and state crop maps that could better account for pesticide drift from nearby farms. Application quantity, application methods, pesticide formulations, or acreage of crops treated may be more accurate predictors of pesticide concentrations in house dust [34]; however, adding these refinements would require re-contacting applicators and obtaining additional information. Meteorological conditions (e.g., wind direction, wind speed, temperature) are important factors predicting agricultural drift, but accurately incorporating these factors would require historical data on meteorological conditions in conjunction with knowledge about where and when pesticides were applied in relation to the home, which is not available for this population.

The residential use pathway accounted for the observed increased pesticide concentration in house dust from multiple pest treatment types. Although information on active ingredients in these treatments was not collected in the AHS questionnaires, an active ingredient-specific estimate was made possible by using the NCI pesticide exposure probability matrix to link treatment types to probable active ingredients. The nation-wide population-based weights from the pesticide matrix applied to these agricultural residents may overestimate or underestimate exposure to specific active ingredients at the individual level. Differences in residential pesticide exposure intensities based on whether the spouse or another household member performed the treatment was only partially taken into account with applicator-specific (e.g., household member vs. commercial applicator) probabilities of use in the pesticide exposure matrix of a specific active ingredient. We did not have consistent, detailed information on the frequency and duration of residential pest treatments and therefore assumed the reported treatment was representative of usual usage patterns over the years the spouse and applicator lived together. Our approach averages probabilities and treatment practices over time and therefore may not capture changes in formulations due to economic changes or regulatory actions. In addition, this approach may not capture a pesticide if its usage does not fit into one of the survey categories. Finally, the weights derived for this pathway focused on patterns observed in rural populations, which may not be applicable to urban settings. The Years Together component of the residential use pathway will require

customization for each active ingredient based on when it was approved for in-home use. For instance, many organochlorine pesticides were banned in the United States in the 1970s and 1980s and therefore, the Years Together variable should be truncated to the year the use of a specific organochlorine active ingredient of interest was restricted. The impacts of the assumptions made for the residential use pathway and its contributions to cancer risk can be examined in sensitivity analyses.

Literature-based pathway-specific weights

The framework's pathway weights were data-driven, with confidence intervals, and were based on a novel application of meta-regression models to synthesize published environmental data to capture broad contrasts in relative contributions [30]. An additional advantage of this approach is that the distribution of pathway weights could be used to estimate uncertainties in the exposure assessment. The weights reflect summary effects across a range of different pesticide compounds, thereby increasing the generalizability of the equations. However, these data-driven weights did not identify finer nuances in exposure intensity because, at this time, data were too sparse to confirm quantitative differences in pathway weights for various subgroups, such as pesticide type, active ingredient, formulation type, crop type, application method, geographic location, or time period. Some subgroup-specific differences were previously identified, but were based on small numbers [30]. For example, the relative increase in house dust pesticide concentrations in homes of farmers with more frequent vs. less frequent pesticide contact was higher for atrazine compared to chlorpyrifos, though the difference was not statistically significant. Also, the decrease in house dust pesticide concentrations with increasing distance between homes and treated fields was greater for herbicides compared to insecticides. The weights might be able to be adjusted for specific chemicals based on their unique physicochemical properties, such as their transport potential or persistence. The transparency of this approach allows for weights to be recalculated and sensitivity analyses repeated if new data become available.

The pathway weights were based on an important assumption that pesticide concentrations in house dust serve as a reasonable proxy for adult exposure. Although dust pesticide concentrations are a potentially useful exposure indicator in children [43], their relevance to adult exposure is less established [44]. We used house dust pesticide concentrations as a surrogate for chronic exposure because pesticides in indoor dust resist degradation due to limited sunlight, microbial activity, and moisture (Simcox et al. 1995; [45, 32]). Biological monitoring data from spouses/

women in agricultural areas, a more direct measure of exposure, was not used here to predict exposure contrast, because pesticide biomarkers tend to have low percent detection and limited variability within study populations, and generally reflect recent exposures [1, 16, 46, 47]. More research is needed on the relevance of house dust for predicting cumulative, adult exposure to pesticides in agricultural populations. Another challenge with dust pesticide measurements is that they integrate the three exposure pathways. We strove to disentangle the relative contribution or weight of each of the non-occupational pathway wherever possible by basing the data-driven weights on published exposure estimates that adjusted for the other exposure pathways [30].

Application of algorithm and correlations among pathways

Application of our algorithm to two pesticides illustrated some important pathway-specific distinctions in estimated exposure intensity by active ingredients. For atrazine, the take-home and drift pathways were dominant, whereas the residential pesticide use pathway was diminutive. In contrast, for chlorpyrifos, the residential pesticide use pathway was a greater contributor to cumulative, non-occupational exposure than the pathways related to the applicator's pesticide use.

The take-home and agricultural drift pathways were highly correlated with each other for both active ingredients (≥ 0.98), due to the common equation components related to the applicators' duration and frequency of pesticide use. Though users of the algorithm may retain only one of these pathways in the algorithm, in this analysis we chose fidelity to the exposure literature, which suggests an independent contribution of each pathway, versus parsimony and simplicity. The impacts of this choice can be examined in sensitivity analyses. New evidence may provide improved information for disentangling these two pathways.

Omitted pathways and time periods

The algorithm captures the three exposure pathways with the strongest empirical evidence. However, at this time the algorithm does not capture all potential non-occupational exposure sources or time periods for the spouses and is expected to non-differentially underestimate non-occupational cumulative exposure intensity. Possibly the most important underestimation of exposure occurs from our inability to characterize exposure for the time window prior to the spouse's marriage to the applicator, because active ingredient-specific information for this earlier time period was not included in the AHS questionnaires. This

may be important for the 58% of the AHS spouses who reported living most of their life on a farm prior to age 18 [48], particularly given the mounting evidence that early-life exposures to environmental chemicals can lead to increased risk of chronic diseases in adult life [49]. An accounting of these earlier-life exposures would require re-contacting the spouses, and even then, they may not be able to reliably report use of specific active ingredients on their childhood farms.

Underestimations of non-occupational exposure may also occur from omitting the dietary exposure pathway; however, our previous literature review found only limited evidence that dietary factors were linked to increased pesticide exposure in farming populations [1]. We also omitted the bystander pathway (incidental exposure from being present but engaged in an unrelated activity at the time of pesticide application) because of limited published data quantifying its contribution [1] and lack of information on the timing of pesticide applications during AHS spouses' field-based tasks. For instance, we previously identified only three publications with relevant data that demonstrated a short-term increase in urinary pesticide biomarker concentrations among a limited number of female spouses who were "outside" or "nearby" when their husbands conducted liquid spray applications [14–16]. Additional studies that more precisely define bystander exposure and examine a greater number of spouses over a longer time period and different application scenarios would be informative. Further, we were not able to identify a suitable surrogate for bystander potential from the AHS questionnaire data. We considered using days spent in the field or data on spouse engagement in farm activities as proxies for the probability of being a bystander; however, we concluded that many of the queried activities would not likely coincide with timing of pesticide applications.

Additional underestimations of exposure may occur within the three pathways evaluated here because the estimates of drift and take-home exposure intensity do not include pesticide usage of the other residents in the home (including the spouses) or neighboring farmers/farmworkers. The spouses' usage could be included in sub-analyses or sensitivity analyses. Incorporation of other household member applications for the take-home pathway would require re-contacting participants or their family members. Neighbors' pesticide usage may be incorporated eventually into the drift pathway using crop maps to predict which active ingredients were applied to neighboring farms. Underestimation of exposure may also occur because the algorithm omits non-occupational pesticide exposure at the spouse's off-farm jobs. No information was collected on active ingredient-specific pesticide applications in or near the spouse's work place. Finally, the model did not modify exposure intensity for hygiene-related factors, such as

removing work clothing and shoes prior to entering the home or in-home cleaning practices; however, we previously identified only limited support for the impact of these behaviors on house dust pesticide levels [1].

Use of algorithm and future efforts

This algorithm was designed for the AHS spouses; however, it may have relevance to a broader population, including children living on farms or residents of agricultural areas. Approximately 2% of the U.S. population lives on farms, while 15% live in rural areas, potentially in proximity to crops or orchards where pesticides are applied [50, 51]. In addition, ~80–90% of households in both urban and rural areas report using pesticides for home and yard pest treatments [36, 52–54]. The framework may be extended with some modification for the population of interest and the study-specific information available. The algorithm is intended to be used separately for individual active ingredients; use of the algorithm to assess health impacts from combining exposure intensities from the algorithm across multiple pesticides would require consideration of the relative toxicokinetic and toxicological properties of the different chemicals.

Ideally, the algorithm's intensity scores require validation, but this is difficult because of a lack of a gold standard over the time frame of interest; as such any current validation studies would require extrapolation or generalization to historical exposures. However, there may be informative value to relate estimates of exposure intensity with shorter-term measurements of exposures to provide an indication of the degree of exposure misclassification associated with surrogate indicators [55]. In future efforts, we will evaluate the validity of the algorithm by determining whether the intensity estimates are associated with pesticide dust concentrations. These evaluations will be similar to those conducted to examine the validity of the applicator's occupational algorithm [22–24]. In addition, we plan to conduct indirect validation of these metrics by evaluating how exposure–response relationships change when using the newly developed algorithm as compared to simpler metrics. We will also examine consistency between exposure–response relationships in the AHS spouses and associations that were previously observed in the applicators based on occupational exposure. Finally, we will investigate the impact on the exposure–response relationships when certain pathways are included or excluded. For instance, because it is uncertain whether the spouse's self-reported employment status is a good surrogate for the potential for take-home exposure, we can examine the impact on pesticide exposure intensity estimates that include or exclude the take-home pathway.

Conclusions

In summary, we developed a deterministic algorithm to provide a metric of cumulative, active ingredient-specific pesticide exposure intensity for AHS spouses that represents additional exposure refinement over using questionnaire variables singly or accounting only for their personal use of pesticides. Our algorithm captured substantial contrast in exposure intensities within the study population and demonstrated important pathway-specific differences for two active ingredients. The transparency of our assumptions and systematic analysis of the published data will allow for future sensitivity analyses that examine the influence of our assumptions on the rank ordering of participants. The approach allows for the algorithm to be readily revised as new information becomes available, and for possible adaptations to other study populations. The algorithm represents an important step forward in identifying pesticide exposure contrast in the AHS spouses. Subsequent etiologic analyses based on our exposure assessment approach have the potential to advance greatly our understanding of the potential health risks pesticides pose to women.

Disclaimer

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health. The United States Environmental Protection Agency through its Office of Research and Development collaborated in the research described here. It has been subjected to Agency review and approved for publication.

Acknowledgements This work was supported by the Intramural Research Program of the National Institutes of Health, National Cancer Institute, Division of Cancer Epidemiology and Genetics (Z01CP010119) and the National Institute of Environmental Health Sciences (Z01-ES049030). NCD was supported, in part, through contract (HHSN261201400231P) and the American Cancer Society grant 127509-MRSG-15-147-01-CNE.

Compliance with ethical standards

Conflict of interest For transparency, NCD discloses that her spouse is an employee of the Dow Chemical Company, a producer of hundreds of chemicals and products, including pesticides. The remaining authors declare that they have no conflict of interest.

References

1. Deziel NC, Friesen MC, Hoppin JA, Hines CJ, Thomas K, Beane Freeman LE. A review of non-occupational pathways for pesticide exposure in women living in agricultural areas. *Environ Health Perspect.* 2015;123:515–24.
2. Blair A, Zahm SH. Methodologic issues in exposure assessment for case-control studies of cancer and herbicides. *Am J Ind Med.* 1990;18:285–93.
3. Hoppin JA, Adgate JL, Eberhart M, Nishioka M, Ryan PB. Environmental exposure assessment of pesticides in farmworker homes. *Environ Health Perspect.* 2006;114:929–35.
4. Carreon T, Butler MA, Ruder AM, Waters MA, Davis-King KE, Calvert GM, et al. Gliomas and farm pesticide exposure in women: the upper midwest health study. *Environ Health Perspect.* 2005;113:546–51.
5. Folsom AR, Zhang S, Sellers TA, Zheng W, Kushi LH, Cerhan JR. Cancer incidence among women living on farms: findings from the Iowa women's health study. *J Occ Env Med.* 1996;38:1171–6.
6. Aschengrau A, Ozonoff D, Coogan P, Vezina R, Heeren T, Zhang Y. Cancer risk and residential proximity to cranberry cultivation in massachusetts. *Am J Public Health.* 1996;86:1289–96.
7. Glorennec P, Serrano T, Fravallo M, Warembourg C, Monfort C, Cordier S, et al. Determinants of children's exposure to pyrethroid insecticides in western france. *Environ Int.* 2017;104:76–82.
8. Duell EJ, Millikan RC, Savitz DA, Newman B, Smith JC, Schell MJ, et al. A population-based case-control study of farming and breast cancer in North Carolina. *Epidemiology.* 2000;11:523–31.
9. Cockburn M, Mills P, Zhang X, Zadnick J, Goldberg D, Ritz B. Prostate cancer and ambient pesticide exposure in agriculturally intensive areas in california. *Am J Epidemiol.* 2011;173:1280–8.
10. Rull RP and Ritz B. 2003. Historical pesticide exposure in California using pesticide use reports and land-use surveys: an assessment of misclassification error and bias. *Send to Environ Health Perspect.* 13:1582–9.
11. Ward MH, Lubin J, Giglierano J, Colt JS, Wolter C, Bekiroglu N, et al. Proximity to crops and residential exposure to agricultural herbicides in Iowa. *Environ Health Perspect.* 2006;114:893–7.
12. Maxwell SK, Airola M, Nuckols JR. Using landsat satellite data to support pesticide exposure assessment in california. *Int J Health Geogr.* 2010;9:1.
13. Curwin BD, Hein MJ, Sanderson WT, Striley C, Heederik D, Kromhout H, et al. Urinary pesticide concentrations among children mothers and fathers living in farm and non-farm households in Iowa. *Ann Occup Hyg.* 2007;511:53–65.
14. Alexander BH, Burns CJ, Bartels MJ, Acquavella JF, Mandel JS, Gustin C, et al. Chlorpyrifos exposure in farm families: results from the farm family exposure study. *J Expo Sci Environ Epidemiol.* 2006;16:447–56.
15. Alexander BH, Mandel JS, Baker BA, Burns CJ, Bartels MJ, Acquavella JF, et al. Biomonitoring of 2,4-dichlorophenoxyacetic acid exposure and dose in farm families. *Environ Health Perspect.* 2007;115:370–6.
16. Arbuckle TE, Ritter L. Phenoxyacetic acid herbicide exposure for women on Ontario farms. *J Toxicol Environ Health A.* 2005;68:1359–70.
17. Barr DB, Thomas K, Curwin B, Landsittel D, Raymer J, Lu C, et al. Biomonitoring of exposure in farmworker studies. *Environ Health Perspect.* 2006;114:936–42.
18. Dopart PJ and Friesen MC. New opportunities in exposure assessment of occupational epidemiology: use of measurements to aid exposure reconstruction in population-based studies. *Curr Environ Health Rep.* 2017. 4:355–363. <https://doi.org/10.1007/s40572-017-0153-0>.
19. Negatu B, Vermeulen R, Mekonnen Y, Kromhout H. A method for semi-quantitative assessment of exposure to pesticides of applicators and re-entry workers: an application in three farming systems in ethiopia. *Ann Occup Hyg.* 2016;60:669–83.

20. Coble J, Thomas KW, Hines CJ, Hoppin JA, Dosemeci M, Curwin B, et al. An updated algorithm for estimation of pesticide exposure intensity in the agricultural health study. *Int J Environ Res Public Health*. 2011;8:4608–22.
21. Dosemeci M, Alavanja MC, Rowland AS, Mage D, Zahm SH, Rothman N, et al. A quantitative approach for estimating exposure to pesticides in the agricultural health study. *Ann Occ Hyg*. 2002;46:245–60.
22. Hines CJ, Deddens JA, Jaycox LB, Andrews RN, Striley CA, Alavanja MC. Captan exposure and evaluation of a pesticide exposure algorithm among orchard pesticide applicators in the Agricultural Health Study. *Ann Occup Hyg*. 2008;52:153–66.
23. Coble J, Arbuckle T, Lee W, Alavanja M, Dosemeci M. The validation of a pesticide exposure algorithm using biological monitoring results. *J Occup Environ Hyg*. 2005;2:194–201.
24. Thomas KW, Dosemeci M, Coble JB, Hoppin JA, Sheldon LS, Chapa G, et al. Assessment of a pesticide exposure intensity algorithm in the agricultural health study. *J Expo Sci Environ Epidemiol*. 2010;20:559–69.
25. Gladen BC, Sandler DP, Zahm SH, Kamel F, Rowland AS, Alavanja MC. Exposure opportunities of families of farmer pesticide applicators. *Am J Ind Med*. 1998;34:581–7.
26. Louis LM, Lerro CC, Friesen MC, Andreotti G, Koutros S, Sandler DP, et al. A prospective study of cancer risk among Agricultural Health Study farm spouses associated with personal use of organochlorine insecticides. *Environ Health*. 2017;16:95
27. Alavanja MC, Aklund G, Baird D, Blair A, Bond A, Dosemeci M, et al. Cancer and noncancer risk to women in agriculture and pest control: the Agricultural Health Study. *J Occ Med*. 1994;6:1247–50.
28. Kirrane EF, Hoppin JA, Umbach DM, Samanic C, Sandler DP. Patterns of pesticide use and their determinants among wives of farmer pesticide applicators in the agricultural health study. *J Occ Env Med*. 2004;46:856–69.
29. Freeman LE, Rusiecki JA, Hoppin JA, Lubin JH, Koutros S, Andreotti G, et al. Atrazine and cancer incidence among pesticide applicators in the Agricultural Health Study (1994–2007). *Environ Health Perspect*. 2011;119:1253–9.
30. Deziel NC, Beane Freeman LE, Graubard BI, Jones RR, Hoppin JA, Thomas K, et al. Relative contributions of agricultural drift, para-occupational, and residential use exposure pathways to house dust pesticide concentrations: meta-regression models of published data. *Environ Health Perspect*. 2017;125:296–305.
31. Deziel NC, Ward MH, Bell EM, Whitehead TP, Gunier RB, Friesen MC, et al. Temporal variability of pesticide concentrations in homes and implications for attenuation bias in epidemiologic studies. *Environ Health Perspect*. 2013;121:565–71.
32. Lewis RG, Fortmann RC, Camann DE. Evaluation of methods for monitoring the potential exposure of small children to pesticides in the residential environment. *Arch Environ Contam Toxicol*. 1994;26:37–46.
33. Bureau of Labor Statistics. American time use survey. 2003. <http://www.bls.gov/tus/data.htm>. Accessed October 2014.
34. Gunier RB, Ward MH, Airola M, Bell EM, Colt J, Nishioka M, et al. Determinants of agricultural pesticide concentrations in carpet dust. *Environ Health Perspect*. 2011;119:970–6.
35. Harnly ME, Bradman A, Nishioka M, McKone TE, Smith D, McLaughlin R, et al. Pesticides in dust from homes in an agricultural area. *Environ Sci Technol*. 2009;43:8767–74.
36. Deziel NC, Colt JS, Kent EE, Gunier RB, Reynolds P, Booth B, et al. Associations between self-reported pest treatments and pesticide concentrations in carpet dust. *Environ Health*. 2015;14:27.
37. Trunnelle KJ, Bennett DH, Tancredi DJ, Gee SJ, Stoecklin-Marois MT, Hennessy-Burt TE, et al. Pyrethroids in house dust from the homes of farm worker families in the micasa study. *Environ Int*. 2013;61:57–63.
38. Colt JS, Cyr MJ, Zahm SH, Tobias GS, Hartge P. Inferring past pesticide exposures: a matrix of individual active ingredients in home and garden pesticides used in past decades. *Environ Health Perspect*. 2007;115:248–54.
39. USEPA (U.S. Environmental Protection Agency). Interim Reregistration Eligibility Decision for Atrazine. Case No. 0062 Decision Documents for Atrazine. 2003. https://archive.epa.gov/pesticides/reregistration/web/pdf/atrazine_combined_docs.pdf.
40. Lerro CC, Koutros S, Andreotti G, Friesen MC, Alavanja MC, Blair A et al. Organophosphate insecticide use and cancer incidence among spouses of pesticide applicators in the Agricultural Health Study. *Occup Environ Med*. 2005;72:736–744.
41. Engel LS, Hill DA, Hoppin JA, Lubin JH, Lynch CF, Pierce J, et al. Pesticide use and breast cancer risk among farmers' wives in the agricultural health study. *Am J Epidemiol* 15. 2005;161:121–35.
42. Engel LS, Werder E, Satagopan J, Blair A, Hoppin JA, Koutros S, et al. Insecticide use and breast cancer risk among farmers' wives in the Agricultural Health Study. *Environ Health Perspect* 6. 2017;125:097002.
43. Bradman A, Eskenazi B, Barr DB, Bravo R, Castorina R, Chevrier J, et al. Organophosphate urinary metabolite levels during pregnancy and after delivery in women living in an agricultural community. *Environ Health Perspect*. 2005;113:1802–7.
44. Butte W, Heinzow B. Pollutants in house dust as indicators of indoor contamination. *Rev Env Contam Tox*. 2002;175:1–46.
45. Simcox NJ, Fenske RA, Wolz SA, Lee IC, Kalman DA. 1995. Pesticides in household dust and soil: exposure pathways for children of agricultural families. *Environ Health Perspect* 10312:1126–1134.
46. Mandel JS, Alexander BH, Baker BA, Acquavella JF, Chapman P, Honeycutt R. Biomonitoring for farm families in the farm family exposure study. *Scand J Work Environ Health*. 2005;31 (Suppl 1):98–104.
47. Arbuckle TE, Bruce D, Ritter L, Hall JC. Indirect sources of herbicide exposure for families on ontario farms. *J Expo Sci Environ Epidemiol*. 2006;16:98–104.
48. Hofmann JN, Hoppin JA, Lynch CF, Poole JA, Purdue MP, Blair A, et al. Farm characteristics, allergy symptoms, and risk of non-hodgkin lymphoid neoplasms in the agricultural health study. *Cancer Epidemiol Biomark Prev*. 2015;24:587–94.
49. Barouki R, Gluckman PD, Grandjean P, Hanson M, Heindel JJ. Developmental origins of non-communicable disease: implications for research and public health. *Environ Health*. 2012;11:42.
50. USDA (U.S. Department of Agriculture). 2007 Census of Agriculture. Document number AC-07-A-51. 2009. http://www.agcensus.usda.gov/Publications/2007/Full_Report/usv1.pdf. Accessed March 2015.
51. USDA (U.S. Department of Agriculture). Population & migration. 2015. <http://www.ers.usda.gov/topics/rural-economy-population/population-migration.aspx>. Accessed May 2015.
52. Adgate JL, Kukowski A, Stroebel C, Shubat PJ, Morrell S, Quackenboss JJ, et al. Pesticide storage and use patterns in minnesota households with children. *J Expo Sci Environ Epidemiol*. 2000;10:159–67.
53. Colt J, Lubin J, Camann D, Davis S, Cerhan J, Severson R. Comparison of pesticide levels in carpet dust and self-reported pest treatment practices in four us sites. *J Expo Sci Environ Epidemiol*. 2004;14:74–83.
54. Wu XM, Bennett DH, Ritz B, Frost J, Cassady D, Lee K, et al. Residential insecticide usage in northern california homes with young children. *J Expo Sci Environ Epidemiol*. 2011;21:427–36.
55. Blair A, Thomas K, Coble J, Sandler DP, Hines CH, Lynch CF, et al. Impact of pesticide exposure misclassification on estimates

- of relative risks in the Agricultural Health Study. *J Occ Env Med.* 2011;68:537–41.
56. Fitzgerald D, Chanasyk DS, Neilson RD, Kiely D, Audette R. Farm well water quality in alberta. *Water Qual Res J Can.* 2001;36:565–88.
57. Friesen MC, Lavoue J, Teschke K, van Tongeren M. Occupational exposure assessment in industry- and population-based epidemiological studies. In: Nieuwenhuijsen M, editor. *Exposure Assessment in Environmental Epidemiology.* 2nd Edition. New York, NY: Oxford University Press; 2015. p. 139–62.
58. Friesen MC, Park DU, Colt JS, Baris D, Schwenn M, Karagas MR, et al. Developing estimates of frequency and intensity of exposure to three types of metalworking fluids in a population-based case-control study of bladder cancer. *Am J Ind Med.* 2014;57:915–27.
59. Goldman L, Eskenazi B, Bradman A, Jewell NP. Risk behaviors for pesticide exposure among pregnant women living in farmworker households in salinas, california. *Am J Ind Med.* 2004;45:491–9.
60. Gunier RB, Nuckols JR, Whitehead TP, Colt JH, Deziel NC, Metayer C, et al. Temporal trends of insecticide concentrations in carpet dust in California from 2001-6. *Environ Sci Technol.* 2016;50:7761–9.
61. Hartge P, Colt JS, Severson RK, Cerhan JR, Cozen W, Camann D, et al. Residential herbicide use and risk of non-hodgkin lymphoma. *Cancer Epidemiol Biomark Prev.* 2005;14:934–7.
62. Jones RR, DellaValle CT, Flory AR, Nordan A, Hoppin JA, Hofmann JN, et al. Accuracy of residential geocoding in the agricultural health study. *Int J Health Geogr.* 2014;13:37.
63. Jones RR, Yu CL, Nuckols JR, Cerhan JR, Airola M, Ross JA, et al. Farm residence and lymphohematopoietic cancers in the Iowa women's health study. *Environ Res.* 2014;133:353–61.
64. McCauley LA, Michaels S, Rothlein J, Muniz J, Lasarev M, Ebbert C. Pesticide exposure and self reported home hygiene practices in agricultural families. *AAOHNJ.* 2003;51:113–9.
65. Peters S, Carey RN, Driscoll TR, Glass DC, Benke G, Reid A, et al. The Australian Work Exposures Study: prevalence of occupational exposure to diesel engine exhaust. *Ann Occup Hyg.* 2015;59:600–8.
66. Pronk A, Stewart PA, Coble JB, Katki HA, Wheeler DC, Colt JS, et al. Comparison of two expert-based assessments of diesel exhaust exposure in a case-control study: programmable decision rules versus expert review of individual jobs. *Occ Env Med.* 2012;69:752–8.
67. Quandt SA, Arcury TA, Rao P, Snively BM, Camann DE, Doran AM, et al. Agricultural and residential pesticides in wipe samples from farmworker family residences in North Carolina and Virginia. *Environ Health Perspect.* 2004;112:382–7.
68. Teitelbaum SL, Gammon MD, Britton JA, Neugut AI, Levin B, Stellman SD. Reported residential pesticide use and breast cancer risk on long island, new york. *Am J Epidemiol.* 2007;165: 643–51.