

# Effects of posture on dynamic back loading during a cable lifting task

Sean Gallagher , William S. Marras , Kermit G. Davis & Kimberly Kovacs

To cite this article: Sean Gallagher , William S. Marras , Kermit G. Davis & Kimberly Kovacs (2002) Effects of posture on dynamic back loading during a cable lifting task, Ergonomics, 45:5, 380-398, DOI: [10.1080/00140130210127639](https://doi.org/10.1080/00140130210127639)

To link to this article: <https://doi.org/10.1080/00140130210127639>



Published online: 09 Nov 2010.



Submit your article to this journal [↗](#)



Article views: 188



View related articles [↗](#)



Citing articles: 21 View citing articles [↗](#)

## Effects of posture on dynamic back loading during a cable lifting task

SEAN GALLAGHER<sup>†\*</sup>, WILLIAM S. MARRAS<sup>‡</sup>, KERMIT G. DAVIS<sup>§</sup> and KIMBERLY KOVACS<sup>‡</sup>

<sup>†</sup>National Institute for Occupational Safety and Health, Pittsburgh Research Laboratory, Pittsburgh, PA 15236-0070, USA

<sup>‡</sup>Biodynamics Laboratory, Department of Industrial, Welding, and Systems Engineering, The Ohio State University, 1971 Neil Avenue, Columbus, OH 43210, USA

<sup>§</sup>Department of Environmental Health, Industrial Hygiene Division, 330 Kettering Laboratory, 3223 Eden Avenue, University of Cincinnati, Cincinnati, OH 45267, USA

*Keywords:* Posture; Low-back pain; Electromyography; Biomechanics; Lifting; Spinal loads; Trunk muscles.

This study evaluated spinal loads associated with lifting and hanging heavy mining cable in a variety of postures. This electrical cable can weigh up to 10 kg per metre and is often lifted in restricted spaces in underground coal mines. Seven male subjects performed eight cable lifting and hanging tasks, while trunk kinematic data and trunk muscle electromyograms (EMGs) were obtained. The eight tasks were combinations of four postures (standing, stooping, kneeling on one knee, or kneeling on both knees) and two levels of cable load (0 N or 100 N load added to the existing cable weight). An EMG-assisted model was used to calculate forces and moments acting on the lumbar spine. A two-way split-plot ANOVA showed that increased load ( $p < 0.05$ ) and changes in lifting posture ( $p < 0.05$ ) independently affected trunk muscle recruitment and spinal loading. The increase in cable load resulted in higher EMG activity of all trunk muscles and increased axial and lateral bending moments on the spine ( $p < 0.05$ ). Changes in posture caused more selective adjustments in muscle recruitment and affected the sagittal plane moment ( $p < 0.05$ ). Despite the more selective nature of trunk EMG changes due to posture, the magnitude of changes in spinal loading was often quite dramatic. However, average compression values exceeded 3400 N for all cable lifting tasks.

### 1. Introduction

The posture adopted by the body during manual materials handling tasks is thought to have a profound impact on spinal loading. The preponderance of manual lifting research in ergonomics has concentrated on analysis of loads experienced during standing postures; however, certain occupations require that workers adopt postures that may significantly alter muscle recruitment patterns and biomechanical loads on the body. For example, coal miners often work in vertically confined workspaces where standing erect is not possible. In such an environment, the miner is obliged to

---

\*Author for correspondence. e-mail: sfg9@cdc.gov

select a lifting posture from a menu of unpalatable alternatives, typically stooping (standing with a fully flexed trunk) or kneeling (on one or both knees). Each of these postures engenders significant biomechanical disadvantages. For example, psychophysical lifting capacity and strength are significantly reduced in the kneeling posture (Gallagher *et al.* 1988, Smith *et al.* 1989, Gallagher and Unger 1990, Gallagher and Hamrick 1992, Gallagher 1997). Stooping requires severe trunk flexion resulting in a number of biomechanical disadvantages, including higher shear loading on the spine, reliance on passive tissues for spinal support, and reduced force output from the back muscles as a function of the muscle length-strength relationship (Floyd and Silver 1955, Potvin *et al.* 1991, Yingling and McGill 1999). However, the stooping posture often appears to be favoured by mineworkers, particularly when high forces need to be exerted in restricted spaces, or when enhanced mobility is needed. However, it is quite conceivable that workers who adopt the stooping posture to take advantage of its strength benefits may, at the same time, be subjecting themselves to potentially damaging spinal loads.

In addition to the unusual postural demands of their job, miners sometimes have to handle materials possessing rather unique characteristics. One example is the large diameter electrical cable used to power heavy mining equipment. These cables are extremely heavy due to the large amounts of copper wire and to the insulation requirements associated with the electrical current demands of this massive equipment. It is not uncommon for this cable to have a diameter of 7.5 cm and to weigh 10 kg per metre. Several metres of cable may have to be handled during a cable lifting task, and the higher the cable is lifted the greater is the load that must be supported by the worker. This progressive increase in the load being supported is unusual in manual handling activities and must place a higher demand on the muscular agonists compared to a similar lift involving a constant load. The flexibility of cable may cause additional concerns due to load instability, especially given the heavy weight of the material. In the mining environment, additional stressors may exist when lifting cable. For example, the cable may become quite muddy, adding significantly to the weight and force requirements of the lift. Not surprisingly, handling of heavy cable has been associated with a large number of lost-time back injuries in the mining industry (Randolph 1991).

Changes in body posture (from standing to kneeling, for instance) are likely to have significant impacts on the synergism of muscular activity, and it is presumed that the sensorimotor cortex would have to adjust muscle activation programmes to accomplish a specified manual lifting task (Dul 1986). Previous investigations have confirmed that significant changes in trunk muscle activity do occur in restricted postures (Gallagher *et al.* 1988, 1994, Gallagher and Unger 1990). Increases in the weight (or load) lifted have also been shown to impact on trunk muscle recruitment (Fathallah *et al.* 1998, De Looze *et al.* 1999, Davis and Marras 2000). However, it is not well understood whether the trunk muscle activity changes due to posture and those due to increased load activity act independently or whether these influences might interact. Since trunk muscle activity is thought to be indicative of spinal load (McGill and Norman 1986, Chaffin and Andersson 1991, Granata and Marras 1993), which is (in turn) associated with development of low-back disorders (Marras *et al.* 1993), understanding such relationships is of critical importance to those who must handle materials in restricted working postures.

Therefore, the current experiment was undertaken to address the following hypotheses.

- (1) *Hypothesis 1.* Changes in posture would alter trunk muscle activity and estimates of lumbar spine loading.
- (2) *Hypothesis 2.* Increased cable load would alter trunk muscle activity and lumbar spine loading.
- (3) *Hypothesis 3.* The main effects described above might interact with one another.

These effects were investigated in a task involving the lifting and hanging of large diameter electrical cables, which are commonly handled in underground coal mines.

## 2. Methods

### 2.1. Subjects

Seven male subjects volunteered to perform a series of lifts using a heavy mining cable used to power heavy underground equipment. Subjects read and signed an informed consent form prior to the initiation of testing. None had a prior history of low-back disorders (LBD). The subjects' ages ranged from 22 to 36 years, and none had prior experience with cable hanging tasks. Anthropometric data are provided in table 1. Comparison of these data with the mining population suggests that these subjects were approximately the same stature, but weighed less and were somewhat younger than underground coal miners (Gallagher 1999).

### 2.2. Experimental design

The study evaluated spinal loads during the lifting and hanging of the cable using an EMG-assisted biomechanical model. The independent variables consisted of lifting posture (with four levels) and cable load (two levels). Subjects served as blocks within which experimental conditions were randomized. The following lifting postures were manipulated in this study: kneeling on one knee (1KNEE), kneeling on two knees (2KNEE), stooping (STOOP), and standing (STAND). The weight of the entire length of cable (7.6 m) used in this study was approximately 367 N. However, the amount of this weight actually supported by the subject during a lift was only a portion of the total load and varied according to the length lifted during the task. Previous research (Gallagher *et al.* 2001), also employing continuous mining cable, indicated a linear increase in load according to the following equation:

$$\text{Cable load (N)} = 24.7 + 0.083 \times \text{Height of centre of cable (mm)} \quad (1)$$

Table 1. Anthropometric data of the subjects.

Subject	Age (yrs)	Weight (kg)	Height (cm)
1	24	76.2	194.4
2	21	72.3	189.9
3	25	64.5	174.0
4	30	92.7	178.6
5	36	95.5	197.5
6	22	80.9	165.0
7	22	79.5	181.3
Mean	25.7	80.2	182.9
Standard deviation	5.4	10.9	11.6

An additional 100 N (50 N load attached to both ends) was placed on the cable to simulate the effect of an additional load (from additional lengths of cable, mud, etc.) as may be experienced in underground cable lifting tasks. While the subjects did not directly lift this additional load, the weight added to either end increased the force required to draw the cable ends together when lifting the centre portion of the cable to the hook. The added load increased the force required to lift the cable by an estimated 20–25%. The magnitude of the additional load was selected such that the load would provide an estimate of the effect without being so large as to create an undue risk of injury to the subject. Each lift was performed three times in each load posture/load condition, for a total of 24 lifts per subject.

### 2.3. Dependent measures

Spine moments and forces were estimated using an EMG-assisted biomechanical model which has been under development in the Ohio State University Biodynamics Laboratory over the past 18 years (Marras and Reilly 1988, Reilly and Marras 1989, Marras and Sommerich 1991a, b, Granata and Marras 1993, 1995, Mirka and Marras 1993, Marras and Granata 1995, 1997, Davis *et al.* 1998). This model provides estimates of spine loading parameters based upon measured activity of ten trunk muscles, from which estimates of muscle force and subsequent spine loading are determined (Marras and Granata 1997). Forces for each muscle are estimated using the following equation.

$$F_j = \text{PCSA}_j \times \text{Gain} \times \frac{\text{EMG}_j(t)}{\text{EMG}_{\text{max-}j}} \times f(\text{L} - \text{S}) \times f(\text{F} - \text{V}) \quad (2)$$

where:

$F_j$  = predicted muscle force for muscle  $j$ ;

$\text{PCSA}_j$  = physiological cross-sectional area for muscle  $j$ ;

Gain = estimate of muscle stress (force/area);

$\text{EMG}_j(t)$  = instantaneous integrated EMG for muscle  $j$  at time  $t$ ;

$\text{EMG}_{\text{Max-}j}$  = integrated EMG from maximum voluntary contraction for muscle  $j$ ;

$f(\text{L-S})$  = muscle length-strength relationship modulation factor; and

$f(\text{F-V})$  = muscle force-velocity relationship modulation factor.

The muscle generated internal moments about the axis of rotation are predicted from the sum vector products combining the tensile muscle force for each muscle and the moment arms of each respective muscle as given by equation (2):

$$M_i = \sum r_{ij} \times F_{ij} \quad (3)$$

where:

$M_i$  = predicted internal moment for the  $i$ th plane;

$r_{ij}$  = moment-arm for muscle  $j$  in the  $i$ th plane; and

$F_{ij}$  = vector force component for muscle  $j$  in the  $i$ th plane.

The model has been validated under forward trunk bending motions (Marras and Sommerich 1991a, b, Granata and Marras 1993, 1995), trunk twisting motions (Marras and Granata 1995), lateral bending motions (Marras and Granata 1997), and lowering tasks (Davis *et al.* 1998).

The spinal loads evaluated in this study include compression, anterior-posterior (A-P) shear force and lateral shear force, as well as moments acting about the

lumbosacral joint ( $L_5/S_1$ ) during the lift. Maximum values (over the entire lift) were obtained for each of these parameters for each of the three repetitions within each experimental condition. The average value of the three repetitions was taken as the best estimate for these spinal load variables for each subject. Normalized electromyographic (EMG) activity of the 10 trunk muscles was also evaluated as a function of posture and load. The values presented represent the mean of the maximum EMGs obtained for the three repetitions in each treatment combination.

#### 2.4. Apparatus

The Lumbar Motion Monitor (LMM) was used to collect the trunk motion variables. The LMM is essentially an exoskeleton of the spine in the form of a tri-axial electrogoniometer that measures instantaneous three-dimensional position, velocity, and acceleration of the trunk. The lightweight design of the LMM allowed the data to be collected with minimal obstruction to the subject's movements. More detailed information on the design, accuracy, and application of the LMM can be found in Marras *et al.* (1992).

EMG activity was collected through the use of bi-polar electrodes spaced 3 cm apart at the 10 major trunk muscle sites. The ten muscles of interest were the right and left pairs of the erectors spinae (RES, LES), latissimus dorsi (RLD, LLD), internal obliques (RIO, LIO), external obliques (REO, LEO), and rectus abdominis (RRA, LRA). The standard locations of electrode placement of muscles used in conjunction with the Ohio State University EMG-assisted biomechanical model are described in Mirka and Marras (1993).

A Bertec 4060A force plate (Bertec Corp., Columbus, OH, U.S.A.) and a set of electrogoniometers measured the external loads and moments placed on  $L_5/S_1$  during calibration exertions. The purpose of the calibration exertions was to determine the subject-specific gain value to be used with the model in an 'open-loop' fashion, as required to model the current set lifting tasks. The term 'open-loop' refers to exertions that use a predetermined gain to calculate internal moments and forces, rather than calculating a specific gain for each exertion. The electrogoniometers measured the relative position of  $L_5/S_1$  with respect to the centre of the force plate, along with the subject's pelvic angle. The forces and moments were translated and rotated from the centre of the force plate to  $L_5/S_1$  by this means as described in Fathallah *et al.* (1997). The internal moments were adjusted to equal the external moments through the use of this gain factor. The value of the gain represented the maximum force output of the muscles per cross-sectional unit area for the particular subject. Gains are highly variable between subjects, depending on the degree of conditioning and natural ability, but have been found to remain stable on a within-subjects basis for this EMG-assisted model (Granata and Marras 1995).

All signals from the aforementioned equipment were collected simultaneously through customized Windows-based software developed in the Biodynamics Laboratory. The signals were collected at 100 Hz and recorded on a personal computer via an analogue-to-digital board.

#### 2.5. Procedure

Upon arrival at the Biodynamics Laboratory, subjects were given a brief description of the study and the tasks that they would be asked to perform. Next, anthropometric measurements were taken. The surface electrodes were applied using standard placement procedures to sample the muscles of interest in accordance

with Marras (1990) and NIOSH (1991). Each subject was then placed into a structure that allowed maximal exertions to be performed in six directions, while a constant resistance was held against the subject (Marras and Mirka 1993). These maxima were performed to allow subsequent EMG data to be normalized. The six exertions consisted of the following: sagittal extension with the trunk at a 20° forward flexion angle, sagittal flexion at 0° flexion, right lateral bending at 0° flexion, left lateral bending at 0° flexion, right twist at 0° flexion, and left twist at 0° flexion. After each maximal exertion, 2 min of rest was given to reduce the effects of fatigue (Caldwell *et al.* 1974).

Before beginning the experimental tasks, subjects completed a set of calibration lifts. These lifts allowed the gain of the individual to be determined for the 'open-loop' exertions. During the calibration exertions, the subject lifted a 30 lb (13.6 kg) case from a sagittally symmetric position at a slow, smooth pace (controlled by the subject). The lift started with the case at the subject's knee height and ended when the subject reached an upright position. The calibration lifts were run under 'closed-loop' conditions; that is, internal moments were validated with measured external moments. Before and after each set of calibrations, data were collected to determine the position of the LMM and the relative position of the subject's L<sub>5</sub>/S<sub>1</sub> joint to the centre of the force plate as measured by electrogoniometers when the subject was standing erect. Along with the LMM and goniometer neutral values, the initial readings from the force plate were recorded. Figure 1 shows a subject lifting the case during a calibration exertion.

### 2.6. Experimental task

In each posture, the subject was required to lift the cable from the floor and hang it on a hook located above the cable. The centre of cable was located on the floor in front of the subject for each lift, with equal lengths of the cable extending laterally to the left and right. The subject faced the cable and grabbed it with both hands and lifted the cable and hung the centre of the cable from a hook located above the cable and slightly in front of the subject. The speed of the lift was left to the subject's discretion.

For restricted lifting postures (kneeling and stooping), the height of the hook was located 137 cm above the floor. For unrestricted (standing) lifts the hook was located 178 cm above the floor. These hook locations reflect the nature of cable hanging tasks in the underground environment. It must be recalled that the cable must always be hung from hooks attached to the 'ceiling' of the mine (or 'mine roof' as it is called). In coal mines with restricted vertical space (1.4 m, for example), a kneeling or stooping lift is compulsory (as most adults cannot stand upright in such a space). In such circumstances, the vertical excursion of the lift will necessarily be limited by the lower height of the mine roof. However, when the coal seam is thicker (say 1.8 m), hanging the cable from the mine roof must be done using a more traditional lift, ending in the upright standing position, so that the cable can be lifted all the way up to the ceiling. Thus, a 'standing' lift cannot be used in a restricted space, and restricted lifting postures (e.g. kneeling) cannot be used to lift the cable to the mine roof when the seam height is higher (due to the lack of the ability to reach the ceiling). The spine loading and muscle activity in each of the four postures were of interest to the investigators, as all of these situations occur in the mining environment. It should be realized however, when comparing results in these postures, that more cable weight is supported by the subject in standing lifts than in




Figure 1. A subject performing the calibration exertions.

lifts involving restricted postures. The increase in load handled at the end of the lift when standing was estimated to be approximately 3.5 kg (Gallagher *et al.* 2001).

Subjects were given instructions regarding the postural constraints for each lifting task. For the 1KNEE condition, subjects began kneeling with one knee on the floor and the shank of the opposite leg perpendicular to the floor. Each subject was permitted to kneel on either the right or left knee, but was required to maintain that same position for all of the 1KNEE conditions. The 2KNEE condition required the subject to kneel on both knees. For the STOOP condition, subjects maintained a semi-squat position at a comfortable knee angle throughout the lift. The height of the hook also acted as an imaginary ‘ceiling’ in the STOOP condition that the subjects were not permitted to exceed. The STAND condition also began with the cable on the floor; however, the subject was able to fully extend his trunk in order to hang the cable on the hook. Thus, the primary difference between the STAND and STOOP conditions was that the subject had a limited ability to extend his trunk in the STOOP, while complete trunk extension was permitted in the STAND. Subjects

were given a maximum of 5 s to complete each lift. Data were collected for the entire lift, and a laboratory assistant marked the end of the lift by depressing a button (allowing a change in voltage to be recorded in the data file) so that post-lift EMG data could be eliminated during subsequent analyses. Figure 2 shows a subject performing the task in each of the four postures.

### 2.7. Data analysis

Lumbar Motion Monitor (LMM) voltage signals were converted into angles, velocities, and accelerations through customized conversion software. The kinematic data were then used in the EMG-assisted spinal loading model. The raw EMG signals were pre-amplified, high-pass filtered at 30 Hz, low-pass filtered at 1000 Hz, rectified, and integrated via a 20-ms sliding window hardware filter. With the aid of a customized software program, the EMG data were normalized with respect to the maximum output of the muscles and muscle length-strength and force-velocity modulations. Finally, the EMG and kinematic data were imported into the EMG-assisted model to calculate spinal forces and moments at the lumbosacral joint (L<sub>5</sub>/S<sub>1</sub>), which (along with the normalized EMG activity) served as the dependent variables in this study.

Univariate descriptive statistics were obtained and a split-plot analysis of variance (ANOVA) was performed for each dependent measure (Kirk 1982). For all significant posture effects, *post hoc* analyses, in the form of Tukey multiple pairwise comparisons (Honestly Significant Difference [HSD]), were performed to determine the source(s) of the significant effect(s). Alpha levels ( $\alpha$ ) were set at 0.05 for all cases.

## 3. Results

Tables 2 and 3 provide summaries of ANOVA results for EMG activity of the ten trunk muscles and spine loading estimates, respectively. Results of Tukey HSD *post hoc* tests for significant posture effects are also provided in these tables. No significant interactions between posture and load condition were detected for any dependent measures.

### 3.1. Effects due to posture

Changes in posture affected the activation patterns of all trunk muscles with the exception of the rectus abdominis ( $p < 0.05$ ); however, the change in activity for each posture typically involved a subset of trunk muscles. As can be seen in figure 3, the primary spine extensors (erectores spinae and latissimus dorsi) exhibited the greatest amount of activity in the STOOP posture, while the STAND and 2KNEE conditions resulted in the least activation of these muscles. The 1KNEE stance was characterized by an intermediate amount of activation of these extensor muscles. On the other hand, the internal obliques (another extensor muscle) tended to be less active in the kneeling postures than when the subject was standing or stooping. The external obliques exhibited increased activity in the STAND posture, with other postures not significantly different from one another.

Changes in posture also resulted in significant differences in the forces and moments experienced by the lumbar spine during the criterion task, as reported in table 3 and as illustrated in figures 4 and 5. However, in all cases, compression and shear forces were quite high. Compressive forces on the spine were significantly higher in the STOOP posture compared to all other positions ( $p < 0.05$ ), while the 2KNEE lifting condition resulted in significantly lower compressive forces than the




Figure 2. Illustration of the four postures used in the study: (a) kneeling on one knee (1KNEE); (b) kneeling on two knees (2KNEE); (c) stooping (STOOP); and (d) standing (STAND).

Table 2. Summary of *F* statistics from ANOVAs examining the effects of posture and load on normalized EMG activity for 10 trunk muscles. *F* statistics have been tested against a pooled error term with 154 degrees of freedom. Results of Tukey HSD *post hoc* tests are provided for significant posture effects.

Source of variation	df	LLD Left latissimus dorsi	RLD Right latissimus dorsi	LES Left-erectores spinae	RES Right erectores spinae	LEO Left external obliques	REO Right external obliques	LRA Left rectus abdominis	RRA Right rectus abdominis	LIO Left internal obliques	RIO Right internal obliques
Posture	3	5.65** C > BD	3.92** C > D	4.89** C > B	13.33*** C > ABD	9.10*** D > ABC	6.28*** d > ABC	0.61	1.48	9.41*** ACD > B	10.07*** CD > AB
Tukey HSD results											
Load	1	12.68***	29.72***	29.34***	77.08***	41.18***	69.04***	28.73***	38.94***	21.41***	51.67***
Posture × Load	3	0.43	0.65	0.17	1.29	1.35	0.12	0.57	0.66	1.61	0.37

\*Significant at 0.05; \*\*Significant at 0.01; \*\*\*Significant at 0.001. A = Kneeling on one knee, B = Kneeling on both knees, C = Stooping, D = Standing.

Table 3. ANOVA results for predicted forces and moments acting on the lumbosacral joint. *F* statistics reported for each force or moment have been tested against a pooled error term with 154 degrees of freedom. Results of Tukey HSD *post hoc* tests are provided for significant posture effects.

Source of variation	df	Force/Moment component						
		$F_x$	$F_y$	$F_z$	$M_x$	$M_y$	$M_z$	$M_{\text{resultant}}$
Posture	3	8.59*** D > BC, A > B	5.88*** C > AB	36.81*** C > AD > B	41.13*** C > AD > B	2.37	1.37	11.64*** C > D > B, A > B
Load	1	14.43***	27.15***	7.02**	3.14	5.42*	6.93***	6.96**
Posture × Load	3	0.25	0.28	0.61	0.53	0.38	0.18	0.53

\*Significant at 0.05; \*\*Significant at 0.01; \*\*\*Significant at 0.001. A = Kneeling on one knee, B = Kneeling on both knees, C = Stooping, D = Standing.




Figure 3. Normalized maximum EMG activity of 10 trunk muscles during cable lifts in the four postures studied (LLD = left latissimus dorsi, RLD = right latissimus dorsi, LES = left erector spinae, RES = right erector spinae, LEO = left external obliques, REO = right external obliques, LRA = left rectus abdominis, RRA = right rectus abdominis, LIO = left internal obliques, RIO = right internal obliques).




Figure 4. Predicted peak forces acting on the lumbar spine in the four postures studied ( $F_x$  = lateral shear force,  $F_y$  = anterior-posterior shear forces, and  $F_z$  = compression).




Figure 5. Predicted peak moments acting on the lumbar spine in the four postures studied ( $M_x$  = sagittal bending moment,  $M_y$  = lateral bending moment,  $M_z$  = axial twisting moment,  $M_r$  = resultant moment).

other postures ( $p < 0.05$ ). STAND and 1KNEE lifting conditions resulted in compressive forces that were not significantly different from one another and intermediate in relation to those described above. The STOOP posture also resulted in higher A-P shear forces than the kneeling postures ( $p < 0.05$ ), but A-P shear was not significantly different when stooping and standing were compared. Lateral shear forces were higher in the STAND condition than in the STOOP or 2KNEE conditions ( $p < 0.05$ ). However, lateral shear in 1KNEE lifts was not different from that in the STAND position.

Forward bending and the resultant moments were significantly affected by the posture adopted (as can be seen in table 3); however, neither lateral bending nor twisting moments were significantly affected by posture ( $p > 0.05$ ). The forward bending moment ( $M_x$ ) was significantly higher in the STOOP than in other postures, and significantly lower in the kneeling conditions than in the other postures ( $p < 0.05$ ). As the forward bending moment in lifting tasks dominates the resultant moment, it is not surprising that much the same result is seen with the resultant. The only difference is that the resultant moment experienced when stooping was not significantly different from that observed when lifting on one knee. This appears to be the result of the increased lateral bending moment experienced when lifting on one knee.

### 3.2. Effects due to increased cable load

Figures 6–8 contain data describing the effects of increased cable load on EMG data and predicted spine loading. As reported in tables 2 and 3, the 100 N increase in cable load caused significant increases in the activity of all measured trunk muscles,

as well as in the spinal loading variables ( $p < 0.05$ ). However, in neither case did the increase result in a differential response according to the posture adopted in the performance of the cable-lifting task.




Figure 6. Effects of cable load on normalized maximum EMG activity of 10 trunk muscles (LLD = left latissimus dorsi, RLD = right latissimus dorsi, LES = left erector spinae, RES = right erector spinae, LEO = left external obliques, REO = right external obliques, LRA = left rectus abdominis, RRA = right rectus abdominis, LIO = left internal obliques, RIO = right internal obliques).




Figure 7. Effects of cable load on predicted peak forces acting on the lumbar spine ( $F_x$  = lateral shear force,  $F_y$  = anterior-posterior shear forces, and  $F_z$  = compression).




Figure 8. Effects of cable load on predicted peak spinal moments ( $M_x$  = sagittal bending moment,  $M_y$  = lateral bending moment,  $M_z$  = axial twisting moment,  $M_r$  = resultant moment).

Increased cable load resulted in a generalized increase in all 10 trunk muscles (table 2). The extensor muscles (including the erector spinae, the latissimus dorsi, and the internal obliques) showed increases of approximately 15–20% in their response to the increased load. A greater percentage increase was observed with the flexors (35–45% increase in activity), but these muscles remained at lower levels of normalized activity compared to the extensors.

Analysis of the forces acting on the lumbar spine indicated that the increase in cable load resulted in significant spinal force increases in all directions (as shown in figures 7 and 8). Compressive loading was increased by approximately 435 N, while lateral shear and A-P shear were increased by 138 N and 280 N, respectively. Increases in the forward bending, lateral bending, and axial twisting moments due to the increased cable load were 16, 30, and 21 Nm, respectively. Interestingly, the forward bending moment was not significantly increased as a result of the augmented cable load ( $p = 0.09$ ), while the increases in lateral bending and axial twisting moments were both statistically significant ( $p < 0.05$ ). The peak resultant moment was increased by 36 Nm with the additional load, the difference being significant at the 0.05 level.

#### 4. Discussion

The results of this study show that both posture and cable load significantly affect trunk muscle recruitment, which, in turn, influences the forces and moments imposed on the lumbar spine. However, posture and load appear to affect muscle recruitment (and subsequent spinal loads) in quite different fashions. Increases in the cable load provoked a broad increase in response from all 10 trunk muscles. The result of this change in recruitment was a more complex pattern of spine loading, characterized by increased off-plane moments (axial twisting and lateral bending) in what was a sagittally symmetric lifting task. Changes in posture altered the response of fewer

trunk muscles, although these were often influential extensor muscles that had substantial impact on spine loading estimates. In fact, the increase in spinal compression between the least stressful posture (KNEEL2) and the most stressful (STOOP) was nearly two-fold. However, the increase in spinal loading resulting from posture did not produce higher off-plane moments (axial twisting and lateral bending). Instead, the postural changes affected only moments in the sagittal plane.

It was hypothesized that the demands on the trunk musculature from postural changes and increased load might interact with one another in this study. However, the effects of these variables were remarkably independent from one another (and none of the 17 dependent measures disclosed such an interaction). This implies that, no matter which posture is adopted, the body's response to the increased load is the same, and appears to be characterized by increased activity of all available muscular resources. Similarly, if one handles a given load but changes the posture of the body, the body's response is a more selective adjustment of available muscle resources. Moreover, these effects appear to be additive in terms of both muscular recruitment and the ensuing spinal load.

The effects of increased load on trunk muscle EMG and spine loading in this study are similar to those of increased weight seen in studies of box lifting in the unrestricted standing posture (Fathallah *et al.* 1998, DeLooze *et al.* 1999, Davis and Marras 2000). Results of the current study suggest that the body's trunk muscle recruitment response to increased load is not dependent on posture, but always appears to be associated with a broad increase in activation of both extensor and flexor muscle groups.

It is easy to imagine that higher load conditions will create a situation where small changes in the position of the load may quickly lead to instability of the lumbar spinal column, due to high magnitude and rapidly changing moments affecting the spine. It may be that the broad co-contraction of trunk muscles in this situation provides an environment where muscles can respond more effectively to rapidly changing moments and that this may help to limit excessive movement of the spine under high moment conditions. In any event, the result of the increased load results is a more intricate pattern of loading on the spine, with notably higher shear forces that may be difficult for the spinal column to safely withstand (Yingling and McGill 1999). The abdominal muscles have variously been predicted to account for about 6–45% of the compressive load experienced by the spine (Potvin *et al.* 1991, Granata and Marras 1995, DeLooze *et al.* 1999). Their recruitment may be important in terms of increasing the stability of the spine in response to the increased load (Cholewicki and McGill 1996).

The stooping position was associated with high bilateral EMG activation of the erectores spinae and latissimus dorsi musculature, along with increased internal oblique activity. It should be noted that these are all extensor muscles of the spine, and that all appeared to be highly recruited in these tasks based on the need to balance the large external moment associated with the criterion task. It is apparent that subjects in this posture performed sufficient trunk extension to go beyond the range of the flexion-silence phenomenon (Floyd and Silver 1955), placing the burden of generating a significant restorative moment on the extensor musculature as noted above. The result of this activation pattern was that the stooping posture consistently showed the greatest spinal loads.

The 2KNEE lifting posture in this study resulted in the lowest spinal loading. Forces were lower in all three axes when this posture was adopted. The forward

bending moment was markedly lower than in any of the alternative postures. In addition, this was the only posture where the mean spinal compression was close to the 3400 N maximum acceptable limit recommended by NIOSH (Waters *et al.* 1993); however, shear forces were still high in this posture. Activity of all trunk extensors (erectores spinae, latissimus dorsi, and internal obliques) was diminished in the 2KNEE cable lifts. It appears that this posture may have allowed subjects to keep the load close to the body (thereby reducing the forward bending moment), and may have reduced this moment even further by allowing the subject to maintain a more upright trunk orientation throughout the lift, thereby decreasing the demands on the trunk extensor muscles to develop a high restorative moment. Comparison of the 1KNEE versus 2KNEE lifting postures revealed a sizeable difference in spinal loading, at least for the condition under study. The raised knee presents a barrier so significant as to increase the moment by 43% when compared to the 2KNEE condition.

Results of this study suggest that the stooping posture should be avoided whenever possible when working in restricted workspace. Such a recommendation is difficult to implement in practice, however, due to the higher strength capabilities in this posture when compared to kneeling. When loads are heavy, workers opt to use the stoop posture because it represents a position where the body can impart considerable force in an effort to move an object. Unfortunately, the results of this and other studies indicate that the costs in terms of spinal loading are severe, especially with respect to shear forces. Results of recent studies indicate that if one must adopt a stoop posture, it may be of benefit to avoid the end-range of spinal motion and to attempt to maintain some degree of lordosis in the lumbar spine region (Adams and Hutton 1982, Potvin *et al.* 1991, McGill and Kippers 1994, McGill 1999). Doing this may allow the paraspinal muscles to remain active and reduce the anterior shear forces observed when the interspinous ligaments assume increased reliance in supporting the spine.

The magnitudes of the forces and moments associated with lifting the cable in this experiment were all quite high, and this may help to explain the high incidence of lost-time back injuries in the coal mining industry associated with workers who perform this task. The NIOSH criterion for recommended weight limit (RWL) is based upon a 3400 N maximum compressive load on the spine (Waters *et al.* 1993). However, recent evidence suggests that injuries resulting from anterior shear loadings (primarily to the pars interarticularis, annulus and vertebral endplates) may begin to occur at shear loadings of only 900 N (McGill 1999, Yingling and McGill 1999). The fact that the mean compression for all experimental conditions exceeded recommended values (Waters *et al.* 1993) testifies to the high level of exertion required when handling such large diameter electrical cable. Efforts should be made to provide workers with mechanical assistance when performing this demanding task.

## 5. Conclusions

Results of this experiment support the following conclusions. First, changes in posture and cable load both influenced trunk muscle recruitment and spinal loading; however, these effects were not interactive. Second, increased cable load resulted in significantly increased activity of all trunk muscles, and resulted in increased off-plane (non-sagittal) moments on the spine. Third, changes in posture resulted in modifications in trunk muscle recruitment that were more selective than those

associated with increased load. That is, changes in posture typically affected the activity of only a few muscles, rather than the entire set. However, the muscles affected by changes in posture often involved the powerful extensors and therefore had a large impact in terms of spine loading. Fourth, kneeling on both knees was the least stressful posture in terms of spine loading and stooping was most stressful, while standing and kneeling on one knee involved a level of spinal loading intermediate between these two. Finally, the magnitudes of compression and shear loading on the spine were always quite high when lifting the cable regardless of the load condition or posture. Efforts should be made to provide mechanical assistance in performance of this task.

### References

- ADAMS, M. A. and HUTTON, W. C. 1982, Prolapsed intervertebral disc: a hyperflexion injury, *Spine*, **7**, 184–190.
- CALDWELL, L. S., CHAFFIN, D. B., DUKES-DOBOS, F. N., KROEMER, K. H. E., LAUBACH, L. L., SNOOK, S. H. and WASSERMAN, D. E. 1974, A proposed standard procedure for static muscle strength testing, *American Industrial Hygiene Association Journal*, **35**, 201–206.
- CHAFFIN, D. B. and ANDERSSON, G. B. J. 1991, *Occupational Biomechanics*, 2nd edn (New York: John Wiley).
- CHOLEWICKI, J. and MCGILL, S. M. 1996, Mechanical stability of the *in vivo* lumbar spine: implications for injury and chronic low back pain, *Clinical Biomechanics*, **11**, 1–15.
- DAVIS, K. G. and MARRAS, W. S. 2000, Assessment of the relationship between box weight and trunk kinematics: does a reduction in box weight necessarily correspond to a decrease in spine loading? *Human Factors*, **42**, 195–208.
- DAVIS, K. G., MARRAS, W. S. and WATERS, T. R. 1998, Evaluation of the spinal loading during lowering and lifting, *Clinical Biomechanics*, **13**, 141–152.
- DE LOOZE, M. P., GROEN, H., HOREMANS, H., KINGMA, I. and VAN DIEËN, J. H. 1999, Abdominal muscles contribute in a minor way to peak spinal compression in lifting, *Journal of Biomechanics*, **32**, 655–662.
- DUL, J. 1986, Muscular coordination in working postures, in N. Corlett, J. Wilson. and I. Manenica (eds), *The Ergonomics of Working Postures* (London: Taylor & Francis), 115–125.
- FATHALLAH, F. A., MARRAS, W. S. and PARNIANPOUR, M. 1998, An assessment of complex spinal loads during dynamic lifting tasks, *Spine*, **23**, 706–716.
- FATHALLAH, F. A., MARRAS, W. S., PARNIANPOUR, M. and GRANATA, K. P. 1997, A method for measuring external loads during unconstrained free-dynamic lifting, *Journal of Biomechanics*, **30**, 975–978.
- FLOYD, W. F. and SILVER, P. H. S. 1955, The function of the erector spinae muscles in certain movements and postures in man, *Journal of Physiology*, **129**, 184–203.
- GALLAGHER, S. 1997, Trunk extension strength and trunk muscle activity in standing and kneeling postures, *Spine*, **22**, 1864–1872.
- GALLAGHER, S. 1999, *Ergonomics Issues in Mining*, in W. Karwowski and W. S. Marras (eds), *Handbook of Occupational Ergonomics* (Boca Raton, FL: CRC Press), 1889–1911.
- GALLAGHER, S. and HAMRICK, C. A. 1992, Acceptable workloads for three common mining materials, *Ergonomics*, **35**, 1013–1031.
- GALLAGHER, S. and UNGER, R. L. 1990, Lifting in four restricted lifting conditions, *Applied Ergonomics*, **21**, 237–245.
- GALLAGHER, S., MARRAS, W. S. and BOBICK, T. G. 1988, Lifting in stooped and kneeling postures: effects on lifting capacity, metabolic costs, and electromyography of eight trunk muscles, *International Journal of Industrial Ergonomics*, **3**, 65–76.
- GALLAGHER, S., HAMRICK, C. A., CORNELIUS, K. M. and REDFERN, M. S. 2001, The effects of restricted workspace on lumbar spine loading, *Occupational Ergonomics*, **2**, 201–213.
- GALLAGHER, S., HAMRICK, C. A., LOVE, A. C. and MARRAS, W. S. 1994, Dynamic biomechanical modeling of symmetric and asymmetric lifting tasks in restricted postures, *Ergonomics*, **37**, 1289–1310.

- GRANATA, K. P. and MARRAS, W. S. 1993, An EMG-assisted model of loads on the lumbar spine during asymmetric trunk extensions, *Journal of Biomechanics*, **26**, 1429–1438.
- GRANATA, K. P. and MARRAS, W. S. 1995, An EMG-assisted model of trunk loading during free-dynamic lifting, *Journal of Biomechanics*, **28**, 1309–1317.
- KIRK, R. E. 1982, *Experimental Design: Procedures for the Behavioral Sciences*, 2nd edn (Pacific Grove, CA: Brooks/Cole), 911 pp.
- MARRAS, W. S. 1990, Industrial electromyography, *International Journal of Industrial Ergonomics*, **6**, 89–93.
- MARRAS, W. S. and GRANATA, K. P. 1995, A biomechanical assessment and model of axial twisting in the thoraco-lumbar spine, *Spine*, **20**, 1440–1451.
- MARRAS, W. S. and GRANATA, K. P. 1997, The development of an EMG-assisted model to assess spine loading during whole-body free-dynamic lifting, *Journal of Electromyography and Kinesiology*, **7**, 259–268.
- MARRAS, W. S. and MIRKA, G. A. 1993, Electromyographic studies of the lumbar trunk musculature during the generation of low-level trunk acceleration, *Journal of Orthopaedic Research*, **11**, 811–817.
- MARRAS, W. S. and REILLY, C. H. 1988, Networks of internal trunk loading activities under controlled trunk motion conditions, *Spine*, **13**, 661–667.
- MARRAS, W. S. and SOMMERICH, C. M. 1991a, A three dimensional motion model of loads on the lumbar spine. I. Model structure, *Human Factors*, **33**, 129–137.
- MARRAS, W. S. and SOMMERICH, C. M. 1991b, A three dimensional motion model of loads on the lumbar spine. II. Model validation, *Human Factors*, **33**, 139–149.
- MARRAS, W. S., FATHALLAH, F. A., MILLER, R. J., DAVIS, S. W. and MIRKA, G. A. 1992, Accuracy of a three-dimensional lumbar motion monitor for recording dynamic trunk motion characteristics, *International Journal of Industrial Ergonomics*, **9**, 75–87.
- MARRAS, W. S., LAVENDER, S. A., LEURGANS, S. E., RAJULU, S. L., ALLREAD, W. G., FATHALLAH, F. A. and FERGUSON, S. A. 1993, The role of dynamic three-dimensional motion in occupationally-related low back disorders. The effects of workplace factors, trunk position, and trunk motion characteristics on risk of injury, *Spine*, **18**, 617–628.
- MCGILL, S. M. 1999, Dynamic low back models: theory and relevance in assisting the ergonomist to reduce risk of low back injury, in W. Karwowski and W. S. Marras (eds), *The Occupational Ergonomics Handbook* (Boca Raton, FL: CRC Press), 945–965.
- MCGILL, S. M. and KIPPERS, V. 1994, Transfer of loads between lumbar tissues during the flexion relaxation phenomenon, *Spine*, **19**, 2190–2196.
- MCGILL, S. M. and NORMAN, R. W. 1986, Partitioning of the L4-L5 dynamic moment into disc, ligamentous, and muscular components during lifting, *Spine*, **11**, 666–678.
- MIRKA, G. A. and MARRAS, W. S. 1993, A stochastic model of trunk muscle coactivation during trunk bending, *Spine*, **18**, 1396–1409.
- NIOSH (NATIONAL INSTITUTE FOR OCCUPATIONAL SAFETY AND HEALTH) 1991, *Selected Topics in Surface Electromyography for Use in the Occupational Setting: Expert Perspectives*. DHHS(NIOSH) Publication No. 91-100, NIOSH Technical Report (Cincinnati, OH: NIOSH).
- POTVIN, J. R., MCGILL, S. M. and NORMAN, R. W. 1991, Trunk muscle and lumbar ligament contributions to dynamic lifts with varying degrees of trunk flexion, *Spine*, **16**, 1099–1107.
- RANDOLPH, R. F. 1991, Unpublished data. Available upon request from NIOSH, PO Box 18070, Pittsburgh, PA 15236, USA.
- REILLY, C. H. and MARRAS, W. S. 1989, SIMULIFT: a simulation model of human trunk motion during lifting, *Spine*, **14**, 5–11.
- SMITH, J. L., AYOUB, M. M., SELAN, J. L., KIM, H. K. and CHEN, H. C. 1989, Manual materials handling in unusual postures, in A. Mital (ed.), *Advances in Industrial Ergonomics and Safety I* (London: Taylor & Francis), 685–691.
- WATERS, T. R., PUTZ-ANDERSON, V., GARG, A. and FINE, L. J. 1993, Revised NIOSH equation of the design and evaluation of manual lifting tasks, *Ergonomics*, **36**, 749–776.
- YINGLING, V. R. and MCGILL, S. M. 1999, Anterior shear of spinal motion segments: kinematics, kinetics, and resultant injuries observed in a porcine model, *Spine*, **24**, 1882–1889.