

## Effects of melatonin and age on gene expression in mouse CNS using microarray analysis

Edward H. Sharman<sup>a,\*</sup>, Stephen C. Bondy<sup>a</sup>, Kaizhi G. Sharman<sup>a</sup>,  
Debomoy Lahiri<sup>c</sup>, Carl W. Cotman<sup>b</sup>, Victoria M. Perreau<sup>b,1</sup>

<sup>a</sup> Center for Occupational and Environmental Health, Department of Community and Environmental Medicine,  
University of California, Irvine, CA 92697-1825, USA

<sup>b</sup> Institute for Brain Aging and Dementia, University of California, Irvine, CA 92697, USA

<sup>c</sup> Institute of Psychiatric Research, Indiana University School of Medicine, Indianapolis, IN 46202, USA

Received 13 June 2006; received in revised form 29 August 2006; accepted 6 September 2006

Available online 21 November 2006

### Abstract

The expression levels of a number of genes associated with inflammation and immune function change with advancing age. Melatonin modulates gene expression levels of several of these genes. Therefore the declining levels of melatonin associated with age may play a role in the physiological effects of aging. We used oligonucleotide microarrays to measure age-related changes in mRNA expression in the murine CNS, and to study the effect of prolonged administration of dietary melatonin upon these changes. CB6F1 male mice were fed 40 ppm melatonin for 2.1 months prior to sacrifice at age 26.5 months, and compared with both age-matched controls and young, 4.5-month-old untreated controls. Total RNA was extracted from whole brain (excluding cerebellum and brain stem) and individual samples were hybridized to Affymetrix Mouse 430-2.0 arrays. The expression of a substantial number of genes was modulated by melatonin treatment and changes in selected genes were validated by quantitative reverse transcription polymerase chain reaction (qRT-PCR). A subset of these genes did not change with age. Conversely, some genes modulated by age were also modulated by melatonin treatment. In general, melatonin treatment drove the expression levels of these genes closer to the expression levels detected in the younger animals. Notably, the abundance of lipocalin 2 (Lcn2) mRNA increased with age and was decreased in old animals treated with melatonin. Lcn2 is a member of the acute phase response family of proteins and its mRNA levels in the brain increase in response to inflammation. Many of the genes with expression reduced by melatonin are involved in inflammation and the immune system. This suggests that melatonin treatment may influence the inflammatory responses of old animals, driving them to resemble more closely those occurring in young animals.

© 2006 Elsevier Ltd. All rights reserved.

**Keywords:** Aging; Brain; Diet; Melatonin; mRNA; Immunity; Microarray

### 1. Introduction

Markers of chronic inflammation and oxidative damage increase in the mammalian CNS with age, whereas melatonin has demonstrated antioxidant, anti-inflammatory and immunomodulatory properties (Zhang et al., 1997; Reiter et al., 2000; Sharman et al., 2002a,b, 2004; Pei and Cheung, 2004). The pulsatile nocturnal peak level of melatonin decreases with age

in rats (Pang et al., 1990), mice (Lahiri et al., 2004a) and humans (Tozawa et al., 2003), and is predicted to be involved in regulation of sleep, which is commonly disturbed in the elderly (Foley et al., 1995).

The only well-established means of lengthening the lifespan of animals is caloric restriction, and notably, short-term caloric restriction increases melatonin levels in brain and gut (Bubenik et al., 1992). Moreover, dietary supplementation with melatonin increases the lifespan of mice (Pierpaoli and Regelson, 1994; Anisimov et al., 2000) and rats (Oaknin-Bendahan et al., 1995). Thus it is reasonable to conclude that melatonin may play a role in promoting life extension. However, little is known of the mechanisms by which melatonin exerts these effects.

\* Corresponding author. Tel.: +1 949 824 5432; fax: +1 949 824 2793.

E-mail address: [esharman@uci.edu](mailto:esharman@uci.edu) (E.H. Sharman).

<sup>1</sup> Current address: Centre for Neuroscience, University of Melbourne, Parkville, VIC 3010, Australia.

Melatonin is secreted by the pineal gland, which in mice is located atop the cerebral cortex but outside the blood–brain barrier. Melatonin receptor mRNA is detected in a number of areas within the CNS, including the pituitary pars tuberalis, the suprachiasmatic nucleus of the hypothalamus (von Gall et al., 2002), the hippocampus and cerebellum (Al-Ghoul et al., 1998). In the rat, melatonin receptor density decreases with age in the hypothalamus and the hippocampus (Laudon et al., 1988), and melatonin supplementation increases both melatonin levels (Menendez-Pelaez et al., 1993; Lahiri et al., 2004a) and melatonin binding (Oaknin-Bendahan et al., 1995) in the brains of older animals. In the current study we sought to gain a better understanding of melatonin's possible neuroprotective mechanisms by comparing the age-related CNS gene expression patterns of young and old mice and old mice receiving dietary melatonin. We sought to improve the understanding of the aging process in the brain generally; thus we measured gene expression changes related to aging and melatonin common to the cerebrum as a whole, rather than inquire into detailed changes occurring in specific regions.

## 2. Experimental procedures

### 2.1. Animal treatment

Male CB6F1 mice, a hybrid between C57BL/6JM and BALB/cJF from Harlan Labs (Indianapolis, IN), aged 4.5 months (young group, YC) and 26.5 months (old group, OC), were housed two to four per cage and maintained on a 12-h light/12-h dark cycle in a temperature controlled ( $22 \pm 1^\circ\text{C}$ ) room. The CB6F1 hybrid was used in order to take advantage of the vigor – increased disease resistance, better survival under stress and greater natural longevity – typical of hybrids, while maintaining genetic similarity to the published C57BL/6 mouse genome sequence (Waterston et al., 2002). Food and water were provided *ad libitum*. Young (YC) and old (OC) control animals were fed a pelleted minimal basal diet (AIN-93M, Dyets #100900, Dyets Inc., Bethlehem, PA) consisting of 10% sucrose and 14% casein (w/w) as well as a minimal salt and vitamin mix. This basal diet was supplemented with 40 ppm (w/w) melatonin (Sigma–Aldrich, St. Louis, MO) in a separate group of old mice (OM). The dosage level was confirmed by independent testing (Irvine Nutri-Chemical Laboratories, Irvine, CA). This dosage was chosen because we have shown previously that it lowers levels of reactive oxygen species and improves activity in older mice (Bondy et al., 2002). 9.3 weeks later, young mice were sacrificed at 4.5 months of age, old at 26.5 months of age.

Separately caged sentinel animals were maintained in the same room to monitor for the occurrence of infectious disease; no disease was detected during the treatment period. All experiments were approved by the Institutional Animal Care and Use Committee at the University of California, Irvine, and conformed to the National Institute of Health guide for the care and use of laboratory animals.

### 2.2. RNA extraction and purification

Mice were killed by cervical dislocation between 10:40 and 14:40 h and were visually inspected for signs of disease or other pathology; brain tissues were excised quickly, immediately frozen on dry ice and stored at  $-80^\circ\text{C}$ . Cerebellum and brain stem were removed; total RNA was extracted from the remaining tissue using the TRI Reagent kit (Molecular Research Center Inc., Cincinnati, OH), following the manufacturer's protocol. Aliquots of the total RNA were further purified on an RNeasy column (Qiagen Inc., Valencia, CA) to yield a 260–280 nm absorbance ratio of  $\geq 1.9$ . RNA concentrations were determined by absorption at 260-nm wavelength with an ND-1000 spectrometer (Nanodrop Technologies, Wilmington, DE).

### 2.3. Gene array processing and analysis

A 20  $\mu\text{g}$  aliquot of total RNA from each of three animals per treatment group was processed and applied to a Mouse Genome 430 2.0 GeneChip<sup>®</sup> array (45,037 probe sets) according to the manufacturer's protocol (Affymetrix, Santa Clara, CA). Briefly, total RNA was reverse transcribed using a T7-Oligo(dT) Promoter Primer in the first-strand cDNA synthesis reaction. Following RNase H-mediated second-strand cDNA synthesis, the double-stranded cDNA was purified and served as a template for *in vitro* transcription (IVT). The IVT reaction was carried out in the presence of T7 RNA polymerase and a biotinylated nucleotide analog/ribonucleotide mix for complementary RNA (cRNA) amplification and biotin labeling. The biotinylated cRNA targets were then cleaned up, fragmented, and hybridized to GeneChip<sup>®</sup> expression arrays according to the manufacturer's instructions. All reagents used were as supplied in the manufacturer's One-Cycle Target Labeling and Control Reagents kit. The background, noise, housekeeper gene levels and 3'/5' ratio values of all chips were within the quality control limits set by Affymetrix (Supplementary Table A). Micro-array data are being submitted to GEO. Normalized expression values from the raw data were generated using default settings for the Robust Multi-array Average (RMA) method (Irizarry et al., 2003) within GeneSpring (Silicon Genetics, Redwood City, CA). Subsequent statistical analysis was also performed in GeneSpring. The cross-gene error model was applied with replicates; poor-quality probe sets with *p*-values  $> 0.2$  were eliminated. In order to focus only on genes that could be verified readily by qRT-PCR, probe sets with expression levels in the lowest 5%, or which had expression changes  $< 30\%$  between the OC and OM groups, were dropped from consideration.

### 2.4. qRT-PCR gene expression analysis

qRT-PCR analysis was used to measure the expression levels of selected mRNA transcripts. Methylenetetrahydrofolate dehydrogenase 1 (Mthfd1) was chosen as a control gene as it has moderate expression levels in this tissue and its expression level has previously shown to remain consistent under a variety of experimental manipulations in similar tissues (Adlard et al., 2004). Reactions utilizing equal amounts of total RNA from each sample were carried out on a LightCycler instrument (Roche Diagnostics, Indianapolis, IN) using the QuantiTect SYBR Green RT-PCR reagent kit (Qiagen, Valencia, CA) according to the manufacturers' directions. Product fluorescence was detected at the end of the elongation cycle at  $72^\circ\text{C}$ . Melting curves all exhibited a single, sharp peak at a temperature characteristic of the primer pair used and none of the primer pairs produced amplicons in the absence of sample or reverse transcriptase. Expression levels for each sample, normalized to Young Control expression, were calculated as the average of two measurements, relative to Mthfd1 expression.

### 2.5. Data and statistical analyses

Expression ratios were calculated as the power-2 exponential of the  $\log_2$  differences. The acceptance criterion for gene array expression changes was a one-way Analysis of Variance (ANOVA) *t*-test *p*-value of  $< 0.05$ .

For qRT-PCR mRNA analyses, differences between groups were assessed by one-way ANOVA; this was followed by Gabriel's multiple comparisons procedure test as computed using the ClinStat program (Bland, 2000). In all cases, the acceptance level of significance was  $p < 0.05$  using a two-tailed distribution.

## 3. Results

The entire set of 45,037 probe sets was analyzed for expression changes between young and old animals and old animals fed diets supplemented with melatonin. In order to validate the results from the arrays, three of the genes were further studied by qRT-PCR. The results from these analyses broadly confirmed the arrays (Fig. 1). A representative




Fig. 1. qRT-PCR (gray bars) compared to microarray analysis (open bars) of cortical levels of TiPARP, Lcn2 and Sgk mRNA expression in 4.5- and 26.5-month-old mice (YC and OC, respectively). One group of 26.5-month-old mice (OM) received 40 ppm dietary melatonin for the preceding 9.3 weeks. Bars represent mean  $\pm$  S.E. ( $n = 3$ ) mRNA expression levels relative to YC = 1.0, and as adjusted by Mthfd1 'housekeeper' gene expression (qRT-PCR), or normalized with RMA (microarray).

experimental plot for serum and glucocorticoid-induced kinase (SGK) is shown in Fig. 2.

Analysis within GeneSpring yielded an age- or melatonin-related set of 432 probe sets with expression ratio changes in excess of  $\pm 30\%$  and with  $p < 0.05$  (Supplementary Table B). The 30% value was chosen so as to identify genes with the smallest changes that would have an acceptably small probability of being false positives and could be readily verified by qRT-PCR.

Expression ratios for multiple probe sets representing a single unique gene were averaged. All resulting genes, classified as either changing with melatonin alone, or with both age and melatonin, are listed (Tables 1 and 2). Expression levels of seven genes that increased with age were decreased by melatonin




Fig. 2. Quantitative, real-time RT-PCR plot for serum and glucocorticoid-induced kinase (SGK) mRNA expression. (a) Plot of SYBR green fluorescence intensity (530 nm, arbitrary units) vs. number of PCR cycles. A single, representative curve is shown for each treatment group; YC, young control; OC, old control; OM, old melatonin group. Indicated mRNA expression levels: OM  $\gg$  YC  $>$  OC. (b) Overlapping melting peaks of the above samples resulting from a first derivative plot of final product fluorescence vs. temperature (melting curve). Overlapping peaks and lack of other significant features indicate accurate, single-product amplification and lack of primer-dimer or other spurious product formation. The unresponsive curve at bottom of each plot was produced by a negative (no RNA) control sample.

treatment to approach levels in YC (Table 1); similarly, two genes were decreased with age and increased with melatonin treatment (Table 1). Four genes were decreased by both age and melatonin (Table 2). All of the seven genes whose expression was elevated with age and reduced by melatonin treatment have immune-related functionality; these included lipocalin 2, immunoglobulin heavy chain 6 (Igh-6, heavy chain of IgM), and immunoglobulin  $\kappa$  chain V8 (Igk-V8).

Four genes, including three stress-inducible heat shock proteins, were significantly decreased by age, and decreased further by melatonin treatment. Another set of four genes, including plasma membrane associated protein S3-12 and arrestin domain containing protein 2 (Arrdc2), were increased by both age and melatonin (Table 2).

Table 1

List of probe sets with significant age-related gene expression changes the directions of which were reversed in the melatonin-treated animals

Gene	GenBank accession numbers	Normalized expression			OC/YC	OM/OC	Function	Description
		YC	OC	OM				
Increased with age, decreased by melatonin								
	AV057155	0.29	7.12	1.39	<b>24.7</b>	<b>0.20</b>	Possibly immunity	Transcribed locus, similar to XP-132633.3 immunoglobulin kappa chain (V28)
Igk-V8	BC013496	0.50	4.03	1.26	<b>8.06</b>	<b>0.31</b>	Immunity	Immunoglobulin kappa chain variable 8
	BI107286	0.66	3.50	1.17	<b>5.29</b>	<b>0.34</b>		
	BG966217	0.77	2.19	1.10	<b>2.85</b>	<b>0.50</b>		
Lcn2	X14607	0.74	3.01	1.13	<b>4.09</b> (21.0)	<b>0.38</b> (0.15)	Immunity	Lipocalin 2
Igh-6	BB226392	0.71	2.02	1.04	<b>2.85</b>	<b>0.52</b>	Immunity	Immunoglobulin IgM heavy chain 6
	AI326478	0.78	1.72	1.02	<b>2.21</b>	<b>0.60</b>		
Lrg1	NM_029796	0.59	1.44	0.98	<b>2.44</b>	<b>0.68</b>	Immunity	Leucine-rich $\alpha$ -2-glycoprotein 1
Igj	BC006026	0.83	1.62	1.07	<b>1.96</b>	<b>0.66</b>	Immunity	Immunoglobulin joining chain
Gvin 1	BM243571	0.92	1.20	0.89	<b>1.31</b>	<b>0.75</b>	Immunity	GTPase, very large interferon inducible 1; ligs 1; Vlig-1
Decreased with age, increased by melatonin								
Tiparp	BB762264	1.06	0.72	0.96	<b>0.68</b> (0.65)	<b>1.34</b> (1.18)	Histone ribosylation	TCDD-inducible poly(ADP-ribose) polymerase
Agxt211	BC022644	1.03	0.77	1.03	<b>0.75</b>	<b>1.34</b>	Transaminase	Alanine-glyoxylate aminotransferase 2-like 1

Probe sets are identified by gene abbreviations and GenBank accession numbers. Normalized expression values from the raw microarray data were generated using the Robust Multi-array Average (RMA) method (Irizarry et al., 2003) within GeneSpring. OC/YC, ratio of expression in old control animals to expression in young control animals; OM/OC, ratio of expression in old melatonin-fed animals to expression in old control animals. Bold ratios determined by gene chip analysis; ratios in parentheses determined by qRT-PCR.

Genes whose expression levels were unaffected by age but increased by melatonin, included CCAAT/enhancer binding protein delta (Cebpd), DNA-damage-inducible transcript 4 (Ddit4), and pyruvate dehydrogenase kinase enzyme (Pdk4). Genes depressed by melatonin and unaltered with age included the oncogene, Ets-1 (Table 2).

We measured the mRNA expression of arylalkylamine *N*-acetyltransferase (AANAT), the rate-limiting enzyme involved in the conversion of serotonin to melatonin, and of the melatonin-specific receptors Mtnr1a and RORa by qRT-PCR. Aside from a non-significant tendency for melatonin to lower their expression slightly, none were significantly affected by age or melatonin (data not shown). The data imply that the levels of these genes may have remained unchanged by these treatments throughout the major part of the brain we sampled; it is also possible that substantial changes in one region were counter-balanced by opposing changes in another region.

The mRNA expression for a second melatonin receptor Mtnr1b was not detectable by qRT-PCR (data not shown; the Mouse Genome 430 GeneChip<sup>®</sup> has no probe sets corresponding to this transcript). Two possibilities are consistent with this result. One is that Mtnr1b is expressed at very low levels in a relatively limited brain region; Mtnr1b mRNA has been detected in rat hypothalamus, for example (Sallinen et al., 2005). A second possibility is that Mtnr1b is not expressed at detectable levels anywhere in brain, consonant with its non-detectability in ovine hypothalamus (Migaud et al., 2005).

## 4. Discussion

### 4.1. Genes changed by age and reversed by melatonin

#### 4.1.1. Immunoglobulins

All seven unique genes whose expression was both increased with age and reduced by melatonin treatment in aged animals are related to immune function. Genespring classifies as immune-related only 278 probe sets out of the total of over 40,000 measured. However, melatonin is additionally involved in a number of other age-modulated processes, such as antioxidant status and circadian rhythm signaling. While melatonin supplementation could be predicted to result in reversal of age-related increases in genes associated with any of these processes, only reversals of age-related increases in immunity-related genes were detected. Thus, for all seven genes that were elevated with age and depressed by melatonin treatment to be immune-related, is especially remarkable.

Four of these immune-related genes were immunoglobulins—Igk-V8, Igh-6, Igj, and an immunoglobulin-related Expressed Sequence Tag (EST). The gene array used has probe sets for at least 12 distinct immunoglobulin genes, with many redundancies, indicating that only a minority of immunoglobulin genes may be modulated by age and melatonin.

#### 4.1.2. Lipocalin 2

Lipocalin 2 (Lcn2/NGAL/24p3) sequesters iron by binding to lipophilic iron chelators as part of the innate immune response to bacterial infection (Flo et al., 2004). Lcn2 also plays a role in

Table 2

List of probe sets with significant melatonin-related gene expression changes

Gene name	GenBank accession code	OC/YC	OM/OC	Function	Description
Significantly increased by melatonin					
Sgk	NM_011361	<b>0.95</b> (0.89)	<b>2.12</b> (3.30)	Extracellular stress response	Serum/glucocorticoid regulated kinase
Rbm3	AY052560	<b>0.90</b> (0.88)	<b>1.73</b> (1.71)	Hypoxia and cold stress response	Possible splice variant of RNA-binding motif protein 3mRNA
Nfkbia	NM_010907	<b>0.96</b>	<b>1.70</b>	Inflammation, oxidative stress	Nuclear factor of kappa light chain gene enhancer in B-cells inhibitor
	AI462015	<b>0.92</b>	<b>1.58</b>		
Nfkbia	BB096843	<b>0.95</b>	<b>1.63</b>	Inflammation, oxidative stress	cDNA clone 9430057A10 3' similar to U57524 IκBα chain non-coding mRNA sequence
[EST]	BB427399	<b>1.22</b>	<b>1.59</b>		EST from <i>Mus musculus</i> adult male hippocampus cDNA
Pdk4	NM_013743	<b>0.94</b>	<b>1.54</b>	Fatty acid/carbohydrate metabolic switch	Pyruvate dehydrogenase kinase, isoenzyme 4
Ddit4	AKO17926	<b>0.83</b>	<b>1.54</b>	Stress and ROS response	DNA-damage-inducible transcript 4
[EST]	BE956710	<b>1.29</b>	<b>1.44</b>		<i>Mus musculus</i> transcribed sequence
Cebpd	BB831146	<b>1.19</b>	<b>1.42</b>	Acute phase response; fear modulation	CCAAT/enhancer binding protein (C/EBP), delta
Fkbp5	AI098139	<b>0.98</b>	<b>1.41</b>	Steroid receptor activation	FK506 binding protein 5
Rhou	AF378088	<b>1.15</b>	<b>1.32</b>	Cell shape	Ras homolog gene family, member U
Cirbp	NM_007705	<b>1.09</b>	<b>1.31</b>	Stress	Cold inducible RNA binding protein
[EST]	BI133499	<b>1.15</b>	<b>1.31</b>		EST from <i>Mus musculus</i> adult male corpora quadrigemina cDNA
Significantly decreased by melatonin					
[EST]	BI076714	<b>1.16</b>	<b>0.74</b>		NIA mouse newborn ovary cDNA library
					<i>Mus musculus</i> cDNA clone L0223B10 3', mRNA sequence
[EST]	BG066678	<b>1.04</b>	<b>0.75</b>	EST	NIA mouse 15K cDNA clone set <i>Mus musculus</i> cDNA clone H3044G12 3', mRNA sequence
Ets-1	BB151715	<b>1.14</b>	<b>0.76</b>	Oncogene	E26 avian leukemia oncogene 1, 5' domain
CD93	AV319144	<b>0.93</b>	<b>0.77</b>	Immunity	CD93 antigen; complement component 1, q subcomponent, receptor 1
Significantly increased by both age and melatonin					
S3-12	NM_020568	<b>1.94</b> (1.33)	<b>1.31</b> (3.61)	Lipid sequestration	Plasma membrane associated protein, S3-12
Arrdc2	AW542672	<b>1.38</b>	<b>1.57</b>	Unknown	Arrestin domain containing 2; llad 1
Alas2	M63244	<b>1.35</b>	<b>1.73</b>	Heme biosynthesis	Aminolevulinic acid synthase 2, erythroid
Hbb-b2	BC027434	<b>1.31</b>	<b>1.32</b>	Oxygen transport	Hemoglobin, beta adult minor chain
Significantly decreased by both age and melatonin					
Hspa1b	M12573	<b>0.35</b>	<b>0.69</b>	Stress	Heat shock protein 1B; Hsp70
	M12573	<b>0.39</b>	<b>0.67</b>		
Serpinhi	BI220012	<b>0.53</b> (0.57)	<b>0.76</b> (0.67)	Stress	Serine (or cysteine) proteinase inhibitor, Hsp47
Hspbi	U03561	<b>0.53</b>	<b>0.72</b>	Stress	Heat shock protein 1; Hsp25
Hist2h3c2	BC015270	<b>0.68</b>	<b>0.74</b>		Histone 2, H3c2

Probe sets are identified by gene abbreviations and GenBank accession numbers. Normalized expression values from the raw microarray data were generated using the Robust Multi-array Average (RMA) method (Irizarry et al., 2003) within GeneSpring. OC/YC, ratio of expression in old control animals to expression in young control animals; OM/OC, ratio of expression in old melatonin-fed animals to expression in old control animals. Bold ratios determined by gene chip analysis; ratios in parentheses determined by qRT-PCR.

normal iron metabolism (Elangovan et al., 2004). The proper functioning of Lcn2 may be especially important for myelin and oligodendrocytes, which contain particularly high levels of iron. Apoptosis of oligodendrocytes and myelin degeneration are both associated with brain aging, and are particularly pronounced in Alzheimer's disease (Bartzokis, 2004).

We observed significantly higher Lcn2 mRNA expression in old animals compared to young controls and this may relate to the age-induced accumulation of iron in the CNS (Zecca et al., 2004). Increased Lcn2 levels in other tissues are associated with

numerous pathological conditions; among these are psoriasis and squamous cell carcinoma in skin (Mallbris et al., 2002), cisplatin-induced nephrotoxicity (Mishra et al., 2004), and the hepatic acute phase inflammatory response (Liu and Nilsen-Hamilton, 1995). By analogy the age-related increases in Lcn2 expression seen here may have relevance to neural pathology, and the melatonin-induced Lcn2 reduction may help to maintain CNS functionality. Determining the extent to which Lcn2 protein levels parallel the mRNA changes is an important area for future study.



The promoter region of the murine *Lcn2* gene contains an activation site for NF- $\kappa$ B (Cowland et al., 2003), and levels of activated NF- $\kappa$ B increase in murine cortex with age (Ye and Johnson, 2001). Thus, increased levels of NF- $\kappa$ B activation in the aged brain may explain, at least in part, the increased levels of *Lcn2* mRNA observed. The increased levels of *Lcn2* in turn add support to the notion that the aged brain is in a chronically inflamed state relative to the young.

Two genes, *Tiparp* and *Agxt211*, had the opposite response: their expression levels decreased with age and were restored to young-animal levels by melatonin; in contrast to the genes with expression increased by age, neither is immune-related.

#### 4.2. Genes decreased by both age and melatonin

##### 4.2.1. Heat shock proteins

Heat shock proteins are molecular chaperones that facilitate the correct folding of other proteins. They are highly inducible and their increased CNS expression in response to a variety of pathological conditions confers neuroprotection (Yenari, 2002). Three of the four unique genes whose mRNA expression was decreased by both age and melatonin are heat shock proteins: *Hspa1b* (Hsp70), *Serpinh1* (Hsp47), and *Hspb1* (Hsp25). A fourth heat shock gene—*Dnajb11*, that previously has been reported to decrease significantly in aged mouse cortex (Jiang et al., 2001), declined by 20% in our study (microarray data not shown). However, age does not appear to decrease the expression of heat shock genes in all cases. The heat shock protein *Dnajb13* was reported to increase in C57BL/6 mouse cortex (Lee et al., 2000), paralleling our report of an age-related increase of Hsp70 expression in B6C3F1 mouse cortex (Sharman et al., 2004).

How heat shock gene expression changes affect animal fitness may vary with differing treatments. Given a greater requirement for repair of miss-folded proteins in aged cortex, the age-related decreased expression of these genes could result in a decline in brain function in aged animals. However, Lee et al. (2000) found that caloric restriction, a treatment widely found to result in life extension and improved function in aged animals, decreased the expression of *Dnajb13*; this decrease could be interpreted as a response to a decreased protein repair requirement in aged calorically restricted animals. We propose that the decrease in heat shock gene expression produced by melatonin, as observed in both this and our previous study, may be a response to a similarly improved condition.

#### 4.3. Genes altered by melatonin but not changed with age

##### 4.3.1. Serum/glucocorticoid-regulated kinase (*Sgk*)

Gene array measurements (confirmed by qRT-PCR analysis) revealed that melatonin treatment increased expression of serum/glucocorticoid-regulated kinase (*Sgk*) in aged mice above corresponding levels in young animals. Increased CNS levels of *Sgk* may be beneficial *via* reduction of excitotoxicity through stimulation of the glutamate transporter, EAAT4 (Bohmer et al., 2004). In cultured rat spinal cord neurons, the number of dendrites and dendritic branch points are increased

by increased *Sgk* gene expression (David et al., 2005). Higher hippocampal expression levels in rats correlate with faster spatial learning in the Morris water maze test (Tsai et al., 2002). Thus increased levels of *Sgk* expression in the CNS produced following melatonin supplementation may result in similar beneficial effects. Such effects may also contribute to improved learning in older humans observed with nightly administration of 1 mg of melatonin (Peck et al., 2004).

##### 4.3.2. *Nfkb*

NF- $\kappa$ B inhibitor  $\alpha$  (*Nfkb* $\alpha$ /I $\kappa$ B $\alpha$ ) prevents activation of the transcription factor NF- $\kappa$ B which regulates expression of a number of oxidative-stress-related and inflammatory mediators such as cyclooxygenase-2 (COX-2) (Wu, 2005). Increases in *Nfkb* expression may thus be anti-inflammatory; decreased expression is deleterious as *Nfkb* knockout mice are perinatally lethal (Cheng et al., 1998). Our present results showing an increased *Nfkb* expression with melatonin are consistent with a neuroprotective and anti-inflammatory role for this substance.

##### 4.3.3. DNA-damage-inducible transcript 4

Given melatonin's demonstrated neuroprotective properties in an *in vivo* stroke model (Kilic et al., 2005), upregulation of the stress-responsive gene DNA-damage-inducible transcript 4 (*Ddit4*/REDD1/dig2/RTP801) following melatonin treatment, is paradoxical. *Ddit4* gene expression is upregulated by hypoxia in cell culture and by ischemia in rat brain, but its increased expression is accompanied by increased toxicity in non-dividing neuron-like PC12 cells and protection of MCF7 breast carcinoma cells (Shoshani et al., 2002). Furthermore, *Ddit4* gene expression is increased by exposure of human neuroblastoma CHP134 cells to amyloid beta ( $A\beta$ ) protein, and transient overexpression of the sense gene heightened  $A\beta$  toxicity, while overexpression of the antisense gene reduced susceptibility to  $A\beta$  in these cells (Kim et al., 2003). These results are consonant with our failure to observe any improvement in cortical histopathology or memory after chronic feeding of melatonin in a triple-transgenic mouse model of Alzheimer's disease (Sharman, Bondy and LaFerla, unpublished results). However, in our earlier animal studies on non-transgenic mice, we observed a significant reduction in levels of  $A\beta$  peptides only in middle-aged (12 months old) compared with older mice (27 months old) (Lahiri et al., 2004b). The reason for the age dependency of effects of melatonin supplementation on levels of these proteins is unclear; nevertheless these data are consistent with the very modest benefits of melatonin for Alzheimer's disease patients in a recent clinical trial (Asayama et al., 2003).

##### 4.3.4. *C/EBP- $\delta$*

CCAAT/enhancer binding protein (*C/EBP- $\delta$* ) is a transcription factor widely expressed in mammalian CNS (Ramji and Foka, 2002). *C/EBP- $\delta$*  mRNA levels are elevated by melatonin supplementation and are expressed throughout the CNS, including the hippocampus. In one study, targeted gene deletion of *C/EBP- $\delta$*  produced an enhanced contextual fear

response in mice after 24 h, but not after 30 min (Sterneck et al., 1998). Conversely, hippocampal C/EBP- $\delta$  mRNA expression was increased 20 h after inhibitory avoidance training in the rat (Taubenfeld et al., 2001). These two studies point to an important role for this transcription factor in long-term memory consolidation, and the observed elevation of C/EBP- $\delta$  suggests melatonin supplementation has the potential to benefit memory consolidation.

Overexpression of C/EBP- $\delta$  led to accumulation of myelin protein zero (PO) and myelin basic protein (MBP) mRNAs in a Schwann cell myeloma cell culture model (Kamaraju et al., 2004). Loss of C/EBP- $\delta$  also promotes chromosomal instability in fibroblast cell culture (Huang et al., 2004) implying that it can function as a tumor suppressor. Consistent with this role for C/EBP- $\delta$  is our observation that untreated old animals exhibited a higher incidence of tumors than melatonin-supplemented ones.

#### 4.3.5. Pyruvate dehydrogenase kinase (pdk4)

Melatonin increased the mRNA expression for pyruvate dehydrogenase kinase (pdk4). Pdk4 phosphorylates the pyruvate dehydrogenase complex, resulting in a suppression of complex activity and a shift from carbohydrate to fatty acid metabolism. Pdk4 gene expression is also elevated in several tissues of the ground squirrel *Spermophilus tridecemlineatus* during hibernation (Buck et al., 2002); the parallel elevation of Pdk4 by melatonin is consistent with the finding that melatonin supplementation lengthens hibernation duration in the related species *Spermophilus lateralis* (Saarela and Reiter, 1994).

#### 4.3.6. *Ets-1*

The melatonin-induced decrease in *Ets-1* mRNA expression is consistent with melatonin being oncostatic (Saez et al., 2005). *Ets-1* protein is upregulated in many tumors including brain tumors such as astrocytomas and meningiomas, and the degree of expression upregulation correlates with tumor invasiveness (Dittmer, 2003). The light experienced by night shift workers suppresses their nocturnal melatonin secretion, and female night shift workers have increased breast cancer incidence (Davis et al., 2001). Moreover, urinary levels of melatonin's excretory product 6-sulfatoxymelatonin are decreased in breast cancer patients (Schernhammer et al., 2004).

#### 4.4. Broad relevance of results

The results reported here on immune function are in broad accord with our earlier study using a different gene array, different dietary melatonin schedule and a different mouse strain (Sharman et al., 2004). The generality of these findings is further confirmed by the good correlation obtained when comparing mouse and human genes that change markedly with age (Sharman et al., 2005).

Expression of genes for melatonin receptors was not significantly changed with age. The simplest interpretation of this is that the reduction of melatonergic activity occurring with age, leads to little overall alteration of receptor gene

expression. It is also possible that increased receptor gene expression in one part of the brain is negated by a decrease of similar magnitude in another part.

The protective effects of melatonin, together with its low toxicity and its capacity to increase mRNA levels of immune- and memory-related proteins, suggest that this hormone should be investigated further as a treatment of age-related neurological disorders.

#### Acknowledgment

This work was supported in part by grants from the National Institutes of Health (ES 7992, AG 16794, AG 14882 and AG 18884).

#### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.neuint.2006.09.001](https://doi.org/10.1016/j.neuint.2006.09.001).

#### References

- Adlard, P.A., Perreau, V.M., Engesser-Cesar, C., Cotman, C.W., 2004. The timecourse of induction of brain-derived neurotrophic factor mRNA and protein in the rat hippocampus following voluntary exercise. *Neurosci. Lett.* 363, 43–48.
- Al-Ghoul, W.M., Herman, M.D., Dubocovich, M.L., 1998. Melatonin receptor subtype expression in human cerebellum. *Neuroreport* 9, 4063–4068.
- Anisimov, V.N., Zavarzina, N., Zabezhinskii, M.A., Popovich, I.G., Anikin, I.V., Zimina, O.A., Solov'ev, M.V., Shtylik, A.V., Arutunian, A.V., Oparina, T.I., Prokopenko, V.M., Khavinson, V.Kh., 2000. The effect of melatonin on the indices of biological age, on longevity and on the development of spontaneous tumors in mice. *Vopr. Onkol.* 46, 311–319.
- Asayama, K., Yamadera, H., Ito, T., Suzuki, H., Kudo, Y., Endo, S., 2003. Double blind study of melatonin effects on the sleep–wake rhythm, cognitive and non-cognitive functions in Alzheimer type dementia. *J. Nippon Med. Sch.* 70, 334–341.
- Bartzokis, G., 2004. Age-related myelin breakdown: a developmental model of cognitive decline and Alzheimer's disease. *Neurobiol. Aging* 25, 5–18.
- Bland, M., 2000. *An Introduction to Medical Statistics*, vol. 3. Oxford University Press, Oxford, p. 177.
- Bohmer, C., Philippin, M., Rajamanickam, J., Mack, A., Broer, S., Palmada, M., Lang, F., 2004. Stimulation of the EAAT4 glutamate transporter by SGK protein kinase isoforms and PKB. *Biochem. Biophys. Res. Commun.* 324, 1242–1248.
- Bondy, S.C., Yang, Y.E., Walsh, T.J., Gie, Y.W., Lahiri, D.K., 2002. Dietary modulation of age-related changes in cerebral pro-oxidant status. *Neurochem. Int.* 40, 123–130.
- Bubenik, G.A., Ball, R.O., Pang, S.F., 1992. The effect of food deprivation on brain and gastrointestinal tissue levels of tryptophan, serotonin, 5-hydroxyindoleacetic acid, and melatonin. *J. Pineal Res.* 12, 7–16.
- Buck, M.J., Squire, T.L., Andrews, M.T., 2002. Coordinate expression of the PDK4 gene: a means of regulating fuel selection in a hibernating mammal. *Physiol. Genomics* 8, 5–13.
- Cheng, J.D., Ryseck, R.P., Attar, R.M., Dambach, D., Bravo, R., 1998. Functional redundancy of the nuclear factor kappa B inhibitors I kappa B alpha and I kappa B beta. *J. Exp. Med.* 188, 1055–1062.
- Cowland, J.B., Sorensen, O.E., Sehested, M., Borregaard, N., 2003. Neutrophil gelatinase-associated lipocalin is up-regulated in human epithelial cells by IL-1b, but not by TNF-a. *J. Immunol.* 171, 6630–6639.
- David, S., Stegenga, S.L., Hu, P., Xiong, G., Kerr, E., Becker, K.B., Venkatapathy, S., Warrington, J.A., Kalb, R.G., 2005. Expression of serum- and glucocorticoid-inducible kinase is regulated in an experience-dependent manner and can cause dendrite growth. *J. Neurosci.* 25, 7048–7053.

- Davis, S., Mirick, D.K., Stevens, R.G., 2001. Night shift work, light at night, and risk of breast cancer. *J. Natl. Cancer Inst.* 93, 1557–1562.
- Dittmer, J., 2003. The biology of the Ets1 proto-oncogene. *Mol. Cancer* 2, 29–49.
- Elangovan, N., Lee, Y.C., Tzeng, W.F., Chu, S.T., 2004. Delivery of ferric ion to mouse spermatozoa is mediated by lipocalin internalization. *Biochem. Biophys. Res. Commun.* 319, 1096–1104.
- Flo, T.H., Smith, K.D., Sato, S., Rodriguez, D.J., Holmes, M.A., Strong, R.K., Akira, S., Aderem, A., 2004. Lipocalin 2 mediates an innate immune response to bacterial infection by sequestering iron. *Nature* 432, 917–921.
- Foley, D.J., Monjan, A.A., Brown, S.L., Simonsick, E.M., Wallace, R.B., Blazer, D.G., 1995. Sleep complaints among elderly persons: an epidemiologic study of three communities. *Sleep* 18, 425–432.
- Huang, A.M., Montagna, C., Sharan, S., Ni, Y., Ried, T., Sterneck, E., 2004. Loss of CCAAT/enhancer binding protein delta promotes chromosomal instability. *Oncogene* 23, 1549–1557.
- Irizarry, R.A., Hobbs, B., Collin, F., Beazer-Barclay, Y.D., Antonellis, K.J., Scherf, U., Speed, T.P., 2003. Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4, 249–264.
- Jiang, C.H., Tsien, J.Z., Schultz, P.G., Hu, Y., 2001. The effects of aging on gene expression in the hypothalamus and cortex of mice. *Proc. Natl. Acad. Sci. U.S.A.* 98, 1930–1934.
- Kamaraju, A.K., Adjalley, S., Zhang, P., Chebath, J., Revel, M., 2004. C/EBP-delta induction by gp130 signaling. Role in transition to myelin gene expressing phenotype in a melanoma cell line model. *J. Biol. Chem.* 279, 3852–3861.
- Kilic, U., Kilic, E., Reiter, R.J., Bassetti, C.L., Hermann, D.M., 2005. Signal transduction pathways involved in melatonin-induced neuroprotection after focal cerebral ischemia in mice. *J. Pineal Res.* 38, 67–71.
- Kim, J.R., Lee, S.R., Chung, H.J., Kim, S., Baek, S.H., Kim, J.H., Kim, Y.S., 2003. Identification of amyloid b-peptide responsive genes by cDNA microarray technology: involvement of RTP801 in amyloid b-peptide toxicity. *Exp. Mol. Med.* 35, 403–411.
- Lahiri, D.K., Chen, D., Ge, Y.W., Bondy, S.C., Sharman, E.H., 2004a. Dietary supplementation with melatonin reduced levels of amyloid beta-peptides in the murine cerebral cortex. *J. Pineal Res.* 36, 224–231.
- Lahiri, D.K., Ge, Y.W., Sharman, E.H., Bondy, S.C., 2004b. Age-related changes in serum melatonin in mice: higher levels of combined melatonin and 6-hydroxymelatonin sulfate in the cerebral cortex than serum, heart, liver and kidney tissues. *J. Pineal Res.* 36, 217–223.
- Laudon, M., Nir, I., Zisapel, N., 1988. Melatonin receptors in discrete brain areas of the male rat. Impact of aging on density and on circadian rhythmicity. *Neuroendocrinology* 48, 577–583.
- Lee, C.K., Weindrich, R., Prolla, T.A., 2000. Gene expression profile of the ageing brain. *Nat. Genet.* 25, 294–297.
- Liu, Q., Nilsen-Hamilton, M., 1995. Identification of a new acute phase protein. *J. Biol. Chem.* 270, 22565–22570.
- Mallbris, L., O'Brien, K.P., Hulthen, A., Sandstedt, B., Cowland, J.B., Borregaard, N., Stahle-Backdahl, M., 2002. Neutrophil gelatinase-associated lipocalin is a marker for dysregulated keratinocyte differentiation in human skin. *Exp. Dermatol.* 11, 584–591.
- Menendez-Pelaez, A., Poeggeler, B., Reiter, R.J., Barlow-Walden, L., Pablos, M.I., Tan, D.X., 1993. Nuclear localization of melatonin in different mammalian tissues: immunocytochemical and radioimmunoassay evidence. *J. Cell. Biochem.* 53, 373–382.
- Migaud, M., Daveau, A., Malpaux, B., 2005. MTRN1A melatonin receptors in the ovine pre-mammillary hypothalamus: day–night variation in the expression of the transcripts. *Biol. Reprod.* 72, 393–398.
- Mishra, J., Mori, K., Ma, Q., Kelly, C., Barasch, J., Devarajan, P., 2004. Neutrophil gelatinase-associated lipocalin: a novel early urinary biomarker for cisplatin nephrotoxicity. *Am. J. Nephrol.* 24, 307–315.
- Oaknin-Bendahan, S., Anis, Y., Nir, I., Zisapel, N., 1995. Effects of long-term administration of melatonin and a putative antagonist on the ageing rat. *Neuroreport* 6, 785–788.
- Pang, S.F., Tsang, C.W., Hong, G.X., Yip, P.C., Tang, P.L., Brown, G.M., 1990. Fluctuation of blood melatonin concentrations with age: result of changes in pineal melatonin secretion, body growth, and aging. *J. Pineal Res.* 8, 179–192.
- Peck, J.S., LeGoff, D.B., Ahmed, I., Goebert, D., 2004. Cognitive effects of exogenous melatonin administration in elderly persons: a pilot study. *Am. J. Geriatr. Psychiatry* 12, 432–436.
- Pei, Z., Cheung, R.T., 2004. Pretreatment with melatonin exerts anti-inflammatory effects against ischemia/reperfusion injury in a rat middle cerebral artery occlusion stroke model. *J. Pineal Res.* 37, 85–91.
- Pierpaoli, W., Regelson, W., 1994. Pineal control of aging: effect of melatonin and pineal grafting on aging mice. *Proc. Natl. Acad. Sci. U.S.A.* 91, 787–791.
- Ramji, D.P., Foka, P., 2002. CCAAT/enhancer-binding proteins: structure, function and regulation. *Biochem. J.* 365, 561–575.
- Reiter, R.J., Tan, D.X., Qi, W., Manchester, L.C., Karbownik, M., Calvo, J.R., 2000. Pharmacology and physiology of melatonin in the reduction of oxidative stress in vivo. *Biol. Signals Recept.* 9, 160–171.
- Saarela, S., Reiter, R.J., 1994. Function of melatonin in thermoregulatory processes. *Life Sci.* 54, 295–311.
- Saez, M.C., Barriga, C., Garcia, J.J., Rodriguez, A.B., Ortega, E., 2005. Effect of the preventive-therapeutic administration of melatonin on mammary tumour-bearing animals. *Mol. Cell. Biochem.* 268, 25–31.
- Sallinen, P., Saarela, S., Ilves, M., Vakkuri, O., Leppaluoto, J., 2005. The expression of MT1 and MT2 melatonin receptor mRNA in several rat tissues. *Life Sci.* 76, 1123–1134.
- Schernhammer, E.S., Rosner, B., Willett, W.C., Laden, F., Colditz, G.A., Hankinson, S.E., 2004. Epidemiology of urinary melatonin in women and its relation to other hormones and night work. *Cancer Epidemiol. Biomarkers Prev.* 13, 936–943.
- Sharman, K.G., Sharman, E.H., Bondy, S.C., 2002a. Dietary melatonin selectively reverses age-related changes in cortical basal cytokine mRNA levels, and their responses to an inflammatory stimulus. *Neurobiol. Aging* 23, 633–638.
- Sharman, E.H., Vaziri, N.D., Ni, Z., Sharman, K.G., Bondy, S.C., 2002b. Reversal of biochemical and behavioral parameters of brain aging by melatonin and acetyl L-carnitine. *Brain Res.* 957, 223–230.
- Sharman, E.H., Sharman, K.G., Lahiri, D.K., Bondy, S.C., 2004. Age-related changes in murine CNS mRNA gene expression are modulated by dietary melatonin. *J. Pineal Res.* 36, 165–170.
- Sharman, E.H., Sharman, K.G., Bondy, S.C., 2005. Parallel changes in gene expression in aged human and mouse cortex. *Neurosci. Lett.* 390, 4–8.
- Shoshani, T., Faerman, A., Mett, I., Zelin, E., Tenne, T., Gorodin, S., Moshel, Y., Elbaz, S., Budanov, A., Chajut, A., Kalinski, H., Kamer, I., Rozen, A., Mor, O., Keshet, E., Leshkowitz, D., Einat, P., Skalter, R., Feinstein, E., 2002. Identification of a novel hypoxia-inducible factor 1-responsive gene, RTP801, involved in apoptosis. *Mol. Cell. Biol.* 22, 2283–2293.
- Sterneck, E., Paylor, R., Jackson-Lewis, V., Libbey, M., Przedborski, S., Tessarollo, L., Crawley, J.N., Johnson, P.F., 1998. Selectively enhanced contextual fear conditioning in mice lacking the transcriptional regulator CCAAT/enhancer binding protein delta. *Proc. Natl. Acad. Sci. U.S.A.* 95, 10908–10913.
- Taubenfeld, S.M., Wiig, K.A., Monti, B., Dolan, B., Pollonini, G., Alberini, C.M., 2001. Fornix-dependent induction of hippocampal CCAAT enhancer-binding proteins  $\beta$  and  $\delta$  co-localizes with phosphorylated cAMP response element-binding protein and accompanies long-term memory consolidation. *J. Neurosci.* 21, 84–91.
- Tozawa, T., Mishima, K., Satoh, K., Echizenya, M., Shimizu, T., Hishikawa, Y., 2003. Stability of sleep timing against the melatonin secretion rhythm with advancing age: clinical implications. *J. Clin. Endocrinol. Metab.* 88, 4689–4695.
- Tsai, K.J., Chen, S.K., Ma, Y.L., Hsu, W.L., Lee, E.H., 2002. Sgk, a primary glucocorticoid-induced gene, facilitates memory consolidation of spatial learning in rats. *Proc. Natl. Acad. Sci. U.S.A.* 99, 3990–3995.
- von Gall, C., Stehle, J.H., Weaver, D.R., 2002. Mammalian melatonin receptors: molecular biology and signal transduction. *Cell Tissue Res.* 309, 151–162.
- Waterston, R.H., Lindblad-Toh, K., Birney, E., et al., 2002. Initial sequencing and comparative analysis of the mouse genome. *Nature* 420, 520–562.



- Wu, K.K., 2005. Control of cyclooxygenase-2 transcriptional activation by pro-inflammatory mediators. *Prostaglandins Leukot. Essent. Fatty Acids* 72, 89–93.
- Ye, S.M., Johnson, R.W., 2001. Regulation of interleukin-6 gene expression in brain of aged mice by nuclear factor kappaB. *J. Neuroimmunol.* 117, 87–96.
- Yenari, M.A., 2002. Heat shock proteins and neuroprotection. *Adv. Exp. Med. Biol.* 513, 281–299.
- Zecca, L., Youdim, M.B., Riederer, P., Connor, J.R., Crichton, R.R., 2004. Iron, brain ageing and neurodegenerative disorders. *Nat. Rev. Neurosci.* 5, 863–873.
- Zhang, Z., Inseerra, P.F., Liang, B., Ardestani, S.K., Elliott, K.K., Molitor, M., Watson, R.R., 1997. Melatonin, immune modulation and aging. *Autoimmunity* 26, 43–53.