

support to the Centre for its operations and research. Current donors providing unrestricted support include the following: Government of the People's Republic of Bangladesh; Canadian International Development Agency; the Swedish International Development Cooperation Agency; the Swiss Agency for Development and Cooperation; and the Department for International Development, UK.

**Mustafizur Rahman,  
Shamsun Nahar,  
Mokibul Hassan Afrad,  
Abu S.G. Faruque,  
and Tasnim Azim**

Author affiliations: International Centre for Diarrhoeal Disease Research, Bangladesh, Dhaka, Bangladesh (M. Rahman, M.H. Afrad, A.S.G. Faruque, T. Azim); and Jahangirnagar University, Dhaka (S. Nahar)

DOI: <http://dx.doi.org/10.3201/eid1908.130227>

## References

- Green KY. Caliciviridae: the noroviruses. In: Fields BN, Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, et al, editors. *Fields virology*, 5th ed. Philadelphia: Lippincott, Williams & Wilkins; 2007. p. 949–79.
- van Beek J, Ambert-Balay K, Botteldoorn N, Eden J, Fonager J, Hewitt J, et al. Indications for worldwide increased norovirus activity associated with emergence of a new variant of genotype II.4, late 2012. *Euro Surveill*. 2013;18:8–9.
- Bennett S, Maclean A, Miller R, Aitken C, Gunson R. Increased norovirus activity in Scotland in 2012 is associated with the emergence of a new norovirus GII.4 variant. *Euro Surveill*. 2013;18: pii: 20349.
- Fonager J, Hindbæk LS, Fischer TK. Rapid emergence and antigenic diversification of the norovirus 2012 Sydney variant in Denmark, October to December, 2012. *Euro Surveill*. 2013;18:20413.
- National Institute of Infectious Diseases, Japan. Flash report of norovirus in Japan. Tokyo [cited 10 Jan 2013]. <http://www.nih.go.jp/niid/en/iasr-noro-e.html>
- White PA, Eden JS, Hansman GS. Molecular epidemiology of noroviruses and sapoviruses and their role in Australian outbreaks of acute gastroenteritis. *Microbiology Australia*. 2012;33:70–3 [cited 2013 Jun 5]. <http://journals.cambridge.org/med/Articles/08white409.pdf>
- Centers for Disease Control and Prevention. Emergence of new norovirus strain GII.4 Sydney—United States, 2012. *MMWR Morb Mortal Wkly Rep*. 2013;62:55.
- Trujillo AA, McCaustland KA, Zheng DP, Hadley LA, Vaughn G, Adams SM, et al. Use of TaqMan real-time reverse transcription-PCR for rapid detection, quantification, and typing of norovirus. *J Clin Microbiol*. 2006;44:1405–12. <http://dx.doi.org/10.1128/JCM.44.4.1405-1412.2006>
- Nayak MK, Balasubramanian G, Sahoo GC, Bhattacharya R, Vinje J, Kobayashi N, et al. Detection of a novel intergenogroup recombinant norovirus from Kolkata, India. *Virology*. 2008;377:117–23. <http://dx.doi.org/10.1016/j.virol.2008.04.027>

Address for correspondence: Mustafizur Rahman, International Centre for Diarrhoeal Disease Research, Bangladesh, Mohakhali, Dhaka-1212, Bangladesh; email: [mustafizur@icddr.org](mailto:mustafizur@icddr.org)

## Norovirus GII.4/Sydney/2012 in Italy, Winter 2012–2013

**To the Editor:** Noroviruses (NoVs) are the major cause of acute gastroenteritis in children and adults; they are responsible for sporadic cases and outbreaks of gastroenteritis in various epidemiologic settings. NoVs can be classified genetically into at least 5 genogroups, GI to GV (1). Although >30 genotypes within genogroups GI, GII, and GIV can infect humans (2), a single genotype, GII.4, has been associated with most NoV-related outbreaks and sporadic cases of gastroenteritis worldwide (3).

GII.4 NoV strains continuously undergo genetic/antigenic diversification and periodically generate novel strains through accumulation of punctate mutations or recombination. New GII.4 variants emerge every 2–3 years (4). Increased incidence of NoV-related illness and/or outbreaks in various

countries in late 2012 has been related to the emergence of a novel GII.4 variant, Sydney 2012. This variant was first identified in March 2012 in Australia (5).

The Italian Study Group for Enteric Viruses (ISGEV; <http://isgev.net>) monitors the epidemiology of enteric viruses in children through hospital-based surveillance (6–8). NoVs are monitored and characterized by multi-target analysis in the diagnostic regions A (open reading frame 1, polymerase) and C (open reading frame 2, capsid) of the NoV genome (9) and interrogation of the Norovirus Typing Tool database ([www.rivm.nl/mpf/norovirus/typing-tool](http://www.rivm.nl/mpf/norovirus/typing-tool)). During November 2011–March 2012, the prevalence of sporadic NoV infections detected (in samples from newborns, infants, and children up to 5 years of age) by real-time reverse transcription PCR was 22.2% (121/545). A subset (≈50%) of the NoV-positive samples representative of the whole winter period was selected for sequence analysis, and 48 were successfully characterized in region A and region C.

Among these 48 NoV strains, 20 (41.7%) were characterized as the variant GII.4 New Orleans 2009, a smaller number, 6 (12.5%), displayed a New Orleans 2009 polymerase (pol) but 2 distinct GII.4 capsid sequences, which were not typeable in the Norovirus Typing Tool database, and only 2 (4.2%) GII.4 strains of the variant Den Haag 2006b were detected. Moreover, 4 sporadic cases in November 2011 and January 2012 and a small outbreak in February 2012 were related to a GII.4 Pe\_GII.4 recombinant strain. After the set of sequences of GII.4 variants from the Norovirus Typing Tool database was updated (access to the updated database: April 11, 2013), 5 (10.4%) GII.4 Pe\_GII.4 recombinant strains were characterized as variant Sydney 2012.

From April through October 2012, a total of 56 (7.6%) NoV-positive samples were detected from 737 analyzed samples, of which 34 (60.7%) NoV-positive samples could be sequenced. Of

these, 41.2% were characterized as GII.3 (mostly with a GII.Pb pol), 26.5% as GII. Pg\_GII.1, and 17.6% as GII.4 variants. From spring to fall 2012, the variant New Orleans 2009 became the predominant GII.4 strain, and the variant Sydney 2012 strain apparently disappeared.

During November–December 2012 and January 2013, ISGEV detected NoV infection in 90 (28.9%) of 311 children hospitalized for gastroenteritis. This finding is comparable to a prevalence of 25.2% in the same period (November–January) of the 2011–12 winter season. A representative subset of 45 samples was randomly selected for sequencing, and 26 (74.3%) of 35 fully typed strains were characterized as GII.4 Sydney 2012, which suggested that the new NoV variant had become the predominant strain in Italy.

Our surveillance seem to mirror observations of a report from Denmark that documented the onset and circulation at low prevalence of the variant GII.4 Sydney 2012 at the beginning of 2012 with a marked increase in the prevalence only by the end of 2012 (10). Our surveillance detected the emergence of this variant in Italy at the end of 2011 and provided us with one of the earliest strains of the variant GII.4 Sydney 2012. This novel variant has a common ancestor with the NoV GII.4 variants Apeldoorn 2008 and New Orleans 2009 and has several amino acid changes on the main epitope in the capsid P2 domain (10).

Sequence analysis of these early strains of the GII.4 variant Sydney 2012 could help clarify the mechanisms driving its global emergence and spread. Continued surveillance for NoV infections through ISGEV and additional data on clinical and epidemiologic features will enable further assessment of the public health implications of the new variant GII.4 Sydney 2012 in Italy.

This study was supported by the grants “Studio dei meccanismi evolutivi

dei calicivirus umani” (Italian Scientific Research Fund PRIN 2008), “Caratterizzazione molecolare di norovirus circolanti nella popolazione pediatrica” (University of Palermo, Italy, Fondi di Ateneo 2007), and “Epidemiologia molecolare e studio dei meccanismi evolutivi di norovirus” (University of Parma, Italy, Fondi di Ateneo 2008).

**Giovanni M. Giammanco,  
Simona De Grazia,  
Fabio Tummolo,  
Floriana Bonura,  
Adriana Calderaro,  
Alessio Buonavoglia,  
Vito Martella, and  
Maria C. Medici**

Author affiliations: Università degli Studi di Palermo, Palermo, Italy (G.M. Giammanco, S. De Grazia, F. Bonura); Università degli Studi di Parma, Parma, Italy (F. Tummolo, A. Calderaro, M.C. Medici); Medical Practitioner, Bari, Italy (A. Buonavoglia); and Università Aldo Moro di Bari, Valenzano, Italy (V. Martella)

DOI: <http://dx.doi.org/10.3201.130119>

## References

- Green KY. Caliciviridae. In: Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, et al., editors. *Fields virology*, 5th ed. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 949–79.
- Kroneman A, Vega E, Vennema H, Vinjé J, White PA, Hansman G, et al. Proposal for a unified norovirus nomenclature and genotyping. *Arch Virol*. 2013; April 25. Epub ahead of print.
- Bok K, Abente EJ, Realpe-Quintero M, Mitra T, Sosnovtsev SV, Kapikian AZ, et al. Evolutionary dynamics of GII.4 noroviruses over a 34-year period. *J Virol*. 2009;83:11890–901. <http://dx.doi.org/10.1128/JVI.00864-09>
- Siebenga JJ, Vennema H, Renckens B, de Bruin E, van der Veer B, Siezen RJ, et al. Epochal evolution of GGII.4 norovirus capsid proteins from 1996 to 2006. *J Virol*. 2007; 81:9932–41. <http://dx.doi.org/10.1128/JVI.00674-07>
- van Beek J, Ambert-Balay K, Botteldoorn N, Eden J, Fonager J, Hewitt J, et al. Indications for worldwide increased norovirus activity associated with emergence of a new variant of genotype II.4, late 2012. *Euro Surveill*. 2013;18:8–9.

- Medici MC, Tummolo F, Albonetti V, Abelli LA, Chezzi C, Calderaro A. Molecular detection and epidemiology of astrovirus, bocavirus, and sapovirus in Italian children admitted to hospital with acute gastroenteritis, 2008–2009. *J Med Virol*. 2012;84:643–50. <http://dx.doi.org/10.1002/jmv.23231>
- Giammanco GM, Rotolo V, Medici MC, Tummolo F, Bonura F, Chezzi C, et al. Recombinant norovirus GII.g/GII.12 gastroenteritis in children. *Infect Genet Evol*. 2012;12:169–74. <http://dx.doi.org/10.1016/j.meegid.2011.10.021>
- De Grazia S, Martella V, Chironna M, Bonura F, Tummolo F, Calderaro A, et al. Nationwide surveillance study of human astrovirus infections in an Italian paediatric population. *Epidemiol Infect*. 2013;141:524–8. <http://dx.doi.org/10.1017/S0950268812000945>
- Kroneman A, Vennema H, Deforche K, Avoort HVD, Peñaranda S, Oberste MS, et al. An automated genotyping tool for enteroviruses and noroviruses. *J Clin Virol*. 2011;51:121–5. <http://dx.doi.org/10.1016/j.jcv.2011.03.006>
- Fonager J, Hindbaek L, Fischer T. Rapid emergence and antigenic diversification of the norovirus 2012 Sydney variant in Denmark, October to December, 2012. *Euro Surveill*. 2013;18:pil: 20413.

Address for correspondence: Giovanni M. Giammanco, Dipartimento di Scienze per la Promozione della Salute e Materno Infantile “G. D’Alessandro,” Via del Vespro 133, 90127, Palermo, Italy; email: [giovanni.giammanco@unipa.it](mailto:giovanni.giammanco@unipa.it)

## Group C Betacoronavirus in Bat Guano Fertilizer, Thailand

**To the Editor:** Bats play a critical role in the transmission and origin of zoonotic diseases, primarily viral zoonoses associated with high case-fatality rates, including those caused by Nipah virus (NiV) and severe acute respiratory syndrome (SARS)-like coronavirus (CoV) infections (1). Recently, the World Health Organization (WHO) reported 44 confirmed cases of human infection with Middle East