

Case-Control Study of Paresthesia Among World Trade Center-Exposed Community Members

Michael Marmor, PhD, Sujata Thawani, MD, MPH, Maria Luisa Cotrina, PhD, Yongzhao Shao, PhD, Ericka S. Wong, MD, Mark M. Stecker, MD, PhD, Bin Wang, Alexander Allen, MD, Marc Wilkenfeld, MD, Etta J. Vinik, MA, Aaron I. Vinik, MD, PhD, and Joan Reibman, MD

Objective: To investigate whether paresthesia of the lower extremities following exposure to the World Trade Center (WTC) disaster was associated with signs of neuropathy, metabolic abnormalities, or neurotoxin exposures. **Methods:** Case-control study comparing WTC-exposed paresthesia cases with “clinic controls” (WTC-exposed subjects without paresthesias), and “community controls” (WTC-unexposed persons). **Results:** Neurological histories and examination findings were significantly worse in cases than controls. Intraepidermal nerve fiber densities were below normal in 47% of cases and sural to radial sensory nerve amplitude ratios were less than 0.4 in 29.4%. Neurologic abnormalities were uncommon among WTC-unexposed community controls. Metabolic conditions and neurotoxin exposures did not differ among groups.

From the Department of Population Health, NYU Grossman School of Medicine, New York (Dr Marmor, Dr Shao, Ms Wang, Dr Alexander), Department of Environmental Medicine, NYU Grossman School of Medicine, New York, New York (Dr Marmor, Dr Shao, Ms Wang, Dr Reibman), Department of Medicine, NYU Grossman School of Medicine, New York, New York (Dr Marmor, Dr Cotrina, Dr Wilkenfeld, Dr Reibman), Department of Neurology, NYU Grossman School of Medicine, New York, New York (Dr Thawani, Dr Alexander, Dr Wong), Department of Neurology, Thomas Jefferson University Hospital, Philadelphia, Pennsylvania (Dr Wong), UCSF/Fresno and University Neurology Associates, Fresno, California (Dr Stecker), Division of Occupational and Environmental Medicine, Department of Medicine, NYU Winthrop Hospital, Mineola, New York (Dr Wilkenfeld) and Department of Internal Medicine, Strelitz Diabetes Center, Eastern Virginia Medical School, Norfolk, Virginia (Ms Vinik, Dr Vinik).

Data Availability Statement: De-identified data on the cases, clinic controls, and community controls can be obtained by contacting Yongzhao Shao, PhD (Yongzhao.Shao@nyulangone.org).

Funding: This study was supported by the cooperative agreement U01 OH010395 from the National Institute for Occupational Health and Safety (NIOSH), Centers for Disease Control and Prevention to the NYU School Grossman of Medicine; NYU Clinical and Translational Science Award (CTSA) number UL1TR001445 from the National Center for Advancing Translational Sciences (NCATS), National Institute of Health (NIH); NIOSH contracts 200-2017-93327 (Data Center) and 200-2017-93427 (Clinical Center of Excellence) to NYC Health + Hospitals, Bellevue Hospital Center; and Center Grant 5P30ES000260 from the National Institute of Environmental Health Sciences to NYU Grossman School of Medicine.

Disclosures: This study was sponsored by the National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention.

M.M. has been funded by CDC/NIOSH cooperative agreement 1U01-OH011305-01 and U01OH010394-01A1, and NIH grant R01EY015520. He has no other disclosures.

S.T. has been funded by CDC/NIOSH cooperative agreement 1U01-OH011305-01. She has no other disclosures.

M.L.C. has been funded by cooperative agreement 1U01-OH011305-01. She has no other disclosures.

Y.S. has been funded by the cooperative agreement 1U01-OH011305-01 and U01OH010394-01A1, U01 OH010395 from the National Institute for Occupational Health and Safety (NIOSH), Centers for Disease Control and Prevention to the NYU Grossman School of Medicine; Center Grant 5P30AG008051 and SPORE grant 1P50CA225450 from the National Institute of Health to NYU Grossman School of Medicine. NIOSH contracts 200-2017-93327 (Data Center) to NYC Health Hospitals, Bellevue Hospital Center; and Center Grant 5P30ES000260 from the National Institute of Environmental Health Sciences to NYU Grossman School of Medicine. He has no other disclosures.

E.S.W. has been funded by CDC/NIOSH cooperative agreement 1U01-OH011305-01. She has no other disclosures.

Conclusions: Paresthesias among WTC-exposed individuals were associated with signs of neuropathy, small and large fiber disease. The data support WTC-related exposures as risk factors for neuropathy, and do not support non-WTC etiologies.

Keywords: electrodiagnostic neurologic measurements, neuropathic symptoms, paresthesia, peripheral neuropathy, small nerve fiber density

Lower extremity paresthesias have been reported among first-responders and community members exposed to the World Trade Center (WTC) disaster of September 11, 2001 (referred to herein as 9/11). We previously reported paresthesias among members of the WTC Environmental Health Center (EHC), a treatment

M.M.S. has received royalties from UPTo Date and research support (not related to the present project) from the Campos Family Farms. He has no other disclosures.

B.W. has been funded by cooperative agreement 1U01-OH011305-01. She has no other disclosures.

A.A. and M.W. have no disclosures.

E.J.V. has not received study funding in the past 2 years. She and A.I.V. hold a licensing agreement with Eastern Virginia Medical School for the use of the Norfolk QOL-DN to commercial companies. No license or fee is required for its use by academic or nonprofit institutions. She has no other disclosures or conflict of interest.

A.I.V. has been funded by the National Institutes of Health (NIH), National Institute of Neurological Disorders and Stroke (NINOS), NeuroNEXT Clinical Trial (U01)—Topiramate as a disease altering therapy for CSPN; ViroMed Laboratories: A Phase II, Double-Blind, Randomized, Placebo-Controlled, Multicenter Study to Assess the Safety and Efficacy of VM202 in Subjects with Painful Diabetic Peripheral Neuropathy. Veroscience LLC: Impact of Timed Bromocriptine-QR Therapy upon Measures of Sympathetic Tone and Vascular Biology in Type 2 Diabetes Subjects; Novo Nordisk A/S: The Effects of Liraglutide on Sudomotor Function and Inflammation in Type 2 Diabetes; Impeto: Medical, Inc. "Autonomic Neuropathy in Patients in Patients with Psoriasis" Aptinyx Inc.: "A Randomized, Double-Blind, Parallel-Group, Placebo-Controlled, Multiple-Dose Study to Assess the Efficacy and Safety of NYX-2925 in Subjects with Neuropathic Pain Associated with Diabetic Peripheral Neuropathy." A.I.V. also has been a Consultant/Speaker for Merck, Pfizer, Alnylam, and Astellas. He has a patent on Norfolk QOL-DN. He has no other disclosures or conflict of interest.

J.R. has been funded by the National Institute of Occupational Safety and Health (NIOSH) contracts 200-2017-93327 and 200-2017-93427. She has no other disclosures.

Clinical significance: World Trade Center-exposed paresthesia cases had greater prevalence than paresthesia-free controls of signs of neuropathy, small and large fiber nerve disease. A majority of WTC survivor's paresthesia reports thus seem to reflect true disease rather than psychosomatic symptoms or questionnaire artifacts. WTC-related exposures may have contributed to peripheral neuropathy.

Supplemental digital contents are available for this article. Direct URL citation appears in the printed text and is provided in the HTML and PDF versions of this article on the journal's Web site (www.joem.org).

Address correspondence to: Michael Marmor, PhD, Department of Population Health, 180 Madison Avenue, Room 9-53, New York, NY 10016 (Michael.Marmor@nyulangone.org).

Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American College of Occupational and Environmental Medicine. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Clay without permission from the J
DOI: 10.1097/JOM.0000000000001828

program for community members (local workers, local residents, students and passersby) exposed to WTC dust and fumes on and after 9/11 who subsequently developed medical or mental health conditions thought to be due to this exposure.^{1,2} The WTC EHC, now a federal program, provides services based upon exposure to the events of 9/11 and development of conditions certifiable as likely caused by WTC exposure.³ Risk factors for paresthesia included intensity of exposure to WTC dust on 9/11 and employment in a job that required cleaning of WTC dust.² Time to paresthesia was associated with these same variables.⁴ Furthermore, electromyographic studies of 16 WTC-exposed first-responders found 1 case of possible demyelinating neuropathy and 3 cases of motor neuron disease without evidence of compression of the corticospinal tracts.⁵ Rodent nerves exposed *ex vivo* to WTC dust had lower conduction velocities than control nerves.⁶

Heavy metals and complex hydrocarbons (eg, polycyclic aromatic hydrocarbons, polybrominated diphenyl ethers, polychlorinated biphenyls, and polychlorinated naphthalenes) in WTC dust and fumes have potential neurotoxicity and provide biologic plausibility⁷⁻⁹ for the hypothesis that WTC exposures were a risk for neuropathy.⁷⁻⁹ Additionally, cleaning of WTC dust may have exposed workers to neurotoxic cleaning agents. Repetitive motions during cleaning activities may have increased the risk of carpal tunnel syndrome.

We performed a case-control study of lower extremity paresthesia among WTC-exposed individuals to explore whether this subjective symptom was associated with clinical signs, including markers of small or large fiber nerve abnormalities, and metabolic and toxic blood markers for possible non-WTC neuropathic etiologies.

MATERIALS AND METHODS

Subjects and Study Design

We enrolled patients primarily from the Bellevue Hospital WTC EHC. Study enrollment occurred during June 28, 2017 to August 22, 2018. WTC EHC patients completed questionnaires interrogating demographics, health, and WTC exposures at their enrollment visits and at recall, or “monitoring” visits. Potential volunteers for the present study were identified among WTC EHC enrollees who consented to the use of their information for research, and who separately consented to future contact to discuss participation in related research projects. Inclusion criteria included aged 18 to 75 years and enrollment in the WTC EHC between August 22, 2005 and March 30, 2018.

We employed a case-control study design to contrast symptoms, signs and exposures among four groups:

1. *Cases:* WTC EHC patients who reported paresthesias of the lower extremities at their most recent WTC EHC visit and in the screening questionnaire with symptom frequency of “often” or “almost continuous.” We excluded isolated upper-extremity paresthesias to minimize cases of carpal tunnel syndrome.
2. *Clinic controls:* WTC EHC patients who did not report paresthesia of the upper or lower extremities at any study visit, and who continued to be free of paresthesias at the times of screening and enrollment into the present study. This group was enrolled primarily to estimate the prevalence of signs of neuropathy among paresthesia-free WTC survivors.
3. *Community controls:* Individuals without WTC exposure who did not live or work south of Canal Street, Manhattan, New York on 9/11. These subjects included: (1) friends or relatives of cases or clinic controls; (2) participants in other research studies being conducted at Bellevue Hospital Center; or (3) participants in the ResearchMatch, a secure registry at NYU Langone Medical Center open to individuals interested in participating in research studies. This group was enrolled primarily to estimate the prevalences of paresthesia and signs of neuropathy in the general population without severe WTC exposure.

4. *Historical controls for small nerve fiber density measurements:* Normative data on small nerve fiber densities were obtained from Therapath Partners, LLC, New York, NY. Therapath provided a de-identified data set of epidermal small nerve fiber densities, sweat gland nerve fiber densities, sex, and age for 128 individuals who had been classified as free of neuropathic symptoms on the basis of history and physical examination. This study had been conducted with oversight from The Copernicus Group Independent Review Board (study THE4-06-146). All subjects provided written informed consent. Study recruitment occurred from June 14, 2006 to March 28, 2014 (Arthur P. Hays, MD, principal investigator). Subjects were recruited at sites in Arizona, California, Idaho, Maryland, New Jersey, and New York. These proprietary data were used with permission by the owner, Therapath Partners, LLC, which also provided sex- and age-specific cut-points for abnormality.

Cases and clinic controls had to meet their respective group’s paresthesia requirements (see below) during the month prior to enrollment. Exclusion criteria applied to cases, clinic and community controls were: paresthesia before 9/11; cancer chemotherapy at any time; known vitamin B12 deficiency; diabetes mellitus; use of antiretroviral medications at any time; and conditions that would preclude elective skin punch biopsies, including bleeding disorders, use of anticoagulants, and history of keloid scarring.

We used frequency-matched sampling to recruit subjects into sex- and age (18 to 25, 25 to 34, 35 to 44, 45 to 54, 55 to 64, and 65 to 74 years)-defined strata. We planned to recruit one clinic control per case and one community control for every two cases.

Of 5457 potential subjects aged 18 to 75 years enrolled in the Bellevue WTC EHC between August 22, 2005 and March 30, 2018, first monitoring visits were completed by 2373 individuals (43.5% of all enrolled WTC EHC patients); second monitoring visits (conducted from December 5, 2012 on) were completed by 1105 individuals (18.4% of enrollees); and third monitoring (conducted from April 1, 2015) were completed by 384 (5.6% of enrollees).

Study Size

We planned to recruit 40 cases, 40 clinic controls, and 20 community controls, a sample size with adequate power at the two-tailed alpha significance level to show a difference between a 30% or greater prevalence of abnormal small nerve fiber densities in cases compared to an anticipated prevalence of 5% or fewer among historical controls with a power of 97%.¹⁰ For other parameters of interest, such as abnormal neurological examination results, the samples of 40 cases and 40 clinic controls would provide 88% power at a 5% two-tailed alpha significance level if the risk factor were present in 30% or more of cases and 5% or fewer of clinic controls. For comparisons between the planned 40 cases and 20 community controls, the study would have power of 76% at an alpha-significance level of 0.05 to detect the difference between a factor present in 30% of cases and 5% of community controls.

Interview Measures

At the single study visit required for the present study, subjects responded to a questionnaire regarding demographics, health history, and WTC exposures. The questionnaire included a self-administered alcohol assessment (cut-annoyed-guilty-eye questionnaire¹¹) modified to inquire into alcohol issues within the past year and the PCL-17, used to quantify subjects’ degree of posttraumatic stress disorder. We used the OccIDEAS to elicit histories of exposure to potential neurotoxins during work or hobbies (see Appendix 1 for a list of these agents).¹² Interview data from first monitoring visits were used to complete data on WTC exposures that were missing from the present study’s

questionnaire responses for several subjects. We also mailed a 5-question supplemental questionnaire to 13 subjects to further supplement missing data on WTC exposures (returned by 8 [62%] of these subjects).

Body mass index (BMI) and vital signs were obtained using standard equipment.

Neurological Examination

All subjects underwent neurologic examination with trained neurologists. Subjects completed additional self-administered questionnaires, including: the Utah Early Neuropathy Scale,¹³ the University of Michigan Neuropathy Screening Instrument—Patient Version,¹⁹ a modification of the neurological history and examination used by the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications Study (personal communication, Brian Callaghan, February 2, 2017),^{14,15} and the Norfolk Quality of Life—Diabetic Neuropathy (QOL-DN).¹⁶

Nerve Conduction Studies

We evaluated large fiber neuropathy via nerve conduction studies using methods recommended for the early identification of diabetic symmetrical peripheral neuropathy.¹⁵ Controls were not subjected to nerve conduction studies out of concern that there would be no benefit to most controls and that anticipated discomfort associated with these studies would reduce participation rates. We measured distal latencies and amplitudes, and calculated conduction velocities of the peroneal, tibial and ulnar motor nerves; and of the sural sensory and radial sensory nerves. These nerves were chosen because they were included in the nerve impairment score + 7 reported by Suanprasert et al¹⁷ and have previously been shown to be useful in describing diabetic neuropathy.¹⁸ Studies were performed whenever possible on the left side. We also analyzed the sural to radial sensory nerve amplitude ratio and categorized subjects as abnormal if this ratio was less than 0.4, as has been suggested for diagnosis of axonal polyneuropathy.¹⁹ In a study of Sjögren syndrome, 20.4% of subjects had sural to radial sensory nerve amplitude ratios less than 0.4 compared to 6% of healthy controls.²⁰ Nerve conduction studies were performed after subjects had been in temperature-controlled spaces maintained at 32°C for at least 1 hour. We did not conduct electromyography out of concern that the required insertion of needles might reduce participation rates. Cut-points for abnormality of electrodiagnostic test results were provided by the Bellevue Hospital Center Neurology Laboratory where the testing was performed.

Epidermal Nerve and Sweat Glands Nerve Fiber Densities

We evaluated small fiber neuropathy by measuring the density of small nerve fibers in skin biopsies.^{21,22} Each case was asked to undergo a single, circular 3 mm skin punch biopsy of the distal left leg, approximately 10 cm above the lateral malleolus. A commercial laboratory (Therapath Partners, LLC) stained and evaluated the biopsy specimens for epidermal nerve fiber density. If the epidermal nerve fiber density was normal, sweat gland nerve fiber density was evaluated. This testing resulted in two measures: the quantitative epidermal nerve fiber density per mm of epidermal length, and a dichotomous characterization of the epidermal or sweat gland small nerve fiber density as “normal” or “abnormal” when compared with age- and gender-specific normative values. Some samples obtained did not have any visible sweat glands due to either sampling or compression of the glands during transfer to the specimen vial.

Study neurologists performed the skin punch biopsies and the nerve conduction velocity studies. Due to their involvement in these procedures and the scheduling thereof, we were unable to mask the neurologists to subjects' case or control status.

TABLE 1. Blood Tests Performed to Investigate Known Causes of Paresthesia or Neuropathic Injury

System	Blood Test
Endocrine	Fasting glucose and hemoglobin A1C
Infectious disorders	Syphilis (<i>Treponema pallidum</i> IgG Ab), Lyme (<i>B. burgdorferi</i>)
Nutritional	B12, methylmalonic acid, folate
Autoimmune disorders	Antinuclear antibody, antibodies to deamidated gliadin peptide (associated with celiac disease)
Hematologic	SPEP with immunofixation
Inflammatory reactions	C-reactive protein
Toxic exposures	Blood lead concentration
	SPEP, serum protein electrophoresis.

Alternative Metabolic, Infectious, Nutritional, or Toxic Causes of Paresthesia and Peripheral Neuropathy

Venous peripheral blood samples were evaluated by the NYU Langone Medical Center Clinical Laboratory, New York, NY to evaluate possible metabolic, infectious, nutritional, or toxic causes of peripheral neuropathy (Table 1).²³ Syphilis testing followed the reverse algorithm.²⁴ Blood lead concentrations were measured by a commercial laboratory (ARUP Laboratories, Salt Lake City, UT). Blood test results were analyzed as continuous variables and also were dichotomized into normal versus abnormal with abnormal defined by the participating laboratories as hemoglobin A1c greater than or equal to 6%, C-reactive protein greater than or equal to three units, methylmalonic acid greater than 0.4 μ mol/L, vitamin B12 less than 213 pg/mL, folate less than 6 ng/mL, hemoglobin A1c greater than or equal to 6%, fasting glucose > 100 mg/dL, ANA inverse titers greater than or equal to 40, C-reactive protein greater than or equal to 3 units, gliadin peptide IgA antibodies greater than or equal to 19 units, gliadin peptide IgG antibodies greater than or equal to 19 units, lead greater than or equal to 4.9 μ g/dL, and *Borrelia burgdorferi* antibodies greater than or equal to 0.91 units.

Data Management

We managed study data using the REDCap system hosted by NYU School of Medicine.²⁵

Statistical Methods

Categorical variables were summarized using counts and proportions, and statistically evaluated using the chi-squared or Fisher exact tests. Continuous variables were summarized using means and standard deviations and differences tested using the *t* test, analysis of variance or linear regression; or for variables that were not normally distributed, medians, interquartile ranges (IQRs) and the Wilcoxon rank-sum or Kruskal-Wallis test. The Mantel-Haenszel technique was used to calculate summary odds ratios after stratifying on the potential confounders of sex and categorized age used in the frequency-matched enrollment process.²⁶ Counts for some variables did not sum to the total numbers of subjects in each group due to missing data.

Ethical Oversight

All study procedures were approved by the NYU School of Medicine Institutional Review Board (study number 016-1490 and WTC EHC research database NCT00404898). Written informed consent was provided by all study participants. Data on historical controls were provided by Therapath, Inc., in a Health Insurance Portability and Accountability Act-compliant, de-identified manner.




FIGURE 1. Flowchart of recruitment for case-control study of paresthesia.

RESULTS

Study Enrollment

We identified 1302 patients who met the paresthesia case definition, 1493 who met the clinic control definition, and 2586 who met neither definition. Patients were excluded based on paresthesias prior to 9/11 ($n = 47$), diabetes ($n = 370$) or cancer chemotherapy ($n = 711$). There resulted 849 potential cases and 1235 potential controls (Fig. 1). Two enrolled cases were referrals from a collaborating hospital to the WTC EHC for certification of illnesses. Participation rates were 11% among cases and clinic controls and 43% among community controls. We enrolled 41 cases, 38 clinic controls, and 20 community controls.

Enrollment Demographics

Due to frequency-matching during enrollment, the age and gender distributions were similar across groups (Table 2). There were no significant differences in race or ethnicity. Cases reported lower annual household incomes compared with clinic or community controls. The distributions of BMI did not differ among groups. cut-annoyed-guilty-eye scores did not differ significantly among groups.

WTC Exposure Histories

Around 27% (11/41) of cases reported being “heavily covered and unable to dust most of the dust off with their hands before returning home on 9/11” compared to 13% (5/38) of clinic controls ($P = 0.17$, Fisher exact test). The Mantel-Haenszel age- and sex-

adjusted summary odds ratio (OR) for paresthesia associated with being heavily exposed to WTC dust on 9/11 was 2.3 (95% CI = 0.7 to 7.8, $P = 0.18$). There also was a suggestion that paresthesia was associated with having worked in a job requiring cleaning of WTC dust (22%, 9/41) compared to controls (11%, 4/38, $P = 0.23$, Fisher exact test; age- and sex-adjusted OR = 2.4, 95% confidence interval = 0.7 to 8.3, $P = 0.16$).

Standardized Neurological Questionnaires and Objective Tests

Scores on the Norfolk Diabetic Neuropathy QoL—Diabetic Neuropathy instrument were significantly higher (worse) in paresthesia cases than in both control groups (Table 3). High total QoL scores as well high large- and small-fiber nerves scores indicated impairment.

The Utah Early Neuropathy Scale indicated significant impairments among cases (median score = 6, IQR = 4 to 10) compared to both clinic controls (median score = 0, IQR = 0 to 2) and community controls (median score = 0, IQR = 0 to 2), $P < 0.0001$ (Table 3). The greatest impairments of cases compared to controls among the components of the Utah Early Neuropathy Scale were in pin sensation and deep tendon reflex, and great toe joint position (Supplemental Table 1, <http://links.lww.com/JOM/A712>).

The Michigan Neuropathy Screening Instrument indicated significantly higher scores for neuropathic symptoms among paresthesia cases than controls (Table 3). The neurological history and examination also indicated significantly higher (worse) scores among paresthesia cases compared to controls (Supplemental

TABLE 2. Demographic Description of Subjects Enrolled in a Case-Control Study of Paresthesia and World Trade Center Exposures^a

Variable	Cases	Clinic Controls	Community Controls	P Value*
Value				
Number enrolled	41	38	20	
Gender				
Male	15 (37)	13 (34)	8 (40)	0.9
Female	26 (63)	25 (66)	12 (60)	
Age				
25–34	2 (5)	1 (3)	1 (5)	1.0
35–44	4 (10)	4 (11)	2 (10)	
45–54	8 (20)	7 (18)	4 (20)	
55–64	16 (39)	15 (39)	7 (35)	
≥65	11 (27)	11 (30)	6 (30)	
Race				
Caucasian	15 (37)	21 (55)	8 (40)	0.16
African-American	11 (37)	8 (21)	9 (45)	
American Indian, Asian and other	3 (7)	4 (11)	0 (0)	
Do not know/Refused to answer	12 (29)	5 (13)	3 (15)	
Ethnicity				
Not Hispanic	27 (66)	29 (76)	14 (70)	0.8
Hispanic	13 (32)	8 (21)	6 (30)	
Refused to answer	1 (2)	1 (3)	0 (0)	
Annual household income (USD)				
<\$25,000	15 (37)	6 (16)	7 (35)	0.04¶
\$25,001–\$75,000	17 (41)	15 (39)	7 (35)	
> \$75,000	3 (12)	16 (42)	6 (30)	
Refused to answer	4 (10)	1 (3)	0 (0)	
BMI (kg/m ²)				
< 25	6 (15)	12 (32)	7 (35)	0.2
25 to < 30	15 (37)	12 (32)	7 (35)	
30 to < 35	10 (24)	6 (16)	5 (25)	
≥ 35	10 (24)	8 (21)	1 (5)	
CAGE alcoholism score				
< 2	39 (98)	31 (86)	18 (95)	0.3
≥ 2	1 (3)	4 (11)	1 (5)	
Weekday alcohol consumption outside mealtimes				
No	37 (95)	31 (89)	14 (74)	0.06
Yes	2 (5)	4 (11)	5 (26)	
Days per week consumed alcohol				
0–1	31 (82)	23 (66)	11 (61)	0.3
2	4 (11)	4 (11)	4 (22)	
≥3	3 (8)	8 (22)	4 (17)	
Caught in WTC dust cloud on 9/11				
No	17 (41)	17 (45)		0.8
Yes	24 (59)	21 (55)		
Heavily covered with dust on 9/11				
No	30 (73)	33 (87)		0.17¶
Yes	11 (27)	5 (13)		
Worked in a job requiring cleaning of WTC dust				
No	32 (78)	34 (89)		0.23¶
Yes	9 (22)	4 (11)		

BMI, body mass index; WTC, World Trade Center.

^aCounts not summing to the total number of subjects in each group reflect missing data.

*P values from the chi-squared test unless otherwise indicated.

¶P value from Fisher exact test.

Table 2, <http://links.lww.com/JOM/A712>). The physician-administered neurological examinations showed that 39% of cases had multiple examination findings consistent with distal symmetric polyneuropathy versus 8% of clinic controls and 0% of community controls. Considering examination findings and symptoms, the study's neurologists rated 58% of cases as "definitely yes" and 34% as "possible yes" for having clinically evident distal symmetric polyneuropathy, which was significantly greater than the 3% and 21%, respectively, among clinic controls, and the 0% and 0% among community controls.

Total PCL total scores were higher among paresthesia cases (median = 44, IQR = 32 to 62) than clinic controls (median = 25, IQR = 21 to 34) or community controls (median = 18.5, IQR = 17 to 20; $P < 0.0001$).

Nerve Conduction Studies

Peroneal and ulnar motor nerve conduction velocities were below the cutoff for normality in 3% of paresthesia cases, a prevalence that did not differ from expected (Table 4).¹⁴ However, delayed latencies were present in 40% of subjects for the tibial

TABLE 3. Findings from the Norfolk Quality of Life—Diabetic Neuropathy (QOL-DN) instrument, the Utah Early Neuropathy Scale, and the Michigan Neuropathy Screening Instrument Among Paresthesia Cases, Clinic Controls, and Community Controls (Higher Scores Indicate Greater Neuropathic Impairment)

Scale	Cases [Median, IQR] (n = 41)	Clinic Controls [Median, IQR] (n = 38)	Community Controls [Median, IQR] (n = 20)	P Value*
Norfolk physical functioning/large fiber score	31 (16,41)	0 (-1,9)	0 (-1,9)	<0.0001
Norfolk small fiber score	2 (1,4)	0 (0,0)	0 (0,0)	<0.0001
Norfolk total score	53 (32,70)	2 (-1,9)	0 (-1.5,3.5)	<0.0001
Utah early neuropathy scale	6 (4,10)	0 (0,2)	0 (0,2)	<0.0001
Michigan neuropathy screening instrument	5 (4,7)	1 (0,2)	0 (0,1)	<0.0001

DN, diabetic neuropathy; IQR, interquartile range; QOL, quality of life.

*P from the Kruskall-Wallis test.

motor nerve and low amplitude nerve action potentials were present in 0 to 36% of nerves tested. The sural to radial sensory nerve amplitude ratio was abnormal (less than 0.4) in 29.4% (10/34) of subjects. Of the 10 paresthesia cases with abnormal sural to radial sensory nerve amplitude ratios, 60% (6/10) had reduced epidermal skin or sweat gland fiber densities compared to 52% (11/23) of those with normal sural to radial sensory nerve amplitude ratios. Quantitative sural to radial sensory nerve amplitude ratios also were not correlated with epidermal nerve fiber densities (Spearman $r=0.07$, $P=0.7$). The Wilcoxon rank-sum test revealed no significant differences in the blood measures shown in Table 5 between subjects with reduced (less than 0.4) sural to radial sensory nerve amplitude ratios and those with normal ratios. None of the subjects had clinical or nerve conduction evidence of chronic inflammatory demyelinating polyneuropathy.

Skin Biopsy Examination for Epidermal Nerve and Sweat Gland Nerve Fiber Density

Epidermal or sweat gland nerve fiber densities were significantly reduced among cases compared to historical controls (Table 6, Part A). This difference remained significant after adjusting for age and gender by linear regression. The composite measure of epidermal or sweat gland small nerve fiber density was abnormally low in 57% of cases compared to an expected prevalence of abnormal findings of 4 to 5%, as estimated from the age- and gender-similar historical controls (Table 6, Part B, $P<0.0001$). Heights of cases with reduced intraepidermal nerve fiber densities were greater (median = 166.5 cm, min = 149.3, max = 202) than heights of cases with normal intraepidermal nerve fiber densities (median = 159.3 cm, min = 146, max = 170.5; Wilcoxon ranked sum $P=0.01$). The difference in heights remained significant after adjustment for age and gender.

TABLE 4. Frequencies of Abnormal Nerve Conduction Velocity Findings Among Paresthesia Cases Enrolled in the World Trade Center Environmental Health Center

Nerve	Latency (% [n Abnormal/n Tested])	Amplitude (% [n Abnormal/ n Tested])	Conduction Velocity (% [n Abnormal/n Tested])
Motor nerves			
Peroneal motor nerve (left)			
Ankle—extensor digitorum brevis	8.3 (3/36)*	13.9 (5/36)†	
Fibular head—ankle		13.9 (5/36)†	3.0 (1/33)‡
Tibial motor nerve (Left)			
Ankle—abductor hallucis brevis muscle	40.0 (14/35)§	8.8 (3/34)	
Ulnar motor nerve (left)			
Wrist—abductor digiti minimi muscle	0 (0/36)¶	25.0 (9/36)♯	
Below elbow—Wrist		36.1 (13/36)♯	2.9 (1/35)**
Sensory nerves:			
Radial sensory nerve (Left)			
Forearm—extensor pollicis longus tendon	22.9 (8/35)††	20.0 (7/35)††	0 (0/35)**
Sural sensory nerve (Left)			
Calf—lateral malleolus	2.9 (1/34)§§	0 (0/34)¶¶	0 (0/34)‡

*Abnormal if >6.0 ms.

†Abnormal if <2.0 mV.

‡Abnormal if <40 m/s.

§Abnormal if >5.6 ms.

||Abnormal if <3 mV.

¶Abnormal if >3.5 ms.

♯Abnormal if <6 mV.

**Abnormal if <48 m/s.

††Abnormal if >2.7 ms.

††Abnormal if <20 µV.

§§Abnormal if >4.2 ms.

¶¶Abnormal if <5 µV.

TABLE 5. Blood Tests Investigating Factors Potentially Associated With Paresthesias Among Cases, Clinic Controls, and Community Controls

Blood Measure	Cases				Clinic Controls				Community Controls				P Value*	P Value**
	N	Median	Min	Max	N	Median	Min	Max	N	Median	Min	Max		
Vitamin B12 (pg/mL)	41	546	230	2500	38	439	222	2500	19	493	209	1150	0.8	0.5
Methylmalonic acid (μmol/L)	40	0.145	0.05	0.33	38	0.15	0.05	0.29	20	0.155	0.11	0.47	0.4	0.4
Folate (ng/mL)	40	12.8	4.6	60	38	12.1	5	60	19	13.5	9.2	60	0.2	0.8
Hemoglobin A1c (%)	41	5.4	4.8	6.7	38	5.45	4.3	6.4	18	5.35	5	9.4	0.7	0.9
Glucose, fasting (mg/dL)	40	86.5	70	107	38	85	64	111	18	83	59	196	0.5	0.4
ANA (inverse titers)	32	20	20	1280	25	20	20	640	15	20	20	160	0.5	0.2
C-reactive protein (mg/L)	41	2.3	0.1	11	38	2.05	0.1	10	19	1.8	0.4	9.8	0.6	1
Gliadin peptide antibody IgA (units)	39	5	2	37	36	5	2	15	20	5	3	15	0.4	0.3
Gliadin peptide antibody IgG (units)	39	3	2	28	36	2	2	16	20	3	1	5	0.4	0.2
Lead concentration (μg/dL)	39	1	1	2.9	37	1	1	2.4	18	1	1	3	0.4	0.2
Lyme antibody IgG/IgM (Western blot [normal < 0.91])	40	0.32	0.11	1	37	0.34	0.08	1.1	20	0.31	0.11	2.14	0.9	0.8

*P from the Kruskall-Wallis test comparing cases, clinic controls and community controls.

**P from the Wilcoxon rank-sum test comparing cases and clinic controls.

*Lead levels shown by the laboratory as less than 2.0 were assigned a value of 1.0.

When categorized by frequency of paresthesia symptoms, there was a suggestion ($P=0.10$, Fisher exact test) of a trend of increasing proportion of subjects with abnormal epidermal nerve fiber density with increasing frequency of paresthesia (0% [0/2] of those with frequency of "occasional" versus 50% [8/16] of those with frequency of "often" versus 68% [13/19] of those with frequency of "almost continuous;" $P=0.10$, Cochran-Armitage exact trend test). Scores on the Utah Early Neuropathy scale also were higher (worse) among persons with abnormally reduced small nerve densities compared to those with normal epidermal nerve fiber densities; this difference remained significant after adjustment for age ($P=0.01$).

Association of Paresthesias With Known Causes of Neuropathy

None of the blood tests for factors commonly associated with peripheral neuropathy differed significantly across groups (Table 5). Syphilis testing revealed one positive individual who was referred for treatment and five individuals (all cases) whose syphilis tests

indicated either false-positive screening tests or syphilis that had been treated and resolved. When blood tests were dichotomized into abnormal versus normal, the only blood test that showed significant heterogeneity across groups was methylmalonic acid, with community controls having a higher prevalence of abnormally high levels (suggestive of vitamin B12 deficiency, found in 2/20, 10%) compared to cases (0/40, 0%) or clinic controls (0/38, 0%) ($P=0.04$, Fisher exact test). There were no significant differences among groups in the prevalence of high hemoglobin A1c ($P=0.5$), high fasting glucose ($P=0.9$), or other dichotomized blood values. When blood lead concentrations were dichotomized at less than 2 μg/dL versus 2 to 4.9 μg/dL, there was a suggestion of an excess of higher lead concentrations in cases (15.4%, 6/39, in the range of 2 to 4.9 μg/dL) compared to clinic controls (5.4%, 2/37, in the range of 2 to 4.9 μg/dL, $P=0.26$, Fisher exact test). Community controls had an intermediate prevalence (11.1%, 2/18) resulting in an P value comparing all 3 groups of 0.4 (Fisher exact test).

ANA titers did not differ among groups, and ANA patterns among ANA positivity also did not differ significantly across the

TABLE 6. Epidermal or Sweat Gland Nerve Fiber Densities in Cases and Historical Controls

Part A. Quantitative Comparison of Epidermal Nerve Fiber Densities

Group	Mean Density (Fibers/mm of Epidermal Length)	95% Confidence Interval	P Value*
Paresthesia cases ($n=37$)	4.70	3.95–5.45	<0.0001
Historical controls ($n=128$)	8.32	7.86–8.77	

Part B. Categorical Comparison of the Prevalence of Abnormally Reduced Nerve Fiber Densities Based on a Composite Evaluation of Epidermal and Sweat Gland Nerve Fiber Densities

Group	Composite Nerve Fiber Density Measure			P Value, Fisher Exact Test
	Abnormal (n, [%])	Normal	Total	
Paresthesia cases	21 (57%)	16 (43%)	37	< 0.0001
Historical controls	5 (3.9%)	123 (96%)	128	

*From generalized linear models regression analysis adjusted for age and gender.

three groups. Among the 14 ANA-positive cases, however, 2 (14%) had nucleolar ANA patterns and 1 (7%) had nuclear dots, whereas these patterns were not found in the 9 ANA-positive clinic controls or 6 ANA-positive community controls.

Serum protein electrophoresis (SPEP) results did not reveal important differences across groups: 67% (26/39) of cases, 71% (27/38) of clinic controls and 85% of community controls had normal SPEP findings. Immunofixation electrophoresis results also did not differ among groups; no monoclonal bands were detected in 90% (36/40) of cases, 84% (32/38) of clinic controls, and 85% (17/20) of community controls.

On interview, lead exposure since 9/11 was more often reported by cases (11/41, 26.8%) than clinic controls (2/38, 5.3%) or community controls (0/20, 0%) (Fisher exact $P=0.003$). A majority (64%; 7/11) of the cases who said they had been exposed to lead since 9/11 reported these exposures occurred in the home, while 27% (3/11) reported WTC-related lead exposures, and 9% (1/11) reported hobby-related lead exposures. There also were suggestions of differences in post-9/11 exposures to mercury, with 9.8% (4/41) of cases reporting such exposure compared to 0% (0/38) of clinic controls, and 0% (0/20) of community controls ($P=0.07$, Fisher exact test); and exposure to solvents, with 14.6% (6/41) of cases reporting such exposure compared to 5.3% (2/38) of clinic controls and 0% (0/20) of community controls ($P=0.12$, Fisher exact test).

DISCUSSION

This study shows that the high prevalences of paresthesia we and others have observed among WTC-exposed individuals were associated with abnormal neurological examination findings. Importantly, pathologic evaluation revealed reduced epidermal or sweat gland nerve fiber densities among 57% of cases. Nerve conduction studies showed reduced sural to radial sensory nerve amplitude ratios in 29.4% of cases. Clinical examinations and histories, conducted with standardized neuropathy scales revealed findings consistent with both small and large fiber neuropathy more often in cases than in either of the two control groups. Blood tests did not reveal any other clear associations with non-WTC-related disease, metabolic, or toxic etiologies of peripheral neuropathy.²⁷ Paresthesia and abnormal neurologic findings were infrequent among the community control group. These findings, including objective measures of disease, support an association between WTC exposures and neurological injury or disease in a substantial proportion of WTC-exposed members of the general community.

WTC-associated dust and fumes contained heavy metals,^{7–9} including lead, mercury and arsenic, and complex hydrocarbons.^{7–9} Exposure to these materials occurred both at the time of the disaster when massive dust clouds were released upon collapse of the towers, and subsequently when materials were re-suspended. Workers involved in WTC clean-up activities also may have been exposed to neurotoxic cleaning agents or other substances after 9/11 at work, during hobbies or at home. These exposures provide biologic plausibility for the findings and are consistent with reports of small fiber neuropathies associated with exposure to environmental toxins, including lead, solvents and chemical warfare agents.^{28–37}

The high prevalence of small fiber abnormalities in our sample of paresthesia cases suggests that a small nerve fiber condition may be operative. Intraepidermal nerve fiber density measurement has been recommended as a sensitive and specific tool for diagnosis of small nerve fiber disease.³⁸ Nerve conduction velocities, which evaluate large fiber nerve function, were abnormal in 29.4% of paresthesia cases and reduced nerve action potentials, were observed in up to 36% of subjects depending on the nerve tested, suggesting axonal injury in some subjects.³⁹

The mechanisms by which toxic exposures contribute to the development of paresthesias are unclear. Small fiber neuropathies have been associated with exposure to environmental toxins. Prior

studies have suggested autoimmune factors may play a role in chemically mediated neuropathy.⁴⁰ We observed no significant difference in ANA positivity or SPEP findings among groups. ANA nucleolar and nuclear dot patterns, present in several cases but not in controls, suggests a possible role of autoimmune disorders in a small fraction of WTC-associated paresthesia cases.

The possibility exists that post-9/11 exposure to environmental toxins contributed to the neurologic symptoms. More cases than controls reported exposures to lead since 9/11, and there was a suggestion in the data that blood lead concentrations in the range of 2 to 4.99 $\mu\text{g}/\text{dL}$ were present more often among cases than clinic controls. Lead levels below 5 $\mu\text{g}/\text{dL}$, however, have not been reported to be associated with neuropathy. Furthermore, it is not possible to know whether blood lead concentrations measured during the study might be due to exposures that occurred around September 11, 2001, rather than previous or substantially after that date. An alternative explanation, might be recall bias, with cases more likely to remember and report exposures to neurotoxins than controls, perhaps because of concern for their symptoms. Participation bias also might have been present if persons with post-9/11 neurotoxin exposures were more likely to enroll due to fears of health consequences and out of a desire to obtain related testing and medical advice.

Heights were greater among cases with reduced epidermal or sweat gland nerve fiber densities compared to cases with normal densities, even after adjustment for sex and age. This suggests the possibility of a length-dependent effect. Greater height is a known risk factor for diabetic peripheral neuropathy.^{41,42}

There are potential limitations to our study. We did not conduct skin punch biopsies in some subjects due to extended blood pressure or heart rate excursions well beyond normal. Some subjects refused the procedure despite initial consent. Nerve conduction studies were not conducted in a few subjects due to equipment problems or refusal because the subjects had previously undergone nerve conduction studies elsewhere. Our dependence on historical controls as a comparison group for the intraepidermal nerve fiber measurements in cases could have led to bias if there had been drift in the evaluating pathologists' methods over time. Observer bias may have occurred due to the lack of masking of the study neurologists,⁴³ a situation necessitated by the neurologists' conduct of the skin punch biopsies and nerve conduction studies that were performed only among cases. Inclusion criteria for the study were based on a screening questionnaire that may have caused under- or over-reporting of paresthesia and the frequency thereof. Two subjects changed their responses to the question on the frequency of paresthesia, saying during screening that they "often" experienced paresthesia while during the main study interview they reported experiencing this symptom "occasionally." It was of interest, however, that neither of these subjects had reduced intraepidermal small nerve density, and that there was an increasing prevalence of abnormal intraepidermal nerve fiber densities as the reported frequency of paresthesia increased from "occasionally" to "often" to "almost continuous."

We enlisted paresthesia cases and clinic controls from the WTC EHC, meaning that subjects in each of those groups were exposed to the WTC disaster. In an ideal case-control study design, cases and controls would be drawn at random from pools of subjects with and without disease, and not from subject pools in which exposure has been in any way predetermined. The WTC EHC patient population is heterogeneous, however, and includes individuals with very different severities of WTC exposures, as well as widely different exposures to clean-up activities following the events of 9/11. This heterogeneity allowed us to ask whether there was any evidence for differential exposures among those with and without paresthesia, although this was not the primary aim of the study. We thus found non-significant, but suggestive exposure differences indicating that more paresthesia cases than clinic

controls had jobs that required cleaning of WTC dust, and more paresthesia cases than controls were heavily covered in dust on 9/11. The study's sampling scheme would be expected to bias the associations with WTC exposure variables towards the null. The lack of significance in these findings may reflect in part the fact that the study size was derived without consideration of analyzing associations between signs of neuropathy and WTC exposures. We have previously demonstrated significant associations between the symptom of paresthesia and WTC exposure variables.^{2,4}

In sum, this study demonstrates increased prevalence of clinical and laboratory-test abnormalities indicative of neuropathy among individuals with WTC exposure and paresthesia of the lower extremities. These abnormalities were observed more often among paresthesia cases than among members of the control groups. Our data are consistent with the hypothesis that WTC exposures or clean-up activities contributed to the development of peripheral neuropathy with paresthesia. Our findings reinforce the need for neurologic evaluation after toxic exposures in WTC-exposed as well as other populations. Future research should seek to identify possible mechanisms and describe the longitudinal course of signs and symptoms. Moreover, the impact of these symptoms on mental health and functional status should be investigated.

ACKNOWLEDGMENTS

The authors thank the research participants for their generosity, time and interest; and the staff and administration of Bellevue Hospital Center, NYC Health + Hospitals for their assistance and oversight.

REFERENCES

- Reibman J, Liu M, Cheng Q, et al. Characteristics of a residential and working community with diverse exposure to World Trade Center dust, gas, and fumes. *J Occup Environ Med*. 2009;51:534–541.
- Marmor M, Shao Y, Bhatt DH, et al. Paresthesias among community members exposed to the World Trade Center disaster. *J Occup Environ Med*. 2017;59:389–396.
- Reibman J, Levy-Carrick N, Miles T, et al. Destruction of the World Trade Center Towers. Lessons learned from an environmental health disaster. *Ann Am Thorac Soc*. 2016;13:577–583.
- Thawani S, Wang B, Shao Y, Reibman J, Marmor M. Time to onset of paresthesia among community members exposed to the World Trade Center Disaster. *Int J Environ Res Public Health*. 2019;16: pii: E1429.
- Stecker MM, Yu H, Barlev R, Marmor M, Wilkenfeld M. Neurologic evaluations of patients exposed to the world trade center disaster. *J Occup Environ Med*. 2016;58:1150–1154.
- Stecker M, Segelnick J, Wilkenfeld M. Analysis of short-term effects of World Trade Center dust on rat sciatic nerve. *J Occup Environ Med*. 2014;56:1024–1028.
- Lioy PJ, Weisel CP, Millette JR, et al. Characterization of the dust/smoke aerosol that settled east of the World Trade Center (WTC) in lower Manhattan after the collapse of the WTC 11 September 2001. *Environ Health Perspect*. 2002;110:703–714.
- Butt CM, Diamond ML, Truong J, Ikonomou MG, Helm PA, Stern GA. Semivolatile organic compounds in window films from lower Manhattan after the September 11th World Trade Center attacks. *Environ Sci Technol*. 2004;38:3514–3524.
- Rayne S, Ikonomou MG, Butt CM, Diamond ML, Truong J. Polychlorinated dioxins and furans from the World Trade Center attacks in exterior window films from lower Manhattan in New York City. *Environ Sci Technol*. 2005;39:1995–2003.
- Hintze J. PASS 2008 Kaysville, UT: NCSS, LLC, 2011.
- Ewing JA. Detecting alcoholism. The CAGE questionnaire. *JAMA*. 1984;252:1905–1907.
- Fritschi L. OccIDEAS – occupational exposure assessment in community-based studies. *Occup Med (Lond)*. 2019;69:156–157.
- Singleton JR, Bixby B, Russell JW, et al. The Utah Early Neuropathy Scale: a sensitive clinical scale for early sensory predominant neuropathy. *J Peripher Nerv Syst*. 2008;13:218–227.
- Hoffken O, Maier C, Richter H, Tegenthoff M, Schwenkreis P. A simplified screening protocol predicts pathological electromyographic results in patients with suspected polyneuropathy. *Int J Neurosci*. 2010;120:28–35.
- Tesfaye S, Boulton AJ, Dyck PJ, et al. Diabetic neuropathies: update on definitions, diagnostic criteria, estimation of severity, and treatments. *Diabetes Care*. 2010;33:2285–2293.
- Vinik AJ, Hayes RP, Oglesby A, et al. The development and validation of the Norfolk QOL-DN, a new measure of patients' perception of the effects of diabetes and diabetic neuropathy. *Diabetes Technol Ther*. 2005;7:497–508.
- Suanprasert N, Berk JL, Benson MD, et al. Retrospective study of a TTR FAP cohort to modify NIS+7 for therapeutic trials. *J Neurol Sci*. 2014;344:121–128.
- Dyck PJ, Carter RE, Litchy WJ. Modeling nerve conduction criteria for diagnosis of diabetic polyneuropathy. *Muscle Nerve*. 2011;44:340–345.
- Rutkove SB, Kothari MJ, Raynor EM, Levy ML, Fadic R, Nardino RA. Sural/radial amplitude ratio in the diagnosis of mild axonal polyneuropathy. *Muscle Nerve*. 1997;20:1236–1241.
- Yasemin E, Gongor YN, Ozisler C. Polyneuropathy and the sural/radial sensory nerve action potential ratio in primary Sjogren's syndrome. *Neurol Res*. 2019;42:17–21.
- Chan AC, Wilder-Smith EP. Small fiber neuropathy: getting bigger! *Muscle Nerve*. 2016;53:671–682.
- Lauria G, Hsieh ST, Johansson O, et al. European Federation of Neurological Societies/Peripheral Nerve Society Guideline on the use of skin biopsy in the diagnosis of small fiber neuropathy. Report of a joint task force of the European Federation of Neurological Societies and the Peripheral Nerve Society. *Eur J Neurol*. 2010;17:903–912. e44–e49.
- England JD, Gronseth GS, Franklin G, et al. Practice Parameter: evaluation of distal symmetric polyneuropathy: role of autonomic testing, nerve biopsy, and skin biopsy (an evidence-based review). Report of the American Academy of Neurology, American Association of Neuromuscular and Electromyography Medicine, and American Academy of Physical Medicine and Rehabilitation. *Neurology*. 2009;72:177–184.
- Binnicker MJ, Jespersen DJ, Rollins LO. Direct comparison of the traditional and reverse syphilis screening algorithms in a population with a low prevalence of syphilis. *J Clin Microbiol*. 2012;50:148–150.
- Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42:377–381.
- Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst*. 1959;22:719–748.
- Callaghan BC, Price RS, Chen KS, Feldman EL. The importance of rare subtypes in diagnosis and treatment of peripheral neuropathy: a review. *JAMA Neurol*. 2015;72:1510–1518.
- Pradhan S, Tandon R. N-hexane neuropathy with vertigo and cold allodynia in a silk screen printer: a case study. *Int J Occup Med Environ Health*. 2015;28:915–919.
- Guimaraes-Costa R, Schoindre Y, Metlaine A, et al. N-hexane exposure: a cause of small fiber neuropathy. *J Peripher Nerv Syst*. 2018;23:143–146.
- Terkelsen AJ, Karlsson P, Lauria G, Freeman R, Finnerup NB, Jensen TS. The diagnostic challenge of small fibre neuropathy: clinical presentations, evaluations, and causes. *Lancet Neurol*. 2017;16:934–944.
- Majersik JJ, Caravati EM, Steffens JD. Severe neurotoxicity associated with exposure to the solvent 1-bromopropane (*n*-propyl bromide). *Clin Toxicol (Phila)*. 2007;45:270–276.
- Sclar G. Encephalomyelradiculoneuropathy following exposure to an industrial solvent. *Clin Neurol Neurosurg*. 1999;101:199–202.
- Samukawa M, Ichihara G, Oka N, Kusunoki S. A case of severe neurotoxicity associated with exposure to 1-bromopropane, an alternative to ozone-depleting or global-warming solvents. *Arch Intern Med*. 2012;172:1257–1260.
- Karam C, Dyck PJ. Toxic neuropathies. *Semin Neurol*. 2015;35:448–457.
- Kushlaf HA. Emerging toxic neuropathies and myopathies. *Psychiatr Clin North Am*. 2013;36:209–218.
- Thomson RM, Parry GJ. Neuropathies associated with excessive exposure to lead. *Muscle Nerve*. 2006;33:732–741.
- Holisaz MT, Rayegani SM, Hafezy R, Khedmat H, Motamed MH. Screening for peripheral neuropathy in chemical warfare victims. *Int J Rehabil Res*. 2007;30:71–74.
- Devigili G, Tugnoli V, Penza P, et al. The diagnostic criteria for small fibre neuropathy: from symptoms to neuropathology. *Brain*. 2008;131:1912–1925.
- McCorquodale D, Smith AG. Clinical electrophysiology of axonal polyneuropathies. *Handb Clin Neurol*. 2019;161:217–240.
- Abou-Donia MB, Lieberman A, Curtis L. Neural autoantibodies in patients with neurological symptoms and histories of chemical/mold exposures. *Toxicol Ind Health*. 2018;34:44–53.

41. Adler AI, Boyko EJ, Ahroni JH, Stensel V, Forsberg RC, Smith DG. Risk factors for diabetic peripheral sensory neuropathy. Results of the Seattle Prospective Diabetic Foot Study. *Diabetes Care*. 1997;20:1162–1167.
42. Unmar Y, Zafar MI, Gao F. Factors associated with peripheral neuropathy in type 2 diabetes: subclinical versus confirmed neuropathy. *J Huazhong Univ Sci Technolog Med Sci*. 2017;37:337–342.
43. Delgado-Rodriguez M, Llorca J, Bias. *J Epidemiol Community Health*. 2004;58:635–641.

Appendix 1. Neurotoxins Queried as Part of the OcclIDEAS Instrument

Subjects were asked about chemical exposures that included arsenic, lead, manganese, mercury, tin, acrylamide, carbon disulfide, ethylene oxide, methyltertiarybutylethane, methylchloride, *n*-hexane, perchlorethylene, toluene, trichloroethylene, pesticides, carbon monoxide, gasoline, and “other chemical exposures.”