

Secondhand Smoke Exposure Among Women and Children: Evidence From 31 Countries

Heather Wipfli, PhD, Erika Avila-Tang, PhD, MHS, Ana Navas-Acien, MD, PhD, MPH, Sungroul Kim, PhD, Georgiana Onicescu, ScM, Jie Yuan, BS, Patrick Breyse, PhD, MHS, and Jonathan M. Samet, MD, for the FAMRI Homes Study Investigators

Extensive research shows that secondhand smoke (SHS) exposure places adults and children at increased risk for premature death, diverse illnesses, and other adverse effects, such as reduced lung-function growth in children.¹ Worldwide, over 40% of men smoke tobacco, compared with only about 12% of women.² This global profile implies that women and children constitute the bulk of the population exposed to SHS. The World Health Organization (WHO) conducted the Global Youth Tobacco Survey between 1999 and 2005 and found that approximately 44% of youths were exposed to tobacco smoke at home and that 47% had at least 1 parent who smoked.³ There is insufficient information, however, regarding levels of SHS in households and the range of exposures among women and children throughout the world. Questionnaires have been the most commonly used tool to assess the prevalence and intensity of SHS exposure at home.^{3–5} Although questionnaires can confirm that SHS exposure is taking place, they are not highly informative as to the specific level of exposure because of inherent limitations of the data collected.¹

For our study, we sought to quantify the levels of SHS exposure among women and children living with smokers in diverse climates and cultures. Measuring air nicotine and hair nicotine concentrations are validated methods of quantifying airborne levels of SHS and the uptake of SHS in the body, respectively.^{6,7} We carried out a multicountry study measuring nicotine concentrations in the air of households with and without smokers and in the hair of women and children living in these households. Survey data were also collected to evaluate smoking behaviors and smoking policies in the households and around the children, as well as perceptions and attitudes about SHS.

Objectives. We sought to describe the range of exposure to secondhand smoke (SHS) among women and children living with smokers around the world and generate locally relevant data to motivate the development of tobacco control policies and interventions in developing countries.

Methods. In 2006, we conducted a cross-sectional exposure survey to measure air nicotine concentrations in households and hair nicotine concentrations among nonsmoking women and children in convenience samples of 40 households in 31 countries.

Results. Median air nicotine concentration was 17 times higher in households with smokers ($0.18 \mu\text{g}/\text{m}^3$) compared with households without smokers ($0.01 \mu\text{g}/\text{m}^3$). Air nicotine and hair nicotine concentrations in women and children increased with the number of smokers in the household. The dose-response relationship was steeper among children. Air nicotine concentrations increased an estimated 12.9 times (95% confidence interval = 9.4, 17.6) in households allowing smoking inside compared with those prohibiting smoking inside.

Conclusions. Our results indicate that women and children living with smokers are at increased risk of premature death and disease from exposure to SHS. Interventions to protect women and children from household SHS need to be strengthened. (*Am J Public Health.* 2008;98:672–679. doi:10.2105/AJPH.2007.126631)

METHODS

Overview

We conducted a cross-sectional exposure survey among adult nonsmoking women and children younger than 11 years in Latin America, Asia, Eastern Europe, and the Middle East. We used passive air monitors to measure household air nicotine concentrations and a hair nicotine test to measure personal exposure to SHS.^{8,9}

Household Sampling

In each country, we selected at least 40 households that had a child younger than 11 years. In 80% of the households, at least 1 male was currently smoking and living in the house at the time of the study; we preferred that this person be the father or primary male caregiver, because he would have the most direct contact with the exposed child. Active smoking was defined as having smoked within the past 30 days. In most of the countries, we selected our convenience sample from the country's capital city, because

sampling frames were not available. Some country teams chose to select the households from previous or ongoing studies (Dominican Republic, Mexico, Nepal, Peru, and the Philippines). Other teams reached out through health clinics (Armenia, Georgia, Guatemala, Hong Kong, Latvia, Panama, and Uruguay), schools (Poland, Romania, Russia, Ukraine, and Venezuela), community workers (Argentina, Cambodia, China, Egypt, India, Indonesia, Laos, Taiwan, Thailand, and Vietnam), and through their organizational staff (Brazil, Malaysia, and Turkey). Whenever possible, households were selected from different socioeconomic levels and different neighborhoods. All sampling strategies and recruitment procedures were reviewed and approved by the Johns Hopkins research team.

Data Collection

Trained field staff administered questionnaires to the primary male and female caregivers in each household. The questionnaire had 4 main sections, including personal and socioeconomic characteristics, smoking his-

tory, exposure to SHS, and attitudes and beliefs about the harms of smoking and tobacco control policies. We considered a woman to be exposed to SHS outside the household if she reported SHS exposure at any of the following places: at work, in any indoor place that was not the household, during transportation, or outdoors. We considered a child to be exposed to SHS outside the household if the child's caregiver reported witnessing the child's SHS exposure at any of the following places: in school, in daycare, in any indoor place that was not the household, during transportation, or outdoors. We classified a household as having a no-smoking policy if the respondents indicated that smoking was not permitted anywhere inside the household.

Sample Collection

Passive air monitors were installed in each household to measure air nicotine for 7 days. Monitors were placed in the main room in which the family congregated and were not highly visible or accessible to the residents.

Hair samples were collected from the primary female caregiver and from 1 child younger than 11 years in each household. In households where the mother or primary female caregiver was a smoker, hair samples from women were not taken. A small sample of hair (approximately 30–50 strands) was cut near the hair root from the back of the scalp where the growth pattern is the most uniform. Hair samples were immediately placed in a labeled plastic bag that was sealed for storage and transportation. Information was also collected on prior chemical treatments of the hair and on time–activity patterns in relation to SHS exposure.

Laboratory Analysis

Air Nicotine. The nicotine collected by each passive sampler was extracted into heptane with an internal standard (isoquinoline; Aldrich Chemical Company, Milwaukee, Wis), and then injected into a gas chromatograph, coupled with a nitrogen phosphorus detector (5890A; Agilent, Santa Clara, Calif) and a capillary column (30 m × 0.32 mm id, 0.5-μm film thickness, [HP-5; Agilent]).^{8,10} We started the gas chromatograph oven temperature at 130°C, increased the temperature to 160°C at a rate of 5°C per min, then again

increased the temperature to 180°C at a rate of 10°C per min. For quality control purposes, 10% of samples were duplicates and blanks. The correlation coefficient between the duplicate and original samples was 0.92 (n=95). Using the analytic results from blank samples, we found the median limit of detection to be 0.001 μg/m³ for a 7-day sample. Final concentrations were provided after subtraction of background levels from the blank samples.

Hair Nicotine. Nicotine in a 30-mg sample of hair taken from within 3 cm of the hair root was extracted using an isotope dilution method with an internal standard (Nicotine-d₃; Supelco, Bellefonte, Pa). Our method was modified from that developed by Kintz.⁹ Hair nicotine analysis was conducted using gas chromatography and mass spectrometry (GC–17/MS-QP5000; Shimadzu, Kyoto, Japan) in SIM and splitless mode. The gas chromatograph oven temperature was maintained at 70°C for 1 minute, increased to 280°C at a rate of 25°C per minute, and held for 1 minute. Nicotine was separated using a capillary column (30 m × 0.25 mm id, 0.25-μm film thickness [Rtx-624, Restek, Bellefonte, Pa]).

For quality control purposes, measurement performance of hair nicotine was examined with respect to bias and precision. Bias was evaluated by calculating recovery (percentage), and precision was determined as the relative standard deviation (percentage) using nonsmokers' hair samples spiked with 2 different concentrations (e.g., 0.67 and 3.33 ng/mg; n=5 within batch; n=30 between batches). The recoveries from within batch were 84.8% and 88.1% with relative standard deviations of 8.5% and 6.3% for the 2 concentrations, respectively; recoveries from between batches were 73% and 83% with relative standard deviations of 21% and 11%, respectively. In addition, 10% of the hair samples from each batch went through duplicate analyses. The coefficient of intersample correlation was 0.97 (n=172). Blank samples were used to determine the limit of detection and blank-corrected hair nicotine concentration. The median limit of detection was 0.02 ng/mg for a 30-mg hair sample.

Data Analysis. Microsoft Access version 2003 (Microsoft Corporation, Redmond, Wash) was provided to each country investi-

gator for data entry using a common format. The data collected in the country databases were checked and reviewed centrally for completeness and consistency and merged with air nicotine and hair nicotine data from the laboratory. We calculated percentages, means, medians, and interquartile ranges for demographic and housing characteristics, smoking behavior, and self-reported SHS exposure. Hair and air nicotine concentrations were skewed and log⁻¹⁰ transformed for statistical analysis. We examined the relationship between air and hair nicotine concentrations using the Pearson correlation coefficient. Multilevel linear models that allowed for country-specific intercepts were used to identify factors associated with differences in concentrations of air nicotine and hair nicotine. Factors of interest were the number of smokers in the household, number of cigarettes smoked per day by the smokers in the household, smokers' behavior (smoking inside their household and smoking in the presence of their child), and women's and children's exposures to SHS outside their households. SAS version 9 was used for data checking and statistical analysis.¹¹

RESULTS

A total of 1284 households from 31 countries were surveyed; these included 208 households with no smokers, 784 households with only 1 male smoker, and 292 households with 2 or more smokers (Table 1). Participants in the households covered a range of socioeconomic levels within and between countries. In Europe, 53% of the male caregivers living in the households had completed some education beyond high school, followed by 39% in Latin America and 16% in Asia. Approximately 50% of women in all regions worked outside of the household. Among smoking households, 38% of households in Europe and the Middle East had 1 or more smokers in that household, compared with 24% in Latin America and 21% in Asia. In Europe, smokers consumed a median of 18 cigarettes a day compared with 14 in Asia and 12 in Latin America. Most smokers indicated that they smoked in their households, which varied little between regions. Across all regions, approximately 82% of smokers

TABLE 1—Characteristics of Households in Which Women and Children are Exposed to Secondhand Smoke (SHS): International Survey, 2006

Country	No. of Participant Households	Households		Smokers			Nonsmoking Women		Children Exposed to SHS Outside the Home, %
		Mean Outdoor Temperature, °C	Smoking Households With ≥ 2 Smokers, %	No. Cigarettes Smoked/Day, Mean	Smokes in the Home, %	Smokes Near Child, %	No. Working Outside the Home	Exposed to SHS Outside the Home, %	
North and South America	364	21	24	12	81	81	48	64	48
Argentina	40	13	22	18	83	92	20	47	20
Brazil	40	22	16	21	94	90	50	78	70
Dominican Republic	40	24	19	11	75	75	22	66	63
Guatemala	40	20	9	11	85	79	69	69	50
Mexico	41	28	27	7	74	83	15	78	61
Panama	40	28	31	9	63	66	45	55	48
Peru	40	16	38	6	83	74	65	57	30
Uruguay	42	13	26	16	86	91	70	53	24
Venezuela	41	24	29	13	91	88	41	73	63
Asia	509	23	21	14	91	82	50	64	46
Cambodia	40	28	33	11	85	68	87	68	73
China	60	−5	5	19	98	98	41	73	43
Hong Kong	40	27	23	15	96	76	50	94	83
India	40	29	13	15	91	97	53	10	8
Indonesia	40	28	15	11	94	79	32	92	73
Laos	46	28	30	14	100	84	50	58	13
Malaysia	40	29	10	12	78	63	70	93	63
Nepal	40	2	34	12	87	80	28	29	13
Philippines	42	29	26	14	83	69	28	92	93
Taiwan	40	25	27	24	100	100	50	33	23
Thailand	40	29	21	9	82	82	68	50	33
Vietnam	41	26	21	12	97	88	54	71	44
Europe and Middle East	411	12	38	18	84	82	47	68	50
Armenia	51	8	41	25	98	98	30	76	65
Egypt	40	19	17	16	100	86	18	60	63
Georgia	40	7	19	19	97	89	35	84	50
Latvia	40	12	57	16	98	85	78	40	35
Poland	40	18	61	17	98	92	81	56	28
Romania	40	1	17	22	90	78	91	74	33
Russia	40	7	52	14	79	72	67	63	58
Syria	40	20	50	17	97	91	19	67	67
Turkey	40	20	26	16	78	62	30	79	55
Ukraine	40	12	31	14	53	69	62	57	40
All Regions	1284	19	27	15	88	82	49	65	48

indicated that they smoked around their children. The majority of women indicated that they were exposed to SHS outside the household (Europe, 68%; Asia, 64%; Latin America, 64%). In all regions, approximately half of the children in the study were reported as being exposed to SHS outside the household.

Air nicotine concentrations were detected in 88% of households with at least 1 smoker and in 60% of households with no smokers.

Median air nicotine concentrations within households with smokers were highest in Europe (0.58 $\mu\text{g}/\text{m}^3$), followed by Latin America (0.16 $\mu\text{g}/\text{m}^3$) and Asia (0.09 $\mu\text{g}/\text{m}^3$; Figure 1). Hair nicotine concentrations were detected in 78% of women and children living with a smoker, and in 59% of women and children not living with a smoker. Median hair nicotine concentrations for both women and children living with smokers were highest

in Asia (0.50 ng/mg and 0.87 ng/mg, respectively), followed by Europe and the Middle East (0.37 ng/mg, 0.72 ng/mg), and Latin America (0.26 ng/mg, 0.50 ng/mg; Figure 2). Air nicotine concentrations in countries with mean outdoor temperatures lower than 20°C at the time of measurement were over 3 times higher (95% confidence interval [CI]=1.35, 8.90) than countries with warmer temperatures.




FIGURE 1—Median and interquartile range of air nicotine concentrations in homes, by country in nonsmoking households (a) and smoking households (b).

Hair nicotine concentrations were positively correlated with the concentration of air nicotine in the households. The correlation was

stronger among children ($r=0.36$, $P<.001$) than among their mothers ($r=0.25$, $P<.001$). Hair nicotine concentrations of women and

children living in the same household were also positively correlated ($r=0.50$, $P<.001$). Median hair nicotine concentrations in children (0.68 ng/mg) were generally higher than those found in women (0.40 ng/mg; Figure 2). Children younger than 5 years living with smokers had hair nicotine concentrations that were nearly twice as high (estimated geometric mean ratio [GMR]=1.93; 95% CI=1.19, 3.12) as children 5 years or older living with smokers.

With a multilevel linear model that allowed for a country-specific random intercept, we evaluated determinants of geometric mean values of air and hair nicotine concentrations (Table 2). A dose-response group relationship was evident between air nicotine concentration and the number of smokers living in a household. We found an estimated 21.6 times increase (95% CI=15.7, 29.6) in air nicotine concentration inside households with 2 or more smokers compared with households with no smokers. Households allowing smoking inside had an estimated 12.9 times increase (95% CI=9.4, 17.6) in air nicotine concentrations compared with households with a no-smoking policy.

A dose-response group relationship was also found between hair nicotine concentrations among women and children and the number of smokers living in a household (Table 3). The increase in hair nicotine concentration among participants living with 2 or more smokers compared to those living with no smokers was greater among children (estimated GMR=3.8; 95% CI=3.0, 4.9) than among women (estimated GMR=2.1; 95% CI=1.7, 2.9). A child whose father and mother smoked had a 2.9 times increase (95% CI=2.1, 4.0) in hair nicotine concentrations compared with a child whose father smoked but whose mother did not.

DISCUSSION

We initiated this project to gain a better perspective of the range of exposures among women and children living within households with smokers, to encourage increased tobacco research capacity in developing countries, and to generate locally relevant data to motivate the development of approaches for reducing household SHS exposure. The resulting




FIGURE 2—Median and interquartile range of hair nicotine concentrations in women and children, by country in nonsmoking households (a) and smoking households (b).

information regarding the level of SHS exposure among women and children and the relative risks associated with that exposure can be used to strengthen and design interventions to protect women and children.

Using a common protocol in 31 countries, we measured concentrations of airborne nicotine in households and concentrations of

nicotine in the hair of women and children living in those households. We found that exposure to SHS is ubiquitous in households with smokers and that smokers smoke around their children with little restraint. Concentrations of air nicotine were higher in households with a smoker and increased with the number of smokers residing in the household.

The hair nicotine biomarker confirmed the uptake of SHS by women and children living with smokers.

Our results expand the body of information on levels of SHS in households, as previous studies were limited largely to developed countries.¹ The countries we surveyed encompassed a wide range of housing types, climates, and smoking profiles, and these differences were reflected in our measurements. Air nicotine concentrations were higher for measurements made at colder temperatures. This inverse association is most likely indicative of housing type. In some of the warmer countries, (e.g., Vietnam, Thailand, and Malaysia), many households had walls and floors that you could see through and through which air could easily pass, so smoke was probably rapidly diluted by natural ventilation. The highest levels of SHS were generally found in countries with high prevalence and high consumption of cigarettes (e.g., Poland, Syria, Turkey, Argentina, and Georgia).¹² The countries with the highest levels of air nicotine positively corresponded with the countries having the highest percentage of youth reporting SHS exposure at home in the Global Youth Tobacco Survey.³

Hair nicotine levels provide information complementary to that obtained from the questionnaire on exposure and reflect direct contact with SHS by children and women, whether at home or elsewhere. In some countries with low concentrations of air nicotine, hair nicotine concentrations were in the same range as in the countries with higher airborne nicotine. Children in Vietnam, for example, had a median hair nicotine concentration of 1.21 ng/mg, in the midrange for the 31 countries. The apparent discrepancy between air and hair nicotine concentrations is likely to reflect the closeness of interactions of the children and women with smoking household members, as well as differences in nicotine uptake and metabolism. Closeness of interactions and differences in nicotine uptake and metabolism are also likely to explain the higher concentrations of nicotine found in younger children.

Several studies in Western countries have measured hair nicotine concentrations in young children or women,^{13–18} and the range of values was similar to our findings. Across

TABLE 2—Geometric Means (GMs) and Geometric Mean Ratios (GMRs) of Air Nicotine Concentrations in Households in Which Women and Children are Exposed to Secondhand Smoke (SHS), by Household Characteristics: International Survey, 2006

Characteristic	Unadjusted GM, $\mu\text{g}/\text{m}^3$	GMR (95% CI)
Smokers in household, no.		
None (Ref)	0.01	1.0
1	0.13	10.3 (7.9,13.6)
≥ 2	0.3	21.6 (15.7, 29.6)
Cigarettes smoked per day, ^a no.		
< 10 (Ref)	0.07	1.0
10–19	0.15	2.1 (1.5, 2.8)
≥ 20	0.34	3.2 (2.4, 4.3)
Policy restricting smoking in the home		
Yes (Ref)	0.01	1.0
No	0.15	12.9 (9.4, 17.6)
Mean outdoor temperature, $^{\circ}\text{C}$		
> 20 (Ref)	0.09	1.0
10–20	0.32	3.47 (1.35, 8.90)
< 10	0.35	3.51 (1.24, 9.93)

Note. CI = confidence interval. GMRs obtained from the regression model that allowed for country-specific random intercepts.

^aAmong respondents who smoked in the household.

the 31 countries we surveyed, median concentrations of hair nicotine in women ranged from 0.02 to 2.5 ng/mg (Figure 2). Similarly, in 2001 Jaakkola et al. found that Finnish women's hair nicotine concentrations ranged from 0.9 ng/mg to about 2 ng/mg, depending on exposure in the household.¹⁸ In children, median concentrations across the 31 countries ranged from 0.02 ng/mg to 4.5 ng/mg. In 2003, a study among Latino children in the United States found a median of 0.5 ng/mg in children exposed to parental smoking.¹⁹ Similar to our findings, studies in New Zealand¹⁴ and Norway¹³ found dose-response group relationships between hair nicotine concentration and the number of smokers in the house and the amount smoked by the mother, respectively.

Our multilevel linear regression model showed a strong effect of smoking on nicotine concentrations in the household and the results of limiting smoking in the household. There was a strong dose-response group relationship between the number of smokers living within a household, the number of cigarettes being smoked, and air nicotine concentration (Table 2). Voluntary smoking bans in households with

smokers were associated with substantial reductions in the concentration of nicotine in the air.

Our findings are potentially limited by the selection of households on a convenience basis and by the number of households included in each country. These aspects of the design reflected our intent of reaching as many countries as was feasible, given the resources available. Other studies, such as the Global Youth Tobacco Survey, provide data pertaining to the percentage of children living with smokers. Although the samples in our study were not necessarily representative of women and children living with smokers within a country as a whole, our sampling approach would not have introduced bias with regard to the relation between smoking in the households and the outcome measures. For assessing airborne nicotine, the passive sampling method we used is well standardized.⁸ The monitors measured nicotine continuously, not just while the household was occupied, so reported concentrations are likely to be lower than those present during the period of actual exposure.

Hair nicotine concentrations have been used increasingly during the past decade to assess personal SHS exposure; hair is easier and less

expensive to sample, store, and transport than is urine, saliva, or serum.⁷ Moreover, hair nicotine has the advantage of characterizing SHS exposure over a longer period of time than does blood or urine cotinine, with each centimeter of hair length representing a month of exposure.⁹ Measuring hair nicotine concentration provides an unbiased indicator of exposure not subject to the reporting bias that may have affected questionnaires or to behavior changes by smokers influenced by active monitoring.

For hair samples, a majority (60%) of the women sampled had applied some form of chemical treatment to their hair (dyes or permanents). The effect of hair treatment on nicotine concentrations is not yet well characterized, although reports from the literature indicate that hair treatments tend to reduce the measured concentration.^{7,20} On the basis of prior studies of the effect of chemical treatments on hair nicotine concentrations, the hair nicotine concentrations measured for women in our study may be underestimations. Nevertheless, we found good reproducibility and the expected relationships between the reported level of smoking in the household, airborne nicotine concentration, and hair nicotine concentration (Table 2).

A growing number of countries have banned smoking in public places, and more are contemplating new smoking restrictions as they meet their commitments under the WHO Framework Convention on Tobacco Control (FCTC).²¹ Article 8 of the FCTC legally binds all ratifying nations to implement effective measures to protect people from exposure to tobacco smoke, to the extent they have the power to do so at the national level. The language further specifies that these measures must apply to "indoor workplaces, public transport, indoor public places, and, as appropriate, other public places."²² However, as with other formal regulatory measures aimed at eliminating SHS exposure in public areas, SHS exposure in households is not covered under the FCTC. If protection from SHS is to be complete, particularly for women and children, SHS exposure should be eliminated in public areas by government authorities through smoke-free legislation and by families in their households.

TABLE 3—Geometric Means (GMs) and Geometric Mean Ratios (GMRs) of Hair Nicotine Concentrations Among Women and Children Living in Households Exposed to Secondhand Smoke, by Household Characteristics: International Survey, 2006

Characteristic	Women		Children	
	Unadjusted GM, ng/mg	GMR (95% CI)	Unadjusted GM, ng/mg	GMR (95% CI)
Smokers in household, no.				
None (Ref)	0.2	1.0	0.26	1.0
1	0.45	1.9 (1.5, 2.3)	0.68	2.3 (1.8, 2.8)
≥ 2	0.43	2.2 (1.7, 2.9)	1.04	3.8 (3.0, 4.9)
Cigarettes smoked per day, ^a no.				
< 10 (Ref)	0.3	1.0	0.49	1.0
10–19	0.44	1.4 (1.1, 1.7)	0.85	1.5 (1.2, 1.9)
≥ 20	0.59	1.7 (1.4, 2.2)	0.97	1.5 (1.2, 1.9)
Policy restricting smoking in the home				
Yes (Ref)	0.18	1.0	0.22	1.0
No	0.44	1.8 (1.4, 2.3)	0.77	2.6 (2.0, 3.3)
At least 1 smoker smokes near child				
No (Ref)	0.64	1.0
Yes	0.77	1.1 (0.9, 1.5)
Smokers sharing child's bedroom, no.				
None (Ref)	0.46	1.0
1	0.91	2.1 (1.6, 2.5)
2	1.6	3.4 (2.3, 5.0)
3	1.91	4.7 (1.5, 14.2)
Mother also smokes				
No (Ref)	0.60	1.0
Yes	1.42	2.9 (2.1, 4.0)
Cigarettes smoked per day, ^b no.				
< 10 (Ref)	0.69	1.0
10–19	1.59	2.6 (1.6, 4.4)
≥ 20	3.17	3.7 (2.3, 6.1)
Child's age, y				
≥ 5 (Ref)	0.61	1.0
< 5	1.08	1.93 (1.19, 3.12)

Note. CI = confidence interval. GMRs obtained from the regression model that allowed for country-specific random intercepts. Ellipses indicate that the characteristic does not apply to women.

^aAmong respondents who smoked in the household.

^bAmong female smokers only.

SHS exposure is a cause of premature death and disease among women and children.¹ As shown in this and other studies, household SHS exposure makes a dominant contribution to the cumulative SHS exposures among women and children. In the United States, the majority of households with smokers now claim to have some policy restricting smoking inside.^{1,23} Such policies, which preferably allow smoking outside only, have been shown to reduce or even eliminate SHS exposure.¹

The data we collected show that household policies are not commonly in place in many countries. Instead, the data demonstrate a nearly universal lack of voluntary restrictions on smoking in households and a high percentage of smokers smoking around their children. Some evidence suggests that legislation for smoke-free public places has a positive effect on voluntary smoke-free households,²⁴ and smoke-free activities mandated by the FCTC may be beneficial for promoting

smoke-free environments. Education is also a necessary and effective strategy in promoting protection from household SHS.²⁵ To protect the health of millions of women and children throughout the world, all individuals should be made aware of the adverse consequences of SHS exposure and encouraged to make their households smoke-free. ■

About the Authors

Heather Wipfli, Erika Avila-Tang, Georgiana Onicescu, and Jonathan M. Samet are with the Department of Epidemiology, Institute for Global Tobacco Control, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD. Ana Navas-Acien, Sungroul Kim, and Patrick Breyse are with the Department of Epidemiology and the Department of Environmental Health Sciences, Johns Hopkins Bloomberg School of Public Health, Baltimore. Jie Yuan is with the Department of Environmental Health Sciences, Johns Hopkins Bloomberg School of Public Health, Baltimore.

Requests for reprints should be sent to: Heather Wipfli, Institute for Global Tobacco Control, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe St, W6027, Baltimore, MD 21205 (e-mail: hwipfli@jhsph.edu; adineva@jhsph.edu)

This article was accepted November 27, 2007.

Contributors

H. Wipfli, A. Navas-Acien, and J.M. Samet designed the study and wrote the article. S. Kim, J. Yuan, and P. Breyse conducted the lab analysis. H. Wipfli, E. Avila-Tang, A. Navas-Acien, and G. Onicescu analyzed the data.

Acknowledgments

Support for this project was provided by the Flight Attendants Medical Research Institute (FAMRI) in the United States through a Dr William Cahan Distinguished Professors award to J.M. Samet, and a Center of Excellence Award to the Johns Hopkins Medical Institutions. Support was also provided by the National Institutes of Health through the Fogarty International Center (grant ROI HL 73699).

The FAMRI Homes Study Investigators were responsible for the acquisition of the data and the analysis and interpretation of the data within their country. The FAMRI Homes Study Investigators includes Marta Angueira (Argentina); Arayik Sargsyan (Armenia); Valeska Figueiredo (Brazil); Sin Sovann (Cambodia); Gonghuan Yang, Shao-jun Ma (China); Sergio Diaz, Deborah Ossip-Klein, Essie Sierra (Dominican Republic); Mostafa Mohamed, Christopher Loffredo (Egypt); Giorgi Magradze, George Bakhturidze (Georgia); Joaquin Barnoya (Guatemala); Sophia S. C. Chan, T.H. Lam, Y.W. Mak (Hong Kong); Mira Aghi (India); Rita Damayanti (Indonesia); Janis Caunitis (Latvia); Maniphanh Vongphosy (Laos); Foong Kin (Malaysia); Luz Myrian Reynales (Mexico); M.R. Pandey (Nepal); Reina Roa (Panama); Alfonso Zavaleta (Peru); Maria Lourdes Reuillida (Philippines); Witold Zatonski, Krzysztof Przewozniak, Jakub Gumkowski, Pawel Polak, Malgorzata Zagroba, Janina Fetlinska (Poland); Florin Mihaltan (Romania); Vladimir Levshin (Russia); Wasim Maziak (Syria); Wayne Kao (Taiwan); Naowarut Charoencra (Thailand); Zeynep Onder (Turkey); Tatiana Andreeva (Ukraine); Adriana Blanco (Uruguay); Natasha Herrera (Venezuela); and Bao Chau (Vietnam).

Human Participant Protection

Participation in the study was voluntary and the study and consent procedures were approved by the Johns Hopkins Bloomberg School of Public Health human subjects institutional review board and by the appropriate ethics committee at the individual country level.

References

1. US Department of Health and Human Services. *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*. Washington, DC: US Dept of Health and Human Services; 2006.
2. Jha P, Chaloupka F. *Curbing the Epidemic: Governments and the Economics of Tobacco Control*. Washington, DC: The World Bank; 1999.
3. The GTSS Collaborative Group. A cross-country comparison of exposure to secondhand smoke among youth. *Tob Control*. 2006;15(suppl 2):ii4–19.
4. Halterman JS, Fagnano M, Conn KM, Szilagyi PG. Do parents of urban children with persistent asthma ban smoking in their homes and cars? *Ambul Pediatr*. 2006;6:115–119.
5. Cobanoglu N, Kiper N, Dilber E, et al. Environmental tobacco smoke exposure and respiratory morbidity in children. *Inhal Toxicol*. 2007;19:779–785.
6. Marbury MC, Hammond SK, Haley NJ. Measuring exposure to environmental tobacco smoke in studies of acute health effects. *Am J Epidemiol*. 1993;137:1089–1097.
7. Al-Delaimy WK. Hair as a biomarker for exposure to tobacco smoke. *Tob Control*. 2002;11:176–182.
8. Hammond S, Leaderer B. A diffusion monitor to measure exposure to passive smoking. *Environ Sci Technol*. 1987;21:494–497.
9. Kintz P. Gas chromatographic analysis of nicotine and cotinine in hair. *J Chromatogr*. 1992;580(1–2):347–353.
10. Hammond S, Leaderer B, Roche A, Schenker M. Collection and analysis of nicotine as a marker for environmental tobacco smoke. *Atmos Environ*. 1987;21:457–462.
11. SAS Software. Version 9, Edition 1. Cary, NC: SAS Institute Inc; 2003.
12. Mackay J, Eriksen M, eds. *The Tobacco Atlas*. Geneva, Switzerland: World Health Organization; 2002.
13. Nafstad P, Jaakkola JJ, Hagen JA, Zahlén K, Magnus P. Hair nicotine concentrations in mothers and children in relation to parental smoking. *J Expo Anal Environ Epidemiol*. 1997;7:235–239.
14. Al-Delaimy WK, Crane J, Woodward A. Questionnaire and hair measurement of exposure to tobacco smoke. *J Expo Anal Environ Epidemiol*. 2000;10:378–384.
15. Kalinic N, Skender L, Karacic V, Brcic I, Vadjic V. Passive exposure to tobacco smoke: hair nicotine levels in preschool children. *Bull Environ Contam Toxicol*. 2003;71:1–5.
16. Eliopoulos C, Klein J, Chitayat D, Greenwald M, Koren G. Nicotine and cotinine in maternal and neonatal hair as markers of gestational smoking. *Clin Invest Med*. 1996;19:231–242.
17. Eliopoulos C, Klein J, Phan MK, et al. Hair concentrations of nicotine and cotinine in women and their newborn infants. *JAMA*. 1994;271:621–623.
18. Jaakkola JJ, Jaakkola N, Zahlén K. Fetal growth and length of gestation in relation to prenatal exposure to environmental tobacco smoke assessed by hair nicotine concentration. *Environ Health Perspect*. 2001;109:557–561.
19. Woodruff SI, Conway TL, Edwards CC, Hovell MF. Acceptability and validity of hair collection from Latino children to assess exposure to environmental tobacco smoke. *Nicotine Tob Res*. 2003;5:375–385.
20. Pichini S, Altieri I, Pellegrini M, Pacifici R, Zuccaro P. Hair analysis for nicotine and cotinine: evaluation of extraction procedures, hair treatments, and development of reference material. *Forensic Sci Int*. 1997;84(1–3):243–252.
21. World Health Organization. *WHO Framework Convention on Tobacco Control*. Geneva, Switzerland: World Health Organization; 2003. Available at: http://www.who.int/tobacco/fctc/text/en/fctc_en.pdf. Accessed August 24, 2007.
22. World Health Organization. *WHO Framework Convention on Tobacco Control. Article 8: Protection from Exposure to Tobacco Smoke*. Geneva, Switzerland: World Health Organization; 2003. Available at: http://www.who.int/tobacco/framework/WHO_FCTC_english.pdf. Accessed August 24, 2007.
23. Centers for Disease Control and Prevention. State-specific prevalence of current cigarette smoking among adults and secondhand smoke rules and policies in homes and workplaces—United States, 2007. *MMWR Morb Mort Wkly Rep*. 2007;56(20):497–500.
24. Borland R, Yong H, Cummings K, Hyland A, Anderson S, Fong G. Determinants and consequences of smoke-free homes: findings from the International Tobacco Control (ITC) Four Country Survey. *Tob Control*. 2006;15(suppl 3):iii42–iii50.
25. World Health Organization. *Policy Recommendations on Protection from Exposure to Second-Hand Tobacco Smoke*. Geneva, Switzerland: World Health Organization; 2007.

This article has been cited by:

1. Judith A. Groner, Ana M. Rule, Sharon A. McGrath-Morrow, Joseph M. Collaco, Angela Moss, Susanne E. Tanski, Robert McMillen, Regina M. Whitmore, Jonathan D. Klein, Jonathan P. Winickoff, Karen Wilson. 2018. Assessing pediatric tobacco exposure using parent report: comparison with hair nicotine. *Journal of Exposure Science & Environmental Epidemiology* 24. . [\[Crossref\]](#)
2. Ludmila Sevcikova, Jana Babjakova, Jana Jurkovicova, Martin Samohyl, Zuzana Stefanikova, Erika Machacova, Diana Vondrova, Etela Janekova, Katarina Hirosova, Alexandra Filova, Michael Weitzman, Lubica Argalasova. 2018. Exposure to Environmental Tobacco Smoke in Relation to Behavioral, Emotional, Social and Health Indicators of Slovak School Children. *International Journal of Environmental Research and Public Health* 15:7, 1374. [\[Crossref\]](#)
3. Philip K. Pattemore, Karen M. Silvers, Chris M. Frampton, Kristin Wickens, Tristram Ingham, David Fishwick, Julian Crane, G. Ian Town, Michael J. Epton. 2018. Hair nicotine at 15 months old, tobacco exposure and wheeze or asthma from 15 months to 6 years old. *Pediatric Pulmonology* 53:4, 443-451. [\[Crossref\]](#)
4. Katherine A Moon, Ana M Rule, Hoda S Magid, Jacqueline M Ferguson, Jolie Susan, Zhuolu Sun, Christine Torrey, Salahaddin Abubaker, Vladimir Levshin, Aslı Çarkoğlu, Ghada Nasr Radwan, Maha El-Rabbat, Joanna E Cohen, Paul Strickland, Patrick N Breyse, Ana Navas-Acien. 2018. Biomarkers of Secondhand Smoke Exposure in Waterpipe Tobacco Venue Employees in Istanbul, Moscow, and Cairo. *Nicotine & Tobacco Research* 20:4, 482-491. [\[Crossref\]](#)
5. Hepzibah Muñoz Martínez, Ann Pederson. 2018. Global frameworks, local strategies: Women's rights, health, and the tobacco control movement in Argentina. *Global Public Health* 68, 1-11. [\[Crossref\]](#)
6. Allison J. Burbank, David B. Peden. 2018. Assessing the impact of air pollution on childhood asthma morbidity. *Current Opinion in Allergy and Clinical Immunology* 1. [\[Crossref\]](#)
7. Nancy Chin, Ann Dozier, Zahira Quinones, Sergio Diaz, Emily Weber, Hector Almonte, Arisleyda Bautista, Kiran Raman, Scott McIntosh, Deborah Ossip. 2017. A qualitative study of tobacco use in eight economically disadvantaged Dominican Republic communities. *Global Health Promotion* 24:4, 23-32. [\[Crossref\]](#)
8. Laura J Rosen, Eimi Lev, Nurit Guttman, Efrat Tillinger, Shira Rosenblat, David M Zucker, Vicki Myers. 2017. Parental Perceptions and Misconceptions of Child Tobacco Smoke Exposure. *Nicotine & Tobacco Research* 377. . [\[Crossref\]](#)
9. P. H. Lykhatskiy, L. S. Fira, L. E. Grytcishin. 2017. Biochemical mechanisms of stress development in rats of different age groups under the influence of sodium nitrite with the tobacco intoxication background. *The Animal Biology* 19:1, 65-72. [\[Crossref\]](#)
10. Shanta Pandey, Vithya Murugan, Yagya B. Karki, Amit Mathur. 2017. In-home Smoking in Households with Women of Reproductive Age in Nepal: Does Women's Empowerment Matter?. *Health & Social Work* 42:1, 32-40. [\[Crossref\]](#)
11. Lazarous Mbulo, Krishna Mohan Palipudi, Linda Andes, Jeremy Morton, Rizwan Bashir, Heba Fouad, Nivo Ramanandraibe, Roberta Caixeta, Rula Cavaco Dias, Trudy M A Wijnhoven, Mina Kashiwabara, Dharendra N Sinha, Edouard Tursan d'Espaignet. 2016. Secondhand smoke exposure at home among one billion children in 21 countries: findings from the Global Adult Tobacco Survey (GATS). *Tobacco Control* 25:e2, e95-e100. [\[Crossref\]](#)
12. Mohammad Hajizadeh, Arijit Nandi. 2016. The socioeconomic gradient of secondhand smoke exposure in children: evidence from 26 low-income and middle-income countries. *Tobacco Control* 25:e2, e146-e155. [\[Crossref\]](#)
13. Rachel Mann, Heather Thomson, Becky Reynolds, Amanda Amos, Kamran Siddiqi. 2016. The challenge of mothers learning about secondhand smoke (MLASS): a quasi-experimental, mixed methods feasibility study. *Pilot and Feasibility Studies* 2:1. . [\[Crossref\]](#)
14. John Marsh, Ann McNeill, Sarah Lewis, Tim Coleman, Manpreet Bains, Alexandra Larwood, Jacqueline Purdy, Laura L Jones. 2016. Protecting children from secondhand smoke: a mixed-methods feasibility study of a novel smoke-free home intervention. *Pilot and Feasibility Studies* 2:1. . [\[Crossref\]](#)
15. Bo Xi, Yajun Liang, Yunxia Liu, Yinkun Yan, Min Zhao, Chuanwei Ma, Pascal Bovet. 2016. Tobacco use and second-hand smoke exposure in young adolescents aged 12–15 years: data from 68 low-income and middle-income countries. *The Lancet Global Health* 4:11, e795-e805. [\[Crossref\]](#)
16. M.J. Lopez, X. Contente. 2016. Secondhand smoke exposure in children: It is time to move forward. *Revista Portuguesa de Pneumologia (English Edition)* 22:4, 187-188. [\[Crossref\]](#)
17. Deborah J. Ossip, Sergio Díaz, Zahira Quinones, Scott McIntosh, Ann Dozier, Nancy Chin, Emily Weber, Heather Holderness, Essie Torres, Arisleyda Bautista, José Javier Sánchez, Esteban Avendaño, Timothy De Ver Dye, Paul McDonald, Eduardo Bianco. 2016. Lessons Learned from Twelve Years of Partnered Tobacco Cessation Research in the Dominican Republic. *Journal of Smoking Cessation* 11:02, 99-107. [\[Crossref\]](#)

18. Pallavi Pant, Sarath K. Guttikunda, Richard E. Peltier. 2016. Exposure to particulate matter in India: A synthesis of findings and future directions. *Environmental Research* **147**, 480-496. [[Crossref](#)]
19. Michelle C. Kegler, Regine Haardörfer, Carla Berg, Cam Escoffery, Lucja Bundy, Rebecca Williams, Patricia Dolan Mullen. 2016. Challenges in Enforcing Home Smoking Rules in a Low-Income Population: Implications for Measurement and Intervention Design. *Nicotine & Tobacco Research* **18**:5, 976-981. [[Crossref](#)]
20. Lian Yang, Elisa K. Tong, Zhengzhong Mao, Teh-wei Hu, Anita H. Lee. 2016. A Clustered Randomized Controlled Trial to Reduce Secondhand Smoke Exposure Among Nonsmoking Pregnant Women in Sichuan Province, China. *Nicotine & Tobacco Research* **18**:5, 1163-1170. [[Crossref](#)]
21. Jonathan M. Samet. 2016. Epidemiology and the Tobacco Epidemic: How Research on Tobacco and Health Shaped Epidemiology. *American Journal of Epidemiology* **183**:5, 394-402. [[Crossref](#)]
22. Scott McIntosh, José Pérez-Ramos, Margaret M Demment, Carmen Vélez Vega, Esteban Avendaño, Deborah J Ossip, Timothy D Dye. 2016. Development and Implementation of Culturally Tailored Offline Mobile Health Surveys. *JMIR Public Health and Surveillance* **2**:1, e28. [[Crossref](#)]
23. Michelle C. Kegler, Regine Haardörfer, Lucja T. Bundy, Cam Escoffery, Carla J. Berg, Maria Fernandez, Rebecca Williams, Mel Hovell. 2015. Do partial home smoking bans signal progress toward a smoke-free home?. *Health Education Research* **64**, cyv066. [[Crossref](#)]
24. Laura Rosen, Vicki Myers, Jonathan Winickoff, Jeff Kott. 2015. Effectiveness of Interventions to Reduce Tobacco Smoke Pollution in Homes: A Systematic Review and Meta-Analysis. *International Journal of Environmental Research and Public Health* **12**:12, 16043-16059. [[Crossref](#)]
25. Nipapun Kungskulniti, Naowarut Charoenca, Jintana Peesing, Songwut Trangwatana, Stephen Hamann, Siriwan Pitayarangsarit, Hatai Chitanondh. 2015. Assessment of secondhand smoke in international airports in Thailand, 2013. *Tobacco Control* **24**:6, 532-535. [[Crossref](#)]
26. Anna Ahn, Kathryn M. Edwards, Carlos G. Grijalva, Wesley H. Self, Yuwei Zhu, James D. Chappell, Sandra R. Arnold, Jonathan A. McCullers, Krow Ampofo, Andrew T. Pavia, Anna M. Bramley, Seema Jain, Derek J. Williams. 2015. Secondhand Smoke Exposure and Illness Severity among Children Hospitalized with Pneumonia. *The Journal of Pediatrics* **167**:4, 869-874.e1. [[Crossref](#)]
27. Joseph G. Allen, Piers MacNaughton, Jose Guillermo Cedeno Laurent, Skye S. Flanigan, Erika Sita Eitland, John D. Spengler. 2015. Green Buildings and Health. *Current Environmental Health Reports* **2**:3, 250-258. [[Crossref](#)]
28. Van T. Tong, Paola Morello, Alicia Alemán, Carolyn Johnson, Patricia M. Dietz, Sherry L. Farr, Agustina Mazzoni, Mabel Berrueta, Mercedes Colomar, Alvaro Ciganda, Ana Becú, Maria G. Bittar Gonzalez, Laura Llambi, Luz Gibbons, Ruben A. Smith, Pierre Buekens, José M. Belizán, Fernando Althabe. 2015. Pregnant Women's Secondhand Smoke Exposure and Receipt of Screening and Brief Advice by Prenatal Care Providers in Argentina and Uruguay. *Maternal and Child Health Journal* **19**:6, 1376-1383. [[Crossref](#)]
29. W. Xiong, M.R. Phillips, Y.Q. Xu, X.H. Wang, H. Li. 2015. Closing the knowledge gap in secondhand smoke exposure among children: employment of a five-minute household survey in China. *Public Health* **129**:6, 716-724. [[Crossref](#)]
30. Van T Tong, Patricia M Dietz, Italia V Rolle, Sara M Kennedy, William Thomas, Lucinda J England. 2015. Clinical interventions to reduce secondhand smoke exposure among pregnant women: a systematic review. *Tobacco Control* **24**:3, 217-223. [[Crossref](#)]
31. Michelle C. Kegler, Lucja Bundy, Regine Haardörfer, Cam Escoffery, Carla Berg, Debbie Yembra, Matthew Kreuter, Mel Hovell, Rebecca Williams, Patricia Dolan Mullen, Kurt Ribisl, Donna Burnham. 2015. A Minimal Intervention to Promote Smoke-Free Homes Among 2-1-1 Callers: A Randomized Controlled Trial. *American Journal of Public Health* **105**:3, 530-537. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
32. Bo Zhang, Farzana Haji, Pamela Kaufman, Sarah Muir, Roberta Ferrence. 2015. 'Enter at your own risk': a multimethod study of air quality and biological measures in Canadian waterpipe cafes. *Tobacco Control* **24**:2, 175-181. [[Crossref](#)]
33. Stephen B Gordon, Nigel G Bruce, Jonathan Grigg, Patricia L Hibberd, Om P Kurmi, Kin-bong Hubert Lam, Kevin Mortimer, Kwaku Poku Asante, Kalpana Balakrishnan, John Balmes, Naor Bar-Zeev, Michael N Bates, Patrick N Breyse, Sonia Buist, Zhengming Chen, Deborah Havens, Darby Jack, Surinder Jindal, Haidong Kan, Sumi Mehta, Peter Moschovis, Luke Naeher, Archana Patel, Rogelio Perez-Padilla, Daniel Pope, Jamie Rylance, Sean Semple, William J Martin. 2014. Respiratory risks from household air pollution in low and middle income countries. *The Lancet Respiratory Medicine* **2**:10, 823-860. [[Crossref](#)]
34. Rishi Caleyachetty, Christopher A Tait, Andre P Kengne, Camila Corvalan, Ricardo Uauy, Justin B Echouffo-Tcheugui. 2014. Tobacco use in pregnant women: analysis of data from Demographic and Health Surveys from 54 low-income and middle-income countries. *The Lancet Global Health* **2**:9, e513-e520. [[Crossref](#)]

35. Isaiah "Shaneequa" Brokenleg, Teresa K. Barber, Nancy L. Bennett, Simone Peart Boyce, Valarie Blue Bird Jernigan. 2014. Gambling with Our Health. *American Journal of Preventive Medicine* 47:3, 290-299. [[Crossref](#)]
36. Sherry Zhou, David G. Rosenthal, Scott Sherman, Judith Zelikoff, Terry Gordon, Michael Weitzman. 2014. Physical, Behavioral, and Cognitive Effects of Prenatal Tobacco and Postnatal Secondhand Smoke Exposure. *Current Problems in Pediatric and Adolescent Health Care* 44:8, 219-241. [[Crossref](#)]
37. Sophia Siu Chee Chan, Yee Tak Derek Cheung, Doris Yin Ping Leung, Yim Wah Mak, Gabriel M. Leung, Tai Hing Lam. 2014. Secondhand Smoke Exposure and Maternal Action to Protect Children from Secondhand Smoke: Pre- and Post-Smokefree Legislation in Hong Kong. *PLoS ONE* 9:8, e105781. [[Crossref](#)]
38. Michael Hennessy, Amy Bleakley, Giridhar Mallya, Daniel Romer. 2014. The Effect of Household Smoking Bans on Household Smoking. *American Journal of Public Health* 104:4, 721-727. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
39. Carrie L. Nieman, Ana Navas-Acien, Sandra Y. Lin, Douglas D. Reh. 2014. Physician screening and recommendations on secondhand smoke in chronic rhinosinusitis patients. *International Forum of Allergy & Rhinology* 4:2, 117-123. [[Crossref](#)]
40. J. Mackay, J.M. Samet. Tobacco, Smoking, and Nicotine . [[Crossref](#)]
41. M. Hennessy, A. Bleakley, G. Mallya, D. Romer. 2014. Beliefs Associated With Intention to Ban Smoking in Households With Smokers. *Nicotine & Tobacco Research* 16:1, 69-77. [[Crossref](#)]
42. Dorota Jarosińska, Kinga Polańska, Bogdan Wojtyniak, Wojciech Hanke. 2014. Towards estimating the burden of disease attributable to second-hand smoke exposure in Polish children. *International Journal of Occupational Medicine and Environmental Health* 27:1. . [[Crossref](#)]
43. Sasha G Hutchinson, Ilse Mesters, Gerard van Breukelen, Jean WM Muris, Frans JM Feron, S Katharine Hammond, Constant P van Schayck, Edward Dompeling. 2013. A motivational interviewing intervention to PREvent PASSive Smoke Exposure (PREPASE) in children with a high risk of asthma: design of a randomised controlled trial. *BMC Public Health* 13:1. . [[Crossref](#)]
44. Hayfaa A Wahabi, Rasmieh A Alzeidan, Amel A Fayed, Ahmed Mandil, Ghadeer Al-Shaikh, Samia A Esmaeil. 2013. Effects of secondhand smoke on the birth weight of term infants and the demographic profile of Saudi exposed women. *BMC Public Health* 13:1. . [[Crossref](#)]
45. Miranda R Jones, Heather Wipfli, Shahida Shahrir, Erika Avila-Tang, Jonathan M Samet, Patrick N Breyse, Ana Navas-Acien. 2013. Secondhand tobacco smoke: an occupational hazard for smoking and non-smoking bar and nightclub employees. *Tobacco Control* 22:5, 308-314. [[Crossref](#)]
46. Neil E. Klepeis, Suzanne C. Hughes, Rufus D. Edwards, Tracy Allen, Michael Johnson, Zohir Chowdhury, Kirk R. Smith, Marie Boman-Davis, John Bellettiere, Melbourne F. Hovell. 2013. Promoting Smoke-Free Homes: A Novel Behavioral Intervention Using Real-Time Audio-Visual Feedback on Airborne Particle Levels. *PLoS ONE* 8:8, e73251. [[Crossref](#)]
47. Shanta Pandey, Yuan Lin. 2013. Tobacco Use Among Married Women in Nepal: The Role of Women's Empowerment. *Maternal and Child Health Journal* 17:3, 530-538. [[Crossref](#)]
48. Jere Reijula, Tom Johnsson, Simo Kaleva, Tapani Tuomi, Kari Reijula. 2013. Total prohibition of smoking but not partial restriction effectively reduced exposure to tobacco smoke among restaurant workers in Finland. *International Journal of Occupational Medicine and Environmental Health* 26:5. . [[Crossref](#)]
49. Akram Abusalah, Magda Gavana, Anna-Bettina Haidich, Emmanouil Smyrnakis, Nikos Papadakis, Alexis Papanikolaou, Alexis Benos. 2012. Low Birth Weight and Prenatal Exposure to Indoor Pollution from Tobacco Smoke and Wood Fuel Smoke: A Matched Case-Control Study in Gaza Strip. *Maternal and Child Health Journal* 16:8, 1718-1727. [[Crossref](#)]
50. C. M. Knottenbelt, S. Bawazeer, J. Hammond, D. Mellor, D. G. Watson. 2012. Nicotine hair concentrations in dogs exposed to environmental tobacco smoke: a pilot study. *Journal of Small Animal Practice* 53:11, 623-626. [[Crossref](#)]
51. Veronica Schoj, Lorena Allemandi, Oscar Ianovsky, Manuel Lago, Mariela Alderete. 2012. Smoke-free environments: age, sex, and educational disparity in 25 Argentinean cities. *Cancer Causes & Control* 23:10, 1607-1614. [[Crossref](#)]
52. Laura Jones, Andy McEwen. 2012. Reducing secondhand smoke exposure at home. *British Journal of School Nursing* 7:8, 389-393. [[Crossref](#)]
53. Wendy Nicholson. 2012. Calling for school nurses' views on the new nursing vision. *British Journal of School Nursing* 7:8, 393-393. [[Crossref](#)]
54. Judith A. Groner, Hong Huang, Lisa Nicholson, Jennifer Kuck, Bethany Boettner, John A. Bauer. 2012. Secondhand Smoke Exposure and Hair Nicotine in Children: Age-Dependent Differences. *Nicotine & Tobacco Research* 14:9, 1105-1109. [[Crossref](#)]
55. B. J. Apelberg, L. M. Hepp, E. Avila-Tang, S. Kim, C. Madsen, J. Ma, J. M. Samet, P. N. Breyse. 2012. Racial Differences in Hair Nicotine Concentrations Among Smokers. *Nicotine & Tobacco Research* 14:8, 933-941. [[Crossref](#)]

56. Bradley N. Collins, Jennifer Ibrahim. 2012. Pediatric Secondhand Smoke Exposure: Systematic Multilevel Strategies to Improve Health. *Global Heart* 7:2, 161-165. [[Crossref](#)]
57. Andria Apostolou, Esther Garcia-Esquinas, Jeffrey J. Fadrowski, Pat McLain, RN, Virginia M. Weaver, Ana Navas-Acien. 2012. Secondhand Tobacco Smoke: A Source of Lead Exposure in US Children and Adolescents. *American Journal of Public Health* 102:4, 714-722. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
58. Tingting Yao, Hai-Yen Sung, Zhengzhong Mao, Teh-wei Hu, Wendy Max. 2012. Secondhand smoke exposure at home in rural China. *Cancer Causes & Control* 23:S1, 109-115. [[Crossref](#)]
59. Natalie Hemsing, Lorraine Greaves, Nancy Poole, Joan Bottorff. 2012. Reshuffling and Relocating: The Gendered and Income-Related Differential Effects of Restricting Smoking Locations. *Journal of Environmental and Public Health* 2012, 1-12. [[Crossref](#)]
60. E. Torres, Z. Quinones de Monegro, L. French, D. P. Swanson, J. Guido, D. J. Ossip. 2011. Tobacco Use and Exposure to Secondhand Smoke Among Pregnant Women in the Dominican Republic: An Exploratory Look into Attitudes, Beliefs, Perceptions, and Practices. *Nicotine & Tobacco Research* 13:12, 1220-1227. [[Crossref](#)]
61. Asma B. Jafri. 2011. Aging and Toxins. *Clinics in Geriatric Medicine* 27:4, 609-628. [[Crossref](#)]
62. Huda Gharaibeh, Linda Haddad, Sukaina Alzyoud, Omar El-Shahawy, Nesrin Abu Baker, Mary Umlauf. 2011. Knowledge, Attitudes, and Behavior in Avoiding Secondhand Smoke Exposure Among Non-Smoking Employed Women with Higher Education in Jordan. *International Journal of Environmental Research and Public Health* 8:11, 4207-4219. [[Crossref](#)]
63. Niki Katsiki, Apostolos I. Hatzitolios, Dimitri P. Mikhailidis. 2011. Passive Smoking: The Democratic Right of Nonsmokers to Survive. *Angiology* 62:7, 520-522. [[Crossref](#)]
64. L. L. Jones, O. Atkinson, J. Longman, T. Coleman, A. McNeill, S. A. Lewis. 2011. The Motivators and Barriers to a Smoke-Free Home Among Disadvantaged Caregivers: Identifying the Positive Levers for Change. *Nicotine & Tobacco Research* 13:6, 479-486. [[Crossref](#)]
65. Ding Ding, Dennis R. Wahlgren, Sandy Liles, Georg E. Matt, McKenzie Oliver, Jennifer A. Jones, Melbourne F. Hovell. 2011. A Second Reporter Matters. *American Journal of Preventive Medicine* 40:5, 572-575. [[Crossref](#)]
66. W. K. Al-Delaimy, W. C. Willett. 2011. Toenail Nicotine Level as a Novel Biomarker for Lung Cancer Risk. *American Journal of Epidemiology* 173:7, 822-828. [[Crossref](#)]
67. David G. Rosenthal, Michael Weitzman. 2011. Examining the Effects of Intrauterine and Postnatal Exposure to Tobacco Smoke on Childhood Cognitive and Behavioral Development. *International Journal of Mental Health* 40:1, 39-64. [[Crossref](#)]
68. Christina N. Lessov-Schlaggar, Dennis R. Wahlgren, Sandy Liles, Jennifer A. Jones, Ming Ji, Suzanne C. Hughes, Gary E. Swan, Melbourne F. Hovell. 2011. Sensitivity to Secondhand Smoke Exposure Predicts Smoking Susceptibility in 8-13-Year-Old Never Smokers. *Journal of Adolescent Health* 48:3, 234-240. [[Crossref](#)]
69. Sandra Y. Lin, Douglas D. Reh, Sandra Clipp, Laili Irani, Ana Navas-Acien. 2011. Allergic Rhinitis and Secondhand Tobacco Smoke: A Population-based Study. *American Journal of Rhinology & Allergy* 25:2, e66-e71. [[Crossref](#)]
70. Seth Kligerman, Charles White. 2011. Epidemiology of Lung Cancer in Women: Risk Factors, Survival, and Screening. *American Journal of Roentgenology* 196:2, 287-295. [[Crossref](#)]
71. Mattias Öberg, Maritta S Jaakkola, Alistair Woodward, Armando Peruga, Annette Prüss-Ustün. 2011. Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *The Lancet* 377:9760, 139-146. [[Crossref](#)]
72. Heather L. Wipfli, Jonathan M. Samet. 2011. Second-hand smoke's worldwide disease toll. *The Lancet* 377:9760, 101-102. [[Crossref](#)]
73. S. Kim, J. Sohn, K. Lee. 2010. Exposure to Particulate Matters (PM2.5) and Airborne Nicotine in Computer Game Rooms After Implementation of Smoke-Free Legislation in South Korea. *Nicotine & Tobacco Research* 12:12, 1246-1253. [[Crossref](#)]
74. Amy E Kalkbrenner, Richard W Hornung, John T Bernert, S Katherine Hammond, Joe M Braun, Bruce P Lanphear. 2010. Determinants of serum cotinine and hair cotinine as biomarkers of childhood secondhand smoke exposure. *Journal of Exposure Science & Environmental Epidemiology* 20:7, 615-624. [[Crossref](#)]
75. Douglas D Reh, Ana Navas-Acien. 2010. Relationship between second-hand tobacco smoke exposure and chronic rhinosinusitis: evidence for causality. *Expert Review of Respiratory Medicine* 4:4, 445-449. [[Crossref](#)]
76. Zubair Kabir, Hillel R Alpert, Patrick G Goodman, Sally Haw, Ilan Behm, Gregory N Connolly, Prakash C Gupta, Luke Clancy. 2010. Effect of smoke-free home and workplace policies on second-hand smoke exposure levels in children: an evidence summary. *Pediatric Health* 4:4, 391-403. [[Crossref](#)]
77. N.M. Shara. 2010. Cardiovascular disease in Middle Eastern women. *Nutrition, Metabolism and Cardiovascular Diseases* 20:6, 412-418. [[Crossref](#)]

78. Harry A. Lando, Bethany J. Hipple, Myra Muramoto, Jonathan D. Klein, Alexander V. Prokhorov, Deborah J. Ossip, Jonathan P. Winickoff. 2010. Tobacco Control and Children: An International Perspective. *Pediatric Allergy, Immunology, and Pulmonology* 23:2, 99-103. [[Crossref](#)]
79. Denis Vinnikov, Nurlan Brimkulov, Shahida Shahrir, Patrick Breyse, Ana Navas-Acien. 2010. Excessive Exposure to Secondhand Tobacco Smoke among Hospitality Workers in Kyrgyzstan. *International Journal of Environmental Research and Public Health* 7:3, 966-974. [[Crossref](#)]
80. Heura Llaquet, Simona Pichini, Xavier Joya, Esther Papaseit, Oriol Vall, Julia Klein, Oscar Garcia-Algar. 2010. Biological matrices for the evaluation of exposure to environmental tobacco smoke during prenatal life and childhood. *Analytical and Bioanalytical Chemistry* 396:1, 379-399. [[Crossref](#)]
81. Moon-Woo Seong, Jin Soo Moon, Jong Hee Hwang, Hye-Jung Ryu, Soo Jin Kang, Sun-Young Kong, Tae Hyun Um, Jae-Gahb Park, Jin Soo Lee, Do-Hoon Lee. 2010. Preschool children and their mothers are more exposed to paternal smoking at home than school children and their mothers. *Clinica Chimica Acta* 411:1-2, 72-76. [[Crossref](#)]
82. Laili Irani, Sandra Y. Lin, Sandra L. Clipp, Anthony J. Alberg, Ana Navas-Acien. 2010. Involving Stakeholders to Optimize a Study Protocol on Secondhand Tobacco Smoke and Chronic Rhinosinusitis in Adults. *American Journal of Rhinology & Allergy* 24:1, 39-44. [[Crossref](#)]
83. H Wipfli, J M Samet. 2009. Global Economic and Health Benefits of Tobacco Control: Part 1. *Clinical Pharmacology & Therapeutics* 86:3, 263-271. [[Crossref](#)]
84. Sung Roul Kim, Heather Wipfli, Erika Avila-Tang, Jonathan M. Samet, Patrick N. Breyse. 2009. Method validation for measurement of hair nicotine level in nonsmokers. *Biomedical Chromatography* 23:3, 273-279. [[Crossref](#)]
85. Sungroul Kim, Ther Aung, Emily Berkeley, Gregory B. Diette, Patrick N. Breyse. 2008. Measurement of nicotine in household dust. *Environmental Research* 108:3, 289-293. [[Crossref](#)]
86. Naomi Priest, Rob Roseby, Elizabeth Waters, Adam Polnay, Rona Campbell, Nick Spencer, Premila Webster, Grace Ferguson-Thorne. Family and carer smoking control programmes for reducing children's exposure to environmental tobacco smoke . [[Crossref](#)]
87. John P Pierce, María E León. 2008. Effectiveness of smoke-free policies. *The Lancet Oncology* 9:7, 614-615. [[Crossref](#)]