

## RESEARCH ARTICLE

# Maternal occupational exposure to selected organic and chlorinated solvents and delivery of small-for-gestational age or preterm infants

Kristen W. Van Buren DrPH, MPH<sup>1,2</sup>  | Carissa M. Rocheleau PhD<sup>1</sup> |  
I-Chen Chen PhD<sup>1</sup>  | Tania A. Desrosiers PhD<sup>3</sup> | Wayne T. Sanderson PhD, CIH<sup>4</sup> |  
Maria D. Politis DrPH, MPH<sup>5</sup> | Elizabeth C. Ailes PhD<sup>6</sup> | The National Birth Defects  
Prevention Study

<sup>1</sup>Division of Field Studies and Engineering,  
National Institute for Occupational Safety and  
Health, Cincinnati, Ohio, USA

<sup>2</sup>Department of Epidemiology and  
Environmental Health, College of Public  
Health, University of Kentucky, Lexington,  
Kentucky, USA

<sup>3</sup>Department of Epidemiology, Gillings School  
of Global Public Health, University of North  
Carolina, Chapel Hill, North Carolina, USA

<sup>4</sup>Department of Biosystems and Agricultural  
Engineering, College of Agriculture, Food and  
Environment, University of Kentucky,  
Lexington, Kentucky, USA

<sup>5</sup>Arkansas Center for Birth Defects Research  
and Prevention, University of Arkansas for  
Medical Sciences, Little Rock, Arkansas, USA

<sup>6</sup>Division of Birth Defects and Infant  
Disorders, National Center on Birth Defects  
and Developmental Disabilities, Atlanta,  
Georgia, USA

## Correspondence

Kristen W. Van Buren, DrPH, MPH, Division of  
Field Studies and Engineering, National  
Institute for Occupational Safety and Health,  
Cincinnati, OH, USA.  
Email: [opv7@cdc.gov](mailto:opv7@cdc.gov)

## Funding information

Centers for Disease Control and Prevention  
(CDC), Grant/Award Numbers: PA #96043,  
PA #02081, FOA #DD09-001, FOA #DD13-  
003, NOFO #DD18-001; National Institute  
for Occupational Safety and Health,  
Grant/Award Number: 200-2000-08018

## Abstract

**Background:** Potential reproductive effects of organic solvent exposure during pregnancy remain unclear. We investigated the association between maternal occupational exposure during pregnancy to six chlorinated solvents, three aromatic solvents, and Stoddard solvent, and delivery of preterm infants or those born small-for-gestational age (SGA).

**Methods:** In this case-control study of SGA and preterm birth (PTB) nested within the National Birth Defects Prevention Study (NBDPS) from 1997 to 2011, we analyzed data from 7504 singleton live births without major birth defects and their mothers. Self-reported information on jobs held in the periconceptional period was assessed for solvent exposure. Unconditional logistic regression was used to estimate the association between maternal occupational exposure (any, none) during early pregnancy to organic solvents and PTB and SGA. Linear regression was used to examine changes in mean birthweight potentially associated with maternal occupational solvent exposure.

**Results:** Maternal occupational exposure to any organic solvents overall was not associated with an increased odds of PTB (adjusted odds ratio [aOR] = 0.94; 95% confidence interval [CI] 0.67–1.33) or SGA (aOR = 0.93; 95% CI 0.65–1.34). Point estimates increased modestly for higher estimated exposure versus lower, but confidence intervals were wide and not statistically significant. Maternal exposure to solvents was not associated with a statistically significant change in term birthweight among infants.

**Conclusions:** Occupational exposure to organic solvents at the frequency and intensity levels found in a population-based sample of pregnant workers was not

**Abbreviations:** CATI, computer-assisted telephone interview; DAG, directed acyclic graph; IH, industrial hygienist; NAICS, North American Industry Classification System; NBDPS, National Birth Defects Prevention Study; PTB, preterm birth; SGA, small-for-gestational age; SOC, Standard Occupational Classification.

Maternal occupational solvent exposure and select birth outcomes

associated with PTB or SGA; however, we cannot rule out any effects among pregnant workers with uncommonly high exposure to organic solvents.

#### KEYWORDS

birth weight, occupation, organic solvents, pregnancy, preterm birth, reproductive health, SGA, worker

## 1 | INTRODUCTION

Organic solvents are carbon-based, lipophilic compounds commonly used in a wide range of occupations and industries. Each year, millions of workers are potentially exposed to organic solvents or solvent mixtures through contact with paints, varnishes, lacquers, adhesives, glues, degreasing, and cleaning agents.<sup>1</sup> The potential health hazards associated with occupational organic solvent exposure vary widely and are influenced heavily by dose, duration, and bioavailability of the chemical compound. Evidence from animal models—as well as occupational epidemiologic studies—have demonstrated that some, but not all, organic solvents are carcinogens, neurotoxins, or reproductive toxicants.<sup>2,3</sup> Although many acute and chronic adverse health effects associated with occupational exposure to organic solvents have been previously described in working adults,<sup>4</sup> the potential for negative effects on fetal development from organic solvent exposure among pregnant workers remains unclear.

Fat-soluble organic solvents can cross the placenta during pregnancy, potentially causing oxidative damage to critical cell structures and disrupting normal fetal development.<sup>5,6</sup> Animal models have for decades established the teratogenic effects of select organic solvents, such as toluene.<sup>2,7,8</sup> Findings from early epidemiological studies among potentially high-exposure populations (e.g., laboratory workers, dry cleaner workers, etc.),<sup>9</sup> as well as several population-based studies,<sup>10–13</sup> have been suggestive of an association between organic solvent exposure and adverse birth outcomes, including small-for-gestational age (SGA) and preterm birth (PTB). However, small sample size has contributed to imprecise (i.e., low statistical power) findings in some studies,<sup>14,15</sup> while others have reported negative or null associations.<sup>16–18</sup> Challenges, including small sample size, limited job history information, and the inability to discern individual solvents from solvent mixtures or other co-exposures in the workplace, have all contributed to the difficulty in accurately quantifying the effects of organic solvent exposure on pregnant workers. Despite these challenges, the identification of potential risk factors, specifically reproductive toxicants, is necessary to mitigate and prevent adverse health outcomes for working mothers and their infants.

In this study, we focus on PTB and SGA, which are two relatively prevalent pregnancy outcomes associated with increased infant mortality and morbidity.<sup>19,20</sup> Preterm birth, defined as birth before 37 weeks' gestation, affects approximately one out of every ten infants born in the United States, while fetuses characterized as SGA—referring to a newborn with a birth weight less than the 10th percentile for gestational age by infant sex—affect approximately

11.1% of all live births in the United States.<sup>21,22</sup> Infants characterized by either condition are at increased risk of intellectual and neurodevelopmental delays in childhood, as well as select chronic diseases in adulthood.<sup>23–30</sup> We also conducted a sensitivity analysis examining the potential effects of maternal occupational exposure to organic solvents on mean change in infant birthweight at term (i.e.,  $\geq 37$  weeks' gestation), as birthweight is a critical component of several birth outcome measurements that can be indicative of intrauterine growth restriction (IUGR), including SGA.

To better understand the potential effects of organic solvent exposure in the workplace on pregnancy outcomes, Desrosiers and colleagues first conducted an analysis using National Birth Defects Prevention Study (NBDPS) data from 1997 to 2002 that evaluated the association between expert-assessed occupational solvent exposure and risk of SGA among infants in a population-based sample of women; elevated but imprecise associations were observed between SGA and exposure to any solvent(s), chlorinated solvents, and aromatic solvents among women with  $\geq 50\%$  probability of exposure.<sup>10</sup> Since these analyses, an additional 9 years of data (2003–2011) were collected and PTB was included as an additional pregnancy outcome of interest. The current study expands the work conducted by Desrosiers and colleagues (2015) by utilizing the full NBDPS analytic data set (1997–2011) to examine associations between maternal occupational exposure during pregnancy to six chlorinated solvents (carbon tetrachloride, chloroform, methylene chloride, perchloroethylene, trichloroethylene, and 1,1,1-trichloroethane), three aromatic solvents (benzene, toluene, and xylene), and a petroleum mixture referred to as Stoddard solvent with delivery of non-malformed liveborn infants characterized as preterm or SGA in a population-based sample of women.

## 2 | MATERIALS AND METHODS

### 2.1 | Study population

The National Birth Defects Prevention Study (NBDPS) was a large, population-based case-control study designed primarily to investigate risk factors for major structural birth defects.<sup>31</sup> The NBDPS was conducted in 10 US states: Arkansas, California, Georgia, Iowa, Massachusetts, New Jersey, New York, North Carolina, Texas, and Utah, although not all study sites contributed for all the study years.

Randomly selected live-born infants without a chromosomal or major structural birth defect identified from birth records and

hospital delivery records were recruited as controls from the same population areas the birth defects registries covered. Because these controls have been demonstrated to be generally representative of the population of non-malformed live births in the geographic regions covered by the NBDPS,<sup>32</sup> they have been used as essentially a birth cohort in other studies of pregnancy-related factors and outcomes.<sup>33–35</sup> Our study is nested within the NBDPS, and our study population consists of mothers of liveborn infants without a birth defect(s) with estimated dates of delivery between October 1, 1997 and December 31, 2011.

Specifically, data were available on 11,829 mothers of infants without birth defects who participated in the computer-assisted telephone interview (CATI) in either English or Spanish (65% of eligible mothers of infants without birth defects participated in the interview). Our eligible study population included mothers who reported employment for at least one calendar month during the periconceptional window of exposure (i.e., one month before conception through the first trimester) (68.9% of those mothers interviewed) and provided enough job history information to assess occupational exposure to organic solvents (99.8% of mothers who reported employment) (Figure 1). Because pregestational diabetes and multiple pregnancies are strong risk factors for both birth outcomes of interest, NBDPS participant mothers with Type 1 or Type 2 diabetes before the index pregnancy ( $N = 52$ ), as well as non-singleton pregnancies ( $N = 260$ ), were excluded from analyses because there were too few exposed pregnancies among these mothers to allow for stratification. In addition, infants with ambiguous or unreported sex at birth ( $N = 4$ ) and those infants born outside the range of 22–44 weeks' gestation ( $N = 3$ ) were also excluded (Figure 1). The final study population consisted of 7504 eligible mother–infant pairs. The NBDPS and this analysis were approved by the Institutional Review Boards of the Centers for Disease Control and Prevention and all participating study centers, and all participating mothers provided informed consent.

## 2.2 | Exposure assessment

Mothers self-reported occupational information during the NBDPS interview, including company name, participant's job title, what the company produced, the participant's main job activities and duties, and chemicals or equipment used. All responses were recorded as open free text. All jobs that were performed for at least 1 month during the periconceptional window were recorded. Jobs were then coded by occupation and industry using the Standard Occupational Classification (SOC) System and the North American Industry Classification System (NAICS).

Maternal occupational exposure assessment to chlorinated, aromatic, and Stoddard solvent(s) took place in two batches over the 15 years of the study's recruitment (first assessment period from 1997 to 2002 and second assessment period from 2003 to 2011); both used the same definitions, source materials, and methods—though staff performing exposure assessments changed between the

assessments. Two industrial hygienists (IH), blinded to case status, reviewed the full job descriptions to assign probability of direct and indirect exposure, intensity of direct and indirect exposure, frequency of direct and indirect exposure, and a subjective rating of the reviewer's confidence in the score based on the strength of data from similar occupations. Inter-rater reliability and validity of the exposure assessment strategy has been previously described.<sup>36</sup>

As some women had multiple jobs during the relevant window during pregnancy, job exposure measures were summarized to assign exposure estimates of probability, intensity, frequency, and duration of exposure for each organic solvent of interest for each participant. Cumulative occupational exposure, as parts per million (ppm)-hours for chlorinated and aromatic solvent classes or milligram per cubic meter ( $\text{mg}/\text{m}^3$ )-hours for Stoddard solvent, was estimated for each job using a calculation that incorporated weighted intensity, frequency, self-reported work frequency, and number of days worked in the exposure window. For participant mothers reporting more than one job during the periconceptional window of exposure, the estimate of total organic solvent exposure was the sum of the job-specific cumulative exposures in that period. For exposure-response analysis, estimated organic solvent exposure categories included no organic solvent exposure, below (low), and at or above (high), the median of the cumulative exposure of organic solvents (any and by class) among liveborn infants without SGA or PTB. For this analysis, we considered a pregnant worker to be exposed to a particular solvent if any of her self-reported job(s) held during the periconceptional period were rated by the IH as 'exposed' (i.e., probability of exposure  $>0$ ); she was considered unexposed to a particular solvent if her self-reported job(s) during this time were rated as 'unexposed' (i.e., probability of exposure = 0).

## 2.3 | Outcome classifications

All variables necessary to determine both primary outcome classifications, including due date, infant birth weight (grams), gestational age at delivery, and infant sex were self-reported during the CATI. Fetal growth restriction was identified based on SGA, defined as birth weight below the 10th percentile for a given gestational age at delivery and infant sex. Infants born SGA were identified using a US-based birth weight for gestational age reference reported by Talge and colleagues.<sup>37</sup> Sex-specific, 10th percentile birth weight values determined SGA and non-SGA infants within the study population from 22 through 44 weeks' gestation. Preterm birth (PTB) was classified as birth occurring before 37 completed weeks of gestation, following the National Center for Health Statistics (NCHS) standards.<sup>21</sup> The increased sample size of the current study was utilized to replicate a sensitivity analysis first conducted by Desrosiers and colleagues (2015) examining changes in mean birthweight (grams) among infants potentially associated with maternal occupational solvent exposure during early pregnancy. Infant birthweight (grams) was analyzed as a continuous outcome and analyses were conducted among 'term' infants (i.e., birth occurring  $\geq 37$  weeks' gestation) to

identify meaningful inferences from the potential influence of maternal occupational solvent exposure on mean change in infant birthweight. We restricted to term infants to reduce the potential for skewed data resulting from infants who weigh less at delivery because they are premature.

## 2.4 | Covariates

Potential confounders were identified through a directed acyclic graph (DAG)<sup>38</sup> analysis (Figure S1)<sup>38,39</sup> based on previous literature of factors known to influence maternal occupational exposure to organic solvents, SGA, and PTB, which included: maternal residence at delivery (NBDPS study location), maternal age at delivery ( $\leq 20$  years, 21–25 years, 26–34 years, and  $\geq 35$  years), maternal race/ethnicity (non-Hispanic White, non-Hispanic Black, Hispanic, Other non-Hispanic), maternal nativity (US-born, foreign-born), maternal education ( $\leq 12$  years,  $>12$  years), maternal pre-pregnancy body mass index (BMI) (underweight range,  $<18.5$  kg/m<sup>2</sup>; healthy weight range, 18.5 to  $<24$  kg/m<sup>2</sup>; overweight range, 25 to  $<30$  kg/m<sup>2</sup>; or obesity range,  $>30$  kg/m<sup>2</sup>), maternal smoking status (any use during periconceptional period, none), maternal alcohol use (any use during periconceptional period, none), and maternal secondhand smoke exposure (any exposure from home or work during periconceptional period, none). The final models were adjusted for maternal education (referent  $\geq 12$  years) and maternal residence at delivery, which were the minimally sufficient set of confounders resulting from the DAG (Figure S1).

## 2.5 | Statistical analysis

Select demographic and behavioral factors of mother/infant pairs were described. Frequencies and percentages were calculated, as well as chi-square tests (excluding missing values) to assess differences between mothers of SGA, non-SGA, PTB, and non-PTB infants. The prevalence of maternal occupational exposure (any vs. no exposure) during the periconceptional window was estimated for any solvent exposure and solvent classes (aromatic, chlorinated, Stoddard), stratified by SGA and PTB. To describe occupational exposure to solvents among our study population, the prevalence of exposure (any vs. no exposure) by job was estimated and presented by SOC major group.


Unconditional logistic regression was used to estimate unadjusted crude (cOR) and adjusted odds ratios (aOR), and 95% confidence intervals (CI) for the associations between periconceptional maternal exposure to organic solvents (any and by class), and SGA and PTB. To assess cumulative occupational exposure to solvents, exposure-response analyses were conducted in three categorical groups (none, low, and high) determined by median cumulative exposure levels among mothers of infants not born prematurely or not characterized as SGA. Linear regression was used to estimate the mean difference in birthweight (grams) among term

infants of exposed mothers compared to term infants of unexposed mothers. We also conducted a sensitivity analysis to account for potential exposure misclassification by restricting the study sample to include only women with an exposure probability of at least 50%, as determined by the expert IHS during exposure assessment. Because this was a continuation of prior analyses by Desrosiers et al. (2015), we also stratified the main analysis by both exposure assessment periods (i.e., 1997–2002 vs. 2003–2011) to identify any notable differences. In all models, women considered “unexposed” are those participant mothers with no assigned exposure to any organic solvent during the periconceptional window exclusively. All statistical analyses were performed using SAS 9.4 (SAS Institute, Inc.) and independently replicated by a second analyst.

## 3 | RESULTS

After eligibility and exclusion criteria were met, a total of 7504 infant/mother pairs were analyzed (Figure 1). Approximately 6.9% (517/7504) of study infants were classified as SGA; 7.7% (577/7504) were born premature. Table 1 displays frequency distributions and chi-square tests of select mother/infant pair demographic and behavioral characteristics described by birth outcome of interest. Overall, study participant mothers were most commonly non-Hispanic White, did not smoke, had greater than a High School education, and were aged between 26 and 34 years of age at delivery. Self-reported jobs that were most frequently rated as incurring some exposure to organic solvents (vs. no exposure) were in the SOC major groups of production ( $n = 134$  exposed jobs; 23.8%), personal care and service ( $n = 83$  exposed jobs; 14.8%), healthcare practitioners and technical ( $n = 80$  exposed jobs; 14.2%) and building and grounds cleaning and maintenance ( $n = 58$  exposed jobs; 10.3%) (Table S1).

Maternal occupational organic solvent exposure among our study population is described in Table 2. Among all study participant mothers, approximately 5.8% were determined to have exposure to chlorinated solvents, while only 1.9% were exposed to aromatic solvents and 1.8% to Stoddard solvent. Exposure variability was observed within the aromatic and chlorinated solvent classes. For example, among mothers occupationally exposed to chlorinated solvents, approximately 4.7% were exposed to 1,1,1-trichloroethane compared with less than 0.5% to carbon tetrachloride. Co-exposures were also common. Over one-third (33%;  $n = 166$ ) of participant mothers were determined to have exposure to more than one solvent class, while over 80% of participant mothers were determined to have co-exposures within solvent classes (93.7% within the aromatic solvents; 81.5% within the chlorinated solvents) (data not shown). The distribution of occupational exposure to any organic solvent, as well as solvent class, did not differ meaningfully across outcomes of interest; however, maternal occupational exposure to both aromatic and Stoddard solvent was higher among mothers of SGA and PTB infants, compared to mothers of non-SGA/non-PTB infants (2.3%–2.5% vs. 1.8%–1.9% for SGA; 2.3%–2.4% vs. 1.8%–1.9% for



**FIGURE 1** Study population eligibility and exclusion criteria, National Birth Defects Prevention Study, USA, 1997–2011.

PTB, respectively). Exposure to chlorinated solvents was slightly lower among SGA infants compared to non-SGA infants (5.4% vs. 5.8%, respectively), while exposure to chlorinated solvents was similar between both preterm and non-preterm infants (5.9% vs. 5.8%, respectively).

Results from the exposure-response analyses are presented in Table 3. After adjusting for maternal residence at delivery and educational attainment, point estimates were elevated for the high exposure groups compared to the low exposure group for both SGA and PTB (aOR for “low” exposure to any solvent = 0.90 95% CI 0.57–1.40; aOR for “high” exposure to any solvent = 1.52 95% CI 0.80–2.87; aOR for “low” exposure to any solvent = 0.97 95% CI 0.64–1.47; aOR for “high” exposure to any solvent = 1.20 95% CI 0.62–2.33); however, confidence intervals were wide and we cannot rule out that this finding was spurious. A similarly suggestive—but not conclusive—dose-response pattern was observed between maternal occupational exposure to chlorinated solvents and PTB (aOR for “low” exposure to chlorinated solvents = 1.07 95% CI 0.69–1.66; aOR for “high” exposure to chlorinated solvents = 1.23 95% CI 0.63–2.39). Crude analysis demonstrated an association between “high” maternal occupational exposure to Stoddard solvent and SGA (cOR for “high” exposure to Stoddard solvent = 2.24 95% CI 1.10–4.56); however, the association was attenuated after adjustment for maternal education and residence (aOR for “high” exposure to Stoddard solvent = 1.85 95% CI 0.90–3.81).

Table 4 shows the crude and adjusted associations between estimated maternal occupational exposure to organic solvents and SGA and PTB. After adjusting for maternal education and residence at delivery, no associations were observed between maternal exposure to any organic solvent and SGA (aOR = 0.93; 95% CI 0.65–1.34) or PTB (aOR = 0.94; 95% CI 0.67–1.33). Similarly, no associations were observed by solvent class and SGA or PTB or when restricting to mothers where the probability of workplace exposure to organic solvents was highest ( $\geq 50\%$ ).

Maternal exposure to any solvent, as well as exposure by solvent class, was not associated with a difference in term birthweight (Table 5). After adjusting for maternal education and residence at delivery, the difference in mean birthweight at term between infants of mothers exposed to any organic solvent and infants of unexposed mothers was 47.3 g, but the 95% CI included the null value of 0 [95% CI –31.5 to 126.0].

Stratifying by exposure assessment periods demonstrated an elevated, but imprecise adjusted OR for any occupational maternal solvent exposure and SGA (aOR = 1.11; 95% CI 0.66–1.89) in the first exposure assessment period (1997–2002) (Table S2). Among women in the second exposure assessment period (2003–2011), we observed a possible inverse association between occupational maternal exposure to any solvent and SGA (aOR = 0.83; 95% CI 0.50–1.37) (Table S2).



**TABLE 1** Distribution of maternal and infant characteristics by classification of small-for-gestational age (SGA) and preterm birth (PTB) among infants without birth defects, National Birth Defects Prevention Study, USA, 1997–2011.

Maternal characteristics	SGA (n = 517)		non-SGA (n = 6987)		$\chi^2$ p-Value	PTB (n = 577)		non-PTB (n = 6927)		$\chi^2$ p-Value
	n	(%)	n	(%)		n	(%)	n	(%)	
Occupational exposure to any organic solvents during the periconceptual period					NS					NS
Exposed <sup>a</sup>	34	6.6	467	6.7		38	6.6	463	6.7	
Unexposed	483	93.4	6520	93.3		539	93.4	6,464	93.3	
Maternal residence at delivery					<0.01					<0.01
Arkansas	83	16.1	881	12.6		100	17.3	864	12.5	
California	35	6.8	642	9.2		45	7.8	632	9.1	
Georgia	74	14.3	763	10.9		71	12.3	766	11.1	
Iowa	54	10.4	936	13.4		86	14.9	904	13.1	
Massachusetts	61	11.8	941	13.5		70	12.1	932	13.5	
New Jersey	29	5.6	338	4.8		22	3.8	345	5.0	
New York	47	9.1	619	8.9		33	5.7	633	9.1	
North Carolina	41	7.9	606	8.7		47	8.2	600	8.7	
Texas	58	11.2	617	8.8		70	12.1	605	8.7	
Utah	35	6.8	644	9.2		33	5.7	646	9.3	
Maternal age at delivery					<0.01					NS
≤20 years	83	16.1	691	9.9		70	12.1	704	10.2	
21–25 years	154	29.8	1663	23.8		141	24.4	1676	24.2	
26–34 years	201	38.9	3615	51.7		270	46.8	3546	51.2	
≥35 years	79	15.3	1018	14.6		96	16.6	1001	14.5	
Maternal race/ethnicity					<0.01					<0.01
Non-Hispanic White	266	51.5	4529	64.8		332	57.5	4463	64.4	
Non-Hispanic Black	98	19.0	760	10.9		97	16.8	761	11.0	
Hispanic	109	21.1	1277	18.3		112	19.4	1274	18.4	
Other Non-Hispanic	44	8.5	421	6.0		36	6.2	429	6.2	
Maternal nativity					<0.05					NS
US-born	415	80.3	5885	84.2		491	85.1	5809	83.9	
Foreign-born	102	19.7	1094	15.7		86	14.9	1110	16.0	
Missing	0	0.0	8	0.1		0	0.0	8	0.1	
Maternal education					<0.01					<0.05
<12 years	69	13.4	642	9.2		67	11.6	644	9.3	
High school diploma or equivalent	146	28.2	1563	22.4		152	26.3	1557	22.5	
>12 years	302	58.4	4769	68.3		358	62.1	4713	68.0	
Missing	0	0.0	13	0.2		0	0.0	13	0.2	
Maternal BMI					<0.01					NS
Underweight range (BMI < 18.5)	39	7.5	301	4.3		32	5.6	308	4.5	
Healthy weight range (18.5 ≤ BMI < 25)	293	56.7	3691	52.8		289	50.1	3695	53.3	
Overweight range (25 ≤ BMI < 30)	95	18.4	1604	23.0		130	22.5	1569	22.7	
Obesity range (BMI ≥ 30)	75	14.5	1255	18.0		118	20.5	1212	17.5	
Missing	15	2.9	136	1.9		8	1.4	143	2.1	

(Continues)

TABLE 1 (Continued)

Maternal characteristics	SGA (n = 517)		non-SGA (n = 6987)		$\chi^2$ p-Value	PTB (n = 577)		non-PTB (n = 6927)		$\chi^2$ p-Value
	n	(%)	n	(%)		n	(%)	n	(%)	
Maternal smoking					<0.05					NS
Yes <sup>b</sup>	121	23.4	1271	18.2		118	20.5	1274	18.4	
No	396	76.6	5714	81.8		459	79.6	5651	81.6	
Missing	0	0.0	2	0.0		0	0.0	2	0.0	
Maternal alcohol use					NS					<0.01
Yes <sup>b</sup>	219	42.4	2981	42.7		212	36.7	2988	43.1	
No	295	57.1	3986	57.1		361	62.6	3920	56.6	
Missing	3	0.6	20	0.3		4	0.7	19	0.3	
Secondhand smoke exposure					<0.05					<0.05
Any <sup>b</sup>	154	29.8	1685	24.1		168	29.1	1671	24.1	
None	362	70.2	5284	75.6		407	70.5	5239	75.6	
Missing	1	0.2	18	0.3		2	0.3	17	0.2	
Infant characteristics										
Infant birthweight					<0.01					<0.01
Low (<2500 g)	146	28.2	212	3.0		242	41.9	116	1.7	
Normal (2500–4000 g)	363	70.2	5892	84.3		318	55.1	5937	85.7	
Macrosomic (>4000 g)	0	0.0	795	11.4		8	1.4	787	11.4	
Missing	8	1.5	88	1.3		9	1.6	87	1.3	
Infant sex					NS					NS
Female	233	45.1	3434	49.2		276	47.8	3391	49.0	
Male	284	54.9	3553	50.9		301	52.2	3536	51.1	

Abbreviation: BMI, body mass index.

<sup>a</sup>Exposed to any solvent.

<sup>b</sup>Any use/exposure during the periconceptional period.

## 4 | DISCUSSION

In the current study, we did not observe any definitive associations between maternal occupational exposure to organic solvents during the periconceptional period and SGA or PTB after adjusting for maternal education and residence. This was true for general exposure (any vs. none), as well as by solvent class (chlorinated, aromatic, and Stoddard solvent). Point estimates were slightly elevated for the highest category of any solvent exposure and SGA (aOR = 1.52; 95% CI 0.80–2.87) and PTB (aOR = 1.20; 95% CI 0.62–2.33), but confidence intervals were wide. Solvent exposure was somewhat limited among our study population for both birth outcomes of interest ( $n = 24$  exposed mothers of infants born SGA;  $n = 38$  exposed mothers of preterm infants); however, overall, our analyses do not suggest any substantial association between low-dose, infrequent periconceptional solvent exposure among expectant workers and SGA or PTB.

Our findings are consistent with other studies of maternal occupational exposure to organic solvents and adverse birth outcomes. Burdorf et al.<sup>40</sup> observed no associations between self-reported maternal occupational exposure to industrial solvents and

PTB with similar sample size ( $N = 8880$  participant mothers) (aOR = 1.02; 95% CI 0.58–1.81). In the same study, using a job exposure matrix (JEM) assessment strategy to determine exposure ( $n = 227$  exposed) or restricting the analysis to only mothers with probable occupational exposure to solvents ( $n = 69$  exposed) still resulted in no significant associations between maternal occupational solvent exposure and PTB.<sup>40</sup> More recently, Shirangi et al.<sup>41</sup> observed no association between maternal occupational exposure to organic solvents (any) and SGA or percentage of optimal birthweight (POBW) among infants in a population-based, prospective birth cohort study of 4142 pregnant workers from 2007 to 2011 in the United Kingdom (SGA aRR = 1.08; 95% CI 0.36–3.29; POBW aRR = 0.89; 95% CI 0.44–1.80). In that same study, there were similar number of infants with SGA in the study sample ( $n = 451$ ) compared to the NBDPS ( $n = 517$ ) and slightly higher study participation rates (approximately 80% vs. 65% among NBDPS controls).<sup>41</sup> Even studies of nonoccupational exposure to solvents have demonstrated similar findings. For example, Sorenson et al.<sup>42</sup> observed that residential exposure to paint fumes during pregnancy was not associated with reduced birth weight or PTB.

**TABLE 2** Prevalence of assessed occupational exposure to organic solvents during the periconceptional period among working mothers, stratified by SGA and PTB among infants without birth defects, National Birth Defect Prevention Study, USA, 1997–2011.

	SGA (n = 517) n (%)	Non-SGA (6987) n (%)	PTB (n = 577) n (%)	Non-PTB (n = 6927) n (%)	Total (n = 7504) n (%)
Any solvent	34 (6.6)	467 (6.7)	38 (6.6)	463 (6.7)	501 (6.7)
Chlorinated solvents	28 (5.4)	404 (5.8)	34 (5.9)	398 (5.8)	432 (5.8)
Carbon tetrachloride <sup>a</sup>	-	11 (0.2)	-	13 (0.2)	13 (0.2)
Chloroform	13 (2.5)	183 (2.6)	15 (2.6)	181 (2.6)	196 (2.6)
Methylene chloride	20 (3.9)	320 (4.6)	25 (4.3)	315 (4.6)	340 (4.5)
Perchloroethylene	18 (3.5)	244 (3.5)	20 (3.5)	242 (3.5)	262 (3.5)
Trichloroethylene	11 (2.1)	217 (3.1)	13 (2.3)	215 (3.1)	228 (3.0)
1,1,1-Trichloroethane	22 (4.3)	319 (4.6)	27 (4.7)	314 (4.5)	341 (4.5)
Stoddard solvent	12 (2.3)	126 (1.8)	14 (2.4)	124 (1.8)	138 (1.8)
Aromatic solvents	13 (2.5)	130 (1.9)	13 (2.3)	130 (1.9)	143 (1.9)
Benzene <sup>a</sup>	-	30 (0.4)	-	30 (0.4)	31 (0.4)
Toluene	13 (2.5)	122 (1.8)	13 (2.3)	122 (1.8)	135 (1.8)
Xylene	13 (2.5)	127 (1.8)	13 (2.3)	127 (1.8)	140 (1.9)

Abbreviations: PTB, preterm birth; SGA, small-for-gestational age.

<sup>a</sup>Exposure among participant mothers of  $n \leq 3$  not reported.**TABLE 3** Organic solvent exposure-response analysis for all jobs held by mothers of SGA, non-SGA, PTB, and non-PTB infants, National Birth Defects Prevention Study 1997–2011.

	non-SGA Controls n	SGA Cases n	cOR	(95% CI)	aOR <sup>a</sup>	(95% CI)	non-PTB Controls n	PTB Cases n	cOR	(95% CI)	aOR <sup>a</sup>	(95% CI)
Occupational organic solvent exposure												
No exposure	5493	395	1.00	-	1.00	-	5441	447	1.00	-	1.00	-
Low exposure	321	22	0.95	(0.61–1.49)	0.90	(0.57–1.40)	320	26	0.99	(0.66–1.49)	0.97	(0.64–1.47)
High exposure	95	11	1.61	(0.86–3.03)	1.52	(0.80–2.87)	93	10	1.31	(0.68–2.53)	1.20	(0.62–2.33)
Chlorinated solvents												
No exposure	5551	401	1.00	-	1.00	-	5502	450	1.00	-	1.00	-
Low exposure	264	16	0.84	(0.50–1.40)	0.79	(0.47–1.32)	261	23	1.08	(0.70–1.67)	1.07	(0.69–1.66)
High exposure	94	11	1.62	(0.86–3.05)	1.52	(0.80–2.87)	91	10	1.34	(0.70–2.60)	1.23	(0.63–2.39)
Aromatic solvents												
No exposure	5793	416	1.00	-	1.00	-	5738	471	1.00	-	1.00	-
Low exposure	70	8	1.59	(0.76–3.33)	1.68	(0.80–3.53)	73	5	0.83	(0.34–2.08)	0.83	(0.33–2.08)
High exposure	46	4	1.21	(0.43–3.38)	1.02	(0.36–2.88)	43	7	1.98	(0.89–4.43)	1.72	(0.77–3.87)
Stoddard solvent												
No exposure	5800	416	1.00	-	1.00	-	5747	469	1.00	-	1.00	-
Low exposure	53	3	0.79	(0.25–2.54)	0.65	(0.20–2.09)	52	8	1.89	(0.89–3.99)	1.73	(0.81–3.69)
High exposure	56	9	2.24	(1.10–4.56)	1.85	(0.90–3.81)	55	6	1.34	(0.57–3.12)	1.12	(0.47–2.63)

Abbreviations: aOR, adjusted odds ratio; CI, confidence interval; cOR, unadjusted crude odds ratio; PTB, preterm birth; SGA, small-for-gestational age.

<sup>a</sup>Model adjusted for maternal residence at delivery and maternal education (referent = >12 years).



**TABLE 4** Association between assessed probability of maternal occupational exposure to organic solvents during pregnancy and small-for-gestational age (SGA) and preterm birth (PTB) among infants without birth defects, National Birth Defects Prevention Study, USA, 1997–2011.

	Any probability of exposure					Probability of exposure $\geq 50\%$		
	n	cOR	(95% CI)	aOR <sup>a</sup>	(95% CI)	n	aOR <sup>a</sup>	(95% CI)
<b>SGA</b>								
Unexposed to any solvent	483	-	-	-	-	158	-	-
Exposed to any solvent	34	0.98	(0.69–1.41)	0.93	(0.65–1.34)	13	1.01	(0.57–1.81)
Chlorinated solvent(s)	28	0.93	(0.63–1.38)	0.88	(0.59–1.31)	11	1.06	(0.56–1.98)
Stoddard solvent	12	1.29	(0.71–2.36)	1.08	(0.59–1.99)	6	1.04	(0.45–2.43)
Aromatic solvent(s)	13	1.36	(0.77–2.43)	1.33	(0.74–2.37)	7	0.92	(0.42–2.00)
<b>PTB</b>								
Unexposed to any solvent	539	-	-	-	-	157	-	-
Exposed to any solvent	38	0.98	(0.70–1.39)	0.94	(0.67–1.33)	14	0.96	(0.55–1.68)
Chlorinated solvent(s)	34	1.03	(0.72–1.47)	0.99	(0.69–1.42)	14	1.23	(0.70–2.16)
Stoddard solvent	14	1.37	(0.78–2.39)	1.20	(0.68–2.11)	9	1.49	(0.73–3.03)
Aromatic solvent(s)	13	1.21	(0.68–2.15)	1.13	(0.63–2.02)	8	0.91	(0.44–1.89)

Abbreviations: aOR, adjusted odds ratio; CI, confidence interval; cOR, unadjusted crude odds ratio.

<sup>a</sup>Model adjusted for maternal residence at delivery and maternal education (referent = >12 years).**TABLE 5** Mean change in infant birthweight at term ( $\geq 37$  weeks' gestation) association with any estimated maternal occupational exposure to organic solvents during pregnancy, National Birth Defects Prevention Study, USA, 1997–2011.

	n	(%)	Mean	(SD)	Difference in birthweight (g) estimate (95% CI)			
					Crude		Adjusted <sup>a</sup>	
Total number of infants	6927	-	-	-	-	-	-	-
Unexposed	6464	93.3	3517.2	845.2	Reference	-	Reference	-
Exposed to any solvent	463	6.7	3559.5	917.7	42.3	(-37.9 to 122.4)	47.3	(-31.5 to 126.0)
Any chlorinated solvent	398	5.8	3567.3	915.1	50.1	(-35.9 to 136.2)	54.5	(-30.0 to 139.0)
Stoddard solvent	124	1.8	3509.9	717.6	-10.4	(-161.4 to 140.7)	36.2	(-112.5 to 184.9)
Any aromatic solvent	130	1.9	3545.7	749.1	26.2	(-121.4 to 173.7)	39.5	(-105.1 to 184.0)

Abbreviations: CI, confidence interval; SD, standard deviation.

<sup>a</sup>Model adjusted for maternal residence at delivery and maternal education (referent = >12 years).

The initial analysis by Desrosiers et al.<sup>10</sup> observed elevated but imprecise associations between fetal growth restriction (measured by SGA) and occupational periconceptional exposure to organic solvents (any and by chlorinated and aromatic solvent classes) among pregnant workers with a  $\geq 50\%$  probability of exposure; however, study findings were limited by relatively small numbers of exposed workers. The current study adds 9 years of analytic data ( $n = 4618$  mother/infant pairs from 2003 to 2011) and PTB as an additional outcome of interest. Increased sample size in the current study allowed for logistic regression analyses on the association between SGA and Stoddard solvent exposure among participant mothers with  $\geq 50\%$  probability of exposure, as well as dose/response analyses for all solvents (any and by class), though the classification criteria for SGA was slightly different between the two studies. Findings from stratifying the analysis by exposure assessment period (1997–2002 vs. 2003–2011) echo the comparisons between the

two studies as well (Table S2). Though we also observed an elevated, but imprecise adjusted OR for any solvent exposure and SGA among women in the first exposure assessment period (1997–2002), we observed a non-statistically significant inverse association between maternal occupational exposure to any solvent and SGA among women in the second exposure assessment period (2003–2011). It is possible that decreases in both occupational uses of organic solvents, as well as workplace substitutions of solvents with known and suspected health hazards, may have contributed to overall decreases in exposure among our study population and reduced our study's statistical power. Uncontrolled confounding by factors related to employment quality (e.g., stable contract, predictable scheduling, paid leave) cannot be ruled out as contributing to this finding, particularly if the jobs with higher solvent exposure also tended to have other more favorable workplace conditions associated with reduced PTB or SGA risk. Despite predominantly null findings from the entire study

period (1997–2011), we did observe elevated, nonsignificant ORs for estimated solvent exposure above the median versus below for both outcomes of interest in the current study. However, maternal exposure—particularly to Stoddard solvent—was still limited despite the increase in sample size and our findings may warrant further research. Most women in this population-based sample (even in the higher exposure group) were still exposed to these solvents infrequently and in small quantities, so these results should not be extrapolated to workplaces where very substantial and frequent exposure to these solvents occurs. Overall, though, these results are generally reassuring that infrequent and small-dose maternal periconceptional exposure to organic solvents in the workplace do not appear to be significantly associated with PTB or infants characterized as SGA.

The NBDPS provides a unique opportunity to examine the influence of a wide range of occupational exposures on reproductive outcomes. Because our study population of singleton infant/mother pairs is generally representative of infants from the 10 included US states during the study years,<sup>32</sup> results are more generalizable compared to other types of studies (e.g., hospital- or workplace-based studies). Details of the exposure assessment strategy have been previously published,<sup>36</sup> but the use of expert IH raters (as opposed to a JEM alone)—including corresponding interrater reliability—reduces the potential for misclassification bias. In addition, because the study population was limited only to participant mothers reporting at least one job for at least 1 month during the periconceptional period, confounding was minimized for variables predicting employment among women (e.g., select socioeconomic factors, sources of external stress, family structure characteristics, etc.) were likely not influential in the current study.<sup>43</sup> NBDPS participants' reported employment and job histories also highlight a study strength. Less than 1% ( $n = 14/7837$ ) of participant mothers who reported employment for at least 1 month during the periconceptional period did not provide sufficient job history information to assess occupational exposure to organic solvents; therefore, participant reporting of employment and job histories are an unlikely source of bias in this study.

Though sample size was robust, maternal occupational exposure to organic solvents among mothers of infants characterized as SGA or preterm was still infrequent (6.6%) (exposed mothers of SGA infants,  $n = 34$ ; exposed mothers of preterm infants,  $n = 38$ ), which may have led to imprecise estimates. In addition, the exposure window for the current study was restricted to the periconceptional period and does not account for potential influences of maternal organic solvent exposure later in pregnancy on either SGA or PTB. However, approximately 93.6% of participant mothers in our study sample worked the same or similar job(s) until delivery (data not shown) indicating that participant exposure likely did not change substantially throughout pregnancy. Reliance on theoretical and retrospective occupational exposure to organic solvents—as opposed to real-time quantitative measurements—may have led to some degree of non-differential misclassification. We also cannot completely rule out selection bias—where mothers predisposed to a higher prevalence of occupational organic solvent exposure and with adverse birth outcomes may have been more inclined to take part in the study—

though the participation rates of NBDPS controls (65%) are relatively high.<sup>31</sup> Additionally, while the most recent estimate of SGA prevalence in the US general population is approximately 11.1%, we noted a lower-than-expected prevalence of SGA in our study sample (approximately 6.9%).<sup>22</sup> Study participant mothers reported higher proportions of protective factors for SGA, such as not smoking and higher educational attainment, which may have contributed to the lower prevalence of SGA observed in our study.

Co-exposures in occupational studies of solvents still present a challenge and warrant some caution in finding interpretations. For example, among participant mothers determined to have exposure to aromatic solvents, approximately 73.4% ( $n = 105$ ) had exposure to both xylene and toluene, while approximately 20.3% ( $n = 29$ ) of mothers had exposure to all three aromatic solvents making it difficult to discern individual solvent effects (data not shown). Furthermore, our findings are representative of the unique exposure distributions within solvent class. For instance, among chlorinated solvents of interest, there were approximately 340 (4.5%) participant mothers with exposure to methylene chloride and only 13 (0.2%) participant mothers with exposure to carbon tetrachloride (Table 2). Because we were unable to examine associations between individual solvents and SGA and PTB, our results reflect the specific distribution of exposure within solvent classes among our study population.

A primary strength of the NBDPS is its ability to allow the study the association between select occupational exposures and pregnancy and birth outcomes. Though exposure to organic solvents is common among many industries, exposure was low within our study population. Occupational safety and health concerns over organic solvent exposure and suspected reprotoxicity have led to industry-wide substitutions and the phasing out of several solvents that may—at least in part—explain such infrequent and low-dose exposures within the study period (1997–2011). Most participant mothers who were determined to have occupational exposure to solvents still had estimated exposure intensities and frequencies that were far below the current OSHA permissible exposure limits (PELs). For example, maternal occupational exposure to carbon tetrachloride within our study population was very infrequent (e.g., only 13 out of 7504 mothers with determined exposure), and even where exposure was estimated to occur more frequently, it was often low-dose. For example, among the 135 participant mothers determined to have been occupationally exposed to toluene, approximately 97 (71.9%) had estimated exposure levels well below the current OSHA PEL (200 ppm averaged over an 8-hour shift) (data not shown). Therefore, it is likely that nonsignificant increases in effect observed between maternal exposure to organic solvents and SGA or PTB may be influenced heavily by high, but still infrequent exposures.

Results from the current study contribute to a growing body of literature that suggests maternal periconceptional exposure to organic solvents in the workplace—at levels well below current PELs—do not appear to be associated with premature delivery or infants characterized as SGA. However, because of the predominant occurrence of infrequent and low-dose exposure within our study population, we cannot rule out any effect(s) among pregnant workers with unusually high exposure to organic solvents.

## AUTHORS CONTRIBUTIONS

Kristen W. Van Buren led data analysis and drafting of the manuscript. Carissa M. Rocheleau, I-Chen Chen, Tania A. Desrosiers, Wayne T. Sanderson, and Elizabeth C. Ailes contributed to study conception and design of the analysis. Carissa M. Rocheleau, Tania A. Desrosiers, Wayne T. Sanderson, and Elizabeth C. Ailes contributed subject matter expertise. Carissa M. Rocheleau, I-Chen Chen, Tania A. Desrosiers, Maria D. Politis, and Elizabeth C. Ailes assisted with drafting of the manuscript. I-Chen Chen assisted with statistical analysis and performed duplication. All authors provided a critical review of the manuscript and approved the submission.

## ACKNOWLEDGMENTS

The authors would like to thank all NBDPS collaborators and participants without whom this work would not be possible. This includes the Arkansas Department of Health; California Department of Public Health Maternal Child and Adolescent Health Division; Georgia Department of Public Health and the Metropolitan Atlanta Congenital Defects Program; Iowa Department of Public Health (Iowa Registry for Congenital and Inherited Disorders); Massachusetts Department of Public Health; North Carolina Department of Health and Human Services; New Jersey Department of Health; New York State Department of Health (Congenital Malformations Registry); Texas Department of State Health Services (Birth Defects Epidemiology and Surveillance Branch); and Utah Department of Health (Utah Birth Defect Network). This project was supported through Centers for Disease Control and Prevention (CDC) cooperative agreements under PA #96043, PA #02081, FOA #DD09-001, FOA #DD13-003, and NOFO #DD18-001 to the Centers for Birth Defects Research and Prevention participating in the National Birth Defects Prevention Study (NBDPS) and/or the Birth Defects Study To Evaluate Pregnancy exposureS (BD-STEPS). Additional support for occupational exposure assessment was provided by contract 200-2000-08018 from the Centers for Disease Control and Prevention and the National Institute for Occupational Safety and Health. This work was completed in partial fulfillment of the lead author's Doctor of Public Health degree in Epidemiology in the Department of Epidemiology and Environmental Health, University of Kentucky College of Public Health, through support from the Central Appalachian Regional Education and Research Center (CARERC).

## CONFLICTS OF INTEREST STATEMENT

The authors declare no conflicts of interest.

## DISCLOSURE BY AJIM EDITOR OF RECORD

John Meyer declares that he has no conflict of interest in the review and publication decision regarding this article.

## DATA AVAILABILITY STATEMENT

The study questionnaires and process for accessing the data used in this study is described at <https://www.cdc.gov/ncbddd/birthdefects/nbdps-public-access-procedures.html>. The code book and analytic code may be made available upon request.

## ETHICS APPROVAL AND INFORMED CONSENT

The study protocols were approved by the Human Subjects Review Board for the Centers for Disease Control and Prevention (protocol #2087) and the Institutional Review Board for each participating site. All interviewed study participants provided informed consent.

## DISCLAIMER

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention.

## ORCID

Kristen W. Van Buren  <https://orcid.org/0000-0002-0825-6882>

I-Chen Chen  <http://orcid.org/0000-0001-6764-8395>

## REFERENCES

- Organic Solvents. National Institute for Occupational Safety and Health (NIOSH). Centers for Disease Control and Prevention (CDC). Published November 2, 2018. <https://www.cdc.gov/niosh/topics/organsolv/default.html?msckid=41b0976ca60011ec9a91b332a736eef0>
- Donald JM, Hooper K, Hopenhayn-Rich C. Reproductive and developmental toxicity of toluene: a review. *Environ Health Perspect.* 1991;94:237. doi:10.2307/3431317
- Dietz D. Toxicity studies of acetone administered in the drinking water of rodents. *Fundam Appl Toxicol.* 1991;17(2):347-360. doi:10.1016/0272-0590(91)90224-r
- White RF, Proctor SP. Solvents and neurotoxicity. *The Lancet.* 1997;349:1239-1243. doi:10.1016/s0140-6736(96)07218-2
- Aylward LL, Hays SM, Kirman CR, et al. Relationships of chemical concentrations in maternal and cord blood: a review of available data. *J Toxicology Environ Health, Part B.* 2014;17(3):175-203. doi:10.1080/10937404.2014.884956
- Firestone JA, Gospe Jr. SM. *Organic Solvents*. Clinical Neurotoxicology. WB Saunders; 2009:401-414.
- Maronpot RR. Ovarian toxicity and carcinogenicity in eight recent National Toxicology Program Studies. *Environ Health Perspect.* 1987;73:125-130. doi:10.1289/ehp.8773125
- Wennborg H, Bodin L, Vainio H, Axelsson G. Pregnancy outcome of personnel in Swedish biomedical research laboratories. *J Occup Environ Med.* 2000;42(4):438-446. doi:10.1097/00043764-200004000-00022
- Ha E, Cho S-I, Chen D, et al. Parental exposure to organic solvents and reduced birth weight. *Arch Environ Health.* 2002;57(3):207-214. doi:10.1080/00039890209602938
- Desrosiers TA, Lawson CC, Meyer RE, et al. Assessed occupational exposure to chlorinated, aromatic and Stoddard solvents during pregnancy and risk of fetal growth restriction. *Occup Environ Med.* 2015;72(8):587-593. doi:10.1136/oemed-2015-102835
- Stoltenburg-Didinger G, Altenkirch H, Wagner M. Neurotoxicity of organic solvent mixtures: embryotoxicity and fetotoxicity. *Neurotoxicol Teratol.* 1990;12(6):585-589. doi:10.1016/0892-0362(90)90066-l
- Ahmed P, Jaakkola JJK. Exposure to organic solvents and adverse pregnancy outcomes. *Hum Reprod.* 2007;22(10):2751-2757. doi:10.1093/humrep/dem200
- Zhu JL, Vestergaard M, Hjollund NH, Olsen J. Pregnancy outcomes among female hairdressers who participated in the Danish National Birth Cohort. *Scand J Work Environ Health.* 2006;32(1):61-66. doi:10.5271/sjweh.977
- Olsen J, Hemminki K, Ahlborg G, et al. Low birthweight, congenital malformations, and spontaneous abortions among dry-cleaning

- workers in Scandinavia. *Scand J Work Environ Health*. 1990;16(3):163-168. doi:10.5271/sjweh.1800
15. Khattak S, K-Moghtader G, McMartin K, Barrera M, Kennedy D, Koren G. Pregnancy outcome following gestational exposure to organic solvents. *JAMA*. 1999;281(12):1106. doi:10.1001/jama.281.12.1106
  16. Zhu JL. Laboratory work and pregnancy outcomes: a study within The National Birth Cohort in Denmark. *Occup Environ Med*. 2006;63(1):53-58. doi:10.1136/oem.2005.021204
  17. Schaumburg I, Olsen J. Birth weight and gestational age among children of Danish pharmacy assistants. *J Epidemiol Commun Health*. 1991;45(1):49-51. doi:10.1136/jech.45.1.49
  18. Hewitt JB, Tellier L. Risk of adverse outcomes in pregnant women exposed to solvents. *J Obstet, Gynecol Neonatal Nurs*. 1998;27(5):521-531. doi:10.1111/j.1552-6909.1998.tb02618.x
  19. Muhihi A, Sudfeld CR, Smith ER, et al. Risk factors for small-for-gestational-age and preterm births among 19,269 Tanzanian newborns. *BMC Pregn Childbirth*. 2016;16(1):110. doi:10.1186/s12884-016-0900-5
  20. Callaghan WM, MacDorman MF, Rasmussen SA, Qin C, Lackritz EM. The contribution of preterm birth to infant mortality rates in the United States. *Pediatrics*. 2006;118(4):1566-1573. doi:10.1542/peds.2006-0860
  21. Division of Reproductive Health, National Center for Chronic Disease Prevention and Health Promotion. Preterm Birth. 2021. Accessed May 2021. <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pretermbirth.htm>
  22. Lee AC, Kozuki N, Cousens S, et al. Estimates of burden and consequences of infants born small for gestational age in low and middle income countries with intergrowth-21st standard: analysis of CHERG datasets. *BMJ*. 2017;358:j4229. doi:10.1136/bmj.j4229
  23. Allin M, Rooney M, Griffiths T, et al. Neurological abnormalities in young adults born preterm. *J Neurol Neurosurg Psychiatry*. 2006;77(4):495-499. doi:10.1136/jnnp.2005.075465
  24. O'Shea TM, Allred EN, Dammann O, et al. The Elgan Study of the brain and related disorders in extremely low gestational age newborns. *Early Hum Dev*. 2009;85(11):719-725. doi:10.1016/j.earlhumdev.2009.08.060
  25. Jensen EA, Foglia EE, Dysart KC, et al. Adverse effects of small for gestational age differ by gestational week among very preterm infants. *Arch Dis Childhood - Fetal Neonatal*. 2018;104(2):F192-F198. doi:10.1136/archdischild-2017-314171
  26. Blair E. Paediatric implications of IUGR with special reference to cerebral palsy. *Intrauterine Growth Restriction*. Springer; 2000:351-366. doi:10.1007/978-1-4471-0735-4\_19
  27. van Wassenae A. Neurodevelopmental consequences of being born SGA. *Pediatr Endocrinol Rev*. 2005;2(3):372-377.
  28. von Beckerath A-K, Kollmann M, Rotky-Fast C, Karpf E, Lang U, Klaritsch P. Perinatal complications and long-term neurodevelopmental outcome of infants with intrauterine growth restriction. *Am J Obstet Gynecol*. 2013;208(2):130.e1-130.e6. doi:10.1016/j.ajog.2012.11.014
  29. Luu TM, Katz SL, Leeson P, Thébaud B, Nuyt A-M. Preterm birth: risk factor for early-onset chronic diseases. *Can Med Assoc J*. 2015;188(10):736-746. doi:10.1503/cmaj.150450
  30. Mericq V, Martinez-Aguayo A, Uauy R, Iñiguez G, Van der Steen M, Hokken-Koelega A. Long-term metabolic risk among children born premature or small for gestational age. *Nat Rev Endocrinol*. 2016;13(1):50-62. doi:10.1038/nrendo.2016.127
  31. Reefhuis J, Gilboa SM, Anderka M, et al. The National birth defects prevention study: a review of the methods. *Birth Defects Res Part A: Clin Mol Teratol*. 2015;103(8):656-669. doi:10.1002/bdra.23384
  32. Cogswell ME, Bitsko RH, Anderka M, et al. Control selection and participation in an ongoing, population-based, case-control study of birth defects. *Am J Epidemiol*. 2009;170(8):975-985. doi:10.1093/aje/kwp226
  33. Langlois PH, Hoyt AT, Desrosiers TA, et al. Maternal occupational exposure to polycyclic aromatic hydrocarbons and small for gestational age offspring. *Occup Environ Med*. 2014;71(8):529-535. doi:10.1136/oemed-2013-101833
  34. Chen L, Bell EM, Browne ML, Druschel CM, Romitti PA. Exploring maternal patterns of dietary caffeine consumption before conception and during pregnancy. *Matern Child Health J*. 2014;18(10):2446-2455. doi:10.1007/s10995-014-1483-2
  35. Carmichael S, Yang W, Shaw G. Maternal dietary nutrient intake and risk of preterm delivery. *Am J Perinatol*. 2012;30(07):579-588. doi:10.1055/s-0032-1329686
  36. Rocheleau CM, Lawson CC, Waters MA, et al. Inter-rater reliability of assessed prenatal maternal occupational exposures to solvents, polycyclic aromatic hydrocarbons, and heavy metals. *J Occup Environ Hyg*. 2011;8(12):718-728. doi:10.1080/15459624.2011.627293
  37. Talge NM, Mudd LM, Sikorskii A, Basso O. United States birth weight reference corrected for implausible gestational age estimates. *Pediatrics*. 2014;133(5):844-853. doi:10.1542/peds.2013-3285
  38. Textor J, van der Zander B, Gilthorpe MK, Liskiewicz M, Ellison GT. Robust causal inference using directed acyclic graphs: the R package 'dagitty'. *Int J Epidemiol*. 2016;45(6):1887-1894.
  39. Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. *Epidemiology*. 1999;10(1):37-48. doi:10.1097/00001648-199901000-00008
  40. Burdorf A, Brand T, Jaddoe VW, Hofman A, Mackenbach JP, Steegers EAP. The effects of work-related maternal risk factors on time to pregnancy, preterm birth and birth weight: the generation R study. *Occup Environ Med*. 2010;68(3):197-204. doi:10.1136/oem.2009.046516
  41. Shirangi A, Wright J, Blair EM, McEachan RR, Nieuwenhuijsen MJ. Occupational chemical exposures in pregnancy and fetal growth: evidence from the born in Bradford study. *Scand J Work Environ Health*. 2020;46(4):417-428. doi:10.5271/sjweh.3878
  42. Sørensen M, Andersen A-MN, Raaschou-Nielsen O. Non-occupational exposure to paint fumes during pregnancy and fetal growth in a general population. *Environ Res*. 2010;110(4):383-387. doi:10.1016/j.envres.2010.02.011
  43. Rocheleau CM, Bertke SJ, Lawson CC, et al. Factors associated with employment status before and during pregnancy: implications for studies of pregnancy outcomes. *Am J Ind Med*. 2017;60(4):329-341. doi:10.1002/ajim.22700

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Van Buren KW, Rocheleau CM, Chen I-C, et al. Maternal occupational exposure to selected organic and chlorinated solvents and delivery of small-for-gestational age or preterm infants. *Am J Ind Med*. 2023;66:842-853. doi:10.1002/ajim.23516