



Short communication

Lessons learned from 9/11: Mental health perspectives on the COVID-19 pandemic

Jonathan DePierro^{a,1,*}, Sandra Lowe^a, Craig Katz^b^a Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA^b Departments of Psychiatry, Medical Education, and Health System Design & Global Health, Icahn School of Medicine at Mount Sinai, New York, USA

ARTICLE INFO

Keywords:

Disaster mental health

PTSD

Emergency responders

ABSTRACT

The COVID-19 pandemic will likely lead to high rates of PTSD, depression, and substance misuse among survivors, victims' families, medical workers, and other essential personnel. The mental health response to the 9/11/01 terrorist attacks, culminating in a federally-funded health program, provides a template for how providers may serve affected individuals. Drawing on the 9/11 experience, we highlight effective prevention measures, likely short and long-term treatment needs, vulnerable subgroups, and important points of divergence between 9/11 and the COVID-19 pandemic. Mental health monitoring, early identification of at-risk individuals, and treatment irrespective of financial barriers are essential for minimizing chronic distress.

At the time of this writing (April 14, 2020), over 1.9 million people across the world have been infected with COVID-19 and there have been approximately 120,000 deaths (Johns Hopkins University, 2020). In New York State, which has become one of the ground zeroes for this pandemic, there have been more than 196,000 infections and 10,000 deaths, far exceeding the number of lives lost on 9/11/01. Several lessons learned from the mental health response to 9/11 should inform how providers and institutions meet the treatment needs of those affected by COVID-19.

1. Prevention

The mental health risk posed to 9/11 responders by the level of trauma to which they were exposed was likely worsened by lack of adequate monitoring of that exposure, especially the duration of their time at Ground Zero. Efforts should be made to ensure that the exposure of healthcare workers responding in the many ground zeros that our hospitals have become is monitored, even as the demand for their services rises with the infection rates.

It is also important to note that 9/11 responders who were part of traditional first responder populations such as the police and fire service also likely had the mental health benefit of robust social networks provided by their affiliation with firehouses, stationhouses, and unions. Such social support offered them the protection that the many

thousands of non-traditional responders, including unaffiliated volunteers who spontaneously came forward to help, had available to them less reliably (Pietrzak et al., 2014). In the COVID-19 crisis, hospitals in particular should ensure that all possible efforts are made to provide social support to their staff, whether through hospital communications, supervisors, or formalized peer support groups. Social support is a largely free, natural, and abundant resource that only needs to be channeled.

2. Short-term treatment needs and barriers

Mental health support for 9/11 responders and survivors started soon after the attacks, as many volunteer mental health professions flooded makeshift family support centers and responder aid stations. This initial support has been described as a "trauma tent," a safe, supportive space surrounding affected individuals (Katz and Nathaniel, 2002). These efforts became formalized within hospital-based clinics and charity-subsidized private treatment. Now, in the midst of COVID-19, again thousands of providers in New York have volunteered for mental health support services, many participating in city and state-organized efforts.

With the benefit of technological advances, this aid can be deployed as effectively via video telehealth platforms. Significant issues were encountered securing space to see patients as the initial Mount Sinai 9/

* Corresponding author at: Mount Sinai World Trade Center Health Program Clinical Center of Excellence, One Gustave L. Levy Place, Box 1230, New York, USA, 10029

E-mail address: jonathan.depierro@mssm.edu (J. DePierro).

¹ Permanent address: Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, Box 1230, New York, NY 10029, USA.

11 mental health treatment and monitoring program in New York was housed within the confines of an already bustling medical center, and modern advances in telepsychiatry can remedy this in the case of COVID-19. To address immediate COVID-19 related mental health needs for staff, the Mount Sinai Hospital System is currently deploying a range of mostly telehealth-based supportive mental health services, including support groups, individual therapy, and crisis lines.

All of these efforts are likely to be fruitful and will alleviate much distress. However, several barriers should be noted. Many individuals may not make use of these short-term services, due to mental health stigma, denial, dissociative defenses, conflicting time demands, or general discomfort with emotional disclosure (particularly in group support/debriefing). Thus, individuals with the highest level of need may not engage when initial services are offered. The Mount Sinai-based mental health program for 9/11 responders lowered these barriers by partnering and co-locating with medical colleagues who were addressing 9/11 related physical health issues, and finding comparable integration will likely be essential to ensure COVID-19 related mental health programming reaches the greatest possible number of affected people.

3. Long-term health monitoring and treatment

The mental health treatment of World Trade Center rescue, recovery and cleanup workers has highlighted the need for long-term monitoring of COVID-19 pandemic survivors, healthcare workers, other essential personnel (e.g. delivery, postal, and grocery store workers) and surviving family members. Health monitoring and treatment efforts for 9/11 survivors and responders were established soon after the attacks and continue to this day, and this work is now funded through the James Zadroga Act. Without similarly unified health registry and treatment services, it is likely that many individuals, particularly those from underserved groups, will experience chronic mental health consequences while being unable to access high-quality healthcare services.

4. Vulnerable populations

Two vulnerable populations should be highlighted. First, high rates of post-traumatic stress disorder (PTSD), clinical depression, and recurrent alcohol use problems are anticipated among healthcare workers, now on the front lines of the COVID-19 pandemic, and serving as the most “visible” responder group during this crisis. Concerns have been raised around moral injury among healthcare workers given that hard choices may need to be made regarding rationing care, including ventilators (Williamson et al., 2020). Moral injury has been associated with increased risk for psychiatric disorders and suicidal ideation (Bryan et al., 2018). This issue recalls the identity conflicts faced by 9/11 uniformed responders: while trained to serve and protect others, many felt that they failed in this mission because so many people were killed and so few were rescued from the “pile” at Ground Zero. Just world assumptions were shattered then, as they likely will be for medical personnel now.

Second, special attention should be paid to the mental health outcomes of non-medical “essential personnel,” including governmental employees, healthcare administrators and support staff, and food delivery workers. This concern arises from striking parallels to the 9/11 experience. Non-traditional 9/11 responders (e.g. construction, cleanup, asbestos workers; city employees; and volunteers) have consistently higher rates of chronic PTSD than uniformed responders (e.g. police), potentially related to having greater pre and post-9/11 life stressors, higher rates of pre-9/11 psychiatric diagnoses, and lower social support around the time of 9/11 (Pietrzak et al., 2014). Further, this group mostly lacked disaster response experience and found themselves taking on tasks well outside the scope of their jobs, often not by choice but due to economic necessity.

5. Contrasts

It is often said of disaster experience, “If you have seen one disaster, you have seen one,” reflecting the uniqueness of every disaster. Despite the many similarities between 9/11 and the COVID-19 pandemic, they must be seen through the lens of differences between two events, differences which suggest the possibility of even greater psychiatric fallout for the pandemic's responders. First, while the daily exposure of responders at Ground Zero took place over months, the acute impact proved to be more or less limited to 9/11/01 itself whereas the acute impact of COVID-19 is unfolding over at least months. Second, the militaristic conceptualization of the COVID-19 response as a “war” wherein healthcare professionals have been “(re)deployed” to work with COVID-19 patients underscores how obligation and immediate personal jeopardy may color this response far more than 9/11 ultimately did. Third, to the extent that problematic pre-disaster psychosocial circumstances are a risk factor for survivors' future risk of psychopathology, the heightened levels of political and social discord in the United States and elsewhere in the last several years is of concern.

6. Concluding thoughts

The reverberations of the 2020 COVID-19 pandemic will likely be felt for many decades, as the impact of 9/11 is still felt by many, including New Yorkers, nearly 20 years after the attacks. The tremendous loss of life will invariably lead to adverse mental health effects in the population, including family members of victims, health workers who tried valiantly to save the ill, and other essential personnel who continued to serve the public in the face of ongoing invisible life threat. Rather than the sudden jolt of fear and horror that accompanied the 9/11 attacks, the COVID-19 pandemic will likely bring a more insidious wave of anxiety, anger, and grief as casualties increase. Widely-accessible short and long-term mental health treatment services, such as those provided following 9/11, may prove necessary to respond to this need.

Funding sources

Drs. Lowe and DePierro are supported by CDC/NIOSH Contract 200-2017-93428. The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of CDC/NIOSH.

CRediT authorship contribution statement

Jonathan DePierro: Conceptualization, Writing - original draft, Writing - review & editing. **Sandra Lowe:** Writing - original draft, Writing - review & editing. **Craig Katz:** Writing - original draft, Writing - review & editing.

Declaration of Competing Interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Craig Katz is a paid consultant to the International Association of Firefighters Center of Excellence for Behavioral Health Treatment and Recovery. Jonathan DePierro and Sandra Lowe have no competing financial or personal relationships that could have appeared influence the work reported in this paper.

Acknowledgments

The authors would like to thank Drs. Adriana Feder and Robert Pietrzak, who provided comments on an earlier version of this manuscript. The authors would also like to thank all of the rescue, recovery

and cleanup workers and survivors who participate in the World Trade Center Health Program.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.psychres.2020.113024](https://doi.org/10.1016/j.psychres.2020.113024).

References

- Bryan, C.J., Bryan, A.O., Roberge, E., Leifker, F.R., Rozek, D.C., 2018. Moral injury, posttraumatic stress disorder, and suicidal behavior among National Guard personnel. *Psychol. Trauma* 10, 36–45. <https://doi.org/10.1037/tra0000290>.
- Johns Hopkins University, 2020. Coronavirus Global Cases by the Center for Systems Science and Engineering [WWW Document]. URL <https://coronavirus.jhu.edu/map.html>.
- Katz, C.L., Nathaniel, R., 2002. Disasters, psychiatry, and psychodynamics. *J. Am. Acad. Psychoanal.* 30, 519–529. <https://doi.org/10.1521/jaap.30.4.519.24188>.
- Pietrzak, R.H., Feder, A., Singh, R., Schechter, C.B., Bromet, E.J., Katz, C.L., Reissman, D.B., Ozbay, F., Sharma, V., Crane, M., Harrison, D., Herbert, R., Levin, S.M., Luft, B.J., Moline, J.M., Stellman, J.M., Udasin, I.G., Landrigan, P.J., Southwick, S.M., 2014. Trajectories of PTSD risk and resilience in World Trade Center responders: an 8-year prospective cohort study. *Psychol. Med.* 44, 205–219. <https://doi.org/10.1017/S0033291713000597>.
- Williamson, V., Murphy, D., Greenberg, N., 2020. COVID-19 and experiences of moral injury in front-line key workers. *Occup. Med.* <https://doi.org/10.1093/occmed/kqaa052>.
- Bryan, C.J., Bryan, A.O., Roberge, E., Leifker, F.R., Rozek, D.C., 2018. Moral injury, posttraumatic stress disorder, and suicidal behavior among National Guard