

REVIEW ARTICLE

Biomarkers of heatstroke-induced organ injury and repair

Zachary J. Schlader¹  | Michael S. Davis² | Abderrezak Bouchama³ 

¹Department of Kinesiology, School of Public Health, Indiana University, Bloomington, IN, USA

²Department of Physiological Sciences, College of Veterinary Medicine, Oklahoma State University, Stillwater, OK, USA

³Department of Experimental Medicine, King Abdullah International Medical Research Center, King Saud bin Abdulaziz University for Health Sciences, Ministry of National Guard – Health Affairs, Riyadh, Saudi Arabia

Correspondence

Zachary J. Schlader, Department of Kinesiology, School of Public Health, Indiana University, Bloomington, IN, USA.
Email: zschlade@iu.edu

Funding information

National Institute of Occupational Safety and Health; R01OH011528; Indiana Clinical and Translational Science Institute, Grant/Award Number: UL1TR002529

Edited by: Jeremy Ward

Abstract

Classic and exertional heatstroke cause acute injury and damage across numerous organ systems. Moreover, heatstroke survivors may sustain long-term neurological, cardiovascular and renal complications with a persistent risk of death. In this context, biomarkers, defined as biological samples obtained from heatstroke patients, are needed to detect early organ injury, and predict outcomes to develop novel organ preservation therapeutic strategies. This narrative review provides preliminary insights that will guide the development and future utilization of these biomarkers. To this end, we have identified numerous biomarkers of widespread heatstroke-associated cellular injury, tissue damage and repair (extracellular heat shock proteins 72 and 60, high mobility group box protein 1, histone H3, and interleukin-1 α), and other organ-specific biomarkers including those related to the cardiovascular system (cardiac troponin I, endothelium-derived factors, circulation endothelial cells, adhesion molecules, thrombomodulin and von Willebrand factor antigen), the kidneys (plasma and urinary neutrophil gelatinase-associated lipocalin), the intestines (intestinal fatty acid-binding protein 2), the brain (serum S100 β and neuron-specific enolase) and skeletal muscle (creatin kinase, myoglobin). No specific biomarkers have been identified so far for liver or lung injury in heatstroke. Before translating the identified biomarkers into clinical practice, additional preclinical and clinical prospective studies are required to further understand their clinical utility, particularly for the biomarkers related to long-term post-heatstroke health outcomes.

KEYWORDS

biomarkers, classic heatstroke, exertional heatstroke, hyperthermia, multiorgan failure, organ systems

1 | INTRODUCTION

With global temperatures rising, heatwaves have become more frequent, intense and longer in duration (IPCC, 2021), thereby exposing a greater proportion of the population to the deadly risk of extreme heat (Mora et al., 2017). Heat exposure elevates the risk of developing heat illness, which describes a continuum of pathologies ranging from relatively mild conditions associated with

alterations in cardiovascular functioning (e.g., heat syncope, heat exhaustion) to potentially lethal heatstroke (Armstrong et al., 2007). Heatstroke is characterized by a rapidly rising core temperature (usually >40.1°C), central nervous system dysfunction, systemic inflammation and organ system injury that can culminate in death (Bouchama et al., 2022). Heatstroke manifests in two distinct forms, classic heatstroke (CHS) and exertional heatstroke (EHS) (Bouchama et al., 2022). CHS typically occurs during resting exposures to hot and

humid conditions, which limit the capacity for heat loss, and more frequently affects the very young, older adults and populations with comorbidities (Bouchama et al., 2022). EHS more often occurs in younger, physically active, healthy adults, such as military personnel and competitive or recreational athletes, and is characterized by the inability to adequately dissipate heat sufficient to offset the heat load caused by physical exertion (i.e., elevations in metabolic heat production) (Bouchama et al., 2022). As a result, EHS can occur across a comparatively wide range of environmental conditions (Rae et al., 2008; Stacey et al., 2022). Therefore, the threat of heatstroke can impact the health and well-being of millions of people around the globe, and this risk will likely continue to increase in the coming decades (IPCC, 2021).

Heatstroke is diagnosed mainly through clinical signs and symptoms. Hallmarks of heatstroke include an elevated core temperature (often $>40.1^{\circ}\text{C}$) and central nervous system alterations (e.g., confusion, irritability, altered consciousness) (Bouchama et al., 2022; Casa et al., 2015) combined with a relevant patient history of exposure to hot and humid conditions and/or strenuous physical exertion (Leon & Bouchama, 2011). Notably, however, a core temperature less than 40.1°C should never rule out the diagnosis of heatstroke as the measurement of core temperature is often delayed, usually after the patient's removal from environmental heat and/or cessation of the exercise and arrival of paramedics. With early diagnosis and rapid cooling, the prognosis for heatstroke victims is generally good (Demartini et al., 2015). In contrast, the risk of prolonged morbidity and mortality increases when heatstroke diagnosis and treatment are delayed, resulting in tissue damage and multiorgan dysfunction/failure, and eventually death. This evolution has been attributed to heat cytotoxicity, excessive systemic inflammation and hypercoagulability (Bouchama et al., 2022; Casa et al., 2012; Leon & Bouchama, 2011). The systemic nature of heatstroke pathology is exemplified in an analysis of 2529 EHS episodes occurring in military personnel from 2008 to 2014, demonstrating abnormal clinical indices of liver, renal and haematological function and muscle damage that peak 0–4 days following diagnosis and persist for up to 16 days post-injury (Ward et al., 2020). These findings corroborate animal literature demonstrating widespread damage across numerous organ systems following CHS (Leon et al., 2006b; Roberts et al., 2008) and EHS (King et al., 2015). Moreover, observational evidence indicates that a single episode of heatstroke could have deleterious long-term all-cause mortality (Wallace et al., 2007), neurological (Argaud et al., 2007; Dematte et al., 1998; Yang et al., 2017), cardiovascular (Wallace et al., 2007; Wang et al., 2019) and renal (Wang et al., 2019; Wu et al., 2021) complications with a continuous risk of death (Bouchama et al., 2022). These observations have raised the need to develop tools that aid in detecting tissue and organ injury, monitoring recovery (or repair failure), and predicting long-term complications following heatstroke (Figure 1). Therefore, this narrative review provides an up-to-date overview of novel biological markers (biomarkers) that could help identify the patients at risk of organ injury or long-term complications following heatstroke. Our goal is not to provide recommendations

New Findings

- **What is the topic of this review?**

The status and potential role of novel biological markers (biomarkers) that can help identify the patients at risk of organ injury or long-term complications following heatstroke.

- **What advances does it highlight?**

Numerous biomarkers were identified related to many aspects of generalized heatstroke-induced cellular injury and tissue damage, and heatstroke-provoked cardiovascular, renal, cerebral, intestinal and skeletal muscle injury. No novel biomarkers were identified for liver or lung injury.

for the clinical deployment of biomarkers because the literature base remains too underdeveloped for this to be reasonably accomplished. Rather, we will provide preliminary insights on biomarkers that could aid the prediction of short-term health outcomes in heatstroke and ultimately help develop novel preservation strategies to reduce organ damage and potentially prevent long-term sequelae (Figure 2).

For this review, a *biomarker* is defined as a measurable substance derived from a bodily fluid that does not require assessment of clinical signs and symptoms. Notably, this definition purposely excludes standard clinical and biochemical tests that are helpful for patient care but are not specific to heatstroke. Furthermore, in this review, the term *heatstroke* will refer to observations that are consistent between CHS and EHS. At the same time, *CHS* and *EHS* will be used when referring to specific classic or EHS observations or pathophysiology. Moreover, we will focus this review on tissues, organs and organ systems impacted by heatstroke. In the interest of conciseness, we will not provide an in-depth overview of the aetiology of heatstroke, except where it is relevant to understanding injury biomarkers. Likewise, we will not be addressing risk predictive or diagnostic biomarkers of heatstroke, but we invite the interested reader to explore studies on these topics (e.g., Alele et al., 2021; Lu, Wang et al., 2004; Stacey et al., 2022).

2 | GENERALIZED CELLULAR INJURY AND TISSUE DAMAGE

Heatstroke is associated with a complex innate immune response characterized by neutrophil and complement protein activation, and production of pro- and anti-inflammatory cytokines, chemokines and acute phase proteins (Bouchama et al., 2022). The activation of the immune response has been attributed to endotoxin leaking from the heat-damaged gastrointestinal tract (Hall et al., 1999) and danger signal molecules or alarmins released into the circulation from damaged or dying cells (Gallucci & Matzinger, 2001). Alarmins, also




FIGURE 1 Potential utility of biomarkers to detect tissue and organ injury, monitor recovery and predict long-term complications following heatstroke. The purpose of this narrative review is to identify biomarkers of heatstroke induced organ injury and repair

known as damage-associated molecular patterns, play a crucial role in removing necrotic tissue debris and other repair mechanisms (Chen & Nuñez, 2010). Alarmins have been implicated in the excessive and sustained inflammatory response leading to the production of reactive oxygen species and proteolytic enzymes that cause further tissue damage and the progression to multiorgan failure (Chen & Nuñez, 2010). Several alarmins have been identified in heatstroke patients (Huisse et al., 2008; Ruell et al., 2014) and preclinical models (Bruchim et al., 2015; Dehbi et al., 2010). These include extracellular heat shock protein (eHSP) 72 (Bruchim et al., 2015; Dehbi et al., 2010; Huisse et al., 2008; Ruell et al., 2014) and eHSP60 (Dehbi et al., 2010; Huisse et al., 2008), high mobility group box protein 1 (HMGB1) (Dehbi et al., 2012; Tong et al., 2011b), histone H3 (Bruchim et al., 2017; Li et al., 2021) and interleukin (IL)-1 α (Bouchama et al., 1991).

Heat shock proteins are a family of proteins produced by cells when exposed to stressful conditions, including extreme heat. They are produced intracellularly and function as chaperones to prevent heat-induced protein misfolding or aggregation (Kultz, 2005). eHSPs do not exert a chaperone function but act as cytokines engaging

pattern-recognition receptors, such as toll-like receptors, to elicit an inflammatory response (Chen & Nuñez, 2010). eHSP72 is released in a sustained manner for up to 72 h in non-human primate models of severe heatstroke and was associated with multiorgan damage and death, suggesting a role as a prognostic biomarker (Dehbi et al., 2010). Moreover, high eHSP70 and eHSP60 were documented in the plasma of heatstroke patients, and the former correlated with plasma IL-8 concentrations (Huisse et al., 2008). HMGB1, histone, and IL-1 α are chromatin-associated nuclear proteins that are released by injured or dying cells (Cohen et al., 2010). HMGB1 was demonstrated as an early mediator of inflammation, tissue injury and lethality in rodent model of heatstroke (Dehbi et al., 2012), and plasma HMGB1 is associated with heatstroke severity and mortality (Huisse et al., 2008). Likewise, histone H3 was linked to coagulopathy, and heatstroke severity and lethality (Bruchim et al., 2017; Li et al., 2021). Finally, plasma concentrations of IL-1 α were increased in a small cohort of CHS patients (Bouchama et al., 1991). Notably, IL-1 α functions intracellularly as a transcription factor in healthy cells, but upon release into the circulation behaves as a proinflammatory cytokine and contributes




FIGURE 2 Identified biomarkers of heatstroke induced organ injury and/or recovery

to tissue remodelling and repair (Cohen et al., 2010). Overall, alarmins are promising biomarkers of heatstroke-associated tissue damage and repair that deserve further validation studies.

3 | CARDIOVASCULAR SYSTEM

3.1 | Heart

Elevations in body temperature have profound effects on the cardiovascular system to support the increases in skin blood flow necessary to promote heat loss (Crandall & Wilson, 2014). The heart responds by increasing cardiac output, mediated mainly through increases in heart rate and the maintenance of stroke volume, which is made possible by increases in cardiac contractility (Crandall & Wilson, 2014). Given these cardiovascular effects, particularly when paired with the profuse haematological and vascular effects during and following heatstroke, it is not surprising that a single episode of heatstroke increases the risk of all-cause mortality by 40% (Wallace et al., 2007). More specifically, patients with EHS display a higher incidence of cardiovascular disease, myocardial infarction, ischaemic heart disease, and acute ischaemic stroke as early as 14 years following the initial heatstroke event (Wallace et al., 2007; Wang et al., 2019).

Evidence of potential myocardial injury during heatstroke was preliminarily identified in a case report published in 2006 of a dog presenting with suspected EHS and elevations in cardiac troponin I (Mellor et al., 2006), which increases when myocardial cells become damaged. These findings were corroborated in a subsequent case report in humans (Whiticar et al., 2008), including one case report of evidence of cardiomyopathy following heatstroke (Chen et al., 2012). Indeed, EHS has recently been shown to provoke myocardial metabolic alterations consistent with the presence of myocardial injury that persists for up to 2 weeks following heatstroke (Laitano et al., 2020) and marked immunosuppression that lasts at least 1 month but likely longer (Murray et al., 2021). Interestingly, these observations were specific to female mice (Laitano et al., 2020; Murray et al., 2021), which was attributed to greater heat tolerance resulting in a higher net heat load than male mice before EHS occurred. Given the consistency of cardiac ramifications during and after heatstroke, a cardiac-specific biomarker likely has utility. To this end, there is a growing body of evidence supporting cardiac troponin I as a sensitive biomarker of myocardial injury during and following heatstroke. For instance, and consistent with the aforementioned case reports (Mellor et al., 2006; Whiticar et al., 2008), graded increases in circulating cardiac troponin I were observed with increased severity of heatstroke 24 h following heatstroke in a rodent model of CHS (Quinn et al., 2014). Moreover,

a point-of-care assessment of cardiac troponin I predicted heatstroke severity ($r^2 = 0.83$) in a rodent model of CHS (Audet et al., 2015). Finally, circulating cardiac troponin I may have longer-term prognostic utility. For example, higher levels of cardiac troponin I are associated with a higher death rate following emergency department admission during heatwaves (Hausfater et al., 2010). Particularly relevant for EHS is that cardiac troponin I can be elevated during prolonged exercise (e.g., a marathon) (Shave et al., 2007). Thus, the specificity of cardiac troponin I to heatstroke-induced cardiac injury remains to be determined.

3.2 | Blood vessels

Clinical and experimental studies suggest that changes in vascular endothelium and coagulation functions are characteristic features of heatstroke (Bouchama, Hammami et al., 1996; Roberts et al., 2008; Shieh et al., 1995; Sohal et al., 1968). Moreover, functional alterations in these systems are also strongly associated with an increased risk of cardiovascular morbidity and mortality (Hadi et al., 2005; Lowe & Rumley, 2014). Thus, it is not surprising that the risk of cardiovascular disease is elevated following a single heatstroke occurrence (Wang et al., 2019). Postmortem studies in heatstroke patients revealed extensive endothelial damage and haemorrhagic thrombosis in most body organs (Chao et al., 1981; Malamud et al., 1946a; Sohal et al., 1968). These findings were replicated and expanded in non-human primate models of heatstroke, demonstrating that endothelial cell, inflammation and coagulation activation precede endothelial damage, thrombosis and bleeding, which culminates in death (Roberts et al., 2008; Sohal et al., 1968). Hence, the capacity to detect early changes in endothelial function may have clinical utility (Bouchama et al., 2005; Roberts et al., 2008). Several studies have investigated potential circulating biomarkers of endothelial dysfunction, including their link with heatstroke-induced organ damage, severity and outcomes (Alzeer et al., 1999; Bouchama, Hammami et al., 1996; Hammami, Bouchama & Al Sedairy et al., 1998; Roberts et al., 2008; Shieh et al., 1995; Sohal et al., 1968). These include mediators of vascular tone, haemostasis, leukocyte trafficking, and endothelium barrier integrity.

3.3 | Vascular tone

A series of elegant experimental studies by Hall et al., demonstrated a central role of nitric oxide in the progression from heat stress to heatstroke (Hall et al., 1994, 2001). Indeed, hyperthermia stimulates nitric oxide synthesis and reactive oxygen species production in the rodent splanchnic circulation, resulting in intestinal barrier dysfunction, cardiovascular collapse, and heatstroke. An imbalance consisting of a marked release of endothelin-1 concomitant to reduced nitric oxide production in the liver sinusoid was also demonstrated in rodents (Zhang et al., 2018). High levels of nitric oxide, assessed by nitrate and nitrite, were identified in a small cohort of heatstroke

patients that correlated with the severity of heatstroke on admission and mortality (Alzeer et al., 1999). Endothelin-1, another endothelium-derived factor with dual properties including vasoconstrictive and vasodilating effects at higher and lower concentrations, was also detected in the splanchnic circulation of heatstroke patients although its clinical significance remains unknown (Bouchama, Hammami et al., 1996; Ohlstein et al., 1990). Taken together, these findings suggest that mediators of vascular tone are implicated in the pathogenesis of heatstroke and may warrant further evaluation as biomarkers of disease activity and survival.

3.4 | Haemostasis

The coagulopathy of heatstroke progresses from the initial activation of coagulation and fibrinolysis followed by late fibrinolytic inhibition, resulting in a hypercoagulable state. Clinically, the patients with heatstroke display low platelet count and fibrinogen levels and increasing coagulation times and D-dimer levels (al-Mashhadani et al., 1994; Bouchama, Bridey et al., 1996). However, except for D-dimer, these conventional coagulation parameters are not likely sensitive enough or specific to be used as biomarkers of organ injury in heatstroke. When assessed in research settings using sensitive assays such as thrombin-antithrombin and plasmin-antiplasmin complexes, an early rise in thrombin and plasmin can be detected, with concomitant decrease of natural anticoagulant proteins (Protein C, S and antithrombin III) before the alteration of coagulation tests used in clinical practice. Thrombin ($r = 0.635$, $P = 0.01$), plasmin ($r = 0.577$, $P = 0.02$) and D-dimer ($r = 0.76$, $P = 0.003$) correlated significantly with the magnitude of hyperthermia in CHS (Bouchama, Bridey et al., 1996), whereas the levels of protein C and antithrombin III discriminate outcome severity in heatstroke, with lower concentrations of these biomarkers being associated with poorer outcomes (i.e., requiring therapeutic blood product intervention) (al-Mashhadani et al., 1994). Hence, further research is needed to assess the value of pro- and anticoagulant molecules as prognostic biomarkers of heatstroke-provoked organ injury/recovery using highly sensitive assays.

Von Willebrand factor antigen (vWfAg) and soluble thrombomodulin are likely the most studied biomarkers of endothelial cell injury activation or injury in heatstroke (Bouchama et al., 2005; Bouchama, Hammami et al., 1996; Huisse et al., 2008; Proctor et al., 2020; Shieh et al., 1995; Tong et al., 2014; Zhang et al., 2018). vWfAg is synthesized and stored in endothelial cells and megakaryocytes, and mediates platelet aggregation and adhesion at vascular injury sites and endothelial damage (Lip & Blann, 1997). In an experimental model of heatstroke, immunostaining analysis revealed higher expression of vWfAg in most organs of the body, suggesting that the vascular endothelium activation is systemic (Roberts et al., 2008). However, no association between organ injury and survival was demonstrated in humans with heatstroke ($r^2 = 0.37$, $P > 0.05$) (Bouchama, Hammami et al., 1996).

Thrombomodulin is a transmembrane proteoglycan located on the surface of the vascular endothelium that functions as an anticoagulant (Martin et al., 2013). There is also a soluble form of thrombomodulin released from membrane-bound thrombomodulin that can be released into the circulation through proteolytic cleavage and is considered a marker of endothelial cell damage (Martin et al., 2013). Thrombomodulin activates coagulation and inhibits tissue plasminogen activator-induced fibrinolysis following binding to thrombin and thrombin-activatable fibrinolysis inhibitor, respectively (Bajzar, 2000; Esmon & Owen, 1981). Recombinant thrombomodulin given to heat-stressed rodents prevented heatstroke by inhibition of HMGB1, a damage-associated molecular pattern, indicating that thrombomodulin also has an anti-inflammatory function in heatstroke (Hagiwara et al., 2010). Several clinical and experimental studies demonstrated an increased plasma soluble thrombomodulin concentration in heatstroke associated with severity and survival in some studies, but this was not consistently observed (Bouchama et al., 2005; Huisse et al., 2008; Shieh et al., 1995; Tong et al., 2014; Zhang et al., 2018). Hence, further studies are needed to clarify the prognostic utility of soluble thrombomodulin in heatstroke.

3.5 | Leukocyte trafficking

A central step in the complex crosstalk between inflammation, coagulation, and endothelial cells in heatstroke is the adherence of leukocytes and platelets to the endothelium via cell adhesion molecules. *Ex vivo* study of leukocytes obtained from CHS patients showed that this process is mediated by the up-regulation of $\beta 2$ integrin and down-regulation of L-selectin (Huisse et al., 2008). Changes in lymphocyte $\beta 2$ integrin expression were also demonstrated *in vivo* in both CHS and EHS (Hammami, Bouchama, Shail et al., 1998; Lu, Lin et al., 2004). Notably, soluble forms of adhesion molecules are released into the circulation and correlate with the concentration of molecules expressed on the cells and various disease activity. Thus, soluble adhesion molecules may constitute more practical biomarkers in clinical settings (Gearing & Newman, 1993). In CHS patients, elevated plasma concentrations of soluble intercellular adhesion molecule 1, E-selectin, and L-selectin adhesion molecules have been documented (Bouchama, Hammami et al., 1996; Hammami, Bouchama & Al Sedairy et al., 1998). L-Selectin correlated significantly with E-selectin ($r = 0.68, P = 0.0002$) and the level of consciousness ($r = -0.45, P = 0.03$) in heatstroke patients (Hammami, Bouchama & Al Sedairy et al., 1998). Compared with other soluble adhesion molecules, E-selectin is probably the most specific among these biomarkers of endothelial cell dysfunction and hence merits further study (Gearing & Newman, 1993).

3.6 | Endothelial barrier integrity

Damage to the endothelium resulting in loss of its barrier function and vascular leaks was demonstrated in human and non-human primate

models of heatstroke (Roberts et al., 2008; Sohal et al., 1968). The luminal surface of the endothelium is covered by the glycocalyx, a network of membrane-bound proteoglycans and glycoproteins that binds and incorporates soluble molecules derived from the plasma and endothelium. Heatstroke disrupts the endothelial glycocalyx as assessed by plasma levels of syndecan-1 and hyaluronan in laboratory animals (Kobayashi et al., 2018; Umemura et al., 2018; Zhang et al., 2018). Syndecan-1 is the principal component of the endothelial glycocalyx, and hyaluronan maintains vascular integrity through endothelial glycocalyx modulation (Lennon & Singleton, 2011). Other biomarkers of endothelial cell activation and loss of integrity include the detachment of endothelial cells and their release into the peripheral circulation. An ~ 2.5 -fold increase in circulating endothelial cells in an experimental heatstroke rat model has been demonstrated, suggesting it can be useful as a marker of endothelial injury (Tong et al., 2014). Overall, biomarkers of vascular barrier function in heatstroke were primarily established experimentally and deserve further translational studies

4 | BRAIN

Heatstroke encephalopathy is a universal primary manifestation of heatstroke (Leon & Bouchama, 2011). Heatstroke-induced encephalopathy is characteristically an early event in the natural course of this condition, preceding the dysfunction of other organ systems. Further, heatstroke encephalopathy is associated with long-term cognitive and motor disability in heatstroke survivors (Argaud et al., 2007; Dematte et al., 1998; Yang et al., 2017). These findings suggest that the brain is extraordinary vulnerable to heatstroke. The neurological signs and symptoms are not specific and include altered mental status, delirium and coma (Bouchama et al., 2022; Casa et al., 2012; Leon & Bouchama, 2011). There are no lateralizing signs on physical examination and brain imaging at presentation is normal with no evidence of structural damage (Leon & Bouchama, 2011). In contrast, follow-up brain imaging performed months after heatstroke in patients with persistent central nervous system dysfunction can reveal damage to the cerebellum, prefrontal cortex, or hippocampus (Leon & Bouchama, 2011). Hence, this indicates that early brain injury leading to delayed brain damage may not be readily detectable with standard diagnostic approaches. In this context, biomarkers that would aid in the early detection of injury and predict delayed brain damage would be crucial. Notably, S100 calcium-binding protein β (S100 β) (Chun et al., 2019; Li et al., 2020) and neuron-specific enolase (NSE) (Li et al., 2020) have been recently proposed as potential biomarkers of heatstroke encephalopathy.

S100 β , a glia-specific protein expressed by astrocytes and Schwann cells, functions as a neurotrophic and neuronal survival factor in the developing brain. S100 β can be released into the circulation from injured cells (Van Eldik et al., 1986). Accordingly, S100 β levels in serum and cerebrospinal fluid have been used as biomarkers of brain and blood-brain barrier injury in many conditions, including traumatic brain injury, subarachnoid haemorrhage and stroke

(Persson et al., 1987). Exercise in the heat increases circulating S100 β indicating that heat stress may alter blood–brain permeability (Watson et al., 2005). However, additional work refutes these findings during exercise in the heat (Cheuvront et al., 2008) and passive heat stress (Shepley et al., 2021) models in humans. Nonetheless, two recent studies in heatstroke patients demonstrated elevated S100 β protein in the serum and cerebrospinal fluid of heatstroke patients (Chun et al., 2019; Li et al., 2020), with one study finding that the heatstroke patients with poor prognosis (defined as not being able to live without assistance at hospital discharge) having serum S100 β concentrations ~5 times higher than in heatstroke patients with a good prognosis (Chun et al., 2019). NSE is a dimeric isoform of the glycolytic enzyme enolase expressed mainly in neurons (Kaiser et al., 1989). NSE is upregulated in injured neuronal cells and can spill over into the circulation from injured or dying cells (Persson et al., 1987). Elevated NSE levels in the serum and cerebrospinal fluid have been reported in various brain pathologies, such as head trauma and vascular stroke (Persson et al., 1987). In heatstroke patients, NSE levels strongly correlated with neurological outcomes for up to 7 days post-heatstroke (all days $r \geq -0.624$, all days $P < 0.05$), suggesting that NSE could be helpful as a diagnostic and prognostic biomarker of heatstroke encephalopathy (Li et al., 2020). Overall, these observations indicate that S100 β and NSE are promising biomarkers for early detection or follow-up of heatstroke-associated brain injury and thus merit larger prospective studies to determine their sensitivity and specificity.

5 | KIDNEYS

Kidney damage is well recognized as one of the hallmarks of multiorgan injury caused by heatstroke (King et al., 2015; Leon et al., 2006b; Roberts et al., 2008). Indeed, patients diagnosed with heatstroke often suffer from acute kidney injury (e.g., Gauss and Meyer, 1917; Kew et al., 1967; Schrier et al., 1970), characterized by elevations in serum creatinine, which is indicative of reductions in glomerular filtration rate, measured as part of the basic clinical bloodwork (Ward et al., 2020). A recent analysis of 187 EHS cases in southern China identified that ~44% of patients developed acute kidney injury and that ~27% of the patients with acute kidney injury died within 90 days (Wu et al., 2021). Thus, the management of acute kidney injury following the diagnosis of heatstroke is an essential consideration. It is important to note, however, that acute kidney injury is not ubiquitous such that other observations indicate that in most heatstroke cases, elevations in serum creatinine (Ward et al., 2020) and other markers of renal dysfunction (Kew et al., 1967; Schrier et al., 1970) subsided as soon as 16 days following heatstroke.

Data in animal models of heatstroke indicate that kidney injury is contributed to by vascular congestion, haemorrhage and thrombi distributed throughout renal tissues (King et al., 2015; Leon et al., 2006b; Roberts et al., 2008). This likely occurs secondarily to the heatstroke-induced systemic inflammation and coagulation activation that catalyse haemorrhagic and thrombotic events within the kidneys

(and other organs), similar to sepsis (Ma et al., 2019). Recent findings support that the aetiology of kidney injury during heatstroke also includes injury to the renal tubules (Lin & Zhang, 2019). This study additionally identified serum neutrophil gelatinase-associated lipocalin (NGAL) as a potential early biomarker of kidney injury secondary to EHS, such that elevations in serum NGAL were almost 10 times higher post-EHS compared to a non-heated control group (Lin & Zhang, 2019). Such observations are consistent with findings in humans undertaking exercise in the heat (Schlader et al., 2017). While NGAL, which functions as a bacteriostatic agent secondary to tissue injury, is primarily expressed in renal tissues, it is also expressed in liver and heart tissues, suggesting that elevations in circulating NGAL may not be specific to the kidneys (Chapman, Johnson, Parker et al., 2020). This issue may be overcome by examining urinary NGAL or other markers of urinary tubular injury, such as kidney injury molecule-1, insulin-like growth factor binding protein-7, and/or tissue inhibitor metalloproteinase-2 (Chapman, Johnson, Parker et al., 2020). Indeed, urinary NGAL is elevated in dogs suffering from heatstroke (Segev et al., 2015). Whether this observation translates to humans and whether urinary markers of kidney injury can be used as biomarkers of kidney injury during or following heatstroke are unknown and warrant further study. It is also worth highlighting that obtaining urine samples during or following heatstroke may be challenging because urine production is reduced by elevations in body temperature and hypovolaemia (Chapman, Johnson, Parker et al., 2020). Thus, urinary kidney injury biomarkers may only be available upon urinary catheterization. Moreover, the specificity of plasma and urinary NGAL in heatstroke should be examined, as they are both temporarily elevated during heat stress in humans without heatstroke and these increases resolve within 24 h (Chapman, Johnson, Vargas et al., 2020; Schlader et al., 2017).

6 | GASTROINTESTINAL SYSTEM

6.1 | Intestines

The systemic immune response to heatstroke has been partially attributed to the development of intestinal cellular dysfunction, making the gastrointestinal tract leaky, and allowing for the translocation of endotoxin into the systemic circulation (Hall et al., 1999; Novosad et al., 2013). The mechanisms by which the gastrointestinal tract may contribute to heatstroke aetiology are outside of the scope of this review. Here, however, we consider the consequences of heatstroke on biomarkers of intestinal injury and recovery. To this end, it is notable that diarrhoea is consistently observed in heatstroke (Hart et al., 1980), which typically subsides with cooling (Bouchama et al., 2022; Leon & Bouchama, 2011). Data from preclinical models of CHS show severe intestinal injury in the duodenum, jejunum and ileum (Novosad et al., 2013) that is characterized by cellular apoptosis (Roberts et al., 2008) and interstitial oedema (Miyamoto et al., 2021). Whether these findings are consistent in the clinical setting remains unknown (Bouchama et al., 2022). However, circulating intestinal fatty acid-binding protein 2 (I-FABP) is a protein found in enterocytes of

the small intestinal epithelium that when found in the circulation is often interpreted as evidence of increased intestinal permeability and/or injury (Wells et al., 2017). I-FABP is elevated in patients with heatstroke, with peak responses occurring 3 days post-heatstroke and reduced with treatment (Zhang et al., 2015), suggesting that I-FABP may be a biomarker of gastrointestinal injury associated with heatstroke. The experimental demonstration of an increased I-FABP gene expression in intestinal mucosa up to 7 h post-passive heat exposure (Miyamoto et al., 2021) and elevations in plasma I-FABP protein levels for 24 h after EHS (King et al., 2015), all of which were associated with intestinal histopathological damage, lend support to this interpretation. Given these I-FABP observations, other biomarkers of increased intestinal permeability (e.g., urinary lactulose/rhamnose ratio) may also be considered as intestinal injury biomarkers following heatstroke. However, data from recent meta-analyses indicate that hyperthermia (Pires et al., 2017) and exercise (Chantler et al., 2021), in the absence of heatstroke, consistently increase markers of intestinal permeability, including I-FABP, casting doubt on the value of I-FABP and other markers of intestinal permeability for the specificity of heatstroke-induced intestinal injury or recovery.

6.2 | Liver

Liver dysfunction and injury are relatively common following heatstroke (Leon & Bouchama, 2011). Indeed, increases in circulating alanine aminotransferase (ALT) and aspartate aminotransferase (AST), standard clinical tests of liver function, are ubiquitous in heatstroke patients (Ward et al., 2020). Moreover, data from a mouse model of EHS (King et al., 2015) and a baboon model of CHS (Roberts et al., 2008) indicate that heatstroke induces vascular congestion and haemorrhagic thrombosis in liver tissues, while postmortem clinical findings provide support for liver tissue ischaemia and necrosis (Malamud et al., 1946b). Notably, these pathological findings also identified that the extent of liver damage was greater when survival following heatstroke exceeded 30 h (Malamud et al., 1946b), suggesting that the aetiology of liver damage initiated by heat cytotoxicity is likely amplified (or worsened) by ischaemia-reperfusion injury and an exacerbated systemic inflammatory response. This interpretation is supported by the timing of changes in ALT and AST following heatstroke, with peak concentrations being observed 2–3 days post-heatstroke (Ji et al., 2021; Ward et al., 2020). There are no formally identified biomarkers, as defined herein, quantifying the extent of liver damage or recovery post-heatstroke. Nevertheless, it is important to recognize that recent evidence supports that increased circulating clinical markers of hepatic dysfunction, including ALT and AST, are associated with an increased likelihood of EHS mortality (Ji et al., 2021; Li et al., 2022). However, it is unknown whether this is due to hepatic causes per se or simply reflects the more severe clinical condition of these patients (Ji et al., 2021). This conclusion is supported by evidence that heatstroke elicits liver injury via IL-1 β mechanisms (Geng et al., 2015) and that pretreatment with specific antibodies that inhibit circulating HMGB1 protein, which is at least

partially implicated in the inflammatory response during EHS (Tong et al., 2011a), alleviates CHS-provoked liver pathology (Tong et al., 2013).

7 | LUNGS

Acute lung injury (ALI) is common and is associated with excessive systemic inflammation and coagulation activation in heatstroke (El-Kassimi et al., 1986). However, whether ALI is a direct consequence of excessive heat or a secondary consequence of systemic inflammation and coagulation activation remains unclear. Several studies in rats used ambient heating to induce CHS and the associated multiorgan failure syndrome that is commonly observed in clinical patients (Hsi-Hsing et al., 2010; Yang et al., 2009). The specific experimental manipulation included exposure of the rats to a thermally stressful environment of 42°C and 80% relative humidity, in which the ambient air had a heat content of approximately 153 kJ/kg. With 146 kJ/kg of heat content at the respiratory surface (assuming 100% relative humidity and a normal body temperature of 37.9°C), this experimental design not only effectively prevents any heat dissipation from the experimental subjects but may also result in *net heat absorption* from the environment into the respiratory tissues. This pattern of preferentially heating the respiratory tract, with the rest of the body accumulating heat due to perfusion of heated blood from the respiratory tract, is likely consistent with the instances of CHS but differs from EHS. Particularly relevant in animal models of heatstroke is that the mechanism of selective brain cooling relies on cooling blood in the upper respiratory tract. Thus, it can be speculated that this arrangement may lead to faster central nervous system heating and signs of heatstroke than in typical clinical situations in which the primary source of heat is other tissues in the body. In contrast, studies of CHS using a lower ambient temperature failed to show any histological evidence of lung injury (Leon et al., 2006a), supporting the contention that early ALI in the development of heatstroke may be a function of selective injury to the lung during conditions favouring heat uptake through that organ. In contrast, ALI developing later in the disease process may be secondary to systemic inflammatory processes and not specific to heat injury. As a result, biomarkers specific to ALI are probably relevant only in this context.

8 | SKELETAL MUSCLE

As the principal source of excess heat, skeletal muscle plays a central role in EHS. Biomarkers of skeletal muscle injury, such as myoglobin and creatine kinase, are prominent in cases of EHS, with reported increases of 2 and 4.5 times, respectively, above normal limits in a retrospective study of cases of EHS (Laitano et al., 2021). However, cases of exertional rhabdomyolysis without associated heatstroke can demonstrate similar or higher serum concentrations of these biomarkers (Lippi & Ceriotti, 2019), making these biomarkers specific to muscle injury but not specific for the underlying cause of muscle injury. Similarly, serum cytokine concentrations lack specificity

regarding skeletal muscle injury in heatstroke since it has been well-established that skeletal muscle has considerable secretory activity during exercise (Pedersen & Febbraio, 2008). Given that skeletal muscle activity is inherently associated with increased skeletal muscle temperature due to the metabolic activity needed to support skeletal muscle contraction, it is impossible to separate the effects of exercise from the thermal consequences in this tissue. Furthermore, the same cytokines released by active skeletal muscles can also be released by other tissues, making it less certain that the cascade of cytokines commonly identified in cases of heatstroke is of muscular origin. Thus, these biomarkers may be more specific to exertion (and perhaps the heating of the skeletal muscle tissue per se) than whole-body hyperthermia secondary to an inability to dissipate the heat produced by the skeletal muscle. That said, an understanding of the extent of skeletal muscle injury may have important implications regarding the management of kidney injury during or following heatstroke.

Skeletal muscle also can suffer injury and presumably release biomarkers of that injury during CHS. A specific role for mitochondria in heat-induced muscle injury is suggested by the fact that lesions induced by passive heating are more prominent in muscles with higher oxidative capacity (Sharma et al., 2021). Increased temperature has been shown to increase mitochondrial leak respiration in skeletal muscle in rats (Jarmuszkiewicz et al., 2015), dogs (Davis & Barrett, 2021), horses (Davis et al., 2020) and humans (Fiorenza et al., 2019). Increased leak respiration will decrease the efficiency of ATP synthesis and as a result increased heat production through oxidative metabolism is necessary to maintain the desired rate of ATP synthesis. If heat dissipation from the tissue is not increased, a cycle will develop resulting in skeletal muscle overheating and metabolic failure. Additional possible consequences of skeletal muscle heating that may result in damage are the production of reactive oxygen species, which is slightly increased during phosphorylating respiration in hyperthermic rat muscle (Jarmuszkiewicz et al., 2015). Finally, tissue hyperthermia may cause failure of cytoplasmic calcium regulation due to heat-induced dysfunction of the sarcoplasmic calcium-mediated calcium channel (aka, ryanodine receptor RyR1) (Capacchione & Muldoon, 2009), contributing to a syndrome of malignant hyperthermia. Due to these mechanisms, the presence of biomarkers of skeletal muscle injury in the absence of a history of recent exertion could support the passive accumulation of heat – and by extension, systemic heat stress or heatstroke – as an important differential diagnosis.

9 | SUMMARY

Based on observational evidence indicating that a single episode of heatstroke could have harmful acute and chronic long-term health outcomes across numerous organ systems, this narrative review aimed to identify biomarkers of heatstroke-induced organ injury and repair that could be used to guide further development and utilization. This includes as a potential aid to initiate specific organ-protective strategies and prognostication (Figure 1). To this end, we

have identified numerous biomarkers related to many aspects of generalized heatstroke-induced cellular injury and tissue damage, and heatstroke provoked cardiovascular, renal, cerebral, intestinal and skeletal muscle injury (Figure 2). The literature supporting biomarkers related to vascular endothelium dysfunction following heatstroke appears to be the most extensive, which may be related to the uniquely multifunctional nature of the endothelium. Moreover, no novel biomarkers were identified for liver or lung injury, which may be because it remains unclear whether injury to these organs is specific to heatstroke or generalized systemic inflammation. In general, there was evidence that the identified biomarkers were specific to acute organ injury. However, the kinetic profile of these biomarkers post-heatstroke, and their reproducibility and specificity to predict organ recovery or failure and long-term outcomes remain relatively unexplored. Collectively, before the translation of the identified biomarkers into clinical practice, additional preclinical and clinical prospective studies are required to better understand the clinical utility of these biomarkers. These studies should likely focus on the relations between the identified biomarkers and how they relate to long-term post-heatstroke health outcomes.

COMPETING INTERESTS

Z.J.S. has received consultant fees from Otsuka Holdings Co., Ltd. Otherwise, no other potential conflicts of interest, financial or otherwise, are declared by the authors.

AUTHOR CONTRIBUTIONS

All authors conceived and designed the manuscript, prepared the figures, drafted the manuscript, and edited and revised the manuscript. All authors have read and approved the final version of this manuscript and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

ORCID

Zachary J. Schlader  <https://orcid.org/0000-0003-3590-3958>

Abderrezak Bouchama  <https://orcid.org/0000-0003-3089-5936>

REFERENCES

- al-Mashhadani, S. A., Gader, A. G., al Harthi, S. S., Kangav, D., Shaheen, F. A., & Bogus, F. (1994). The coagulopathy of heat stroke: Alterations in coagulation and fibrinolysis in heat stroke patients during the pilgrimage (Haj) to Makkah. *Blood Coagulation & Fibrinolysis*, 5, 731–736.
- Alele, F. O., Malau-Aduli, B. S., Malau-Aduli, A. E., & Crowe, M. J. (2021). Haematological, biochemical and hormonal biomarkers of heat intolerance in military personnel. *Biology*, 10(10), 1068.
- Alzeer, A. H., Al-Arifi, A., Warsy, A. S., Ansari, Z., Zhang, H., & Vincent, J. L. (1999). Nitric oxide production is enhanced in patients with heat stroke. *Intensive Care Medicine*, 25, 58–62.
- Argaud, L., Ferry, T., Le, Q.-H., Marfisi, A., Ciorba, D., Achache, P., Ducluzeau, R., & Robert, D. (2007). Short- and long-term outcomes of heatstroke following the 2003 heat wave in Lyon, France. *Archives of Internal Medicine*, 167(20), 2177–2183.

- Armstrong, L. E., Casa, D. J., Millard-Stafford, M., Moran, D. S., Pyne, S. W., & Roberts, W. O. (2007). Exertional heat illness during training and competition. *Medicine & Science in Sports & Exercise*, *39*, 556–572.
- Audet, G. N., Quinn, C. M., & Leon, L. R. (2015). Point-of-care cardiac troponin test accurately predicts heat stroke severity in rats. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, *309*(10), R1264–R1272.
- Bajzar, L. (2000). Thrombin activatable fibrinolysis inhibitor and an anti-fibrinolytic pathway. *Arteriosclerosis, Thrombosis, and Vascular Biology*, *20*(12), 2511–2518.
- Bouchama, A., Abuyassin, B., Lehe, C., Laitano, O., Jay, O., O'Connor, F. G., & Leon, L. R. (2022). Classic and exertional heatstroke. *Nature Reviews. Disease Primers*, *8*(1), 8.
- Bouchama, A., Bridey, F., Hammami, M. M., Lacombe, C., al-Shail, E., al-Ohali, Y., Combe, F., al-Sedairy, S., & de Prost, D. (1996). Activation of coagulation and fibrinolysis in heatstroke. *Thromb Haemost*, *76*, 909–915.
- Bouchama, A., Hammami, M. M., Haq, A., Jackson, J., & al-Sedairy, S. (1996). Evidence for endothelial cell activation/injury in heatstroke. *Critical Care Medicine*, *24*(7), 1173–1178.
- Bouchama, A., Parhar, R. S., el-Yazigi, A., Sheth, K., & al-Sedairy, S. (1991). Endotoxemia and release of tumor necrosis factor and interleukin 1 alpha in acute heatstroke. *Journal of Applied Physiology*, *70*(6), 2640–2644.
- Bouchama, A., Roberts, G., Al Mohanna, F., El-Sayed, R., Lach, B., Chollet-Martin, S., Ollivier, V., Al Baradei, R., Loualich, A., Nakeeb, S., Eldali, A., & de Prost, D. (2005). Inflammatory, hemostatic, and clinical changes in a baboon experimental model for heatstroke. *Journal of Applied Physiology*, *98*(2), 697–705.
- Bruchim, Y., Ginsburg, I., Segev, G., Mreizat, A., Avital, Y., Aroch, I., & Horowitz, M. (2017). Serum histones as biomarkers of the severity of heatstroke in dogs. *Cell Stress & Chaperones*, *22*, 903–910.
- Bruchim, Y., Segev, G., Kelmer, E., Codner, C., Marisat, A., & Horowitz, M. (2015). Hospitalized dogs recovery from naturally occurring heatstroke; does serum heat shock protein 72 can provide prognostic biomarker? *Cell Stress and Chaperones*, *21*(1), 123–130.
- Capacchione, J. F., & Muldoon, S. M. (2009). The relationship between exertional heat illness, exertional rhabdomyolysis, and malignant hyperthermia. *Anesthesia & Analgesia*, *109*, 1065–1069.
- Casa, D. J., Armstrong, L. E., Kenny, G. P., O'Connor, F. G., & Huggins, R. A. (2012). Exertional heat stroke: New concepts regarding cause and care. *Current Sports Medicine Reports*, *11*(3), 115–123.
- Casa, D. J., DeMartini, J. K., Bergeron, M. F., Csillan, D., Eichner, E. R., Lopez, R. M., Ferrara, M. S., Miller, K. C., O'Connor, F., & Sawka, M. N. (2015). National Athletic Trainers' Association position statement: Exertional heat illnesses. *Journal of Athletic Training*, *50*(9), 986–1000.
- Chantler, S., Griffiths, A., Matu, J., Davison, G., Jones, B., & Deighton, K. (2021). The effects of exercise on indirect markers of gut damage and permeability: A systematic review and meta-analysis. *Sports Medicine*, *51*(1), 113–124.
- Chao, T. C., Sinniah, R., & Pakiam, J. E. (1981). Acute heat stroke deaths. *Pathology*, *13*(1), 145–156.
- Chapman, C. L., Johnson, B. D., Parker, M. D., Hostler, D., Pryor, R. R., & Schlader, Z. J. (2020). Kidney physiology and pathophysiology during heat stress and the modification by exercise, dehydration, heat acclimation and aging. *Temperature*, In Press.
- Chapman, C. L., Johnson, B. D., Vargas, N. T., Hostler, D., Parker, M. D., & Schlader, Z. J. (2020). Both hyperthermia and dehydration during physical work in the heat contribute to the risk of acute kidney injury. *Journal of Applied Physiology*, *128*(4), 715–728.
- Chen, G. Y., & Nuñez, G. (2010). Sterile inflammation: Sensing and reacting to damage. *Nature Reviews. Immunology*, *10*(12), 826–837.
- Chen, W.-T., Lin, C.-H., Hsieh, M.-H., Huang, C.-Y., & Yeh, J.-S. (2012). Stress-induced cardiomyopathy caused by heat stroke. *Annals of Emergency Medicine*, *60*(1), 63–66.
- Cheuvront, S. N., Chivever, T. D., Ely, B. R., Kenefick, R. W., Goodman, D. A., McClung, J. P., & Sawka, M. N. (2008). Serum S-100 β response to exercise-heat strain before and after acclimation. *Medicine & Science in Sports & Exercise*, *40*(8), 1477–1482.
- Chun, J.-K., Choi, S., Kim, H.-H., Yang, H. W., & Kim, C. S. (2019). Predictors of poor prognosis in patients with heat stroke. *Clinical and Experimental Emergency Medicine*, *6*(4), 345.
- Cohen, I., Rider, P., Carmi, Y., Braiman, A., Dotan, S., White, M. R., Voronov, E., Martin, M. U., Dinarello, C. A., & Apte, R. N. (2010). Differential release of chromatin-bound IL-1 α discriminates between necrotic and apoptotic cell death by the ability to induce sterile inflammation. *Proceedings of the National Academy of Sciences, USA*, *107*(6), 2574–2579.
- Crandall, C. G., & Wilson, T. E. (2014). Human cardiovascular responses to passive heat stress. *Comparative Physiology*, *5*, 17–43.
- Davis, M. S., & Barrett, M. R. (2021). Effect of conditioning and physiological hyperthermia on canine skeletal muscle mitochondrial oxygen consumption. *Journal of Applied Physiology*, *130*(5), 1317–1325.
- Davis, M. S., Fulton, M. R., & Popken, A. A. (2020). Effect of hyperthermia and acidosis on equine skeletal muscle mitochondrial oxygen consumption. *Comparative Exercise Physiology*, *17*(2), 171–179.
- Dehbi, M., Baturcam, E., Eldali, A., Ahmed, M., Kwaasi, A., Chishti, M. A., & Bouchama, A. (2010). Hsp-72, a candidate prognostic indicator of heatstroke. *Cell Stress & Chaperones*, *15*(5), 593–603.
- Dehbi, M., Uzzaman, T., Baturcam, E., Eldali, A., Ventura, W., & Bouchama, A. (2012). Toll-like receptor 4 and high-mobility group box 1 are critical mediators of tissue injury and survival in a mouse model for heatstroke. *PLoS One*, *7*(9), e44100.
- Demartini, J. K., Casa, D. J., Stearns, R., Belval, L., Crago, A., Davis, R., & Jardine, J. (2015). Effectiveness of cold water immersion in the treatment of exertional heat stroke at the Falmouth Road Race. *Medicine and Science in Sports and Exercise*, *47*(2), 240–245.
- Dematte, J. E., O'Mara, K., Buescher, J., Whitney, C. G., Forsythe, S., McNamee, T., Adiga, R. B., & Ndukwu, I. M. (1998). Near-fatal heat stroke during the 1995 heat wave in Chicago. *Annals of Internal Medicine*, *129*(3), 173–181.
- El-Kassimi, F. A., Al-Mashhadani, S., Abdullah, A. K., & Akhtar, J. (1986). Adult respiratory distress syndrome and disseminated intravascular coagulation complicating heat stroke. *Chest*, *90*(4), 571–574.
- Esmon, C. T., & Owen, W. G. (1981). Identification of an endothelial cell cofactor for thrombin-catalyzed activation of protein C. *Proceedings of the National Academy of Sciences*, *78*(4), 2249–2252.
- Fiorenza, M., Lemming, A. K., Marker, M., Eibye, K., Iaia, F. M., Bangsbo, J., & Hostrup, M. (2019). High-intensity exercise training enhances mitochondrial oxidative phosphorylation efficiency in a temperature-dependent manner in human skeletal muscle: Implications for exercise performance. *FASEB Journal*, *33*(8), 8976–8989.
- Gallucci, S., & Matzinger, P. (2001). Danger signals: SOS to the immune system. *Current Opinion in Immunology*, *13*(1), 114–119.
- Gauss, H., & Meyer, K. (1917). Heat stroke: Report of one hundred and fifty-eight cases from Cook County hospital, Chicago. *The American Journal of the Medical Sciences (1827-1924)*, *154*(4), 554.
- Gearing, A. J. H., & Newman, W. (1993). Circulating adhesion molecules in disease. *Immunology Today*, *14*(10), 506–512.
- Geng, Y., Ma, Q., Liu, Y.-N., Peng, N., Yuan, F.-F., Li, X.-G., Li, M., Wu, Y.-S., Li, B.-L., & Song, W.-B. (2015). Heatstroke induces liver injury via IL-1 β and HMGB1-induced pyroptosis. *Journal of Hepatology*, *63*(3), 622–633.
- Hadi, H. A., Carr, C. S., & Al Suwaidi, J. (2005). Endothelial dysfunction: Cardiovascular risk factors, therapy, and outcome. *Vascular Health and Risk Management*, *1*, 183.

- Hagiwara, S., Iwasaka, H., Goto, K., Ochi, Y., Mizunaga, S., Saikawa, T., & Noguchi, T. (2010). Recombinant thrombomodulin prevents heatstroke by inhibition of high-mobility group box 1 protein in sera of rats. *Shock*, 34(4), 402–406.
- Hall, D. M., Baumgardner, K. R., Oberley, T. D., & Gisolfi, C. V. (1999). Splanchnic tissues undergo hypoxic stress during whole body hyperthermia. *American Journal of Physiology*, 276, G1195–1203.
- Hall, D. M., Buettner, G. R., Matthes, R. D., & Gisolfi, C. V. (1994). Hyperthermia stimulates nitric oxide formation: Electron paramagnetic resonance detection of .NO-heme in blood. *Journal of Applied Physiology*, 77(2), 548–553.
- Hall, D. M., Buettner, G. R., Oberley, L. W., Xu, L., Matthes, R. D., & Gisolfi, C. V. (2001). Mechanisms of circulatory and intestinal barrier dysfunction during whole body hyperthermia. *American Journal of Physiology. Heart and Circulatory Physiology*, 280(2), H509–H521.
- Hammami, M. M., Bouchama, A., & Al Sedairy, S. (1998). Levels of soluble L-selectin and E-selectin in heatstroke and heatstress. *Chest*, 114(3), 949–950.
- Hammami, M. M., Bouchama, A., Shail, E., Aboul-Enein, H. Y., & Al-Sedairy, S. (1998). Lymphocyte subsets and adhesion molecules expression in heatstroke and heat stress. *Journal of Applied Physiology*, 84(5), 1615–1621.
- Hart, L., Egier, B., Shimizu, A., Tandan, P., & Sutton, J. (1980). Exertional heat stroke: The runner's nemesis. *Canadian Medical Association Journal*, 122, 1144.
- Hausfater, P., Doumenc, B., Chopin, S., Le Manach, Y., Santin, A., Dautheville, S., Patzak, A., Hericord, P., Mégarbane, B., & Andronikof, M. (2010). Elevation of cardiac troponin I during non-exertional heat-related illnesses in the context of a heatwave. *Critical Care*, 14(3), 1–9.
- Hsi-Hsing, Y., Ching-Ping, C., Juei-Tang, C., & Lin, M. T. (2010). Inhibition of acute lung inflammation and injury is a target of brain cooling after heatstroke injury. *Journal of Trauma*, 69, 805–812.
- Huisse, M. G., Pease, S., Hurtado-Nedelec, M., Arnaud, B., Malaquin, C., Wolff, M., Gougerot-Pocidallo, M. A., Kermarrec, N., Bezeaud, A., Guillin, M. C., Paoletti, X., & Chollet-Martin, S. (2008). Leukocyte activation: The link between inflammation and coagulation during heatstroke. A study of patients during the 2003 heat wave in Paris. *Critical Care Medicine*, 36(8), 2288–2295.
- IPCC (2021). *Climate change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change*. Cambridge University Press.
- Jarmuszkiwicz, W., Woyda-PloSzczyca, A., Koziel, A., Majerczak, J., & Zoladz, J. A. (2015). Temperature controls oxidative phosphorylation and reactive oxygen species production through uncoupling in rat skeletal muscle mitochondria. *Free Radical Biology & Medicine*, 83, 12–20.
- Ji, J., Gao, J., Wang, C., Ouyang, L., Liu, Z., & Liu, Z. (2021). Characteristics and outcome of exertional heatstroke patients complicated by acute hepatic injury: A cohort study. *Journal of Clinical and Translational Hepatology*, 9, 655.
- Kaiser, E., Kuzmits, R., Pregant, P., Burghuber, O., & Worofka, W. (1989). Clinical biochemistry of neuron specific enolase. *Clinica Chimica Acta*, 183(1), 13–31.
- Kew, M., Abrahams, C., Levin, N., Seftel, H., Rubenstein, A., & Bersohn, I. (1967). The effects of heatstroke on the function and structure of the kidney. *QJM*, 36, 277–300.
- King, M. A., Leon, L. R., Mustico, D. L., Haines, J. M., & Clanton, T. L. (2015). Biomarkers of multiorgan injury in a preclinical model of exertional heat stroke. *Journal of Applied Physiology*, 118(10), 1207–1220.
- Kobayashi, K., Mimuro, S., Sato, T., Kobayashi, A., Kawashima, S., Makino, H., Doi, M., Katoh, T., & Nakajima, Y. (2018). Dexmedetomidine preserves the endothelial glycocalyx and improves survival in a rat heatstroke model. *Journal of Anesthesia*, 32(6), 880–885.
- Kultz, D. (2005). Molecular and evolutionary basis of the cellular stress response. *Annual Review of Physiology*, 67(1), 225–257.
- Laitano, O., Garcia, C. K., Mattingly, A. J., Robinson, G. P., Murray, K. O., King, M. A., Ingram, B., Ramamoorthy, S., Leon, L. R., & Clanton, T. L. (2020). Delayed metabolic dysfunction in myocardium following exertional heat stroke in mice. *The Journal of Physiology*, 598(5), 967–985.
- Laitano, O., Oki, K., & Leon, L. R. (2021). The role of skeletal muscles in exertional heat stroke pathophysiology. *International Journal of Sports Medicine*, 42, 673–681.
- Lennon, F. E., & Singleton, P. A. (2011). Hyaluronan regulation of vascular integrity. *American Journal of Cardiovascular Disease*, 1, 200–213.
- Leon, L. R., Blaha, M. D., & DuBose, D. A. (2006a). Time course of cytokine, corticosterone, and tissue injury responses in mice during heat strain recovery. *Journal of Applied Physiology*, 100(4), 1400–1409.
- Leon, L. R., Blaha, M. D., & DuBose, D. A. (2006b). Time course of cytokine, corticosterone, and tissue injury responses in mice during heat strain recovery. *Journal of Applied Physiology*, 100(4), 1400–1409.
- Leon, L. R., & Bouchama, A. (2011). Heat stroke. *Comprehensive Physiology*, 5, 611–647.
- Li, B., Jia, Y. -R., Gao, W., Li, H. -P., Tao, W. -H., & Zhang, H. -X. (2020). The expression and clinical significance of neuron specific enolase and S100B protein in patients of severe heatstroke-induced brain injury. *Medical Journal of Chinese People's Liberation Army*, 45, 1282–1287.
- Li, C., Su, H. -B., Li, H., Li, X., Wang, H. -M., Song, Q., & Hu, J. -H. (2022). Severe acute liver injury in patients with exertional heat stroke associated with poor short-term prognosis. *World Journal of Emergency Medicine*, 13(2), 124.
- Li, Y., Liu, Z., Shi, X., Tong, H., & Su, L. (2021). Prognostic value of plasma exosomal levels of histone H3 protein in patients with heat stroke. *Experimental and Therapeutic Medicine*, 22(3), 922.
- Lin, Y., & Zhang, Y. (2019). Renoprotective effect of oral rehydration solution III in exertional heatstroke rats. *Renal Failure*, 41(1), 190–196.
- Lip, G. Y., & Blann, A. (1997). von Willebrand factor: A marker of endothelial dysfunction in vascular disorders? *Cardiovascular Research*, 34(2), 255–265.
- Lippi, G. S. F., & Ceriotti, F. (2019). Diagnostic biomarkers of muscle injury and exertional rhabdomyolysis. *Clinical Chemistry and Laboratory Medicine*, 57, 7.
- Lowe, G., & Rumley, A. (2014). The relevance of coagulation in cardiovascular disease: What do the biomarkers tell us? *Thrombosis and Haemostasis*, 112, 860–867.
- Lu, K. -C., Wang, J. -Y., Lin, S. -H., Chu, P., & Lin, Y. -F. (2004). Role of circulating cytokines and chemokines in exertional heatstroke. *Critical Care Medicine*, 32(2), 399–403.
- Lu, K. C., Lin, S. H., Chu, P., Tsai, W. S., & Lin, Y. F. (2004). Correlation of neutrophil phagocytosis and lymphocyte adhesion molecules in exertional heat stroke. *American Journal of the Medical Sciences*, 327(2), 68–72.
- Ma, S., Evans, R. G., Iguchi, N., Tare, M., Parkington, H. C., Bellomo, R., May, C. N., & Lankadeva, Y. R. (2019). Sepsis-induced acute kidney injury: A disease of the microcirculation. *Microcirculation*, 26(2), e12483.
- Malamud, N., Haymaker, W., & Custer, R. (1946a). Heatstroke: A clinicopathologic study of 125 fatal cases. *Military Surgery*, 99(5), 397–449.
- Malamud, N., Haymaker, W., & Custer, R. P. (1946b). Heat stroke. A clinicopathologic study of 125 fatal cases. *Military Surgeon*, 99(5), 397–449.
- Martin, F. A., Murphy, R. P., & Cummins, P. M. (2013). Thrombomodulin and the vascular endothelium: Insights into functional, regulatory, and therapeutic aspects. *American Journal of Physiology. Heart and Circulatory Physiology*, 304(12), H1585–H1597.
- Mellor, P. J., Mellanby, R. J., Baines, E. A., Villiers, E. J., Archer, J., & Herrtage, M. E. (2006). High serum troponin I concentration as a marker of severe myocardial damage in a case of suspected exertional heatstroke in a dog. *Journal of Veterinary Cardiology*, 8(1), 55–62.
- Miyamoto, K., Suzuki, K., Ohtaki, H., Nakamura, M., Yamaga, H., Yagi, M., Honda, K., Hayashi, M., & Dohi, K. (2021). A novel mouse model of

- heatstroke accounting for ambient temperature and relative humidity. *Journal of Intensive Care*, 9(1), 1–11.
- Mora, C., Dousset, B., Caldwell, I. R., Powell, F. E., Geronimo, R. C., Bielecki, C. R., Counsell, C. W., Dietrich, B. S., Johnston, E. T., & Louis, L. V. (2017). Global risk of deadly heat. *Nature Climate Change*, 7(7), 501–506.
- Murray, K. O., Brant, J. O., Iwaniec, J. D., Sheikh, L. H., de Carvalho, L., Garcia, C. K., Robinson, G. P., Alzahrani, J. M., Riva, A., & Laitano, O. (2021). Exertional heat stroke leads to concurrent long-term epigenetic memory, immunosuppression and altered heat shock response in female mice. *The Journal of Physiology*, 599(1), 119–141.
- Novosad, V. L., Richards, J. L., Phillips, N. A., King, M. A., & Clanton, T. L. (2013). Regional susceptibility to stress-induced intestinal injury in the mouse. *American Journal of Physiology. Gastrointestinal and Liver Physiology*, 305(6), G418–G426.
- Ohlstein, E. H., Vickery, L., Sauermech, C., & Willette, R. N. (1990). Vasodilation induced by endothelin: Role of EDRF and prostanoids in rat hindquarters. *American Journal of Physiology*, 259, H1835–1841.
- Pedersen, B. K., & Febbraio, M. A. (2008). Muscle as an endocrine organ: Focus on muscle-derived interleukin-6. *Physiological Reviews*, 88(4), 1379–1406.
- Persson, L., Hårdemark, H., Gustafsson, J., Rundström, G., Mendel-Hartvig, I., Esscher, T., & Pålman, S. (1987). S-100 protein and neuron-specific enolase in cerebrospinal fluid and serum: Markers of cell damage in human central nervous system. *Stroke*, 18(5), 911–918.
- Pires, W., Veneroso, C. E., Wanner, S. P., Pacheco, D. A., Vaz, G. C., Amorim, F. T., Tonoli, C., Soares, D. D., & Coimbra, C. C. (2017). Association between exercise-induced hyperthermia and intestinal permeability: A systematic review. *Sports Medicine*, 47(7), 1389–1403.
- Proctor, E. A., Dineen, S. M., Van Nostrand, S. C., Kuhn, M. K., Barrett, C. D., Brubaker, D. K., Yaffe, M. B., Lauffenburger, D. A., & Leon, L. R. (2020). Coagulopathy signature precedes and predicts severity of end-organ heat stroke pathology in a mouse model. *Journal of Thrombosis and Haemostasis*, 18(8), 1900–1910.
- Quinn, C. M., Duran, R. M., Audet, G. N., Charkoudian, N., & Leon, L. R. (2014). Cardiovascular and thermoregulatory biomarkers of heat stroke severity in a conscious rat model. *Journal of Applied Physiology*, 117(9), 971–978.
- Rae, D. E., Knobel, G. J., Mann, T., Swart, J., Tucker, R., & Noakes, T. D. (2008). Heatstroke during endurance exercise: Is there evidence for excessive endothermy? *Medicine and Science in Sports and Exercise*, 40(7), 1193–1204.
- Roberts, G. T., Ghebeh, H., Chishti, M. A., Al-Mohanna, F., El-Sayed, R., Al-Mohanna, F., & Bouchama, A. (2008). Microvascular injury, thrombosis, inflammation, and apoptosis in the pathogenesis of heatstroke: A study in baboon model. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 28(6), 1130–1136.
- Ruell, P. A., Simar, D., Périard, J. D., Best, S., Caillaud, C., & Thompson, M. W. (2014). Plasma and lymphocyte Hsp72 responses to exercise in athletes with prior exertional heat illness. *Amino Acids*, 46(6), 1491–1499.
- Schlader, Z. J., Chapman, C. L., Sarkar, S., Russo, L., Rideout, T. C., Parker, M. D., Johnson, B. D., & Hostler, D. (2017). Firefighter work duration influences the extent of acute kidney injury. *Medicine & Science in Sports & Exercise*, 49, 1745–1753.
- Schrier, R. W., Hano, J., Keller, H. I., Finkel, R. M., Gilliland, P. F., Cirksena, W. J., & Teschan, P. E. (1970). Renal, metabolic, and circulatory responses to heat and exercise. Studies in military recruits during summer training, with implications for acute renal failure. *Annals of Internal Medicine*, 73(2), 213–223.
- Segev, G., Daminet, S., Meyer, E., De Loor, J., Cohen, A., Aroch, I., & Bruchim, Y. (2015). Characterization of kidney damage using several renal biomarkers in dogs with naturally occurring heatstroke. *The Veterinary Journal*, 206(2), 231–235.
- Sharma, S., Chaudhary, P., Sandhir, R., Bharadwaj, A., Gupta, R. K., Khatri, R., Bajaj, A. C., Baburaj, T. P., Kumar, S., Pal, M. S., Reddy, P. K., & Kumar, B. (2021). Heat-induced endoplasmic reticulum stress in soleus and gastrocnemius muscles and differential response to UPR pathway in rats. *Cell Stress & Chaperones*, 26, 323–339.
- Shave, R., George, K. P., Atkinson, G., Hart, E., Middleton, N., Whyte, G., Gaze, D., & Collinson, P. O. (2007). Exercise-induced cardiac troponin T release: A meta-analysis. *Medicine and Science in Sports and Exercise*, 39(12), 2099–2106.
- Shepley, B. R., Ainslie, P. N., Hoiland, R. L., Donnelly, J., Sekhon, M. S., Zetterberg, H., Blennow, K., & Bain, A. R. (2021). Negligible influence of moderate to severe hyperthermia on blood-brain barrier permeability and neuronal parenchymal integrity in healthy men. *Journal of Applied Physiology*, 130(3), 792–800.
- Shieh, S. D., Shiang, J. C., Lin, Y. F., Shiao, W. Y., & Wang, J. Y. (1995). Circulating angiotensin-converting enzyme, von Willebrand factor antigen and thrombomodulin in exertional heat stroke. *Clinical Science*, 89(3), 261–265.
- Sohal, R. S., Sun, S. C., Colcolough, H. L., & Burch, G. E. (1968). Heat stroke. An electron microscopic study of endothelial cell damage and disseminated intravascular coagulation. *Archives of Internal Medicine*, 122(1), 43–47.
- Stacey, M. J., Hill, N. E., Parsons, I. T., Wallace, J., Taylor, N., Grimaldi, R., Shah, N., Marshall, A., House, C., & O'Hara, J. P. (2022). Relative changes in brain and kidney biomarkers with Exertional Heat Illness during a cool weather marathon. *PLoS One*, 17(2), e0263873.
- Tong, H. -S., Tang, Y. -Q., Chen, Y., Qiu, J. -M., Wen, Q., & Su, L. (2011a). Early elevated HMGB1 level predicting the outcome in exertional heatstroke. *Journal of Trauma and Acute Care Surgery*, 71(4), 808–814.
- Tong, H., Tang, Y., Chen, Y., Yuan, F., Liu, Z., Peng, N., Tang, L., & Su, L. (2013). HMGB1 activity inhibition alleviating liver injury in heatstroke. *Journal of Trauma and Acute Care Surgery*, 74(3), 801–807.
- Tong, H., Wan, P., Zhang, X., Duan, P., Tang, Y., Chen, Y., Tang, L., & Su, L. (2014). Vascular endothelial cell injury partly induced by mesenteric lymph in heat stroke. *Inflammation*, 37(1), 27–34.
- Tong, H. S., Tang, Y. Q., Chen, Y., Qiu, J. M., Wen, Q., & Su, L. (2011b). Early elevated HMGB1 level predicting the outcome in exertional heatstroke. *Journal of Trauma*, 71, 808–814.
- Umemura, Y., Ogura, H., Matsuura, H., Ebihara, T., Shimizu, K., & Shimazu, T. (2018). Bone marrow-derived mononuclear cell therapy can attenuate systemic inflammation in rat heatstroke. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine*, 26(1), 97.
- Van Eldik, L., Jensen, R., Ehrenfried, B. A., & Whetsell Jr, W. (1986). Immunohistochemical localization of S100 beta in human nervous system tumors by using monoclonal antibodies with specificity for the S100 beta polypeptide. *Journal of Histochemistry & Cytochemistry*, 34, 977–982.
- Wallace, R. F., Kriebel, D., Punnett, L., Wegman, D. H., & Amoroso, P. J. (2007). Prior heat illness hospitalization and risk of early death. *Environmental Research*, 104(2), 290–295.
- Wang, J. -C., Chien, W. -C., Chu, P., Chung, C. -H., Lin, C. -Y., & Tsai, S.-H. (2019). The association between heat stroke and subsequent cardiovascular diseases. *PLoS One*, 14(2), e0211386.
- Ward, M. D., King, M. A., Gabrial, C., Kenefick, R. W., & Leon, L. R. (2020). Biochemical recovery from exertional heat stroke follows a 16-day time course. *PLoS One*, 15(3), e0229616.
- Watson, P., Shirreffs, S. M., & Maughan, R. J. (2005). Blood-brain barrier integrity may be threatened by exercise in a warm environment. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 288(6), R1689–R1694.
- Wells, J. M., Brummer, R. J., Derrien, M., MacDonald, T. T., Troost, F., Cani, P. D., Theodorou, V., Dekker, J., Méheust, A., & De Vos, W. M. (2017). Homeostasis of the gut barrier and potential biomarkers. *American Journal of Physiology. Gastrointestinal and Liver Physiology*, 312(3), G171–G193.
- Whiticar, R., Laba, D., & Smith, S. (2008). Exertional heat stroke in a young man with a documented rise in troponin I. *Emergency Medicine Journal*, 25(5), 283–284.
- Wu, M., Wang, C., Liu, Z., Zhong, L., Yu, B., Cheng, B., & Liu, Z. (2021). Clinical characteristics and risk factors associated with acute kidney

- injury inpatient with exertional heatstroke: An over 10-year intensive care survey. *Frontiers in Medicine*, 8, 692.
- Yang, H.-H., Chang, C.-P., Cheng, J.-T., & Lin, M.-T. (2009). Attenuation of acute lung inflammation and injury by whole body cooling in a rat heatstroke model. *BioMed Research International*, 2009, 768086.
- Yang, M., Li, Z., Zhao, Y., Zhou, F., Zhang, Y., Gao, J., Yin, T., Hu, X., Mao, Z., & Xiao, J. (2017). Outcome and risk factors associated with extent of central nervous system injury due to exertional heat stroke. *Medicine*, 96(44), e8417.
- Zhang, L., Fan, X., Zhong, Z., Xu, G., & Shen, J. (2015). Association of plasma diamine oxidase and intestinal fatty acid-binding protein with severity of disease in patient with heat stroke. *The American Journal of Emergency Medicine*, 33(7), 867–871.
- Zhang, X., Chen, Y., Tang, L., Zhang, Y., Duan, P., Su, L., & Tong, H. (2018). The liver sinusoidal endothelial cell damage in rats caused by heatstroke. *European Journal of Inflammation*, 16, 205873921879432.

How to cite this article: Schlader, Z. J., Davis, M. S., & Bouchama, A. (2022). Biomarkers of heatstroke-induced organ injury and repair. *Experimental Physiology*, 107, 1159–1171. <https://doi.org/10.1113/EP090142>