



OPEN

Author Correction: Predicting dissolution and transformation of inhaled nanoparticles in the lung using abiotic flow cells: The case of barium sulphate

Johannes G. Keller, Uschi M. Graham, Johanna Koltermann-Jülly, Robert Gelein, Lan Ma-Hock, Robert Landsiedel, Martin Wiemann, Günter Oberdörster, Alison Elder & Wendel Wohlleben

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-019-56872-3>, published online 16 January 2020

This Article contains errors. In the Methods section, under subheading ‘Flow-through abiotic dissolution and transformation’,

“For the lower flow rate, this corresponds to a ratio, $SA/V = 0.02 \text{ h/cm}$ ”.

should read:

“For the lower flow rate, this corresponds to a ratio, $SA/V = 0.02 \text{ h/}\mu\text{m}$ ”.

In the Results section, under subheading ‘Dynamic abiotic dissolution’,

“If we determine for each sampling interval the instantaneous rates k (in units of $\text{ng/cm}^2/\text{h}$, Eq. 3) and the instantaneous surface area per volume flow SA/V (in units of h/cm , Eq. 4), hundreds of instantaneous release rates collapse on a single linear relationship, regardless if SA/V was modulated by initial surface area or by flow rate or by gradual dissolution (Fig. 3)”.

should read:

“If we determine for each sampling interval the instantaneous rates k (in units of $\text{ng/cm}^2/\text{h}$, Eq. 3) and the instantaneous surface area per volume flow SA/V (in units of $\text{h/}\mu\text{m}$, Eq. 4), hundreds of instantaneous release rates collapse on a single linear relationship, regardless if SA/V was modulated by initial surface area or by flow rate or by gradual dissolution (Fig. 3)”.

And,

“The best match of the predicted halftime with the *in vivo* halftime is obtained for SA/V ratios around 0.01 to 0.03 h/cm ”.

should read:

“The best match of the predicted halftime with the *in vivo* halftime is obtained for SA/V ratios around 0.01 to 0.03 $\text{h/}\mu\text{m}$ ”.

Furthermore, the x-axis of Figure 3 is incorrectly labeled as “ $SA/V \text{ h/cm}$ ”, whereas the correct unit is “ $SA/V \text{ h/}\mu\text{m}$ ”. The correct Figure 3 appears below as Figure 1.

Published online: 19 April 2021




Figure 1. Instantaneous rate evaluation of biodissolution of BaSO_4 in flow-through cells with pH 4.5 PSF media. Each cloud of stepwise rates stems from separate experiment of initial mass M_0 and volume flow V . Five experiments for BaSO_4 (orange) and two for CuO (black). See Table 2 for conventional evaluation (cumulative rates) of the same raw data.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021