

POSTER PRESENTATIONS - PROFFERED ABSTRACTS | JUNE 01 2020

## Abstract A093: Employment and work experiences after breast cancer treatment *⊗*

Christine C. Ekenga; Cora McElwain; Maria Perez; Donna B. Jeffe



+ Author & Article Information

Cancer Epidemiol Biomarkers Prev (2020) 29 (6\_Supplement\_1): A093.

https://doi.org/10.1158/1538-7755.DISP18-A093

Split-Screen	Share ∨	Tools 🗸	Versions ∨	

## **Abstract**

**Background:** Employment is a social determinant of health. However, few longitudinal studies have examined employment outcomes in African-American women with breast cancer. We examined factors associated with return to work over 2-year follow-up in a sample of African-American breast cancer patients participating in a randomized controlled trial of a cancer-information intervention's impact (vs. standard of care) on quality-of-life and treatment adherence outcomes.


**Methods:** Interview and medical-record data from 227 newly diagnosed African-American breast cancer patients (stage 0-III), who enrolled a mean 6 days from surgical post-op visit or start of neoadjuvant therapy, were analyzed in association with return to work; four more interviews were conducted over two years. Potential predictors included sociodemographic variables (age, marital status, income, education, insurance status), treatment(s) received (surgery type, chemotherapy, radiation), comorbidity, and elevated depressed mood (Center for Epidemiologic Studies Depression Scale [CES-D] score > 15). Multivariable logistic regression models were used to identify factors independently associated with return to work.

**Results:** At enrollment, 100 patients (44%) were employed part- or full-time; 71 of employed patients returned to work during 2-year follow-up. Study arm and other treatment and sociodemographic variables were not significantly associated with return to work and was not included in the final model. Patients with elevated depressed mood at baseline were less likely to return to work than nondepressed patients (adjusted odds ratio = 5.8, 95% CI = 1.7-9.3).

**Conclusions:** Patients with elevated depressed mood were less likely to return to work over 2-year follow-up. Screening for depressed mood at diagnosis and providing treatment might be an effective strategy to improve continued workforce participation in African-American breast cancer patients.

**Citation Format:** Christine C. Ekenga, Cora McElwain, Maria Perez, Donna B. Jeffe. Employment and work experiences after breast cancer treatment [abstract]. In: Proceedings of the Eleventh AACR Conference on the Science of Cancer Health Disparities in Racial/Ethnic Minorities and the Medically Underserved; 2018 Nov 2-5; New Orleans, LA. Philadelphia (PA): AACR; Cancer Epidemiol Biomarkers Prev 2020;29(6 Suppl):Abstract nr A093.

©2020 American Association for Cancer Research.



## **Citing Articles Via**

Google Scholar

CrossRef



Skip to Main Content

Advertisement

Issues

Online First

Collections

News

**Twitter** 

Online ISSN 1538-7755 Print ISSN 1055-9965

## AACR Journals

**Blood Cancer Discovery** 

Cancer Discovery

Cancer Epidemiology, Biomarkers, & Prevention

Cancer Immunology Research

Cancer Prevention Research

Cancer Research

**Cancer Research Communications** 

Clinical Cancer Research

Molecular Cancer Research

Molecular Cancer Therapeutics

Info for Advertisers

Info for Librarians

Privacy Policy

Skin to Main Content





Copyright © 2022 by the American Association for Cancer Research.