

SUPPLEMENTARY TABLE 2. Onset, offset, and peak respiratory syncytial virus detections and percentage of tests positive at peak by surveillance week and U.S. Department of Health and Human Services Region* — National Respiratory and Enteric Virus Surveillance System, United States, June 30, 2024–June 28, 2025

Region [†]	Onset [§] (surveillance week)	Offset [¶] (surveillance week)	Peak ^{**} (surveillance week)	Percentage of tests positive at peak
National	Nov 9, 2024 (45)	Mar 29, 2025 (13)	Dec 21, 2024 (51)	11
<i>HHS Region 1</i>	Nov 16, 2024 (46)	Mar 8, 2025 (10)	Dec 21, 2024 (51)	17.3
<i>HHS Region 2</i>	Nov 9, 2024 (45)	Feb 8, 2025 (06)	Dec 21, 2024 (51)	13.7
<i>HHS Region 3</i>	Oct 26, 2024 (43)	Mar 15, 2025 (11)	Dec 21, 2024 (51)	16.9
<i>HHS Region 4</i>	Oct 19, 2024 (42)	Mar 1, 2025 (09)	Dec 21, 2024 (51)	15.8
<i>HHS Region 5</i>	Nov 23, 2024 (47)	Apr 5, 2025 (14)	Jan 4, 2024 (01)	10.7
<i>HHS Region 6</i>	Oct 5, 2024 (40)	Mar 29, 2025 (13)	Nov 30, 2024 (48)	19.3
<i>HHS Region 7</i>	Dec 7, 2024 (49)	Apr 19, 2025 (16)	Jan 18, 2024 (03)	9.7
<i>HHS Region 8</i>	Dec 14, 2024 (50)	Apr 26, 2025 (17)	Feb 22, 2024 (08)	14.1
<i>HHS Region 9</i>	Nov 16, 2024 (46)	Mar 29, 2025 (13)	Jan 4, 2024 (01)	8.8
<i>HHS Region 10</i>	Nov 30, 2024 (48)	Mar 29, 2025 (13)	Jan 25, 2025 (04)	7.9
<i>Alaska*</i>	††	††	††	††
<i>Florida*</i>	Sep 28, 2024 (39)	Mar 22, 2025 (12)	Dec 7, 2024 (49)	23.0
<i>Hawaii*</i>	Sep 28, 2024 (39)	Feb 8, 2025 (06)	Nov 30, 2025 (48)	15.2

Abbreviation: HHS = Department of Health and Human Services

* To account for notable subregional variations in seasonal RSV activity patterns, HHS Region 4 excludes Florida, HHS Region 9 excludes Hawaii, and HHS Region 10 excludes Alaska; those 3 states are described individually.

[†] <https://www.hhs.gov/about/agencies/iea/regional-offices/index.html>

[§] Week ending date corresponding to the first of 2 consecutive weeks when the percentage of weekly tests positive for RSV reaches or exceeds 3%.

[¶] Week ending date corresponding to the last of 2 consecutive weeks when the percentage of weekly tests positive for RSV reaches or exceeds 3%.

^{**} Week ending date corresponding to the week between onsets and offsets when the highest percentage of weekly tests are positive for RSV.

^{††} Fewer than two consistently reporting laboratories within the state. Laboratories with consistent reporting are defined as those that reported during ≥30 weeks and an average of ≥10 RSV polymerase chain reaction tests during the surveillance year.