



Emergency Preparedness and Response

# Monkeypox Caused by Human-to-Human Transmission of *Monkeypox Virus* in the Democratic Republic of the Congo with Spread to Neighboring Countries



Distributed via the CDC Health Alert Network

August 7, 2024, 3:15 PM ET

CDCHAN-00513

## Summary

The Centers for Disease Control and Prevention (CDC) is issuing this Health Alert Network (HAN) Health Update to provide additional information about the outbreak of monkeypox virus (MPXV) in the Democratic Republic of the Congo (DRC); the first [Health Advisory](#) about this outbreak was released in December 2023.

Since January 2023, the DRC has reported the largest number of yearly suspected clade I monkeypox cases on record. While clade I MPXV is endemic, or naturally occurring, in DRC, the current outbreak is more widespread than any previous DRC outbreak and has resulted in clade I monkeypox transmission to some neighboring countries. The Republic of the Congo (ROC), which borders DRC to the west, declared a clade I monkeypox outbreak in April 2024, and there have been confirmed cases in the Central African Republic (CAR). While clade I monkeypox is endemic in ROC and CAR, the epidemiologic pattern of recent cases suggests a possible link to DRC.

In late July 2024, Burundi, Rwanda, and Uganda, which sit on the eastern border of DRC, reported confirmed cases of monkeypox, with some cases having linkages to DRC. Rwanda and Uganda have confirmed these cases are due to clade I MPXV; in Burundi, clade-specific testing is underway, but cases are presumed to be clade I due to DRC's proximity. Monkeypox is not known to be endemic in these countries.

No cases of clade I monkeypox have been reported outside central and eastern Africa at this time. Because there is a risk of additional spread, CDC recommends clinicians and jurisdictions in the United States maintain a heightened index of suspicion for monkeypox in patients who have recently been in DRC or to any country sharing a border with DRC (ROC, Angola, Zambia, Rwanda, Burundi, Uganda, South Sudan, CAR) and present with [signs and symptoms consistent with](#)

**monkeypox.** These can include: rash that may be located on the hands, feet, chest, face, mouth, or near the genitals; fever; chills; swollen lymph nodes; fatigue; myalgia (muscle aches and backache); headache; and respiratory symptoms like sore throat, nasal congestion, and cough.

## Background

MPXV has two distinct genetic clades (subtypes of MPXV), I and II, which are endemic to central and west Africa, respectively. Clade I MPXV has previously been observed to be more transmissible and to cause a higher proportion of severe infections than clade II MPXV. The [ongoing global monkeypox outbreak](#) that began in 2022 is caused by clade II MPXV, and cases continue to be reported worldwide.

Clade I MPXV is endemic in DRC and several other Central African countries, and cases are reported annually. More than 22,000 suspect cases, with more than 1,200 suspected deaths, have been reported in DRC since January 1, 2023, a substantial increase from the median 3,767 suspect [clade I monkeypox cases reported annually in DRC](#) during 2016–2021. Clade I monkeypox cases have been reported from every DRC province, including areas where clade I monkeypox does not normally occur, such as the capital city Kinshasa. Outbreaks of clade I MPXV associated with sexual contact among men who have sex with men and female sex workers and their contacts have been reported in some provinces. In other provinces, patients have acquired infection through contact with infected dead or live wild animals, household transmission, or patient care (transmitted in the absence of appropriate personal protective equipment); a high proportion of cases have been reported in children younger than 15 years of age. Monkeypox vaccine, which is expected to be effective against both clades, is not generally available in DRC at this time. However, the country is actively working on a plan to vaccinate.

Confirmed clade I monkeypox cases were reported in April in CAR and ROC. In late July 2024, clade I cases were confirmed in Rwanda and Uganda. Cases were also confirmed in Burundi; due to Burundi's proximity to DRC and Rwanda, these cases are presumed to be clade I while clade-specific testing is conducted. Clade I MPXV is not known to be endemic in Burundi, Rwanda, and Uganda.

Due to the limited number of travelers and lack of direct commercial flights from DRC or its neighboring countries to the United States, the [risk of clade I monkeypox importation](#) to the United States is considered to be very low.

The United States has robust monkeypox testing capacity in state public health laboratories and several commercial laboratories, including clade-specific testing, sequencing, and/or flagging high-likelihood clade I MPXV samples (i.e., negative for clade II MPXV but positive for orthopoxvirus). In addition, CDC continues to receive a subset of MPXV samples from across the United States that were not differentiated during the initial diagnosis to test for MPXV clade and to look for mutations using genetic sequencing. CDC is helping communities monitor the presence of both clades of [MPXV in wastewater samples](#), including from select airports. Data from samples can provide an early warning of monkeypox activity and spread in communities.

## Recommendations for Clinicians

### *Evaluation and Diagnosis*

- Follow CDC guidance on [infection prevention and control](#) for monkeypox to minimize transmission risk when evaluating and providing care to patients with suspected monkeypox.
- Consider monkeypox as a possible diagnosis in patients with [epidemiologic characteristics](#) and [lesions or other clinical signs and symptoms](#) consistent with monkeypox. This includes persons who have been in DRC or, due to the demonstrated risks of regional spread, any of its neighboring countries (ROC, CAR, Rwanda, Burundi, Uganda, Zambia, Angola, Tanzania, and South Sudan) in the previous 21 days.
- Ask patients with signs and symptoms of monkeypox but no recent travel whether they have had contact with people who had recently been in any of the above countries and who were symptomatic for monkeypox.
- Consider monkeypox as a possible diagnosis if a clinically consistent presentation occurs, even in people vaccinated for or previously diagnosed with monkeypox.
- Advise all patients suspected of having monkeypox to [isolate themselves](#) from others.

- Evaluate all suspected cases related to DRC or its neighboring countries with laboratory testing (rather than clinical diagnosis alone). In most situations, specimens should be sent to the appropriate state public health laboratory or a commercial laboratory for initial testing. If you are authorized by your health department to send specimens directly to CDC for testing, contact CDC at [poxviruslab@cdc.gov](mailto:poxviruslab@cdc.gov) for information regarding specimen types accepted, labeling, specimen storage, and shipping timeframes.
- Follow [specimen collection guidelines](#) (including collecting two swabs per ~2-3 lesions) to ensure specimen availability for clade-specific testing. This testing will help distinguish between cases that are part of the ongoing clade II monkeypox global outbreak and those associated with this clade I outbreak.
- Avoid unroofing or aspiration of lesions or otherwise using sharp instruments for monkeypox testing to minimize the risk of a sharps injury.

### **Treatment and Prevention**

- Recommend [monkeypox vaccine to people exposed to MPXV](#) to help prevent the spread of monkeypox.
- Offer monkeypox vaccination to people  $\geq 18$  years of age with risk factors for monkeypox, following [the Advisory Committee on Immunization Practices \(ACIP\)](#) recommendation for vaccination before an exposure with two doses of the JYNNEOS vaccine 28 days apart.
  - Two doses of JYNNEOS vaccine [offer substantial protection against monkeypox](#), and is expected to offer protection regardless of clade.
  - Additional JYNNEOS vaccine doses (“boosters,” more than two doses) are not currently recommended.
- Consider vaccinating patients [eligible for monkeypox vaccination](#) and planning travel to affected countries, with two doses of JYNNEOS vaccine. Eligible patients who received one dose of the JYNNEOS vaccine more than 28 days ago should receive the second dose as soon as possible.
- There is no vaccination recommendation for travelers who do not meet current vaccine eligibility.
- Consult your health department or CDC ([poxvirus@cdc.gov](mailto:poxvirus@cdc.gov)) promptly about any monkeypox cases for which severe manifestations might occur (e.g., those with advanced HIV infection). Medical countermeasures (e.g., tecovirimat, brincidofovir, and vaccinia immune globulin intravenous) used during the ongoing clade II monkeypox outbreak are expected to be effective for clade I MPXV infections.
- Inform all patients with monkeypox, including those with mild disease, about the STOMP Trial and recommend that they enroll. Oral tecovirimat (TPOXX) is available through the STOMP Trial. To enroll in STOMP, call 1-855-876-9997.
- Contact your state, local, or territorial health departments to see if oral TPOXX remains available from prior prepositioned supplies for patients who are ineligible for STOMP’s open-label arm (e.g., illness  $\geq 14$  days or prior TPOXX receipt) but [meet expanded use Investigational New Drug \(EA-IND\) eligibility](#) for tecovirimat treatment for monkeypox.
- Clinicians should counsel patients about [what to do if they are sick](#) to prevent household transmission, if they have monkeypox symptoms; staying away from other people and not sharing things they have touched with others; and cleaning and disinfecting the spaces they occupy regularly to limit household contamination.

### **Recommendations for Health Departments**

- Promote monkeypox vaccination in your community to protect as many [eligible people](#) as possible from monkeypox.
- Report monkeypox cases to CDC within 24 hours. Initial reports can be submitted with only the minimum required data elements of a local record ID and case jurisdiction of residence.
  - Enter the necessary sCRF data directly in the CDC DCIPHER platform. Data can also be entered into an existing jurisdictional case surveillance system configured for monkeypox reporting, with data uploaded to the CDC DCIPHER platform as a CSV file.
- Collect the data listed in the [2022/2023 U.S. Monkeypox Outbreak Short Case Report Form \(sCRF\)](#) for patients who meet the probable or confirmed monkeypox case definition. Local health departments should check with state or territorial health authorities to verify their jurisdiction’s preferred case reporting process.

### **Recommendations for Laboratories**

- Follow CDC guidance on [infection prevention and control](#) for monkeypox to minimize transmission risk when working with suspected monkeypox specimens.
- Send clinical specimens collected from patients who traveled from DRC or its neighboring countries, or had close or intimate contact with symptomatic people from these countries, to a laboratory that can perform clade-specific testing **as quickly as possible**.
- If clade-specific testing is warranted based on epidemiologic criteria but is not available in a jurisdiction, [specimen submission](#) to a public health laboratory with this capability or to CDC is encouraged; specimen submission to CDC can be coordinated through your state or local health department. Specimens that cannot be accepted at CDC for clinical testing under [Clinical Laboratory Improvement Amendments \(CLIA\)](#) will be redirected for surveillance purposes and tested, providing critical data on MPXV clade(s) circulating in the United States.
- [Laboratory Response Network](#) laboratories and commercial laboratories using CDC's non-variola orthopoxvirus (NVO) polymerase chain reaction (PCR) test should continue submitting duplicate specimens to CDC from all patients with positive NVO PCR test results for routine MPXV clade-specific testing. This testing will assist with national surveillance.
- Some non-CDC laboratories may also have other options available for clade-specific testing, (g., molecular testing or genetic sequencing). These laboratories should alert their [state health department](#) [↗](#) and CDC ([poxvirus@cdc.gov](mailto:poxvirus@cdc.gov)) if results from such tests indicate detection of clade I MPXV.

## Recommendations for the Public

- The [risk of clade I monkeypox spreading to the United States](#) is very low at this time.
- Seek medical care immediately and avoid contact with others if you have been in the DRC or its neighboring countries in the last 21 days and develop a new, [unexplained skin rash \(lesions on any part of the body\), with or without fever and chills](#).
- Consider getting vaccinated against monkeypox if you have [risk factors and are eligible for vaccination](#). CDC continues to recommend that people who are eligible for vaccination receive two doses of the JYNNEOS vaccine for the best protection. People at risk for monkeypox who have only received one dose more than 28 days prior should receive a second dose as soon as possible. JYNNEOS vaccine is believed to protect against both monkeypox clades.
- Review [CDC Travel Health Notices for the DRC and neighboring countries](#) before traveling. People with risk factors for MPXV infection who are not able to be vaccinated or (e.g., pregnant women, infants less than 1 year, people with eczema or active skin conditions, and people who are immunocompromised) should avoid situations that might increase their risk for monkeypox.
- All travelers to areas with monkeypox cases should [protect themselves](#) by avoiding close contact with people who have skin or genital lesions; avoiding contact with dead or live wild animals; avoiding contact with materials used by sick people like clothing, bedding, or in health care; avoiding materials that came into contact with wild animals; and avoiding eating or preparing meat from wild animals (bushmeat), or using products made from wild animals in countries where monkeypox occurs in animals.

## For More Information

### For clinicians and laboratory staff

- Monkeypox Clinical Recognition and Vaccine Information for Healthcare Providers: [Information For Healthcare Professionals | Monkeypox | Poxvirus | CDC](#)
- Biosafety and Select Agent Considerations: [Monkeypox Laboratory Information](#)
- Diagnostic Specimen Packaging and Shipping: [Transporting Infectious Substances Safely.pdf \(dot.gov\)](#) [📄](#) [↗](#)
- CDC Poxvirus and Rabies Branch: [poxvirus@cdc.gov](mailto:poxvirus@cdc.gov) or for emergencies, CDC's 24/7 Emergency Operations Center (EOC): 770-488-7100.
- State and Local Health Department Contacts: [After Hours/Epi-on-Call Contact Lists – Council of State and Territorial Epidemiologists \(cste.org\)](#) [↗](#)

## For the public

- General inquiries: CDC-INFO (1-800-232-4636)
- Monkeypox Information for the Public: [About Monkeypox](#)
- [August 2024 Travel Health Notice: Monkeypox in DRC and Neighboring Countries](#)

## References

- Dalton AF, Diallo AO, Chard AN, et al. Estimated Effectiveness of JYNNEOS Vaccine in Preventing Monkeypox: A Multijurisdictional Case-Control Study — United States, August 19, 2022–March 31, 2023. *MMWR Morb Mortal Wkly Rep.* 2023;72:553–558. DOI: <http://dx.doi.org/10.15585/mmwr.mm7220a3> [↗](#)
- Kibungu EM, Vakaniaki EH, Kinganda-Lusamaki E, et al. Clade I-Associated Monkeypox Cases Associated with Sexual Contact, the Democratic Republic of the Congo. *Emerg Infect Dis.* Published online November 29, 2023. [doi:10.3201/eid3001.231164](https://doi.org/10.3201/eid3001.231164)

---

*The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national and international organizations.*

## DEPARTMENT OF HEALTH AND HUMAN SERVICES

---

### HAN Message Types

- **Health Alert:** Conveys the highest level of importance about a public health incident.
- **Health Advisory:** Provides important information about a public health incident.
- **Health Update:** Provides updated information about a public health incident.

###

This message was distributed to state and local health officers, state and local epidemiologists, state and local laboratory directors, public information officers, HAN coordinators, and clinician organizations.


###

[Top of Page](#)

### Additional Resources

- [HAN Archive By Year](#)
- [HAN Types](#)
- [Sign Up for HAN Email Updates](#)
- [HAN Jurisdictions](#)

Last Reviewed: August 7, 2024

 Was this page helpful?

Yes    Partly    No