

Age-Adjusted Colorectal Cancer Death Rates, by State — United States, 2024

Jurisdiction	Deaths	Age-adjusted rate (per 100,000 standard population)
United States (overall)	55,898	12.9
Alabama	969	14.6
Alaska	121	16.1
Arizona	1,270	12.5
Arkansas	631	16.0
California	5,817	12.2
Colorado	753	11.0
Connecticut	527	10.3
Delaware	190	12.9
District of Columbia	93	12.9
Florida	4,212	11.9
Georgia	1,847	14.3
Hawaii	233	10.9
Idaho	308	12.6
Illinois	2,098	12.9
Indiana	1,285	14.8
Iowa	587	13.8
Kansas	511	13.7
Kentucky	899	15.5
Louisiana	826	14.6
Maine	253	12.0
Maryland	998	12.5
Massachusetts	986	10.5
Michigan	1,723	12.8
Minnesota	879	12.0
Mississippi	642	17.2
Missouri	1,266	15.5
Montana	195	12.6
Nebraska	349	14.3
Nevada	564	14.0
New Hampshire	227	11.4
New Jersey	1,377	11.0
New Mexico	382	13.4
New York	2,895	10.9
North Carolina	1,866	13.5
North Dakota	119	12.1
Ohio	2,213	14.2
Oklahoma	866	17.8
Oregon	733	12.9
Pennsylvania	2,427	13.2

Rhode Island	153	10.1
South Carolina	939	13.0
South Dakota	191	16.0
Tennessee	1,366	15.1
Texas	4,611	14.1
Utah	334	10.6
Vermont	117	12.2
Virginia	1,400	12.6
Washington	1,143	11.8
West Virginia	445	16.8
Wisconsin	962	12.1
Wyoming	100	12.9

NOTES: Colorectal cancer deaths were identified using International Classification of Diseases, Tenth Revision underlying cause-of-death codes C18–C21. Age-adjusted colorectal cancer deaths were calculated using the direct method and the 2000 U.S. standard population.

SOURCE: National Center for Health Statistics, National Vital Statistics System, Mortality Data.