

Supplemental Table 1. Everyday Discrimination Scale ^a

	Overall (N=245)	White (n=84)	Black (n=25)	Hispanic (n=59)	Asian (n=77)
You are treated with less courtesy than other people are	1 [1.1]	0.8 [1.1]	1.4 [1.2]	0.8 [1.1]	1.2 [1]
You are treated with less respect than other people are	1 [1.1]	0.8 [1]	1.4 [1.2]	0.8 [1.1]	1.1 [1]
You receive poorer service than other people at restaurants or stores	0.8 [1.1]	0.4 [0.7]	1.4 [1.4]	0.7 [1.1]	1.1 [1.2]
People act as if they think you are not smart	0.8 [1.1]	0.6 [0.9]	1.6 [1.7]	0.8 [1.1]	1 [1]
People act as if they are afraid of you	0.5 [0.9]	0.3 [0.7]	1.2 [1.4]	0.5 [1]	0.5 [0.8]
People act as if they think you are dishonest	0.2 [0.5]	0.1 [0.3]	0.3 [0.5]	0.3 [0.7]	0.2 [0.6]
People act as if they're better than you are	1.1 [1.2]	0.7 [1]	1.9 [1.7]	1.2 [1.3]	1.3 [1.2]
You are called names or insulted	0.5 [0.8]	0.3 [0.6]	0.6 [1.2]	0.5 [0.9]	0.6 [0.8]
You are threatened or harassed	0.3 [0.7]	0.2 [0.5]	0.6 [1.2]	0.2 [0.5]	0.4 [0.7]
You are followed around in stores	0.2 [0.7]	0.05 [0.3]	1.1 [1.4]	0.1 [0.4]	0.2 [0.5]

Tabled values are mean [standard deviation]

^a Questions asked verbatim as above on a Likert scale (never, less than once a year, a few times a year, a few times a month, at least once a week, almost every day, and don't know). Likert scale responses were coded (never = 0, less than once a year = 1, a few times a year = 2, a few times a month = 3, at least once a week = 4, and almost every day = 5) and summed to create a score for each type of discrimination experience.

Supplemental Table 2. Immunosuppressive Medication Use within the Past Month ^a

Medications	N=245
Imuran or Azathioprine	28 (11.4%)
CellCept, Mycophenolate Mofetil, or Myfortic	69 (28.2%)
Methotrexate, Rheumatrex, or Trexall	13 (5.3%)
Cyclosporine, Sandimmune, Neoral, Gengraf, Prograf, Tacrolimus, or Sirolimus	14 (5.7%)
Leflunomide or Arava	4 (1.6%)
Cytosan or Cyclophosphamide	0 (0%)
Enbrel, Etanercept, Remicade Infliximab, Humira, Adalimumab, Simponi, Golimumab, Cimzia, or Certolizumab	2 (0.8%)
Rituxan or Rituximab	0 (0%)
Orencia or Abatacept	3 (1.2%)
Benlysta or Belimumab	5 (2.0%)

Tabled values are n (%)

^a Participants were asked if they had taken any of the following medications in the past month. Each row corresponds to a separate question.