

Table S4. Adherence to recommended follow-up for the subset of MI-SIGHT program participants who screened positive for glaucoma and were included in the RCT (n=490)

Follow-up Adherence	Standard Glaucoma Education	Personalized Glaucoma Education	P-value**
	n (column %)	n (column %)	
Analysis 1*			
Overall (n=490)	(n=243)	(n=247)	
Non-Adherent	99 (40.7)	97 (39.2)	0.7399
Adherent	144 (59.3)	150 (60.7)	
Free Clinic (n=180)	(n=89)	(n=91)	
Non-Adherent	26 (29.2)	21 (23.1)	0.3487
Adherent	63 (70.8)	70 (76.9)	
FQHC (n=310)	(n=81)	(n=80)	
Non-Adherent	73 (47.4)	76 (48.7)	0.8167
Adherent	81 (52.6)	80 (51.3)	
Analysis 2*			
Overall (n=448)	(n=219)	(n=229)	
Non-Adherent	75 (34.3)	79 (34.5)	0.9554
Adherent	144 (65.8)	150 (65.5)	
Free Clinic (n=169)	(n=81)	(n=88)	
Non-Adherent	18 (22.2)	18 (20.5)	0.7792
Adherent	63 (77.8)	70 (79.6)	
FQHC (n=279)	(n=138)	(n=141)	
Non-Adherent	57 (41.3)	61 (43.3)	0.7406
Adherent	81 (58.7)	80 (56.7)	

MI-SIGHT, Michigan Screening & Intervention for Glaucoma and Eye Health through
Telemedicine; RCT, Randomized Controlled Trial; FQHC, Federally Qualified Health Center

* Analysis 1 assumes those with unknown adherence are non-adherent; Analysis 2 removes
those with unknown adherence

**Chi-square test