

486

487 Supplemental Table 2: Length of Stay, Hospitalization Charges, and Costs for Neonatal Herpes Infections by Survival Status at Discharge, United
488 States, 2019

	Non-survivors	Survivors
Length of stay, days		
Range (IQR)	4-15	12-23
Mean (95% CI)	18 (2-34)	24 (22-26)
Median (95% CI)	7 (3-11)	16 (14-18)
Hospitalization charges^{a,b}, in 2019 US\$ thousands		
Range (IQR)	102-354	59-225
Mean (95% CI)	423 (167-789)	199 (170-227)
Median (95% CI)	134 (45-223)	107 (96-118)
Hospitalization cost^{b,c} (adjusted by CCR), in 2019 US\$ thousands		
Range (IQR)	25-68	14-53
Mean (95% CI)	117 (50-227)	49 (42-55)
Median (95% CI)	40 (20-60)	27 (23-30)

489

490 Abbreviations: CI, confidence interval; IQR, Interquartile Range; CCR, Cost-to-Charge Ratio; HCUP, Healthcare Cost and Utilization Project; KID, Kids' Inpatient Database

491 ^aHospitalization charge estimated using total charge in HCUP KID

492 ^bRounded to the nearest 1,000

493 ^cHospitalization cost is determined by adjusting the hospitalization charge using the HCUP CCR¹¹

494

495

496

497 Supplemental Table 3: Comparison of 2019 Neonatal Herpes Infections Incidence Rates: Live Birth vs. Hospital Birth as Denominator

	Incidence Rate per 100 000		Relative Difference (%)
	Live Birth ^a	Hospital Birth ^b	(Live Birth as denominator compared to Hospital Birth as Denominator)
United States	15.0	15.7	4.7
Northeast	13.6	14.1	3.7
Midwest	13.4	14.1	5.2
South	20.4	21.3	4.4
West	8.1	8.7	7.4

498

499 ^aCenters for Disease Control and Prevention, National Center for Health Statistics. National Vital Statistics System, Natality on CDC WONDER Online Database. Data are from the Natality Records 2016-2021, as compiled
500 from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/natality-expanded-current.html> on May 10, 2023 8:34:21 AM

501 ^bHospital births estimated from Project Kids' Inpatient Database(KID)

502