

Supplemental Table 1. Distribution of cases by multiple risk categories

	Total^a	+MSM	+Injection drug use	+Homelessness	+Incarceration	+International travel	+Epi-linkage
MSM	347 (6%)	MSM only=220 (63%)	53 (15%)	44 (13%)	10 (3%)	25 (7%)	34 (10%)
Injection drug use	1823 (33%)	53 (3%)	Injection drug use only=779 (43%)	632 (35%)	237 (13%)	4 (2%)	421 (23%)
Homelessness	1007 (18%)	44 (4%)	632 (63%)	Homeless only=242 (24%)	131 (13%)	7 (<1%)	223 (22%)
Incarceration	392 (7%)	10 (3%)	237 (60%)	131 (33%)	Incarceration only=91 (23%)	4 (1%)	83 (21%)
International travel	222 (4%)	25 (11%)	4 (2%)	7 (3%)	4 (2%)	International travel only=173 (78%)	14 (6%)
Epidemiologic linkage	1224 (22%)	34 (3%)	421 (34%)	223 (18%)	83 (7%)	14 (1%)	Epi-linkage only=656 (54%)
No risk factor identified	2059 (37%)	NA	NA	NA	NA	NA	NA
Total	5491						

^aTotal column indicates the number and percentage of cases in each risk category. Percentages do not add to 100%.

For each row, the denominator used to calculate percentages in the 6 right-hand columns is the number in the total column for each row. The highlighted cells indicate the number and percentage of cases that had only that risk category identified. Other cells in the row indicate the number and percentage of cases that had the primary risk category for that row, plus one other risk category indicated by the column. Combinations with more than 2 risk categories per case are not indicated in the table.

Example for row 1: Total column, 347 of 5491 cases (6%) identified as MSM. Of these, 220 of 347 (63%) identified as MSM only, without other risk categories identified; 53 of 347 (15%) who identified as MSM also reported injection drug use; 44 of 347 (13%) who identified as MSM also experienced homelessness, etc.

NA = not applicable