

MMWR™

MORBIDITY AND MORTALITY WEEKLY REPORT

- 1065 Mycobacterial Infection Following Liposuction and Liposculpture — Caracas, Venezuela, 1996–1998
- 1067 Impact of Closure of a Sexually Transmitted Disease Clinic — Washington, D.C., 1995
- 1070 Outbreak of Primary and Secondary Syphilis — 1996–1997
- 1073 Fatal *Cercopithecine herpesvirus 1* (B Virus) Infection Following a Mucocutaneous Exposure
- 1083 Notice to Readers



Rapidly Growing Mycobacterial Infection Following Liposuction and Liposculpture — Caracas, Venezuela, 1996–1998

During October 1996–March 1998, nine patients in eight hospitals in Caracas, Venezuela, acquired surgical-site infections (SSI) caused by rapidly growing mycobacteria (RGM). All episodes of RGM infection occurred within 2 months after liposuction or liposculpture (aesthetic surgical procedures). This report describes the findings of an epidemiologic investigation of this cluster by the Venezuelan Ministry of Health and underscores the importance of sterilizing surgical equipment to prevent nosocomial infections.

A confirmed case was defined as RGM in a patient who underwent liposuction or liposculpture during October 1996–March 1998 (study period) in a surgical facility in Caracas, in whom local signs of SSI were present and for whom cultures of surgical site drainage grew RGM. A probable case was defined as RGM in a patient who underwent liposuction or liposculpture in a surgical facility in Caracas during the study period, who had local signs of infection at the surgical site, and for whom microscopic examination of stained smears of surgical site drainage were positive for acid-fast bacilli.

Seven confirmed and two probable cases from eight hospitals were identified. All case-patients were previously healthy women aged 28–49 years (median: 37.5 years). Eight surgeons and surgical teams performed the cosmetic surgery on the women. All nine case-patients underwent general anesthesia during their surgical procedure; procedures consisted of abdominal liposuction (seven patients), anterior and posterior thigh liposuction (three), or bilateral nasolabial fold liposculpture (two). The median time from surgical procedure to onset of infection was 15 days (range: 4–45 days). Clinical findings included fever, local inflammation, microabscesses, purulent drainage from the wound, or fistulae.

Seven case-patients had culture-confirmed RGM; species identified were *Mycobacterium chelonae* (four patients), *M. fortuitum* (two), and *M. abscessus* (one). Molecular typing of RGM isolates were not performed.

All hospitals cleaned surgical instruments (i.e., liposuction and liposculpture cannulae) with tap water and soap followed by low-level disinfection with a commercial quaternary ammonium solution. Environmental cultures, including cultures of tap water, at two surgical units did not yield bacteria or mycobacteria. The epidemiologic

Rapidly Growing Mycobacterial Infection — Continued

investigation did not reveal risk factors such as exposure to certain persons, cleaning solutions, medical supplies, or contaminated quaternary ammonium compounds.

Following the outbreak in Caracas, two of the affected surgical facilities modified their reprocessing procedures for surgical instruments (including suction cannulae) used in cosmetic surgical procedures by replacing quaternary ammonium compounds used for low-level disinfection with either high-level disinfection using 2% glutaraldehyde or ethylene oxide gas sterilization. No further cases of RGM infections complicating cosmetic surgical procedures in Caracas have been reported.

Reported by: J Torres, J Murillo, L Bofill, A Ríos, E Irausquin, R Isturiz, M Guzmán, L Rubino, M Cordido, Venezuelan Collaborative Infectious Diseases and Tropical Medicine Study Group, Caracas; Ministry of Health, Venezuela. Hospital Infections Program, National Center for Infectious Diseases; and an EIS Officer, CDC.

Editorial Note: This is the first official report to CDC of SSI caused by RGM following liposuction or liposculpture. Both surgical procedures involve small surgical wounds with limited environmental exposure; both require using cannulae for tissue suction. The underlying mechanism for the cluster of SSI described in this report was not determined. However, potential causes included contaminated tap water used in cleaning cannulae during liposuction or liposculpture or contamination of the quaternary ammonium solution used to disinfect these instruments.

Nosocomial infections associated with contaminated quaternary ammonium compounds that were used to disinfect patient-care supplies or equipment (e.g., cystoscopes, cardiac catheters, or surgical instruments) have been reported; none of these infections were caused by RGM. Quaternary ammonium compounds are used widely as low-level disinfectants (1). Surgical instruments used in liposuction and liposculpture procedures are critical items (i.e., intended to enter a normally sterile environment, sterile tissue, or the vasculature) according to the Spaulding Classification (2). Critical items should be sterilized between patient procedures.

Based on the risk for contamination of postsurgical wounds, aesthetic surgical procedures such as liposuction or liposculpture are considered clean wounds according to the classification system developed by the National Research Council (3). National Nosocomial Infections Surveillance (NNIS) system data indicate that among 5652 integumental surgical procedures (including aesthetic surgical procedures with risk index=0) performed during 1986–1996 in the United States, only 1.4% were complicated by SSI.

SSI caused by RGM following aesthetic surgical procedures is rare. Reports include infection following augmentation mammoplasty procedures (4,5) and an outbreak of infection following either face-lift or augmentation mammoplasty procedures that implicated using contaminated gentian violet skin-marking solution as the source of infection (6).

The Venezuelan Ministry of Health operates a national program for surveillance of antimicrobial resistance, and some of the large university hospitals occasionally provide rates of specific hospital-acquired infections. However, no active surveillance programs exist for SSI or systematic monitoring of tap water for microorganisms in health-care settings.

To prevent SSI in health-care settings, all surgical instruments used in liposuction or liposculpture procedures should be cleaned carefully after the procedure and sterilized in accordance with a validated reprocessing protocol provided by the medical

Rapidly Growing Mycobacterial Infection — Continued

device manufacturer. The exclusive use of low- or intermediate-level disinfectants to reprocess surgical instruments between patient procedures is inconsistent with the Food and Drug Administration guidance and recommended standards of practice (1,2).

References

1. Rutala W. Disinfection and sterilization of patient-care items. *Infect Control Hosp Epidemiol* 1996;17:377–84.
2. Spaulding EH. Chemical disinfection of medical and surgical materials. In: Lawrence CA, Block SS, eds. *Disinfection, sterilization, and preservation*. Philadelphia, Pennsylvania: Lea & Febiger, 1968:517–31.
3. National Academy of Sciences, National Research Council, Division of Medical Sciences, Ad Hoc Committee on Trauma. Postoperative wound infections: the influence of ultraviolet irradiation of the operating room and of various other factors. *Ann Surg* 1964;160:1–192.
4. Clegg HW, Foster MT, Sanders WE Jr, Baine WB. Infection due to organisms of the *Mycobacterium fortuitum* complex after augmentation mammoplasty: clinical and epidemiologic features. *J Infect Dis* 1983;147:427–33.
5. Wallace RJ, Steele LC, Labidi A, Silcox VA. Heterogeneity among isolates of rapidly growing mycobacteria responsible for infections following augmentation mammoplasty despite case clustering in Texas and other southern coastal states. *J Infect Dis* 1989;160:281–8.
6. Safranek TJ, Jarvis WR, Carson LA, et al. *Mycobacterium chelonae* wound infections after plastic surgery employing contaminated gentian violet skin-marking solution. *N Engl J Med* 1987;317:197–201.

Impact of Closure of a Sexually Transmitted Disease Clinic on Public Health Surveillance of Sexually Transmitted Diseases — Washington, D.C., 1995

In April 1995, a public sexually transmitted disease (STD) clinic in the northwest (NW) ward of the District of Columbia closed, leaving one public STD clinic in the southeast (SE) ward to provide public STD services for the entire city. This report summarizes an investigation by CDC following a request from the District of Columbia Department of Health's STD program to evaluate the impact of the NW STD clinic closure on STD case reports. The findings of this investigation indicate that the clinic closure resulted in a marked decrease in reported syphilis in the NW ward, and suggest that NW ward residents with syphilis and their partners may not have received proper diagnostic testing, therapy, and counseling.

To determine the number and characteristics of patients seen at the STD clinics, the health department's STD clinic reports from May 1, 1994, to April 30, 1996 (the 12 months before and the 12 months after the clinic closed) were reviewed. To assess ward- and clinic-specific trends in case reporting, syphilis and gonorrhea case reports in the health department's STD surveillance database were analyzed. For this analysis, data for the year before and the year after the clinic closed were available for primary and secondary (P&S) syphilis cases, and data for 4 months before and 4 months after the clinic closed were available for gonorrhea cases.

Compared with the 12-month period before the NW STD clinic closed, during the 12-month period after the clinic closed the overall number of patient visits at the health department's STD clinics decreased 37%, from 20,155 to 12,759. The reported cases of P&S syphilis decreased 23%, from 143 cases before the clinic closure to 110 cases after the closure. Among those residing in the NW ward, the number of

Impact of Clinic Closure — Continued

reported cases of P&S syphilis decreased 57%, from 44 cases to 19 cases. However, reported cases among persons residing in the SE ward increased 10% during the same period, from 52 cases to 57 cases. The number of reported cases among women residing in the NW ward did not change, whereas reported cases among women residing in the SE ward increased by 41%, from 22 to 31 (Figure 1). However, among men residing in the NW ward, the number of reported cases decreased 78% (from 32 to seven), and reported cases from men residing in the SE ward decreased 13% (from 30 to 26).

Cases of reported gonorrhea in the District of Columbia decreased 26%, from 6935 cases before the clinic closure to 5166 cases after the closure. The decline in reported gonorrhea cases was seen in all wards.

The numbers of P&S syphilis cases reported for NW residents at the SE STD clinic did not change substantially following the NW STD clinic closure. However, the number of gonorrhea cases reported among NW residents increased at the SE STD clinic from 22 cases to 153 cases.

Reported by: M Levy, MD, J Heath, District of Columbia Dept of Health. Epidemiology and Surveillance Br, Div of Sexually Transmitted Disease Prevention, National Center for HIV, STD, and TB Prevention, CDC.

Editorial Note: Although in the United States syphilis has declined to historically low levels (1), it remains a problem in the District of Columbia. In 1996, the city's syphilis rate was 13th highest for U.S. cities of >200,000 population (2). Data from this investigation suggest that closing the NW STD clinic resulted in unreported syphilis cases. Reporting of syphilis cases is essential if health departments are to ensure that patients and their sex partners are treated and counseled properly, that trends in disease are monitored effectively, and that outbreaks are identified and addressed promptly.

FIGURE 1. Number of persons with reported primary and secondary syphilis, by sex and ward of residence before and after closure of the northwest sexually transmitted disease clinic — Washington, D.C., May 1994–April 1996*



*The 12 months before and the 12 months after closure of the northwest clinic.

Impact of Clinic Closure — Continued

The substantial increase in syphilis among women residing in the SE ward following the clinic closure suggests that a simultaneous increase in the NW ward might not have been detected. Cases in the NW ward may have been missed because of limited access to STD care after the NW STD clinic closure. The number of cases reported among women residing in the NW ward did not decrease as it did for men, possibly because a higher proportion of women than men are tested for syphilis in health-care settings other than STD clinics (e.g., family planning and antenatal care).

The elimination of STD care can result in substantial decreases in STD clinic visits, laboratory testing, and chlamydia and gonorrhea case reports (3). In the District of Columbia, gonorrhea case reports declined, but decreases in all wards were similar. Differences between the specific behaviors of syphilis patients and gonorrhea patients may help to explain the differential impact on reporting. Syphilis patients are more likely than gonorrhea patients to have a greater number of unnamed sex partners and to engage in illicit-drug use and exchange of sex for drugs or money (4). Syphilis patients and their partners may have particular difficulty accessing the health-care system. The signs and symptoms of early syphilis in men often are transient and painless compared with the often persistent urethral discharge and dysuria of gonorrhea; thus, persons with syphilis may not seek health care as readily as persons with gonorrhea. The loss of a public STD clinic may have had a greater impact on the likelihood of identifying, locating, and treating syphilis patients than gonorrhea patients.

The findings in this report are subject to at least two limitations. First, patients who would have been served by the NW STD clinic may have sought STD services from other health-care facilities; however, if such cases were not reported to the health department their sex partners probably did not receive adequate follow-up. Second, although the clinic closure appears to be the most likely explanation for the sharp decline in reporting of syphilis cases among NW residents, other unmeasured factors might have affected the syphilis and gonorrhea rates in the NW ward and elsewhere in the city.

When considering closing any public facility providing health-care services, health departments should evaluate the potential impact on populations with high rates of disease. Specifically, they should assess the extent to which these patients can access the remaining health-care facilities and the capacity of these facilities to handle an increase in patient volume. In settings such as the District of Columbia, measures to increase syphilis case finding should be implemented by expanding routine syphilis serologic screening, strengthening partner notification activities, and improving patient education.

References


1. CDC. Primary and secondary syphilis—United States, 1997. *MMWR* 1998;47:493–7.
2. Division of STD Prevention. Sexually transmitted disease surveillance, 1996. Atlanta, Georgia: US Department of Health and Human Services, Public Health Service, CDC, 1997.
3. Kimball AM, Lafferty WE, Kassler WJ, Hundt A, MacCornack R, Bolan G. The impact of health care market changes on local decision making and STD care: experience in three counties. *Am J Prev Med* 1997;13(suppl 6):75–84.
4. Andrus JK, Fleming DW, Harger DR, et al. Partner notification: can it control epidemic syphilis? *Ann Intern Med* 1990;112:539–43.

Outbreak of Primary and Secondary Syphilis — Guilford County, North Carolina, 1996–1997

In 1996 and 1997, 153 cases of primary and secondary (P&S) syphilis were reported each year in Guilford County, North Carolina, a 147% increase from the 62 cases reported in 1994 (Figure 1). The incidence of P&S syphilis in Guilford County during 1996–1997 was 40.5 cases per 100,000 persons, substantially higher than the national health objective for 2000 of four cases per 100,000 (objective 19.3) (1). In comparison, the number of P&S syphilis cases in North Carolina declined 57% from 1994 to 1997 (Figure 1), to a rate of 10.9 per 100,000 in 1997. This report summarizes the results of an investigation conducted by the Guilford County Health Department (GCHD), the North Carolina Division of Epidemiology, and CDC, which suggest this ongoing outbreak has been associated with missed opportunities for syphilis screening and treatment in high-risk settings, increased exchange of sex for money or drugs, and substantial rates of coinfection with syphilis and human immunodeficiency virus (HIV) among those tested.

To assess factors associated with the epidemic, interviews were conducted with P&S syphilis patients, state and local health department staff members, clinicians, and community residents. Demographic data for all residents of Guilford County with reported cases of syphilis from January 1993 (when the present data registry system was initiated) through August 1997 were analyzed to assess trends in factors that might influence syphilis rates (e.g., access to medical care and adequacy of screening and treatment). Also reviewed were the contact index (the number of sex partners for whom information was sufficient to initiate efforts to locate the person divided by the

FIGURE 1. Number of cases of primary and secondary syphilis, by year of report — Guilford County and North Carolina, 1986–1997



Outbreak of Primary and Secondary Syphilis — Continued

number of persons with syphilis interviewed) and the treatment index (the number of persons treated as a result of partner notification divided by the number of persons interviewed). The roles of illicit-drug use and sex worker activity during the epidemic were assessed. HIV screening and prevalence data were used to assess the extent of HIV coinfection among P&S syphilis patients. Syphilis registry data were used to compare risk factors among P&S syphilis patients reported during the pre-epidemic period (January 1993–December 1995) with P&S syphilis patients reported during the epidemic period (January 1996–August 1997). Screening and prevalence data from the local jails were reviewed (2–6).

Seventy-three percent of Guilford County residents reside in two major cities: Greensboro (1990 population: 192,000) and High Point (1990 population: 74,000). Most (96%) reported P&S syphilis patients in Guilford County reside in these two cities. Of patients in Guilford County who had infectious syphilis from January 1996 through August 1997, 55% were men. The mean age of men with P&S syphilis was 34.5 years in 1993 and 37.2 years during January–August 1997 ($p=0.2$). The mean age of women with P&S syphilis increased significantly from 1993 (27.8 years) through August 1997 (33.3 years) ($p=0.01$).

Patients during the epidemic period were more likely to have used illicit drugs at some time since 1978 (odds ratio [OR]=1.9; 95% confidence interval [CI]=1.1–3.3) and to have exchanged sex for drugs or money during the preceding year (OR=2.1; 95% CI=1.4–3.3) and were less likely to have been tested for HIV (18.6%) than patients before the epidemic period (27.8%; OR=0.6; 95% CI=0.4–0.9). Of P&S syphilis patients tested for HIV infection before and during the epidemic, 16% and 13%, respectively, were HIV infected. On the basis of local police records, prostitution arrests did not increase during 1993–1996, but crack cocaine-related arrests increased 69%.

Public sexually transmitted diseases clinical care appeared to meet the needs of persons seeking care during the epidemic in Greensboro. The contact index was 2.0 in 1993 and 1.7 in 1996, indicating fewer sex partners named per patient interviewed in 1996. However, the treatment index was 0.9 in 1993 and 1.0 in 1996, indicating more patients and contacts were treated for syphilis or preventively treated in 1996.

At the Guilford County jail, full health assessments were offered after 10–14 days of detainment. However, because of a rapid turnover and a high refusal rate, most detainees were not screened. In 1996, 9.6% of those detained in the jail system were screened for syphilis and <1% were screened for HIV infection; 7.5% of syphilis tests and 3.3% of HIV tests were positive. During January–August 1997, 8.0% of detained inmates had a history, physical examination, and syphilis serology, of whom 13.3% had reactive syphilis serologic tests.

To control the increase in syphilis cases in Guilford County, the North Carolina HIV/STD Prevention and Care Section and GCHD, in collaboration with local community organizations, conducted a community intervention effort from July through September 1997. This intervention combined sex partner notification strategies, community outreach, and extended local clinical services to find and treat more patients with P&S syphilis and to educate the community about syphilis. Other prevention measures included alerting the local medical community; obtaining help from community-based organizations in identifying locations where at-risk persons are commonly found and increasing education, outreach, and screening at these locations; and increasing screening and treatment for syphilis at local settings where per-

Outbreak of Primary and Secondary Syphilis — Continued

sons at high risk may have been encountered (e.g., jails). Based on reported cases of P&S syphilis in Guilford County through August 1998, P&S syphilis is expected to decrease 38% in 1998 compared with 1997.

Reported by: H Gabel, MD, Guilford County Health Dept, Greensboro; E Foust, MPH, D Ogburn, N Engel, J Owen-O'Dowd, A Howerton-Privott, North Carolina HIV/STD Prevention and Care Section, North Carolina Dept of Health and Human Svcs; Greensboro Police Dept. Div of Sexually Transmitted Disease Prevention, National Center for HIV, STD, and TB Prevention, CDC.

Editorial Note: This investigation identified epidemiologic factors frequently associated with syphilis outbreaks in other urban areas of the United States: increased illicit-drug use and exchange of money or drugs for sex. This investigation also identified missed opportunities for rapid syphilis screening and treatment at the local jails. Previous studies have identified emergency departments (EDs) and jails as sites of high syphilis prevalence during epidemics (2–6). Many arrested persons lack medical insurance or have used hospital EDs at their last medical visit (2). Therefore, jails and EDs are potentially high-impact settings for rapid screening and treatment of patients at high risk for syphilis in areas with endemic or epidemic syphilis (2–6).

Increased cocaine arrests corroborated community perceptions of increased crack cocaine use in Guilford County before the onset of the P&S syphilis epidemic. Also, data on P&S syphilis patients during 1996–1997 document increased exchange of sex for drugs or money and an increase in injecting or other drug use, compared with patients during 1993–1995. The link between crack cocaine and injecting-drug use and high-risk sex behaviors has been reported previously (7).

The sequelae of syphilis are substantial, including facilitation of HIV transmission, congenital syphilis, and advanced syphilis lesions affecting the cardiovascular and central nervous systems. The high frequency of HIV infection among persons tested who also have P&S syphilis underscores the need to make HIV counseling, testing, and prevention a priority for patients with syphilis.

Syphilis elimination is a feasible goal in the United States as syphilis rates continue to decline nationally, but outbreaks of P&S syphilis and persisting endemic foci are major obstacles (8). Outbreaks, such as the one in Guilford County, emphasize the prevention strategies and activities needed to maintain national and local progress toward elimination of syphilis, including innovative public health responses tailored to meet the challenge of shifting community patterns of high-risk behaviors and associated new outbreaks of communicable diseases. In addition, findings from this outbreak suggest that strengthening and maintaining screening in jails may be a useful component of syphilis surveillance and early outbreak detection, even in areas with little or no recognized syphilis transmission.

References

1. US Department of Health and Human Services. Healthy people 2000 review, 1997. Hyattsville, Maryland: US Department of Health and Human Services, CDC, 1997.
2. Beltrami JF, Cohen DA, Hamrick JT, Farley TA. Rapid screening and treatment for sexually transmitted diseases in arrestees: a feasible control measure. *Am J Public Health* 1997;87:1423–6.
3. Blank S, McDonnell DD, Rubin SR, et al. New approaches to syphilis control: finding opportunities for syphilis treatment and congenital syphilis prevention in a women's correctional setting. *Sex Transm Dis* 1997;24:218–26.
4. Heimberger TS, Chang HGH, Birkhead GS, et al. High prevalence of syphilis detected through a jail screening program: a potential public health measure to address the syphilis epidemic. *Arch Intern Med* 1993;153:1799–804.

Outbreak of Primary and Secondary Syphilis — Continued

5. Cohen D, Scribner R, Clark J, Cory D. The potential role of custody facilities in controlling sexually transmitted diseases. *Am J Public Health* 1992;82:552–6.
6. Ernst AA, Martin DH. High syphilis rates among cocaine abusers identified in an emergency department. *Sex Transm Dis* 1993;20:66–9.
7. Wolfe H, Vranizan KM, Gorter RG, et al. Crack use and human immunodeficiency virus infection among San Francisco intravenous drug users. *Sex Transm Dis* 1992;19:111–4.
8. St. Louis M, Farley TA, Aral SO. Untangling the persistence of syphilis in the south. *Sex Transm Dis* 1996;23:1–4.

Fatal *Cercopithecine herpesvirus 1* (B Virus) Infection Following a Mucocutaneous Exposure and Interim Recommendations for Worker Protection

On December 10, 1997, a 22-year-old female worker at a primate center died from *Cercopithecine herpesvirus 1* (B virus) infection 42 days after biologic material (possibly fecal) from a rhesus macaque (*Macaca mulatta*) splashed into her right eye. This report summarizes the clinical features of her illness and the subsequent investigation by CDC in response to a technical assistance request from the Occupational Safety and Health Administration (OSHA) and presents interim recommendations to prevent ocular splash exposures. This investigation documented the hazard of ocular splashes and indicated that dendritic corneal lesions, such as herpetic skin vesicles, are not always present in B virus infection (1).

The exposure occurred on October 29, 1997, while the worker moved the animal within cages during a routine capture of free-ranging monkeys. She was not wearing protective eyewear because the activities in which she was engaged involved caged macaques, and the activities were judged by the primate center to carry a low risk for exposure to B virus. Following the exposure, the worker wiped her eye with a paper towel and, approximately 45 minutes later, irrigated the eye for 2–3 minutes with tap water but did not file an incident report. The monkey involved was not identified.

On November 8, the worker's eye was red and swollen. At the emergency department (ED) of a medical center affiliated with the same university as the primate center, she informed the physician that she worked with nonhuman primates and may have been exposed to B virus. Dendritic corneal lesions typical of ocular herpes infections were not observed by Wood's lamp examination. The ED physician consulted the B virus protocol in place in the ED and then consulted an infectious diseases specialist by telephone. On the basis of the reported circumstances of the contact and the absence of previous recognized transmission of B virus following mucocutaneous exposure, the infectious diseases specialist concluded that B virus infection was unlikely but recommended follow-up with the infectious diseases clinic within the next few days. The ED physician prescribed sulfonamide eye drops.

An appointment at the infectious diseases clinic was not available immediately. On November 11, the worker called her primary-care physician for a referral because her eye symptoms were worsening. The physician referred her to an ophthalmologist, who elicited history of a recent cat scratch and prescribed doxycycline for suspected Parinaud's oculoglandular syndrome secondary to cat-scratch fever. Routine eye cultures were obtained. Confirmatory serologic testing for *Bartonella* species, also ordered during the visit, subsequently was negative.

Fatal Cercopithecine herpesvirus Infection — Continued

On November 13, the worker sought care from another ophthalmologist because of increased right retro-orbital pain and onset of photophobia, anorexia, nausea, and abdominal pain. After reconsultation with the infectious diseases specialist, the worker was immediately hospitalized for suspected B virus infection. The worker's temperature, normal on admission, reached 101.4 F (38.6 C) during the first day of hospitalization. Physical examination identified a swollen right orbit with conjunctivitis and one small tender right preauricular lymph node. Laboratory examination of urine found trace proteinuria. Cerebrospinal fluid (CSF) analysis identified 8 white blood cells per milliliter (83% lymphocytes [normal: 0–10 cells, 100% mononuclear]). Serum for Western blot testing and CSF specimens and eye swabs for B virus culture were sent to the B Virus Research and Resource Laboratory. All previously collected eye cultures were retrieved from commercial laboratories to minimize biosafety hazards to laboratory workers.

Acyclovir therapy (15 mg/kg intravenously every 8 hours) was started within 2 hours of hospital admission. On November 14, therapy was changed to ganciclovir (5 mg/kg every 12 hours) when a vesicular eruption was noted in the distribution of the first and second branches of the right trigeminal nerve. Magnetic resonance imaging (MRI) of the head was normal. The vesicles resolved over the following week. A sharp mid-cervical/high thoracic back discomfort occurred on November 19 but subsided over an 8-hour period. All symptoms resolved, and on November 24 the worker was discharged on outpatient intravenous (IV) ganciclovir therapy.

Despite uninterrupted ganciclovir therapy, on November 25 the worker woke with right foot weakness, inability to urinate, and lower abdominal pain, followed by a rapidly progressive ascending myelitis. The hospital readmission examination found profound right leg weakness, moderate left leg weakness, decreased hand grip strength bilaterally, and urinary retention. MRI revealed abnormalities extending from the cervical spinal cord to the upper thoracic cord. The worker was intubated electively within 13 hours and developed flaccid paralysis from C2 caudad.

The diagnosis of postviral acute demyelinating encephalomyelitis was considered by neurology consultants, and a short course of plasmapheresis and steroids was administered. On November 30 seizure activity (involuntary facial and eye movements) developed, and foscarnet, usually not recommended for B virus infection because of its toxicity, was added to ongoing ganciclovir therapy. During December 1–9, the worker developed nosocomial pneumonia with bacteremia, followed by adult respiratory distress syndrome. Repeat MRI revealed abnormalities extending from midbrain through the thoracic spinal cord. On December 10, the worker died from refractory respiratory failure.

Eye and CSF cultures obtained in the hospital on November 13 and November 14 were negative for B virus when tested at the B Virus Research and Resource Laboratory. Serum collected November 13 and November 21 and tested for reactivity to B virus by Western blot showed indeterminate and positive reactivity, respectively, confirming B virus infection.

Reported by: C Perlino, MD, Emory Univ School of Medicine; J Hilliard, PhD, B Virus Research and Resource Laboratory, Georgia State Univ, Atlanta; J Koehler, DVM, Div of Public Health, Georgia Dept of Human Resources. National Institute for Occupational Safety and Health; National Center for Infectious Diseases; and Office of Health and Safety, CDC.

Fatal Cercopithecine herpesvirus Infection — Continued

Editorial Note: *C. herpesvirus 1* (B virus) causes persistent latent infections in $\geq 70\%$ of captive adult macaques (2) but not other primates. During intermittent reactivations, the macaque may shed B virus from the buccal mucosa, urogenital tract, and in conjunctival fluid (2). Reactivations may be asymptomatic or accompanied by clustered vesicles on an erythematous base.

This is the first report of a worker developing a recognized B virus infection following mucocutaneous exposure without injury. Previously reported human infections usually have been attributed to macaque bites or scratches, injuries from needles used near a macaque's mucous membranes or central nervous system, or contact with infective biologic materials from macaques (3–5). One human-to-human transmission has been identified (6). The incubation period in humans has been as short as 2 days but more frequently is 2–5 weeks. Previously reported patients infected with B virus who were treated aggressively with either IV acyclovir or ganciclovir after onset of symptoms but before respiratory arrest or coma have survived (3). The death of this patient despite aggressive antiviral therapy may have resulted from factors related to the route of virus inoculation, the virulence of the virus infecting the patient, the patient's immune response, or timing of initiation of treatment following the exposure.

Interim Recommendations to Prevent Ocular Splash Exposures

Preventing worker exposure to biohazardous material is the best protection against infection. Reviews of injuries and biohazard exposures among workers exposed to nonhuman primates suggest that mucocutaneous contact with nonhuman primate body fluids is common; 16 (94%) of 17 contacts with primate body fluids in one survey involved ocular exposure (6,7). Each institution working with macaques should develop a written comprehensive personal protective equipment (PPE) program based on thorough hazard assessments of all work procedures, potential routes of exposure (e.g., bites, scratches, or mucosal exposures) and potential adverse health outcomes. This plan should clearly identify the PPE required for each task or working area and address training, inspection, maintenance, and periodic assessment of program effectiveness.

Previous recommendations for preventing B virus infections in humans advise presuming that all macaques are infected with B virus and protecting workers with a faceshield (or surgical mask and goggles or glasses) when handling uncaged active macaques (3,8). The incident described in this report indicates that proper eye protection also should be mandatory during activities such as entering areas containing macaques, conducting captures, and transporting caged macaques. Other activities where eye protection is necessary should be determined by the hazard assessment. All personnel who work in situations determined to be hazardous should wear eyewear conforming to established standards for eye and splash protection (9). Personal eyeglasses are not PPE.

Protective goggles designed for splash protection (available with antifog lenses for humid environments and in models that preserve peripheral vision) should be worn to protect the eyes against splash hazards in combination with a mask designed to protect other mucous membranes. Faceshields are commonly considered secondary eye protectors that are worn in combination with protective goggles (9,10). Although previous guidelines indicate a faceshield may be sufficient, ocular exposures have oc-

Fatal Cercopithecine herpesvirus Infection — Continued

curred to workers wearing faceshields, including to a worker who was wearing a combination surgical mask/faceshield while moving a macaque within cages. To minimize the potential for mucous membrane exposure, faceshields must prevent droplet splashes to the head from running down into the eyes and prevent mucous membrane exposure around the edges (sides, top, and bottom to below the chin) (10). Decisions to use faceshields as the sole means for preventing ocular exposure should only be made after full consideration of both the limitations of faceshields and regulatory (OSHA) considerations.

Exposure Management


If exposure prevention fails, the adequacy and timeliness of wound or exposure decontamination procedures are critical factors determining the risk for infection. Institutions that house or conduct procedures involving nonhuman primates or potentially contaminated tissues should develop institution-specific postexposure procedures (3,8). Such procedures would eliminate institutional barriers to patient access and ensure appropriate diagnostic testing and infection control. First, animal handlers should be instructed to cleanse immediately and thoroughly all bites, scratches, and/or mucosal surfaces or abraded skin exposed to macaque biologic materials and to report these exposures immediately (3). Following an exposure to the eye, existing guidelines recommend immediately flushing the eye with water for at least 15 minutes (3). Second, postexposure procedures also should provide potentially exposed workers with direct and rapid access to a local medical consultant knowledgeable about B virus and other biohazards associated with nonhuman primates. The employer should ensure that direct access to the knowledgeable consultant is available immediately following exposures and at any time the worker is concerned that potential occupational exposure to B virus may be relevant to worker symptoms. Finally, postexposure procedures also should include routing diagnostic specimens to the B Virus Research and Resource Laboratory, now at Georgia State University in Atlanta. These interim recommendations will be reviewed and may be revised or augmented following additional consideration by a working group convened by Office of Health and Safety, CDC.

References

1. Fierer J, Bazely P, Braude AI. Herpes B virus encephalomyelitis presenting as ophthalmic zoster: a possible latent infection reactivated. *Ann Intern Med* 1973;79:225-8.
2. Weigler BJ. Biology of B virus in macaque and human hosts: a review. *Clin Infect Dis* 1992; 14:555-67.
3. Holmes GP, Chapman LE, Stewart JA, Straus SE, Hilliard JK, Davenport DS. Guidelines for the prevention and treatment of B-virus infections in exposed persons. *Clin Infect Dis* 1995; 20:421-39.
4. Wells DL, Lipper SL, Hilliard JK, et al. *Herpesvirus simiae* contamination of primary rhesus monkey kidney cell cultures: CDC recommendations to minimize risks to laboratory personnel. *Diagn Microbiol Infect Dis* 1989;12:333-5.
5. CDC/National Institutes of Health. Biosafety in microbiological and biomedical laboratories. 3rd ed. Atlanta, Georgia: US Department of Health and Human Services, Public Health Service, CDC and the National Institutes of Health, May 1993; DHHS publication no. (CDC)93-8395.
6. Holmes GP, Hilliard JK, Klontz KC, et al. B virus (*Herpesvirus simiae*) infection in humans: epidemiologic investigation of a cluster. *Ann Intern Med* 1990;112:833-9.
7. bin Zakaria M, Lerche NW, Chomel BB, Kass PH. Accidental injuries associated with nonhuman primate exposure at two regional primate research centers (USA): 1988-93. *Lab Anim Sci* 1996;46:298-304.

(Continued on page 1083)

FIGURE I. Selected notifiable disease reports, comparison of provisional 4-week totals ending December 12, 1998, with historical data — United States



*Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

TABLE I. Summary — provisional cases of selected notifiable diseases, United States, cumulative, week ending December 12, 1998 (49th Week)

| | Cum. 1998 | | Cum. 1998 |
|---|-----------|---|-----------|
| Anthrax | - | Plague | 8 |
| Brucellosis | 55 | Poliomyelitis, paralytic | 1 |
| Cholera | 12 | Psittacosis | 48 |
| Congenital rubella syndrome | 3 | Rabies, human | - |
| Cryptosporidiosis* | 3,018 | Rocky Mountain spotted fever (RMSF) | 325 |
| Diphtheria | 1 | Streptococcal disease, invasive Group A | 1,964 |
| Encephalitis: California* | 86 | Streptococcal toxic-shock syndrome* | 49 |
| eastern equine* | 3 | Syphilis, congenital [¶] | 399 |
| St. Louis* | 26 | Tetanus | 34 |
| western equine* | - | Toxic-shock syndrome | 123 |
| Hansen Disease | 100 | Trichinosis | 13 |
| Hantavirus pulmonary syndrome* [†] | 19 | Typhoid fever | 318 |
| Hemolytic uremic syndrome, post-diarrheal* | 80 | Yellow fever | - |
| HIV infection, pediatric* [§] | 243 | | |

-:no reported cases

*Not notifiable in all states.

[†] Updated weekly from reports to the Division of Viral and Rickettsial Diseases, National Center for Infectious Diseases (NCID).

[§] Updated monthly from reports to the Division of HIV/AIDS Prevention—Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention (NCHSTP), last update November 29, 1998.

[¶] Updated from reports to the Division of STD Prevention, NCHSTP.

TABLE II. Provisional cases of selected notifiable diseases, United States, weeks ending December 12, 1998, and December 6, 1997 (49th Week)

| Reporting Area | AIDS | | Chlamydia | | <i>Escherichia coli</i> O157:H7 | | Gonorrhea | | Hepatitis C/NA,NB | |
|----------------|------------|-----------|-----------|-----------|---------------------------------|-----------|-----------|-----------|-------------------|-----------|
| | Cum. 1998* | Cum. 1997 | Cum. 1998 | Cum. 1997 | NETSS† | PHLIS‡ | Cum. 1998 | Cum. 1997 | Cum. 1998 | Cum. 1997 |
| | | | | | Cum. 1998 | Cum. 1998 | | | | |
| UNITED STATES | 42,564 | 53,705 | 539,229 | 443,010 | 2,853 | 1,883 | 321,493 | 280,265 | 4,679 | 3,279 |
| NEW ENGLAND | 1,688 | 2,248 | 17,046 | 17,019 | 328 | 260 | 4,852 | 5,591 | 106 | 53 |
| Maine | 28 | 51 | 963 | 949 | 36 | - | 63 | 63 | - | - |
| N.H. | 40 | 39 | 883 | 770 | 46 | 45 | 83 | 93 | - | - |
| Vt. | 19 | 32 | 378 | 400 | 21 | 17 | 34 | 48 | 3 | 4 |
| Mass. | 862 | 803 | 7,907 | 6,909 | 148 | 147 | 2,167 | 1,984 | 100 | 42 |
| R.I. | 118 | 145 | 2,172 | 1,915 | 13 | 1 | 395 | 404 | 3 | 7 |
| Conn. | 621 | 1,178 | 4,743 | 6,076 | 64 | 50 | 2,110 | 2,999 | - | - |
| MID. ATLANTIC | 11,418 | 16,079 | 63,627 | 53,698 | 284 | 73 | 37,772 | 36,192 | 338 | 305 |
| Upstate N.Y. | 1,323 | 2,379 | N | N | 214 | - | 6,190 | 6,192 | 251 | 228 |
| N.Y. City | 6,564 | 8,583 | 32,149 | 25,759 | 8 | 12 | 14,403 | 13,729 | - | - |
| N.J. | 2,025 | 3,119 | 10,385 | 9,612 | 62 | 51 | 7,137 | 7,097 | - | - |
| Pa. | 1,506 | 1,998 | 21,093 | 18,327 | N | 10 | 10,042 | 9,174 | 87 | 77 |
| E.N. CENTRAL | 3,063 | 4,078 | 94,238 | 60,172 | 446 | 321 | 65,418 | 38,973 | 480 | 509 |
| Ohio | 640 | 837 | 24,190 | 21,177 | 123 | 65 | 15,745 | 13,827 | 8 | 19 |
| Ind. | 472 | 485 | 4,656 | 8,759 | 102 | 49 | 4,704 | 5,722 | 7 | 12 |
| Ill. | 1,195 | 1,710 | 25,448 | U | 109 | 58 | 21,097 | U | 33 | 85 |
| Mich. | 578 | 801 | 20,469 | 19,728 | 112 | 62 | 15,424 | 14,718 | 432 | 367 |
| Wis. | 178 | 245 | 19,475 | 10,508 | N | 87 | 8,448 | 4,706 | - | 26 |
| W.N. CENTRAL | 832 | 1,099 | 29,508 | 30,987 | 486 | 384 | 15,209 | 13,818 | 279 | 58 |
| Minn. | 163 | 191 | 6,123 | 6,286 | 202 | 202 | 2,361 | 2,248 | 11 | 4 |
| Iowa | 63 | 99 | 2,063 | 4,199 | 93 | 58 | 660 | 1,074 | 8 | 27 |
| Mo. | 402 | 557 | 11,550 | 11,421 | 53 | 61 | 8,527 | 7,175 | 250 | 10 |
| N. Dak. | 5 | 10 | 849 | 834 | 12 | 15 | 71 | 69 | - | 3 |
| S. Dak. | 15 | 8 | 1,486 | 1,319 | 35 | 34 | 212 | 156 | - | - |
| Nebr. | 65 | 90 | 2,667 | 2,575 | 58 | - | 1,119 | 1,134 | 5 | 2 |
| Kans. | 119 | 144 | 4,770 | 4,353 | 33 | 14 | 2,259 | 1,962 | 5 | 12 |
| S. ATLANTIC | 11,132 | 13,315 | 107,017 | 88,611 | 257 | 155 | 88,126 | 87,153 | 184 | 237 |
| Del. | 154 | 211 | 2,461 | 61 | - | 2 | 1,454 | 1,205 | - | - |
| Md. | 1,489 | 1,800 | 6,759 | 7,044 | 38 | 14 | 9,024 | 10,827 | 22 | 10 |
| D.C. | 809 | 1,016 | N | N | 1 | - | 3,305 | 4,116 | - | - |
| Va. | 910 | 1,113 | 12,706 | 11,077 | N | 42 | 8,939 | 8,268 | 12 | 25 |
| W. Va. | 79 | 117 | 2,439 | 2,772 | 13 | 10 | 784 | 863 | 7 | 16 |
| N.C. | 752 | 796 | 20,648 | 16,285 | 56 | 46 | 17,850 | 16,190 | 20 | 48 |
| S.C. | 719 | 746 | 16,770 | 11,801 | 17 | 12 | 10,728 | 10,842 | 11 | 37 |
| Ga. | 1,174 | 1,600 | 21,585 | 14,830 | 76 | - | 18,085 | 17,236 | 9 | - |
| Fla. | 5,046 | 5,916 | 23,649 | 24,741 | 56 | 29 | 17,957 | 17,606 | 103 | 101 |
| E.S. CENTRAL | 1,684 | 1,901 | 36,245 | 33,085 | 114 | 39 | 35,609 | 33,126 | 186 | 332 |
| Ky. | 263 | 340 | 6,083 | 5,950 | 33 | - | 3,577 | 3,787 | 20 | 13 |
| Tenn. | 622 | 738 | 12,424 | 11,821 | 53 | 33 | 10,804 | 10,399 | 159 | 222 |
| Ala. | 456 | 511 | 9,705 | 8,037 | 25 | 2 | 12,324 | 11,215 | 5 | 11 |
| Miss. | 343 | 312 | 8,033 | 7,277 | 3 | 4 | 8,904 | 7,725 | 2 | 86 |
| W.S. CENTRAL | 5,140 | 5,650 | 75,283 | 65,006 | 125 | 24 | 45,742 | 42,251 | 414 | 472 |
| Ark. | 189 | 216 | 3,665 | 2,556 | 11 | 10 | 3,640 | 4,323 | 10 | 14 |
| La. | 878 | 1,016 | 14,301 | 9,603 | 5 | 7 | 12,326 | 9,316 | 114 | 213 |
| Okla. | 272 | 274 | 8,749 | 6,942 | 24 | 7 | 4,895 | 4,498 | 16 | 7 |
| Tex. | 3,801 | 4,144 | 48,568 | 45,905 | 85 | - | 24,881 | 24,114 | 274 | 238 |
| MOUNTAIN | 1,479 | 1,548 | 30,763 | 28,234 | 340 | 238 | 8,528 | 7,701 | 336 | 312 |
| Mont. | 28 | 40 | 1,205 | 1,092 | 16 | - | 44 | 60 | 7 | 21 |
| Idaho | 28 | 50 | 1,920 | 1,559 | 40 | 24 | 168 | 147 | 87 | 79 |
| Wyo. | 3 | 14 | 626 | 585 | 53 | 55 | 29 | 50 | 66 | 73 |
| Colo. | 286 | 366 | 8,134 | 6,943 | 90 | 69 | 2,193 | 2,123 | 33 | 34 |
| N. Mex. | 202 | 164 | 3,699 | 3,578 | 19 | 20 | 894 | 812 | 93 | 60 |
| Ariz. | 589 | 375 | 10,243 | 10,132 | 21 | 26 | 3,717 | 3,477 | 8 | 25 |
| Utah | 128 | 140 | 2,050 | 1,656 | 79 | 21 | 217 | 262 | 23 | 5 |
| Nev. | 215 | 399 | 2,886 | 2,689 | 22 | 23 | 1,266 | 770 | 19 | 15 |
| PACIFIC | 6,128 | 7,787 | 85,502 | 66,198 | 473 | 389 | 20,237 | 15,460 | 2,356 | 1,001 |
| Wash. | 390 | 608 | 10,389 | 8,736 | 108 | 127 | 1,861 | 1,803 | 22 | 28 |
| Oreg. | 166 | 284 | 5,610 | 4,701 | 104 | 99 | 829 | 700 | 6 | 3 |
| Calif. | 5,396 | 6,757 | 65,469 | 49,630 | 254 | 147 | 16,801 | 12,125 | 2,273 | 805 |
| Alaska | 17 | 46 | 1,782 | 1,444 | 7 | - | 300 | 357 | 1 | - |
| Hawaii | 159 | 92 | 2,252 | 1,687 | N | 16 | 446 | 475 | 54 | 165 |
| Guam | 1 | 2 | 201 | 193 | N | - | 24 | 27 | - | - |
| P.R. | 1,602 | 1,974 | U | U | 6 | U | 356 | 519 | - | - |
| V.I. | 31 | 94 | N | U | N | U | U | U | U | U |
| Amer. Samoa | - | - | U | U | N | U | U | U | U | U |
| C.N.M.I. | - | 1 | N | N | N | U | 28 | 23 | - | 2 |

N: Not notifiable U: Unavailable -: no reported cases C.N.M.I.: Commonwealth of Northern Mariana Islands

*Updated monthly from reports to the Division of HIV/AIDS Prevention-Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention, last update November 29, 1998.

†National Electronic Telecommunications System for Surveillance.

‡Public Health Laboratory Information System.

TABLE II. (Cont'd.) Provisional cases of selected notifiable diseases, United States, weeks ending December 12, 1998, and December 6, 1997 (49th Week)

| Reporting Area | Legionellosis | | Lyme Disease | | Malaria | | Syphilis (Primary & Secondary) | | Tuberculosis | | Rabies, Animal |
|----------------|---------------|--------------|--------------|--------------|--------------|--------------|-----------------------------------|--------------|---------------|--------------|-------------------|
| | Cum. 1998 | Cum. 1997 | Cum. 1998 | Cum. 1997 | Cum. 1998 | Cum. 1997 | Cum. 1998 | Cum. 1997 | Cum. 1998* | Cum. 1997 | Cum. 1998 |
| UNITED STATES | 1,239 | 1,009 | 12,273 | 11,422 | 1,298 | 1,741 | 6,759 | 7,907 | 13,945 | 16,688 | 6,462 |
| NEW ENGLAND | 83 | 79 | 2,636 | 2,898 | 59 | 85 | 71 | 130 | 433 | 414 | 1,379 |
| Maine | 1 | 3 | 12 | 8 | 5 | 1 | 1 | 2 | 10 | 19 | 214 |
| N.H. | 7 | 7 | 45 | 37 | 5 | 10 | 2 | - | 13 | 15 | 77 |
| Vt. | 7 | 13 | 11 | 8 | 1 | 2 | 4 | - | 4 | 6 | 65 |
| Mass. | 32 | 27 | 749 | 285 | 16 | 31 | 44 | 66 | 248 | 235 | 489 |
| R.I. | 21 | 12 | 654 | 385 | 14 | 10 | 1 | 2 | 52 | 33 | 97 |
| Conn. | 15 | 17 | 1,165 | 2,175 | 18 | 31 | 19 | 60 | 106 | 106 | 437 |
| MID. ATLANTIC | 289 | 226 | 8,080 | 6,699 | 322 | 496 | 278 | 377 | 2,845 | 2,944 | 1,499 |
| Upstate N.Y. | 101 | 70 | 3,974 | 2,818 | 89 | 72 | 35 | 41 | 365 | 424 | 1,034 |
| N.Y. City | 28 | 24 | 29 | 173 | 151 | 303 | 74 | 81 | 1,410 | 1,487 | U |
| N.J. | 15 | 30 | 1,690 | 1,827 | 52 | 84 | 78 | 149 | 574 | 641 | 204 |
| Pa. | 145 | 102 | 2,387 | 1,881 | 30 | 37 | 91 | 106 | 496 | 392 | 261 |
| E.N. CENTRAL | 409 | 327 | 166 | 578 | 126 | 162 | 1,134 | 593 | 1,199 | 1,697 | 130 |
| Ohio | 126 | 116 | 84 | 37 | 15 | 19 | 125 | 204 | 88 | 239 | 57 |
| Ind. | 118 | 53 | 62 | 33 | 11 | 17 | 244 | 162 | 139 | 146 | 12 |
| Ill. | 37 | 34 | 8 | 13 | 41 | 67 | 455 | U | 604 | 898 | 16 |
| Mich. | 78 | 84 | 12 | 27 | 48 | 43 | 194 | 128 | 350 | 297 | 35 |
| Wis. | 50 | 40 | U | 468 | 11 | 16 | 116 | 99 | 18 | 117 | 10 |
| W.N. CENTRAL | 75 | 57 | 214 | 152 | 92 | 59 | 123 | 167 | 391 | 528 | 677 |
| Minn. | 8 | 3 | 174 | 111 | 56 | 29 | 9 | 16 | 147 | 136 | 119 |
| Iowa | 12 | 9 | 23 | 7 | 8 | 10 | - | 7 | 51 | 57 | 147 |
| Mo. | 24 | 21 | 2 | 27 | 15 | 11 | 93 | 111 | 95 | 218 | 27 |
| N. Dak. | - | 2 | - | - | 2 | 3 | - | - | 10 | 12 | 138 |
| S. Dak. | 3 | 2 | - | 1 | 1 | 1 | 1 | 1 | 17 | 10 | 151 |
| Nebr. | 20 | 15 | 5 | 2 | 1 | 1 | 7 | 3 | 28 | 20 | 7 |
| Kans. | 8 | 5 | 10 | 4 | 9 | 4 | 13 | 29 | 43 | 75 | 88 |
| S. ATLANTIC | 140 | 117 | 863 | 736 | 312 | 310 | 2,438 | 3,296 | 1,946 | 3,182 | 1,856 |
| Del. | 13 | 11 | 45 | 109 | 3 | 5 | 21 | 22 | 18 | 32 | 49 |
| Md. | 28 | 21 | 596 | 468 | 86 | 81 | 619 | 870 | 261 | 292 | 423 |
| D.C. | 8 | 4 | 4 | 9 | 19 | 20 | 73 | 105 | 98 | 95 | - |
| Va. | 20 | 26 | 67 | 62 | 56 | 65 | 146 | 225 | 280 | 305 | 534 |
| W. Va. | N | N | 13 | 10 | 2 | 1 | 3 | 3 | 41 | 49 | 76 |
| N.C. | 14 | 14 | 57 | 34 | 29 | 19 | 691 | 949 | 448 | 409 | 136 |
| S.C. | 11 | 8 | 7 | 2 | 6 | 17 | 309 | 348 | 234 | 316 | 143 |
| Ga. | 8 | 1 | 5 | 7 | 37 | 48 | 272 | 493 | 496 | 588 | 288 |
| Fla. | 36 | 32 | 69 | 35 | 74 | 54 | 304 | 281 | 70 | 1,096 | 207 |
| E.S. CENTRAL | 64 | 55 | 93 | 88 | 31 | 38 | 1,128 | 1,597 | 1,038 | 1,226 | 259 |
| Ky. | 25 | 11 | 25 | 16 | 7 | 12 | 103 | 125 | 158 | 181 | 31 |
| Tenn. | 24 | 33 | 44 | 40 | 16 | 10 | 526 | 693 | 396 | 430 | 135 |
| Ala. | 8 | 4 | 20 | 11 | 6 | 10 | 270 | 395 | 302 | 390 | 91 |
| Miss. | 7 | 7 | 4 | 21 | 2 | 6 | 229 | 384 | 182 | 225 | 2 |
| W.S. CENTRAL | 41 | 33 | 36 | 93 | 47 | 56 | 1,000 | 1,253 | 2,100 | 2,405 | 135 |
| Ark. | - | 2 | 7 | 25 | 1 | 5 | 103 | 152 | 143 | 171 | 31 |
| La. | 4 | 6 | 7 | 5 | 15 | 15 | 409 | 344 | 274 | 257 | - |
| Okla. | 12 | 2 | 2 | 28 | 4 | 8 | 121 | 115 | 147 | 190 | 104 |
| Tex. | 25 | 23 | 20 | 35 | 27 | 28 | 367 | 642 | 1,536 | 1,787 | - |
| MOUNTAIN | 74 | 62 | 23 | 15 | 62 | 65 | 211 | 169 | 421 | 509 | 213 |
| Mont. | 2 | 1 | - | - | 1 | 2 | - | - | 19 | 16 | 53 |
| Idaho | 2 | 2 | 6 | 4 | 8 | - | 2 | 1 | 13 | 11 | - |
| Wyo. | 1 | 1 | 1 | 3 | - | 2 | 1 | - | 4 | 2 | 63 |
| Colo. | 19 | 18 | 5 | - | 19 | 30 | 11 | 15 | U | 78 | 39 |
| N. Mex. | 2 | 3 | 4 | 1 | 12 | 8 | 22 | 8 | 64 | 63 | 6 |
| Ariz. | 19 | 12 | 1 | 4 | 9 | 11 | 160 | 130 | 195 | 207 | 19 |
| Utah | 22 | 18 | - | 1 | 1 | 3 | 4 | 5 | 49 | 31 | 27 |
| Nev. | 7 | 7 | 6 | 2 | 12 | 9 | 11 | 10 | 77 | 101 | 6 |
| PACIFIC | 64 | 53 | 162 | 163 | 247 | 470 | 376 | 325 | 3,572 | 3,783 | 314 |
| Wash. | 12 | 8 | 7 | 10 | 20 | 48 | 27 | 10 | 198 | 280 | - |
| Oreg. | 1 | - | 21 | 19 | 16 | 25 | 6 | 9 | 125 | 136 | 7 |
| Calif. | 49 | 44 | 133 | 132 | 203 | 383 | 341 | 304 | 3,046 | 3,141 | 284 |
| Alaska | 1 | - | 1 | 2 | 3 | 3 | 1 | 1 | 49 | 66 | 23 |
| Hawaii | 1 | 1 | - | - | 5 | 11 | 1 | 1 | 154 | 160 | - |
| Guam | 2 | - | - | - | 1 | - | 1 | 3 | 36 | 13 | - |
| P.R. | - | - | - | - | - | 5 | 172 | 236 | 68 | 212 | 51 |
| V.I. | U | U | U | U | U | U | U | U | U | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | - | - | - | - | - | 164 | 11 | 77 | 20 | - |

N: Not notifiable

U: Unavailable

-: no reported cases

TABLE III. Provisional cases of selected notifiable diseases preventable by vaccination, United States, weeks ending December 12, 1998, and December 6, 1997 (49th Week)

| Reporting Area | <i>H. influenzae</i> , invasive | | Hepatitis (Viral), by type | | | | Measles (Rubeola) | | | | | |
|----------------|---------------------------------|-----------|----------------------------|-----------|-----------|-----------|-------------------|-----------|-----------|-----------|-----------|-----------|
| | Cum. 1998* | Cum. 1997 | A | | B | | Indigenous | | Imported† | | Total | |
| | | | Cum. 1998 | Cum. 1997 | Cum. 1998 | Cum. 1997 | 1998 | Cum. 1998 | 1998 | Cum. 1998 | Cum. 1998 | Cum. 1997 |
| UNITED STATES | 956 | 1,008 | 20,803 | 26,591 | 8,195 | 8,912 | 5 | 67 | - | 25 | 92 | 130 |
| NEW ENGLAND | 65 | 59 | 264 | 612 | 179 | 175 | - | 1 | - | 2 | 3 | 19 |
| Maine | 3 | 5 | 20 | 59 | 5 | 6 | - | - | - | - | - | 1 |
| N.H. | 9 | 11 | 14 | 34 | 19 | 17 | - | - | - | - | - | 1 |
| Vt. | 9 | 3 | 16 | 14 | 6 | 11 | - | - | - | 1 | 1 | - |
| Mass. | 37 | 35 | 106 | 248 | 56 | 73 | - | 1 | - | 1 | 2 | 16 |
| R.I. | 6 | 3 | 17 | 127 | 68 | 16 | - | - | - | - | - | - |
| Conn. | 1 | 2 | 91 | 130 | 25 | 52 | - | - | - | - | - | 1 |
| MID. ATLANTIC | 141 | 155 | 1,402 | 2,003 | 1,043 | 1,296 | - | 8 | - | 6 | 14 | 26 |
| Upstate N.Y. | 63 | 51 | 349 | 348 | 280 | 297 | - | 1 | - | 1 | 2 | 5 |
| N.Y. City | 26 | 41 | 362 | 875 | 266 | 439 | - | - | - | - | - | 10 |
| N.J. | 46 | 44 | 321 | 297 | 181 | 235 | U | 7 | U | 1 | 8 | 3 |
| Pa. | 6 | 19 | 370 | 483 | 316 | 325 | - | - | - | 4 | 4 | 8 |
| E.N. CENTRAL | 154 | 158 | 3,452 | 2,846 | 1,507 | 1,417 | - | 12 | - | 3 | 15 | 10 |
| Ohio | 46 | 82 | 312 | 299 | 74 | 87 | - | - | - | 1 | 1 | - |
| Ind. | 40 | 18 | 326 | 310 | 744 | 94 | - | 2 | - | 1 | 3 | - |
| Ill. | 53 | 39 | 650 | 795 | 186 | 267 | - | 1 | - | - | 1 | 7 |
| Mich. | 8 | 18 | 1,993 | 1,269 | 454 | 427 | - | 9 | - | 1 | 10 | 2 |
| Wis. | 7 | 1 | 171 | 173 | 49 | 542 | - | - | - | - | - | 1 |
| W.N. CENTRAL | 89 | 58 | 1,268 | 2,064 | 401 | 459 | - | 1 | - | - | 1 | 17 |
| Minn. | 66 | 44 | 124 | 192 | 49 | 41 | - | - | - | - | - | 8 |
| Iowa | 3 | 6 | 394 | 442 | 56 | 40 | - | 1 | - | - | 1 | - |
| Mo. | 12 | 5 | 575 | 1,058 | 242 | 323 | - | - | - | - | - | 1 |
| N. Dak. | - | - | 3 | 10 | 4 | 5 | - | - | - | - | - | - |
| S. Dak. | 1 | 2 | 32 | 23 | 2 | 1 | - | - | - | - | - | 8 |
| Nebr. | 1 | 1 | 41 | 89 | 22 | 19 | - | - | - | - | - | - |
| Kans. | 6 | - | 99 | 250 | 26 | 30 | - | - | - | - | - | - |
| S. ATLANTIC | 182 | 153 | 1,923 | 1,965 | 1,117 | 1,148 | - | 3 | - | 5 | 8 | 15 |
| Del. | - | - | 6 | 29 | 4 | 6 | - | - | - | 1 | 1 | - |
| Md. | 52 | 56 | 318 | 179 | 151 | 158 | - | - | - | 1 | 1 | 2 |
| D.C. | - | - | 62 | 33 | 18 | 29 | - | - | - | - | - | 1 |
| Va. | 18 | 13 | 200 | 217 | 99 | 121 | - | - | - | 2 | 2 | 1 |
| W. Va. | 5 | 4 | 7 | 11 | 10 | 16 | - | - | - | - | - | - |
| N.C. | 24 | 21 | 123 | 196 | 228 | 245 | - | - | - | - | - | 2 |
| S.C. | 3 | 4 | 38 | 106 | 46 | 94 | - | - | - | - | - | 1 |
| Ga. | 46 | 31 | 647 | 621 | 139 | 126 | - | 1 | - | 1 | 2 | 1 |
| Fla. | 34 | 24 | 522 | 573 | 422 | 353 | - | 2 | - | - | 2 | 7 |
| E.S. CENTRAL | 57 | 54 | 347 | 590 | 381 | 674 | - | - | - | 2 | 2 | 1 |
| Ky. | 7 | 8 | 23 | 71 | 44 | 38 | - | - | - | - | - | - |
| Tenn. | 34 | 30 | 211 | 363 | 264 | 422 | - | - | - | 1 | 1 | - |
| Ala. | 14 | 14 | 70 | 79 | 71 | 74 | - | - | - | 1 | 1 | 1 |
| Miss. | 2 | 2 | 43 | 77 | 2 | 140 | - | - | - | - | - | - |
| W.S. CENTRAL | 56 | 47 | 3,937 | 5,380 | 1,162 | 1,213 | - | 1 | - | - | 1 | 8 |
| Ark. | - | 2 | 87 | 206 | 88 | 83 | U | - | U | - | - | - |
| La. | 23 | 12 | 124 | 220 | 164 | 161 | - | 1 | - | - | 1 | - |
| Okla. | 30 | 30 | 598 | 1,372 | 108 | 49 | - | - | - | - | - | 1 |
| Tex. | 3 | 3 | 3,128 | 3,582 | 802 | 920 | - | - | - | - | - | 7 |
| MOUNTAIN | 108 | 85 | 3,074 | 3,993 | 793 | 814 | 5 | 8 | - | 2 | 10 | 8 |
| Mont. | - | 1 | 94 | 68 | 5 | 12 | - | - | - | - | - | - |
| Idaho | 2 | 1 | 231 | 133 | 46 | 52 | - | - | - | - | - | - |
| Wyo. | 1 | 4 | 36 | 32 | 8 | 24 | U | - | U | - | - | - |
| Colo. | 18 | 21 | 331 | 387 | 107 | 140 | - | - | - | - | - | - |
| N. Mex. | 8 | 8 | 143 | 334 | 310 | 240 | - | - | - | - | - | - |
| Ariz. | 54 | 31 | 1,842 | 2,098 | 173 | 186 | 5 | 8 | - | 2 | 10 | 5 |
| Utah | 6 | 3 | 188 | 527 | 66 | 88 | - | - | - | - | - | 1 |
| Nev. | 19 | 16 | 209 | 414 | 78 | 72 | - | - | - | - | - | 2 |
| PACIFIC | 104 | 239 | 5,136 | 7,138 | 1,612 | 1,716 | - | 33 | - | 5 | 38 | 26 |
| Wash. | 10 | 5 | 900 | 623 | 116 | 77 | - | - | - | 1 | 1 | 2 |
| Oreg. | 39 | 33 | 364 | 356 | 121 | 112 | - | - | - | - | - | - |
| Calif. | 46 | 185 | 3,816 | 5,982 | 1,355 | 1,502 | - | 5 | - | 3 | 8 | 20 |
| Alaska | 1 | 8 | 17 | 33 | 12 | 14 | - | 28 | - | 1 | 29 | - |
| Hawaii | 8 | 8 | 39 | 144 | 8 | 11 | - | - | - | - | - | 4 |
| Guam | - | - | - | - | 2 | 3 | U | - | U | - | - | - |
| P.R. | 2 | - | 49 | 264 | 335 | 764 | - | - | - | - | - | - |
| V.I. | U | U | U | U | U | U | U | U | U | U | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | 6 | 3 | 1 | 53 | 46 | U | - | U | - | - | 1 |

N: Not notifiable U: Unavailable -: no reported cases

*Of 220 cases among children aged <5 years, serotype was reported for 110 and of those, 43 were type b.

†For imported measles, cases include only those resulting from importation from other countries.

TABLE III. (Cont'd.) Provisional cases of selected notifiable diseases preventable by vaccination, United States, weeks ending December 12, 1998, and December 6, 1997 (49th Week)

| Reporting Area | Meningococcal Disease | | Mumps | | | Pertussis | | | Rubella | | |
|----------------|-----------------------|-----------|-------|-----------|-----------|-----------|-----------|-----------|---------|-----------|-----------|
| | Cum. 1998 | Cum. 1997 | 1998 | Cum. 1998 | Cum. 1997 | 1998 | Cum. 1998 | Cum. 1997 | 1998 | Cum. 1998 | Cum. 1997 |
| UNITED STATES | 2,488 | 2,969 | 3 | 580 | 611 | 101 | 5,799 | 5,411 | - | 334 | 158 |
| NEW ENGLAND | 104 | 187 | - | 7 | 12 | 11 | 910 | 981 | - | 38 | 1 |
| Maine | 6 | 18 | - | - | - | - | 5 | 22 | - | - | - |
| N.H. | 4 | 14 | - | - | 1 | 2 | 121 | 130 | - | - | - |
| Vt. | 5 | 4 | - | - | - | 2 | 74 | 244 | - | - | - |
| Mass. | 56 | 93 | - | 4 | 4 | 2 | 651 | 541 | - | 8 | 1 |
| R.I. | 8 | 21 | - | 1 | 6 | 4 | 13 | 16 | - | 1 | - |
| Conn. | 25 | 37 | - | 2 | 1 | 1 | 46 | 28 | - | 29 | - |
| MID. ATLANTIC | 235 | 323 | 1 | 170 | 58 | 9 | 572 | 383 | - | 135 | 34 |
| Upstate N.Y. | 69 | 84 | 1 | 11 | 12 | 7 | 300 | 156 | - | 111 | 6 |
| N.Y. City | 24 | 51 | - | 139 | 3 | - | 39 | 60 | - | 18 | 28 |
| N.J. | 55 | 71 | U | 2 | 8 | U | 5 | 14 | U | 4 | - |
| Pa. | 87 | 117 | - | 18 | 35 | 2 | 228 | 153 | - | 2 | - |
| E.N. CENTRAL | 374 | 463 | - | 72 | 86 | 20 | 624 | 605 | - | - | 6 |
| Ohio | 133 | 156 | - | 28 | 34 | 10 | 279 | 158 | - | - | - |
| Ind. | 70 | 53 | - | 6 | 14 | 1 | 145 | 69 | - | - | - |
| Ill. | 89 | 147 | - | 11 | 12 | 6 | 113 | 108 | - | - | 2 |
| Mich. | 42 | 66 | - | 27 | 22 | 3 | 70 | 62 | - | - | - |
| Wis. | 40 | 41 | - | - | 4 | - | 17 | 208 | - | - | 4 |
| W.N. CENTRAL | 216 | 220 | - | 30 | 18 | 10 | 541 | 527 | - | 33 | - |
| Minn. | 32 | 34 | - | 13 | 6 | 6 | 337 | 306 | - | - | - |
| Iowa | 47 | 46 | - | 11 | 10 | 3 | 76 | 106 | - | - | - |
| Mo. | 79 | 93 | - | 3 | - | - | 32 | 67 | - | 2 | - |
| N. Dak. | 5 | 2 | - | 2 | - | - | 3 | 1 | - | - | - |
| S. Dak. | 8 | 5 | - | - | - | - | 8 | 5 | - | - | - |
| Nebr. | 15 | 18 | - | - | 1 | 1 | 19 | 13 | - | - | - |
| Kans. | 30 | 22 | - | 1 | 1 | - | 66 | 29 | - | 31 | - |
| S. ATLANTIC | 440 | 505 | - | 48 | 73 | 9 | 323 | 411 | - | 19 | 78 |
| Del. | 2 | 5 | - | - | - | - | 5 | 1 | - | - | - |
| Md. | 33 | 42 | - | - | 1 | - | 57 | 113 | - | 1 | - |
| D.C. | 3 | 12 | - | - | - | - | 1 | 3 | - | - | 1 |
| Va. | 45 | 58 | - | 8 | 18 | - | 41 | 52 | - | 1 | 1 |
| W. Va. | 16 | 19 | - | - | - | - | 4 | 6 | - | - | - |
| N.C. | 58 | 88 | - | 11 | 12 | - | 98 | 118 | - | 13 | 59 |
| S.C. | 55 | 55 | - | 7 | 11 | - | 27 | 29 | - | - | 15 |
| Ga. | 97 | 94 | - | 1 | 10 | - | 27 | 13 | - | - | - |
| Fla. | 131 | 132 | - | 21 | 21 | 9 | 63 | 76 | - | 4 | 2 |
| E.S. CENTRAL | 227 | 223 | - | 16 | 31 | 1 | 119 | 143 | - | 2 | 1 |
| Ky. | 36 | 46 | - | 1 | 3 | - | 50 | 63 | - | - | - |
| Tenn. | 69 | 76 | - | 1 | 6 | - | 37 | 37 | - | 2 | - |
| Ala. | 98 | 76 | - | 8 | 9 | 1 | 29 | 32 | - | - | 1 |
| Miss. | 24 | 25 | - | 6 | 13 | - | 3 | 11 | - | - | - |
| W.S. CENTRAL | 292 | 278 | - | 60 | 84 | 9 | 361 | 276 | - | 88 | 4 |
| Ark. | 30 | 34 | U | 12 | 1 | U | 91 | 53 | U | - | - |
| La. | 60 | 48 | - | 10 | 16 | - | 9 | 20 | - | - | - |
| Okla. | 41 | 41 | - | - | - | - | 30 | 51 | - | - | - |
| Tex. | 161 | 155 | - | 38 | 67 | 9 | 231 | 152 | - | 88 | 4 |
| MOUNTAIN | 141 | 169 | - | 38 | 55 | 11 | 1,093 | 1,189 | - | 5 | 7 |
| Mont. | 4 | 8 | - | - | - | - | 13 | 18 | - | - | - |
| Idaho | 13 | 10 | - | 6 | 3 | 5 | 257 | 521 | - | - | 2 |
| Wyo. | 7 | 3 | U | 1 | 1 | U | 8 | 7 | U | - | - |
| Colo. | 27 | 45 | - | 6 | 3 | 4 | 229 | 379 | - | - | - |
| N. Mex. | 26 | 29 | N | N | N | 1 | 97 | 175 | - | 1 | - |
| Ariz. | 41 | 41 | - | 6 | 33 | - | 199 | 36 | - | 1 | 5 |
| Utah | 14 | 15 | - | 5 | 8 | 1 | 249 | 25 | - | 2 | - |
| Nev. | 9 | 18 | - | 14 | 7 | - | 41 | 28 | - | 1 | - |
| PACIFIC | 459 | 601 | 2 | 139 | 194 | 21 | 1,256 | 896 | - | 14 | 27 |
| Wash. | 63 | 87 | - | 11 | 19 | 18 | 329 | 384 | - | 9 | 5 |
| Oreg. | 88 | 120 | N | N | N | 2 | 90 | 47 | - | - | - |
| Calif. | 300 | 384 | 2 | 103 | 142 | - | 803 | 430 | - | 3 | 14 |
| Alaska | 3 | 3 | - | 2 | 8 | - | 15 | 16 | - | - | - |
| Hawaii | 5 | 7 | - | 23 | 25 | 1 | 19 | 19 | - | 2 | 8 |
| Guam | 1 | 1 | U | 2 | 1 | U | - | - | U | - | - |
| P.R. | 7 | 8 | - | 1 | 7 | - | 6 | - | - | - | - |
| V.I. | U | U | U | U | U | U | U | U | U | U | U |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | - | U | 2 | 4 | U | 1 | - | U | - | - |

N: Not notifiable

U: Unavailable

-: no reported cases

**TABLE IV. Deaths in 122 U.S. cities,* week ending
December 12, 1998 (49th Week)**

| Reporting Area | All Causes, By Age (Years) | | | | | | P&J† | Total | Reporting Area | All Causes, By Age (Years) | | | | | | P&J† | Total |
|---------------------|----------------------------|-------|-------|-------|------|----|------|-----------------------|----------------|----------------------------|-------|-------|-------|------|-----|------|-------|
| | All Ages | >65 | 45-64 | 25-44 | 1-24 | <1 | | | | All Ages | >65 | 45-64 | 25-44 | 1-24 | <1 | | |
| NEW ENGLAND | 604 | 434 | 110 | 38 | 14 | 8 | 35 | S. ATLANTIC | 1,170 | 749 | 242 | 128 | 22 | 25 | 67 | | |
| Boston, Mass. | 144 | 85 | 35 | 12 | 9 | 3 | 8 | Atlanta, Ga. | U | U | U | U | U | U | U | | |
| Bridgeport, Conn. | 45 | 39 | 3 | 2 | - | 1 | 2 | Baltimore, Md. | 281 | 171 | 65 | 37 | 5 | 3 | 25 | | |
| Cambridge, Mass. | 16 | 12 | 3 | 1 | - | - | - | Charlotte, N.C. | 94 | 65 | 14 | 9 | 4 | 2 | 11 | | |
| Fall River, Mass. | 24 | 22 | - | 2 | - | - | 2 | Jacksonville, Fla. | 145 | 88 | 30 | 16 | 1 | 9 | 3 | | |
| Hartford, Conn. | 47 | 26 | 13 | 4 | 3 | 1 | 1 | Miami, Fla. | 123 | 61 | 34 | 22 | 1 | 4 | - | | |
| Lowell, Mass. | 24 | 17 | 4 | 2 | 1 | - | 1 | Norfolk, Va. | 47 | 30 | 10 | 5 | 2 | - | 2 | | |
| Lynn, Mass. | 12 | 9 | 2 | 1 | - | - | 1 | Richmond, Va. | 62 | 48 | 9 | 3 | - | 2 | 2 | | |
| New Bedford, Mass. | 36 | 33 | 1 | 2 | - | - | 4 | Savannah, Ga. | 33 | 24 | 6 | 1 | - | 2 | 4 | | |
| New Haven, Conn. | 42 | 26 | 11 | 4 | 1 | - | 1 | St. Petersburg, Fla. | 75 | 54 | 11 | 9 | - | 1 | 8 | | |
| Providence, R.I. | 66 | 53 | 9 | 4 | - | - | - | Tampa, Fla. | 200 | 139 | 40 | 15 | 5 | - | 10 | | |
| Somerville, Mass. | 5 | 4 | 1 | - | - | - | - | Washington, D.C. | 100 | 60 | 22 | 11 | 4 | 2 | 2 | | |
| Springfield, Mass. | 43 | 31 | 9 | 1 | - | 2 | 4 | Wilmington, Del. | 10 | 9 | 1 | - | - | - | - | | |
| Waterbury, Conn. | 34 | 22 | 11 | - | - | 1 | 4 | E.S. CENTRAL | 881 | 590 | 169 | 62 | 24 | 33 | 50 | | |
| Worcester, Mass. | 66 | 55 | 8 | 3 | - | - | 7 | Birmingham, Ala. | 222 | 141 | 46 | 14 | 7 | 11 | 16 | | |
| MID. ATLANTIC | 2,209 | 1,568 | 418 | 159 | 28 | 35 | 116 | Chattanooga, Tenn. | 68 | 51 | 11 | 4 | 1 | 1 | 3 | | |
| Albany, N.Y. | 50 | 40 | 8 | 1 | - | 1 | 6 | Knoxville, Tenn. | 79 | 49 | 22 | 6 | - | 2 | 4 | | |
| Allentown, Pa. | 24 | 21 | 3 | - | - | - | 2 | Lexington, Ky. | 46 | 32 | 6 | 4 | 2 | 2 | 7 | | |
| Buffalo, N.Y. | 89 | 72 | 11 | 5 | - | - | 8 | Memphis, Tenn. | 147 | 89 | 33 | 17 | 4 | 4 | 8 | | |
| Camden, N.J. | 30 | 20 | 5 | 3 | 1 | 1 | 2 | Mobile, Ala. | 81 | 58 | 15 | 3 | 4 | 1 | - | | |
| Elizabeth, N.J. | U | U | U | U | U | U | U | Montgomery, Ala. | 77 | 60 | 10 | 4 | - | 3 | 6 | | |
| Erie, Pa. | 48 | 43 | 4 | - | 1 | - | 3 | Nashville, Tenn. | 161 | 110 | 26 | 10 | 6 | 9 | 6 | | |
| Jersey City, N.J. | 54 | 35 | 10 | 6 | - | 3 | 3 | W.S. CENTRAL | 1,596 | 1,064 | 324 | 120 | 35 | 53 | 104 | | |
| New York City, N.Y. | 1,187 | 809 | 244 | 98 | 19 | 17 | 54 | Austin, Tex. | 94 | 66 | 12 | 10 | 2 | 4 | 4 | | |
| Newark, N.J. | U | U | U | U | U | U | U | Baton Rouge, La. | 41 | 24 | 8 | 6 | 1 | 2 | 2 | | |
| Paterson, N.J. | 29 | 13 | 11 | 5 | - | - | - | Corpus Christi, Tex. | 57 | 36 | 18 | 1 | 1 | 1 | 2 | | |
| Philadelphia, Pa. | 300 | 203 | 61 | 22 | 5 | 9 | 15 | Dallas, Tex. | 221 | 134 | 52 | 18 | 6 | 11 | 3 | | |
| Pittsburgh, Pa.‡ | 58 | 42 | 11 | 2 | 1 | 2 | 3 | El Paso, Tex. | 79 | 55 | 9 | 11 | 4 | - | 3 | | |
| Reading, Pa. | 20 | 16 | 4 | - | - | - | 1 | Ft. Worth, Tex. | 115 | 87 | 20 | 6 | 1 | 1 | 4 | | |
| Rochester, N.Y. | 132 | 101 | 21 | 7 | 1 | 2 | 9 | Houston, Tex. | 408 | 239 | 103 | 35 | 10 | 21 | 50 | | |
| Schenectady, N.Y. | 28 | 24 | 4 | - | - | - | 4 | Little Rock, Ark. | 51 | 35 | 10 | 3 | 1 | 2 | 5 | | |
| Scranton, Pa. | 31 | 24 | 6 | 1 | - | - | - | New Orleans, La. | 86 | 56 | 20 | 7 | 1 | 2 | - | | |
| Syracuse, N.Y. | 81 | 67 | 8 | 6 | - | - | 5 | San Antonio, Tex. | 244 | 174 | 43 | 17 | 4 | 6 | 14 | | |
| Trenton, N.J. | 30 | 22 | 5 | 3 | - | - | 1 | Shreveport, La. | 59 | 53 | 2 | 1 | 1 | 2 | 8 | | |
| Utica, N.Y. | 18 | 16 | 2 | - | - | - | - | Tulsa, Okla. | 141 | 105 | 27 | 5 | 3 | 1 | 9 | | |
| Yonkers, N.Y. | U | U | U | U | U | U | U | MOUNTAIN | 1,086 | 735 | 200 | 104 | 21 | 25 | 66 | | |
| E.N. CENTRAL | 2,251 | 1,553 | 406 | 161 | 63 | 64 | 159 | Albuquerque, N.M. | 96 | 60 | 21 | 13 | 2 | - | 5 | | |
| Akron, Ohio | 68 | 56 | 9 | 1 | 2 | - | 1 | Boise, Idaho | 35 | 25 | 8 | 1 | - | 1 | 2 | | |
| Canton, Ohio | 46 | 34 | 9 | 2 | 1 | - | 6 | Colo. Springs, Colo. | 58 | 42 | 11 | 5 | - | - | 8 | | |
| Chicago, Ill. | 429 | 245 | 90 | 52 | 24 | 14 | 29 | Denver, Colo. | 116 | 61 | 24 | 21 | 4 | 6 | 9 | | |
| Cincinnati, Ohio | 75 | 48 | 14 | 7 | 4 | 2 | 6 | Las Vegas, Nev. | 230 | 154 | 47 | 22 | 4 | 3 | 5 | | |
| Cleveland, Ohio | 152 | 111 | 26 | 10 | 1 | 4 | 4 | Ogden, Utah | 34 | 24 | 5 | 1 | 1 | 3 | 1 | | |
| Columbus, Ohio | 223 | 155 | 38 | 18 | 6 | 6 | 20 | Phoenix, Ariz. | 225 | 152 | 39 | 20 | 6 | 7 | 13 | | |
| Dayton, Ohio | 130 | 93 | 20 | 7 | 5 | 5 | 6 | Pueblo, Colo. | 39 | 27 | 9 | 2 | 1 | - | 5 | | |
| Detroit, Mich. | 206 | 127 | 47 | 18 | 7 | 7 | 14 | Salt Lake City, Utah | 98 | 71 | 11 | 10 | 3 | 3 | 6 | | |
| Evansville, Ind. | 36 | 25 | 8 | 2 | - | 1 | 2 | Tucson, Ariz. | 155 | 119 | 25 | 9 | - | 2 | 12 | | |
| Fort Wayne, Ind. | 67 | 50 | 7 | 6 | 1 | 3 | 1 | PACIFIC | 1,845 | 1,321 | 316 | 138 | 32 | 38 | 155 | | |
| Gary, Ind. | 16 | 9 | 3 | 1 | 2 | 1 | - | Berkeley, Calif. | 23 | 15 | 5 | 2 | - | 1 | 2 | | |
| Grand Rapids, Mich. | 78 | 65 | 6 | 1 | 1 | 5 | 10 | Fresno, Calif. | 112 | 89 | 12 | 6 | 3 | 2 | 11 | | |
| Indianapolis, Ind. | 202 | 137 | 45 | 12 | 4 | 4 | 21 | Glendale, Calif. | 25 | 23 | 1 | 1 | - | - | 3 | | |
| Lansing, Mich. | 40 | 34 | 3 | 2 | - | 1 | 4 | Honolulu, Hawaii | 66 | 52 | 12 | 1 | 1 | - | 6 | | |
| Milwaukee, Wis. | 126 | 89 | 20 | 13 | - | 4 | 16 | Long Beach, Calif. | 54 | 46 | 4 | 3 | - | 1 | 7 | | |
| Peoria, Ill. | 55 | 38 | 12 | 1 | 2 | 2 | 1 | Los Angeles, Calif. | 452 | 319 | 82 | 32 | 8 | 11 | 21 | | |
| Rockford, Ill. | 48 | 41 | 4 | 2 | - | 1 | 2 | Pasadena, Calif. | 36 | 25 | 7 | 3 | 1 | - | 2 | | |
| South Bend, Ind. | 69 | 49 | 17 | 2 | 1 | - | 3 | Portland, Oreg. | 141 | 97 | 26 | 13 | 1 | 4 | 8 | | |
| Toledo, Ohio | 108 | 80 | 21 | 4 | 1 | 2 | 6 | Sacramento, Calif. | 207 | 141 | 35 | 20 | 6 | 5 | 28 | | |
| Youngstown, Ohio | 77 | 67 | 7 | - | 1 | 2 | 7 | San Diego, Calif. | 147 | 105 | 24 | 11 | 3 | 4 | 13 | | |
| W.N. CENTRAL | 777 | 536 | 138 | 62 | 17 | 16 | 50 | San Francisco, Calif. | 83 | 56 | 16 | 9 | 1 | 1 | 11 | | |
| Des Moines, Iowa | 41 | 26 | 11 | 2 | 1 | 1 | 3 | San Jose, Calif. | 222 | 165 | 36 | 15 | 3 | 3 | 26 | | |
| Duluth, Minn. | 43 | 37 | 3 | 3 | - | - | 4 | Santa Cruz, Calif. | 23 | 13 | 6 | 2 | 1 | 1 | 1 | | |
| Kansas City, Kans. | U | U | U | U | U | U | U | Seattle, Wash. | 121 | 71 | 31 | 16 | 2 | 1 | 5 | | |
| Kansas City, Mo. | 109 | 65 | 19 | 13 | 1 | 3 | 2 | Spokane, Wash. | 58 | 49 | 6 | 1 | - | 2 | 1 | | |
| Lincoln, Nebr. | 36 | 24 | 9 | 2 | 1 | - | 3 | Tacoma, Wash. | 75 | 55 | 13 | 3 | 2 | 2 | 10 | | |
| Minneapolis, Minn. | 224 | 175 | 34 | 8 | 3 | 4 | 15 | TOTAL | 12,419‡ | 8,550 | 2,323 | 972 | 256 | 297 | 802 | | |
| Omaha, Nebr. | 85 | 59 | 14 | 6 | 2 | 4 | 9 | | | | | | | | | | |
| St. Louis, Mo. | 103 | 55 | 21 | 17 | 7 | 3 | 8 | | | | | | | | | | |
| St. Paul, Minn. | 136 | 95 | 27 | 11 | 2 | 1 | 6 | | | | | | | | | | |
| Wichita, Kans. | U | U | U | U | U | U | U | | | | | | | | | | |

U: Unavailable - : no reported cases

*Mortality data in this table are voluntarily reported from 122 cities in the United States, most of which have populations of 100,000 or more. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included.

†Pneumonia and influenza.

‡Because of changes in reporting methods in this Pennsylvania city, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks.

¶Total includes unknown ages.

Fatal Cercopithecine herpesvirus Infection — Continued

8. CDC. Guidelines for prevention of *Herpesvirus simiae* (B virus) infection in monkey handlers. MMWR 1987;36:680–2,687–9.
9. American National Standards Institute. Practice for occupational and educational eye and face protection, Z87-1. Des Plaines, Illinois: American Society of Safety Engineers, 1989.
10. Heinsohn P, Jacobs R, Concoby B. Biosafety reference manual. 2nd ed. Fairfax, Virginia: American Industrial Hygiene Association, 1995:79–80.

Notice to Readers**Revising CDC's Guidelines for Evaluating Surveillance Systems**

A surveillance system provides for the ongoing collection, analysis, and dissemination of data to prevent and control disease. Because all surveillance systems should be assessed periodically for their purpose and usefulness, in 1988, CDC published *Guidelines for Evaluating Surveillance Systems* (1). Recent developments in the electronic exchange of health data (2,3), the establishment of data collection standards (2,4), and interest in the integration of health information and surveillance systems (4,5) have resulted in the need to revise CDC's guidelines.

The guidelines will be revised by a working group under the direction of CDC's Surveillance Coordination Group, comprising representatives from each of the program areas at CDC and ATSDR and from state organizations that collaborate with CDC. Because the surveillance systems at CDC and ATSDR are implemented in collaboration with state and local prevention partners, these groups will be included in the development of revised guidelines.

Comments on the revision of the guidelines should be submitted by December 1999 by e-mail to revguide@cdc.gov or by mail to Attention: Revised Guidelines, Epidemiology Program Office, CDC, Mailstop C-08, 1600 Clifton Road, N.E., Atlanta, GA, 30333.

References

1. CDC. Guidelines for evaluating surveillance systems. MMWR 1988;37(no. SS-5).
2. Harman J. Topics for our times: new health care data—new horizons for public health. Am J Public Health 1998;88:1019–21.
3. Duke University Medical Center. Health Level Seven (HL7) application protocol, Medical Center Information Systems, version 3.0. Durham, North Carolina: Duke University Medical Center, 1996.
4. CDC. Integrating public health information and surveillance systems: a report and recommendations from the CDC/ATSDR Steering Committee on Public Health Information and Surveillance System Development. Atlanta, Georgia: US Department of Health and Human Services, CDC, 1995.
5. CDC. Surveillance review and notification policy. Atlanta, Georgia: US Department of Health and Human Services, CDC, 1998.

Erratum: Vol. 47, No. 48

In the notice to readers, "Federal Register Notice on the Draft Guidelines for HIV Case Surveillance, Including Monitoring HIV Infection and AIDS," on page 1056, the closing date for comments is incorrect. The correct deadline for submitting comments on the draft is January 11, 1999.

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/> or from CDC's file transfer protocol server at <ftp.cdc.gov>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (888) 232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Director, Centers for Disease Control
and Prevention
Jeffrey P. Koplan, M.D., M.P.H.
Deputy Director, Centers for Disease
Control and Prevention
Claire V. Broome, M.D.

Director, Epidemiology Program Office
Stephen B. Thacker, M.D., M.Sc.
Editor, *MMWR* Series
John W. Ward, M.D.
Managing Editor,
MMWR (weekly)
Karen L. Foster, M.A.

Writers-Editors,
MMWR (weekly)
David C. Johnson
Teresa F. Rutledge
Caran R. Wilbanks
Desktop Publishing and
Graphics Support
Morie M. Higgins
Peter M. Jenkins

☆ U.S. Government Printing Office: 1998-733-228/87049 Region IV
