



Published in final edited form as:

Prev Med. 2013 May ; 56(5): 288–292. doi:10.1016/j.yjmed.2013.01.018.

A Farmers' Market at a Federally Qualified Health Center Improves Fruit and Vegetable Intake among Low-income Diabetics

Darcy A. Freedman, Ph.D.^{a,b}, Seul Ki Choi, M.P.H.^{b,c}, Thomas Hurley, M.S.^b, Edith Anadu, Ph.D.^d, and James R. Hebert, Sc.D.^{b,f}

Darcy A. Freedman: darcy.freedman@sc.edu; Seul Ki Choi: choi27@email.sc.edu; Thomas Hurley: THURLEY@mailbox.sc.edu; Edith Anadu: edith.anadu@myfhc.org; James R. Hebert: JHEBERT@mailbox.sc.edu

^aUniversity of South Carolina, College of Social Work, DeSaussure Hall, Columbia, SC, 29208, USA

^bUniversity of South Carolina, Cancer Prevention and Control Program, 915 Greene St., Columbia, SC, 29208, USA

^cUniversity of South Carolina, Department of Health Promotion, Education, and Behavior, 800 Sumter St., Columbia, SC, 29208, USA

^dFamily Health Centers, Inc., 3310 Magnolia St., Orangeburg, SC, 29115, USA

^fUniversity of South Carolina, Department of Epidemiology and Biostatistics, 800 Sumter St., Columbia, SC, 29208, USA

Abstract

Objective—A 22-week federally qualified health center (FQHC)-based farmers' market (FM) and personal financial incentive intervention designed to improve access to and consumption of fruits and vegetables (FV) among low-income diabetics in rural South Carolina was evaluated.

Methods—A mixed methods, one-group, repeated-measures design was used. Data were collected in 2011 before (May/June), during (August), and after (November) the intervention with 41 diabetes patients from the FQHC. FV consumption was assessed using a validated National Cancer Institute FV screener modified to include FV sold at the FM. Sales receipts were recorded for all FM transactions. A mixed-model, repeated measures analysis of variance was used to assess intervention effects on FV consumption. Predictors of changes in FV consumption were examined using logistic regression.

© 2012 Elsevier Inc. All rights reserved.

Corresponding Author: Darcy A. Freedman, Ph.D., M.P.H., University of South Carolina, DeSaussure Hall, Columbia, SC 29208, Ph: 1-803-777-1326, Fax: 1-803-777-3498, darcy.freedman@sc.edu.

Conflict of Interest Statement

The authors declare there are no conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Results—A marginally significant ($p=0.07$) average increase of 1.6 servings of total FV consumption per day occurred. The odds of achieving significant improvements in FV consumption increased for diabetics using financial incentives for payment at the FM (OR: 38.8, 95% CI: 3.4–449.6) and for those frequenting the FM more often (OR: 2.1, 95% CI: 1.1–4.0).

Conclusions—Results reveal a dose-response relationship between the intervention and FV improvements and emphasize the importance of addressing economic barriers to food access.

Keywords

Prevention & control; Community health centers; Health promotion; Diabetes Mellitus; Type 2; Obesity; Poverty

Introduction

Obesity rates in the United States are rising (Finkelstein et al., 2012; Flegal et al., 2010) resulting in increases in type 2 diabetes (Hu et al., 2001). Interventions designed to improve fruit and vegetable (FV) consumption are important strategies for preventing and treating obesity and diabetes (Ford et al., 2012; Gillies et al., 2007; Montonen et al., 2004; Yamaoka and Tango, 2005). Improving FV intake among Americans, however, has proven to be difficult (Grimm et al., 2010).

Recently, there has been a focus on increasing individual consumption of FV by improving access, availability, and affordability to FV in communities (Grimm et al., 2010). This focus emerged because populations disproportionately burdened by diet-related health conditions (Pan et al., 2009) are less likely to have healthy food retailers in their community (Dutko et al., 2012). Moreover, behaviorally-based interventions have not resulted in sustained improvements in diet (Jetter and Cassady, 2006).

Farmers' markets (FM) are targeted approaches for improving access to FV (Centers for Disease Control and Prevention, 2011). Primary health care settings such as federally qualified health centers (FQHC) provide an ideal context for establishing FMs because they are situated in underserved communities (Health Resources and Services Administration, 2012). Moreover, locating at a FQHC makes an explicit connection between FMs and preventive medicine.

Only a few studies reported have examined the influence of FM interventions on FV consumption; most have limitations related to study design and measurement of FV intake, are focused on the Women's Infant and Children (WIC) program participants, and are based in urban areas (McCormack et al., 2010). A recent review of the nutritional implications of FMs concluded "...there is limited research assessing the specific health benefits of farmers' markets" (McCormack et al., 2010).

Four FM interventions used repeated, validated measures to examine changes in FV intake (Abusabha et al., 2011; Anderson et al., 2001; Evans et al., 2012; Herman et al., 2008). Not one of these is focused on high-risk sub-populations like people with diabetes, conducted in a rural context, or conducted in a setting that serves medically underserved populations. In this analysis we addressed this gap by evaluating the influence of a FQHC-based FM and

personal financial incentive intervention on FV consumption among low-income, diabetics in a rural context using validated measures of FV intake and objective measures of FM usage.

Methods

Design, Setting, and Participants

We used a community-based participatory research (CBPR) approach that involved a partnership between the University of South Carolina and Family Health Centers, Inc., an FQHC located in a majority minority (63% African American) rural county in South Carolina (U.S. Census Bureau, 2010). A one-group repeated-measures design was used. The study was approved by the university Institutional Review Board.

Adult participants were eligible if they were patients at the FQHC with a diabetes diagnosis as of March 1, 2011 (N=2,306 patients). Health center staff randomly selected 345 diabetics; each received a mailing that described the purpose of the study including information about the chance to receive \$50 in vouchers to shop at the FM. Due to HIPAA, patient names and contact information were not revealed to the research staff; we are unable to track mailings that were undeliverable. It was impossible to determine the total sample that actually received an invitation to participate in the study. Interested potential participants were required to contact the research staff to express interest and determine eligibility. A total of 63 patients expressed interest; 9 could not be reached to schedule a survey, 9 were ineligible (e.g., no longer patient at FQHC, refused to consent). Our resulting analytic sample includes 45 diabetics. Informed consent was obtained from all participants; this involved verbal review of the written consent form to describe the purpose and process of the research prior to voluntary agreement to participate.

Intervention

The intervention was designed using community feedback, which is detailed in a documentary about the market (Murphy and Jacobs, 2011). We conducted a visioning exercise with 50 community members to understand their hopes for a new FM using a modified version of a nominal group process (Johnson et al., 2011). A 10-member Community Advisory Council was formed to guide the development of the FM including representatives from the FQHC, local schools/universities, agricultural extension, faith-based institutions, and community volunteers. Pre-market interest surveys were conducted with patients at the FQHC and local farmers.

Community feedback informed the two components of the FM intervention. First, onsite produce-only FMs operated at the FQHC once per week (10am–2pm) for 22 weeks from June–October 2011. The market was managed by a community member hired through grant funding. Supplemental Nutrition Assistance Program (SNAP) vouchers were accepted at the FM through a central point-of-purchase electronic benefits transfer system. Most of the vendors were certified to accept Senior and WIC Farmers' Market Nutrition Program Vouchers. Second, study participants were enrolled in a personal financial incentive program that provided up to \$50 in vouchers to purchase FV at the FM. Vouchers were

provided after participants completed two study-related surveys (described below) and were paid for through grant resources. Voucher usage at the FM was documented manually; in two instances a study participant exceeded the \$50 maximum due to a recording error.

Instrumentation

We collected data with the diabetic patient sample at three time points: before the FM intervention (T1, May/June 2011), midway through the intervention (T2, August 2011), and immediately after the intervention (T3, November 2011). Data were collected via structured surveys either in-person or over the telephone by trained research assistants. Vouchers to shop at the FM were provided to participants after completing surveys at T1 and T2 (\$25 each time). A stipend of \$40 was provided after the third survey. A total of 45 participants completed the first survey and 44 completed the second and third.

The main outcome, assessed at each time point, was FV intake measured using a modified version of the 10-item NCI FV screener (Greene et al., 2008; Peterson et al., 2008; Thompson et al., 2002). The modified 19-item version included nine additional FV available at the FM (e.g., peach, apple, orange, cantaloupe, cabbage, broccoli, squash/zucchini, sweet potato, tomato). Demographics, social context, and health status information were collected at T1 using close-ended questions from the Behavioral Risk Factor Surveillance System survey (Table 1). We developed one question to assess self-reports of seven diet-related health conditions.

A receipt of each sales transaction (N=3,747) at the FQHC-based FM was recorded on an optically scannable form; 438 of the receipts were related to purchases made by the study participants. Receipts were recorded by trained research assistants and included the following information: date, participant ID, volume of produce purchased measured in units (e.g., 1 unit=1 peach or 1 basket of okra), cost of produce purchased, and form of payment.

Analysis

Descriptive statistics were computed to describe sociodemographic characteristics of participants, FM use, and food items purchased. FV consumption estimation followed NCI's guidelines (2010). Participants' reports of the frequency of intake and portion size of each food item were converted to average daily frequency and MyPyramid servings. Missing values for frequency of intake were considered as "never" for that food item. Missing values for portion size were imputed to individual's most frequently reported portion size for vegetable items or reported portion size of other fruit for fruit items. Daily frequency and servings were multiplied to generate average daily servings of single food items. Total FV consumption was calculated by summing all food items. Participants without surveys at T2 or T3 (n=1) and those reporting greater than three standard deviations above the mean (approximately 12.5 servings/day) at any time point (n=3) were excluded resulting in an analytic sample of 41 participants.

A mixed-model, repeated measures analysis of variance (ANOVA) was used to assess the effects of the FM and personal financial incentive intervention on FV consumption over time. The sample was dichotomized to explore factors associated with increases in FV consumption. "Increasers" were 56.1% of the sample whose average FV consumption at T2

and T3 was at least 0.5 servings greater than their consumption at T1 whereas the “Non-Increaser” group constituted 43.9% of the sample that did not increase FV consumption over time. This level of change was selected because effective behavioral interventions typically improve diet by about 0.5 servings per day (Ammerman et al., 2002). Students’ t-tests were used to compare FV consumption between the two groups at each time point. Logistic regression was conducted to examine potential predictors of changes in FV consumption. Self-reported height and weight were used to calculate body mass index [BMI=weight(kg)/height(m²); continuous], payment type (study voucher only versus voucher+other form of payment), number of FM visits (continuous), total amount of money spent at the FM (continuous), and receipt of food assistance in the past year (yes/no) were included in the regression model. The goodness of fit of the regression model was adequate ($\chi^2=11.30$, $p=0.19$), as assessed by the Hosmer and Lemeshow statistic. All statistical tests were performed using the SAS 9.2 (SAS Institute Inc., Cary, NC). The level of significance was set at $p < 0.05$.

Results

The diabetic sample was majority African American, female, and older (Table 1). Most participants were obese (BMI 30kg/m^2) (World Health Organization, 2000) with an average BMI of $34.9\pm 6.8\text{kg/m}^2$ calculated from self-reports of height and weight at T1. Participants had high rates of economic and food insecurity: most earned $<\$10,000$ per year and 51.2% reported being at least somewhat worried about having enough money to buy nutritious meals during the past year. At T1, 75.6% had not shopped at a FM in the month before the FQHC-based FM opened.

All participants came to the FQHC-based FM on at least 2 dates throughout the 22-week season (average, 4.5 days; range, 2–15). On average, participants made 10.7 (range, 5–28) sales transactions at the FM with an average of 2.5 (range, 1–6) transactions per day indicating participants frequented multiple vendors at the market during their visit. In total, the 41 study participants made 438 sales transactions. The five most popular items purchased were peaches, sweet potatoes, squash/zucchini, corn, and tomatoes. Participants spent an average of \$53.30 (range, \$29–126) throughout the market season and an average of \$5.49 during each sales transaction. Participants purchased food using multiple forms of payment often in the same sales transaction. Most participants (70.7%) paid for purchases with the study vouchers and at least one other form of payment (e.g., cash, SNAP) whereas 29.3% only used the study vouchers for payment.

Total FV intake increased from 5.9 servings per day at T1 to 7.5 and 6.5 servings per day at T2 and T3, respectively (Table 2). The increase of 1.6 servings per day in total FV consumption from T1 to T2 was marginally significant ($p=0.07$). FV intake at T3 was less than at T2, but was not statistically significant.

In the diabetic cohort at T1, we found that Increasers consumed fewer servings of FV per day compared to the Non-Increasers (T1: 4.9 vs. 7.3, $p=0.02$), whereas at T3 Increasers consumed significantly more FV servings per day compared to the Non-Increasers (T3: 7.8 vs. 4.8, $p=0.01$) (Table 3).

The odds of being an Increaser in FV consumption were higher for diabetics who only used vouchers for payment at the market (OR: 38.8, 95% CI: 3.35–445.0) and for those who visited the FM more often (OR: 2.07, 95% CI: 1.09–3.95) (Table 4). Increasing FV intake was not associated with BMI at baseline, receipt of food assistance, or total amount of money spent at the FM.

Discussion

While FMs are beginning to open at a variety of health care delivery sites (Estabrook et al., 2012; George et al., 2011), this is the first FQHC-based FM intervention in the scientific literature. Findings highlight the benefit of a FQHC-based FM and personal financial incentive intervention designed to improve diet among diabetics. FV consumption increased by 1.6 servings per day from baseline to the mid-point (August) of the intervention and remained about half a serving higher than baseline after the market ended (November). Higher FV consumption patterns at the end of the FM intervention compared to baseline is noteworthy because FV consumption patterns tend to be higher in the summer compared to fall/winter months (Locke et al., 2009; Ziegler et al., 1987). Only a few studies have used repeated, validated measures to examine the influence of FMs on FV consumption (McCormack et al., 2010) finding improvements between 0.4 to 2.4 servings per day (Abusabha et al., 2011; Evans et al., 2012; Herman et al., 2008). The FQHC-based FM model contributed to increases in FV consumption at levels equivalent to or better than behaviorally-based interventions (Ammerman et al., 2002).

Results illuminate a dose-response relationship between the FM intervention and increases in FV consumption among the diabetic cohort. More frequent usage of the FM was associated with higher odds of increasing FV consumption. Findings also emphasize the importance of the personal financial incentive program. The relatively small financial incentive (\$50) was quite beneficial to the diabetics. Those who only used the financial vouchers for payment at the FM were significantly more likely to increase FV consumption compared to those who used the voucher and at least one other form of payment. Findings suggest the FM and personal financial incentive intervention was particularly beneficial for those consuming the lowest levels FVs at baseline.

Strengths of this research include the use of a random sample, validated tools for measuring FV intake, and a repeated measures design. The context of the research is another strength; most FM research is focused on urban settings (McCormack et al., 2009) whereas this study occurred in a rural context. The sample is both a strength and limitation. The cohort of diabetics represents a population disparately affected by disease and hard-to-reach. The sample, however, may not be representative given its small size. The study design was enhanced by using a CBPR approach to engage community members in study development and implementation and by using mixed methods. Lack of a control group is a limitation. Future research is warranted that includes a larger sample and a more robust study design. Finally, there are limitations related to the sales transaction data collection process. There is a chance that some sales transactions made by the diabetic cohort were not recorded due to the busyness of the market; thus, findings may underrepresent FM utilization and benefit.

Conclusion

Findings offer evidence for developing FMs at health centers as a strategy for improving patient health. FQHC-based FMs may be instrumental to providing preventive healthcare services to patient populations, particularly in contexts with high rates of diet-related health conditions or limited access to healthy food retailers (i.e., food deserts) or both. FQHC-based FMs have the potential to serve as a “farmacy” for patients to access nutrients fundamental to good health, especially if patients shop at the market on a regular basis. Personal financial incentives to improve economic access to FMs, similar to co-payment programs that facilitate patient access to pharmaceuticals, may further enhance the benefit of a FM intervention.

Acknowledgments

This publication was supported by the South Carolina Cancer Prevention and Control Research Network funded under Cooperative Agreement Number 3U48DP001936-01W1 from the Centers for Disease Control and Prevention and the National Cancer Institute. J.R. Hebert and T. Hurley were supported by an Established Investigator Award in Cancer Prevention and Control from the Cancer Training Branch of the National Cancer Institute (K05 CA136975; JR Hebert, P.I.). We are thankful for our partners at Family Health Centers, Inc. and the South Carolina Primary Healthcare Association and for research assistance from Jason Greene, Kassy Alia, Natalia Carvahlo, Shanna Hastie, Amy Mattison Faye, and Samira Khan. We thank the anonymous reviewers for their guidance in refining this manuscript.

Abbreviations

CBPR	community-based participatory research
FM	Farmers’ market
FQHC	federally qualified health center
FV	Fruit and vegetable
HIPAA	Health Insurance Portability and Accountability Act of 1996
NCI	National Cancer Institute

References

- Abusabha R, Namjoshi D, Klein A. Increased access and affordability of produce improves perceived consumption of vegetables in low-income seniors. *J Am Diet Assoc.* 2011; 111:1548–1555.
- Ammerman AS, Lindquist CH, Lohr KN, Hersey J. The efficacy of behavioral interventions to modify dietary fat and fruit and vegetable intake: a review of the evidence. *Prev Med.* 2002; 35:25–41. [PubMed: 12079438]
- Anderson JV, Bybee DI, Brown RM, et al. 5 A Day fruit and vegetable intervention improves consumption in a low income population. *J Am Diet Assoc.* 2001; 101:195–202. [PubMed: 11271692]
- Centers for Disease Control and Prevention. Strategies to prevent obesity and other chronic diseases: The CDC guide to strategies to increase the consumption of fruits and vegetables. Atlanta, GA: U.S. Department of Health and Human Services; 2011.
- Dutko, P.; Ver Ploeg, M.; Farrigan, T. Economic Research Service. Characteristics and influential factors of food deserts. U.S. Department of Agriculture; Washington, DC: 2012.

- Estabrook B, Zapka J, Lemon SC. Evaluating the implementation of a hospital work-site obesity prevention intervention: Applying the RE-AIM framework. *Health Promot Pract.* 2012; 13:190–197. [PubMed: 21441205]
- Evans AE, Jennings R, Smiley AW, et al. Introduction of farm stands in low-income communities increases fruit and vegetable among community residents. *Health & Place.* 2012; 18:1137–1143. [PubMed: 22608130]
- Finkelstein EA, Khavjou OA, Thompson H, et al. Obesity and severe obesity forecasts through 2030. *Am J Prev Med.* 2012; 42:563–570. [PubMed: 22608371]
- Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA.* 2010; 303:235–241. [PubMed: 20071471]
- Ford ES, Bergmann MM, Boeing H, Li C, Capewell S. Healthy lifestyle behaviors and all-cause mortality among adults in the United States. *Prev Med.* 2012; 55:23–27. [PubMed: 22564893]
- George DR, Kraschnewski JL, Rovniak LS. Public health potential of farmers' markets on medical center campuses: A case study. *Am J Public Health.* 2011; 101:2226–2232. [PubMed: 22021298]
- Gillies CL, Abrams KR, Lambert PC, et al. Pharmacological and lifestyle interventions to prevent or delay type 2 diabetes in people with impaired glucose tolerance: systematic review and meta-analysis. *BMJ.* 2007; 334:299–302. [PubMed: 17237299]
- Greene GW, Resnicow K, Thompson FE, et al. Correspondence of the NCI Fruit and Vegetable Screener to repeat 24-H recalls and serum carotenoids in behavioral intervention trials. *J Nutr.* 2008; 138:200S–204S. [PubMed: 18156425]
- Grimm KA, Blanck HM, Scanlon DS, et al. State-specific trends in fruit and vegetable consumption among adults—United States, 2000–2009. *MMWR.* 2010; 59:1126–1130.
- Health Resources and Services Administration. About health centers. U.S Department of Health and Human Services; Washington, DC: 2012. Available: <http://bphc.hrsa.gov/about/index.html> [Accessed December 15 2012]
- Herman DR, Harrison GG, Afifi AA, Jenks E. Effect of a targeted subsidy on intake of fruits and vegetables among low-income women in the Special Supplemental Nutrition Program for Women, Infants, and Children. *Am J Public Health.* 2008; 98:98–105. [PubMed: 18048803]
- Hu FB, Manson JE, Stampfer MJ, et al. Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *N Engl J Med.* 2001; 345:790–797. [PubMed: 11556298]
- Jetter KM, Cassady DL. The availability and cost of healthier food alternatives. *Am J Prev Med.* 2006; 30:38–44. [PubMed: 16414422]
- Johnson C, Freedman DA, Joosten Y, Duke M. Cultivating an agenda for change: a dynamic model for community food assessments. *Arête.* 2011; 32:61–89.
- Locke E, Coronado GD, Thompson B, Kuniyuki A. Seasonal variation in fruit and vegetable consumption in a rural agricultural community. *J Am Diet Assoc.* 2009; 109:45–51. [PubMed: 19103322]
- McCormack LA, Laska MN, Larson NI, Story M. Review of the nutritional implications of farmers' markets and community gardens: A call for evaluation and research efforts. *J Am Diet Assoc.* 2010; 110:399–408. [PubMed: 20184990]
- Montonen J, Knekt P, Jarvinen R, Reunanen A. Dietary antioxidant intake and risk of type 2 diabetes. *Diabetes Care.* 2004; 27:362–366. [PubMed: 14747214]
- Murphy, T.; Jacobs, T. Planting Healthy Roots: A look at the Right Choice Fresh Start Farmers' Market. Produced by University of South Carolina; Columbia, SC: 2011. Available: <http://www.youtube.com/watch?v=viPtYFwzD6I> [Accessed December 21, 2012]
- National Cancer Institute. Fruit and vegetable screeners in the Eating at America's Table Study (EATS): Scoring the all-day screener. National Institutes of Health; 2010. Available: <http://riskfactor.cancer.gov/diet/screeners/fruitveg/scoring/allday.html> [Accessed June 15, 2012]
- Pan L, Galuska DA, Sherry B, et al. Differences in prevalence of obesity among black, white, and Hispanic adults -- United States, 2006–2008. *MMWR.* 2009; 58:740–744. [PubMed: 19609247]
- Peterson KE, Hebert JR, Hurley TG, et al. Accuracy and precision of two short screeners to assess change in fruit and vegetable consumption among diverse populations participating in health promotion intervention trials. *J Nutr.* 2008; 138:218S–225S. [PubMed: 18156428]

- Thompson FE, Subar AF, Midthune D, et al. Fruit and vegetable assessment: Performance of 2 new short instruments and a food frequency questionnaire. *J Am Diet Assoc.* 2002; 102:1764–1772. [PubMed: 12487538]
- U.S. Census Bureau. Decennial Census. U.S. Census Bureau; Washington, DC: 2010.
- World Health Organization. Obesity: Preventing and managing the global epidemic, WHO Technical Report Series. World Health Organization; Geneva, Switzerland: 2000.
- Yamaoka K, Tango T. Efficacy of lifestyle education to prevent Type 2 diabetes. *Diabetes Care.* 2005; 28:2780–2786. [PubMed: 16249558]
- Ziegler RG, Wilcox HB III, Mason TJ, Bill JS, Virgo PW. Season variation in intake of carotenoids and vegetables and fruits among white men in New Jersey. *Am J Clin Nutr.* 1987; 45:107–114. [PubMed: 3799494]

Highlights

- We evaluated a farmers' market and personal financial incentive intervention.
- The study occurred at a federally qualified health center in a rural context.
- Diabetics frequenting the market more often had greater improvements in diet.
- The financial incentive was critical to improving fruit and vegetable intake.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Characteristics of diabetics enrolled in the farmers' market intervention in rural South Carolina, June–October, 2011 (N=41).

Variables	Mean	Range
Age, years	63.6	34–88
Number of people living in household ^{a)}	2.1	1–8
Body mass index at baseline (T1) ^{b)}	34.9	24.2–51.6
Shopping days at farmers' market	4.5	2–15
Number of sales transactions at farmers' market	10.7	5–28
Amount of money spent at farmers' market, \$	53.30	29.00–125.75
	Frequency	Percent
Gender		
Female	34	82.9
Race		
African American	38	92.7
White	3	7.3
Marital status		
Widowed	17	41.5
Married	10	24.4
Never married	9	22.0
Divorced or separated	5	12.2
Education level		
Less than high school	12	29.3
High school graduate or GED	18	43.9
Some college or technical school	6	14.6
College graduate or more	5	12.2
Annual household income (last year) ^{c)}		
Less than \$10,000	22	53.7
\$10,000 to \$19,999	12	29.3
\$20,000 to \$29,999	5	12.2
Household food assistance (e.g., SNAP, WIC, and/or free or reduced price lunches)		
Yes	22	53.7
Household financial assistance (e.g., TANF, Medicaid, Disability, SSI)		
Yes	15	36.6
Primary form of transportation^{c)}		
Personal vehicle	26	63.4
Ride with friend or family	11	26.8
Bus or taxi	3	7.3
Employment status^{c)}		
Unable to work	18	43.9
Employed for wages	7	17.1

Variables	Mean	Range
Retired	8	19.5
Out of work for 1 year or more	4	9.8
Not employed for wages (e.g., homemaker, student)	3	7.3
Worried about having enough money to buy nutritious meals in past year		
Always or usually	11	26.8
Sometimes	10	24.4
Rarely or never	20	48.8
Self-reported health status^{c)}		
Excellent	1	2.4
Very good	1	2.4
Good	9	22.0
Fair	16	39.0
Poor	13	31.7
Self-reported disease status		
Diabetes	41	100.0
High Blood Pressure	37	90.2
Arthritis	33	80.5
Obesity	14	34.2
Heart Disease	13	31.7
Gallbladder	3	7.3
Cancer (ever)	2	4.9

^{a)}Including the respondent;

^{b)}Based on self-reports of height and weight at T1;

^{c)}Totals do not add up to 100% due to missing data.

Fruit and vegetable consumption before (T1), during (T2), and after (T3) the federally qualified health center-based farmers' market intervention in rural South Carolina, June–October, 2011 (N=41).

Table 2

Time point ¹	Mean (Standard Deviation)	Mean Difference	95% Confidence Interval of the Difference		
			Lower	Upper	p-value for difference
Total fruit ²					
T1	3.23(2.41)	T2-T1 0.95	-0.20	2.09	0.10
T2	4.18(3.06)	T3-T1 0.08	-1.07	1.23	0.89
T3	3.31(2.29)	T3-T2 -0.87	-2.01	0.28	0.14
Total vegetable ³					
T1	2.70(1.44)	T2-T1 0.63	-0.24	1.50	0.16
T2	3.33(2.29)	T3-T1 0.46	-0.41	1.34	0.29
T3	3.17(2.11)	T3-T2 -0.16	-1.03	0.71	0.71
Total fruit & vegetable					
T1	5.94(3.34)	T2-T1 1.58	-0.11	3.26	0.07
T2	7.51(4.22)	T3-T1 0.54	-1.14	2.23	0.52
T3	6.48(3.88)	T3-T2 -1.03	-2.71	0.65	0.23

¹Time points include T1 (May/June 2011), T2 (August 2011), and T3 (November 2011).

²Fruit includes 100% juice, peach, apple, orange, cantaloupe, other fruits.

³Vegetable includes lettuce salad, cabbage, broccoli, squash or zucchini, sweet potatoes, French fries or fried potatoes, white potatoes, cooked dried beans, tomatoes, tomato sauce, vegetable soup, other vegetables.

Fruit and vegetable consumption among diabetics participating in a federally qualified health center-based farmers' market intervention in rural South Carolina, June–October, 2011 who did and did not increase consumption of fruits and vegetables over time.

Table 3

Time point ²	Non-Increaser ¹ (n=18)		Increaser ¹ (n=23)		p-value for difference
	Mean Fruit and Vegetable Consumption ³ (Standard Deviation)	Mean Fruit and Vegetable Consumption ³ (Standard Deviation)	Mean difference	95% Confidence Interval of the difference	
T1	7.28(3.45)	4.88(2.90)	2.40	0.39 – 4.40	0.02
T2	6.34(3.77)	8.43(4.40)	-2.09	-4.73 – 0.55	0.12
T3	4.80(2.30)	7.80(4.37)	-3.01	-5.16 – -0.85	0.01

¹ Increaser is defined as having average fruit and vegetable consumption at T2 and T3 that was at least 0.5 servings per day greater than consumption at T1. Non-increaser is defined as having average fruit and vegetable consumption at T2 and T3 less than 0.5 servings per day greater than consumption at T1.

² Time points include T1 (May/June 2011), T2 (August 2011), and T3 (November 2011).

³ Expressed as servings per day

Table 4

Odds ratios and 95% confidence intervals of increases in fruit and vegetable consumption over time among diabetics frequenting a FQHC-based farmers' market in rural South Carolina, June–October, 2011 (N=41).

Variables	Odds Ratio	95% Confidence Interval
Self-reported BMI	1.04	0.92–1.17
Payment type		
Study voucher only	38.8**	3.35–445.0
Study voucher + other form of payment	1.00	Referent
Number of farmers' market visits	2.07*	1.09–3.95
Total amount of money spent at the farmers' market	1.02	0.94–1.09
Receipt of food assistance in the past year		
Yes	0.39	0.07–2.08
No	1.00	Referent

*
p<0.05

**
p<0.01

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript