



Published in final edited form as:

J Occup Environ Med. 2014 June ; 56(6): 588–594. doi:10.1097/JOM.000000000000179.

Natural history of upper extremity musculoskeletal symptoms and resulting work limitations over 3 years in a newly hired working population

Ms. Bethany T. Gardner, OTD, OTR/L, Dr. Ann Marie Dale, PhD, OTR/L, Dr. Alexis Descatha, MD, and Dr. Bradley Evanoff, MD, MPH

Department of General Medical Sciences (Ms. Gardner, Dr. Dale, Dr. Evanoff) Washington University School of Medicine, St. Louis, MO; from the Inserm U1018- AP-HP, Occupational Health Unit, F92380 Garches, France (Dr. Descatha)

Abstract

Objective—To describe the proportions of workers with upper extremity (UE) symptoms and work limitations due to symptoms in a newly hired working population over a 3-year study period and to describe transitions between various outcome states.

Methods—827 subjects completed repeat self-reported questionnaires including demographics, medical and work history, symptoms and work status. Outcomes of interest were UE symptoms and work limitations due to symptoms.

Results—72% of workers reported symptoms at least once during the study, with 12% reporting persistent symptoms and 27% reporting fluctuating symptoms. 31% reported work limitations at least once, with 3% reporting consistent work limitations and 8% reporting fluctuating limitations.

Conclusions—UE symptoms and work limitations are common among workers and dynamic in their course. A better understanding of the natural course of symptoms is necessary for targeted interventions.

Background

Transiency of symptoms is a characteristic of many health conditions such as rheumatoid arthritis and multiple sclerosis, such that there are periods of increased disease activity alternating with remission or abatement of symptoms. Previous studies have hypothesized that the course of work-related musculoskeletal disorders (MSDs) may be similar, with several stages of symptom severity from mild discomfort to functionally disabling pain^{1–3}. The transience of MSD symptoms may be due either to the nature of the disorder^{1, 2} or to cyclical or seasonal variance in physical work exposures⁴.

Correspondence to: Bethany Gardner, OTD, OTR/L, Washington University School of Medicine, Division of General Medical Sciences, Campus Box 8005, 660 S. Euclid Avenue, Saint Louis, MO 63110; bgardner@dom.wustl.edu. Phone: 314-747-9318; Fax: 314-454-5113.

The authors declare that they have no conflict of interest relevant to the publication of this manuscript..

Its contents are solely the responsibility of the authors and do not necessarily represent the official view of NIOSH, NCRR, or NIH.

Evanoff et al.,⁵(2014) recently described the complex and multi-factorial nature of MSDs in a simple conceptual model showing a pyramid of disability (Figure 1). As Evanoff outlined, epidemiological studies of MSDs have used a wide variety of case definitions with varying degrees of disease severity and related work disability. Yet the factors influencing progression of MSD, and thus potential targets for intervention, may be different at different stages of disease or disability^{2, 5, 6}. Traditionally, duration of lost work time has been the primary measure of work disability due to musculoskeletal injuries. Lost time as the primary measure of work disability largely underestimates the cost of MSDs, to employers, individual workers, and to society as a whole and misses the earliest opportunity for prevention and intervention efforts.

Most lost productivity, and thus cost, is due to workers who are able to continue working but at less than full ability, rather than from workers who are on lost time⁷⁻¹¹. This phenomenon, of decreased work performance due to a health condition, is sometimes known as “presenteeism”¹²⁻¹⁵. Previous studies showing links between employee health and presenteeism have focused on chronic health conditions including migraines, allergic rhinitis, gastrointestinal disorders, arthritis, and depression in single-employer studies, clinical populations, or national telephone surveys¹⁶⁻²⁸.

Despite the breadth of epidemiological studies of the development, prevention, and treatment of MSDs, and return to work interventions, relatively few studies have examined productivity and functional abilities of workers who remain at work while experiencing musculoskeletal pain^{8, 29, 30, 31, 32}. Most existing studies on presenteeism in workers with MSDs have been conducted with clinical populations who were treatment seeking rather than working populations. These studies have focused on the impact of presenteeism in terms of the cost to employers rather than exploring its causes and the experience of the worker³³.

Due to the high prevalence of upper extremity symptoms and the transient nature of both symptoms and resulting disability, studies of the natural history of upper extremity symptoms and work limitations are necessary to understand the experience of individual workers and the individual and occupational characteristics associated with these limitations. Better understanding of the natural course of symptoms and work limitations can lead to more effective preventive and treatment strategies to improve workers’ functional abilities and provide cost savings for employers. The aims of this study were to describe the proportions of workers with upper extremity symptoms and work limitations due to symptoms in a newly hired working population during repeated follow-ups over a 3-year study period and to describe the transitions between different states of symptoms and work limitations.

Methods

The present study was conducted within an ongoing prospective, longitudinal study of carpal tunnel syndrome and upper extremity musculoskeletal disorders, the Predictors of Carpal Tunnel Syndrome study (PrediCTS). From July 2004 to October 2006, 1107 newly hired workers were recruited from various high and low hand-intensive industries. Inclusion

criteria included a minimum age of 18 years, newly hired or becoming benefits eligible within the last 30 days, working at least 30 hours per week, and English-speaking. Exclusion criteria included having a prior diagnosis of CTS or peripheral neuropathy, being pregnant during study recruitment, or having a contraindication to nerve conduction testing. All subjects provided written informed consent to participate and were compensated for participation. The Washington University School of Medicine Institutional Review Board approved this study.

Data collection and study population

At baseline all subjects completed a self-reported questionnaire including demographics, medical and work history, and current symptom status, nerve conduction studies of the bilateral median and ulnar nerves, and a physical exam of the upper extremities. Questionnaires were repeated at 6 months, 18 months, and 36 months after enrollment to collect information on physical and psychosocial work exposures, symptom status, and work and activities of daily living (ADL) limitations. When follow-up questionnaires were not returned, a second questionnaire was mailed. Subjects who did not return mailed questionnaires were called to complete the survey by phone. Subjects were pursued for up to 6 months after the due date of an unreturned questionnaire.

Data analyses for the present study were limited only to subjects who completed all 4 surveys between baseline and 36-month follow-up since the primary aim of the study was to describe the dynamic and transient natural course of symptoms and work limitations with repeated follow-ups. Figure 2 shows the study design and follow-up rates at each survey time-point. Survey response rates ranged from 80 to 88% at 6, 18, and 36 month follow-up. Overall, 827 subjects (75%) completed all four surveys and were included in the present analyses. Mean follow-up times by study survey were as follows: 6 month follow-up: 7.0 months (range 3.8–18.0), 18 month follow-up: 19.2 months (range 9.1–27.0), and 36 month follow-up: 32.5 months (range 26.6–44.8).

Outcome Measures

For the current study, we defined two primary outcomes of interest: 1) upper extremity (UE) symptoms and 2) work limitations resulting from UE symptoms.

1. **UE symptoms:** At baseline and at each study follow-up, subjects were asked about the presence of upper extremity (UE) symptoms in any one of three upper extremity regions: “In the past YEAR, have you had any RECURRING (repeated) symptoms in your (Neck/shoulder/upper arm, Elbow/forearm, or Hand/wrist/fingers) more than 3 times or lasting more than ONE week?”.
2. **Work limitations due to UE symptoms:** Subjects who reported symptoms were asked to complete additional questions about the effect of their symptoms on their work abilities. Six questionnaire items pertaining to limitations in work abilities, productivity, job restrictions, lost time, and job or company changes due to UE symptoms were available from the follow-up questionnaires at 6, 18, and 36 months (See Appendix). Work ability and productivity questions were similar to items from the University of Michigan Upper Extremity Questionnaire^{34, 35}. A

composite outcome of these items was created such that cases were defined as having any limitations in work ability or productivity or a positive response to any one of the other items.

Statistical analysis

Differences on demographic and clinical characteristics and presence of UE symptoms at baseline were analyzed between the groups of subjects with completed surveys at all study time-points versus subjects with at least one missing survey using the chi-square statistic and student's t-test. To describe the natural history of UE symptoms and work limitations in a working population, the percentage of subjects reporting UE symptoms and work limitations due to UE symptoms were calculated at each study time-point as well as the overall percentages of subjects who reported symptoms or work limitations at least once during the study period. We calculated the percentage of workers whose symptom and/or functional status changed between each study time-point. We also examined whether workers became symptomatic or experienced work limitations and remained symptomatic or limited throughout the study, whether symptoms and/or work limitations resolved, or whether workers remained asymptomatic for the remaining time in the study. We explored the effects of job change and unemployment on symptoms and future work limitations. We calculated the percentage of subjects who changed jobs during the study period. Then, we stratified subjects by symptom status to determine if job changes appeared to be more common among subjects who experienced symptoms during the study. Finally, we conducted a sensitivity analysis to determine the effects of unemployment on symptoms and work limitations. Workers with periods of unemployment were included in the study in order to avoid a uni-directional bias in the data that could have potentially excluded workers who were unable or chose not to work due to their symptoms; we also compared the proportion of subjects with symptoms and work limitations among those without any unemployment to workers reporting periods of unemployment concurrent with follow-up at each study time-point. The analyses were performed using SPSS³⁶, and $p < 0.05$ was considered as statistically significant.

Results

The demographic characteristics of the study population at baseline are shown in Table 1. Study subjects were young, with a mean age of 30.4 years and predominately male (64%). The largest proportion of subjects was employed in construction (40%). Subjects who were missing at least 1 follow-up survey were generally less educated and a higher proportion worked in service industry jobs than the study population; however, the proportion of workers reporting UE symptoms at baseline did not differ significantly between workers with missing surveys and the study population ($p=0.15$).

The natural course of symptoms and work limitations in the study population is graphically displayed in Figure 3. The pyramid of disability⁵ has been modified to reflect the type of data available by time-point in the present study. At baseline, all subjects reported the presence of upper extremity symptoms, shown as 2 levels within the pyramid. Since study subjects were newly hired workers and in many cases had not yet begun performing their

regular work duties at baseline, work limitation status for the new job was not available until 6 months. From 6 month through 36 month follow-up, symptom and work limitation status are shown as three levels, between which workers fluctuated over time. The arrows in the figure show the proportions of workers reporting changes in symptom status and work limitations that occurred since the previous study time-point. The length of the arrow indicates a greater degree of change, with longer arrows indicating movement of 2 levels up or down the pyramid of disability.

At baseline, 31% of subjects reported having UE symptoms in the past year. At 6 month follow-up, the proportion of subjects with UE symptoms increased to 44%, but then remained relatively stable throughout the rest of the study period, 46% at 18 months and 45% at 36 months. At 6 months, approximately one-third of those who reported UE symptoms also reported a limitation in their work activities due to their symptoms (15% of all study subjects). The proportion of subjects with symptoms and work limitations and symptoms alone remained stable from 6 month through 36 month follow-up, although a proportion of subjects within these categories changed. The arrows show that the symptom experience for the individual worker is dynamic, and a considerable percentage of subjects, between 33 and 48%, experienced a change of symptoms or work abilities, either worsening or improving, during each time interval between study follow-ups.

Overall, 596 of 827 workers (72%) reported UE symptoms at least once during the study, yet the proportion of subjects with symptoms was less than 50% at each study time-point. As shown in Table 2, 40% of subjects consistently reported no change in their symptom status: 28% remained asymptomatic and 12% had persistent symptoms throughout the entire 3 year study period. Thus, the majority of subjects (60%) experienced at least 1 change in symptom status, either worsening, improving during the study, with a substantial proportion (27%), experiencing symptom fluctuations, defined as 2 or more changes in symptom status during the study period.

Similarly, 253 of 827 workers (31%) reported work limitations due to UE symptoms at least once during the study; however, the overall proportion of work limitations within a single time-point remained stable at 15% throughout the study. Also similar to symptoms, most workers who reported work limitations experienced at least 1 change in status described as worsened or improved, whereas 8% of workers fluctuated, experiencing 2 or more changes in their work abilities.

Job changes and Unemployment during the study period

Nearly half of the 827 workers in the study population (n=399, 48%) reported at least 1 job change during the three year study follow-up. Job changes were defined as a change in either company or job title which constituted a change in work activities. Table 2 describes the proportion of workers who changed jobs during the study according to symptom and functional work status. Among workers with symptoms at any point during the study, there was a significantly higher proportion of job changes among workers who reported worsening of symptoms over the study period (57%) compared to the asymptomatic workers (47%) whose job changes would have been due to reasons other than symptoms ($p=0.010$). On the contrary, workers whose symptoms improved during the study made the fewest

number of job changes although the difference was not statistically significant ($p=0.352$). Similarly, among workers who reported work limitations due to their symptoms, workers with worsening work abilities reported the highest percentage of job changes (62%) over the study period compared to workers with no work limitations (44%) ($p=0.007$), whereas workers with improved work abilities reported the lowest percentage of job changes (49%), although the difference was not statistically significant ($p=0.855$). Workers who had persistent work limitations (61%, $p=0.181$) or whose work abilities fluctuated (57%, $p=0.117$) also made more job changes than workers with no limitations, however, these differences were not statistically significant.

We examined the effects of unemployment on reported symptoms and work limitations over the course of the study. A sizeable proportion of workers in the study population, 14% ($n=113$), reported at least one period of unemployment concurrent with one of the follow-up surveys. The proportion of unemployed workers increased throughout the study period with 3.3% unemployed at 6 month follow-up, 5.7% at 18 months, and 7.4% at 36 months. The increasing proportion of unemployed workers from 2004 to 2009 followed the upward trend of national unemployment during the recession of 2007–2008.

Of the 113 workers who had a period of unemployment during the study, 45% reported symptoms concurrently. A sensitivity analysis that compared workers who had periods of unemployment versus workers who were employed throughout the study period showed no statistically significant differences in the proportions of workers with symptoms assessed at any time-point during the study (6, 18, 36 months); however, a higher proportion of workers with periods of unemployment reported work limitations due to symptoms at 18 months (22% versus 14%, $p=0.030$) and 36 months (21% versus 15%, $p=0.074$) versus workers with no unemployment.

Discussion

This descriptive study showed the natural course of upper extremity (UE) symptoms and work limitations due to symptoms in a newly hired working population over a three year study period. A considerable majority of workers (72%) reported symptoms at least once during the study, yet less than half of workers reported symptoms within any single follow-up period. In addition, nearly a third of workers (31%) reported work limitations due to their symptoms at least once during the study, but only 15% within any single follow-up period. These results provide evidence for the dynamic nature of both symptoms and work abilities over time, which has been theorized but not explicitly described in previous studies. A better understanding of the natural history of symptoms could help identify targets for early intervention to prevent progression of symptoms and disability.

The increase in the proportion of workers reporting UE symptoms from 31% at baseline to 44% at first follow-up is not surprising since subjects were enrolled in to the PrediCTS study at the time of hire in to a new job. Prior to study enrollment, some subjects had been unemployed or had worked in jobs representing very different physical exposures from their job at enrollment. A significant proportion of study subjects (40%) were just beginning apprenticeship training in the construction trades. Similar associations of musculoskeletal

symptoms with increasing job tenure have been observed in other studies³⁷. The relatively stable proportion of workers with UE symptoms and work limitations at each subsequent follow-up in this study would suggest that cross-sectional studies of risk factors in active workers would yield similar results at any point in time. Yet the sizeable proportion of workers whose symptom and work limitation status changed between study follow-ups, ranging from nearly one-third (32%) to one-half (48%), highlights the need to explore MSDs longitudinally. Cross-sectional studies likely oversample workers whose symptoms are persistent and miss those whose symptoms fluctuate³⁸. The result may be differences in the proportions of workers who would meet a clinical case definition and may result in identification of different risk factors. While we only described the proportions of workers with and without symptoms and not those who progressed to meeting a clinical case definition, musculoskeletal symptoms alone are a precursor of developing a clinical disorder^{2, 3, 39, 40}.

Our findings suggest that there could be a non-linear progression of symptoms to meeting a clinical or epidemiological case definition, as seen in the significant proportion of workers whose symptoms fluctuated (27%). These fluctuating symptoms are important to capture as they may represent an early stage of disease, whereas symptoms that persist over time or cause work limitations may parallel later stages of disease⁴¹. Longitudinal studies commonly report on outcomes over multi-year follow-up periods which may underestimate the prevalence of MSDs in working populations^{38, 42}. Frequency of follow-up is an important consideration in future designs, in order to capture these fluctuations and improve predictive models.

As shown in our findings and in other studies, UE symptoms are common, affecting up to half of workers at any point in time^{3, 43-45}. Yet work ability or productivity has more often been studied in relation to chronic and less common conditions such as rheumatoid arthritis, whereas MSD studies have focused on lost time¹⁵. Presenteeism, or decreased work performance due to MSD symptoms, likely occurs both during the onset of an MSD and during recovery or return to work. However, the factors affecting recovery or disease progression may be different during different stages of disease and for transitions between different stages of impairment. Future studies should examine the natural history and temporal sequence of work ability outcomes related to MSD in order to identify potential differences in risk and prognostic factors at various stages.

Limitations of this study included a general definition of any recurrent UE symptoms that was based on self-report. The high prevalence of disease in our population using self-reported symptoms as the outcome (up to 46%) likely captured a much wider range of disease severity from mild symptoms to severe disease than using a more restrictive epidemiological case definition³⁸. The prevalence of symptoms in this population is consistent with previous studies^{3, 34, 44, 45} but is not a reflection of the proportion of workers who are likely to seek treatment, or whose symptoms will result in an accepted workers' compensation claim. The aim of this study was not to describe prevalence or incidence rates by diagnosis or case definition, but rather to show the transiency of symptoms at the person level. The definition of work limitations was broad and included measures of both presenteeism and absenteeism. Future studies will separate these outcomes to identify

differences in predictors of early versus later stages of work disability due to UE symptoms. While we did have nearly complete employment records for all workers with start and end dates of jobs performed during the study period, we did not have an exact date of onset for symptoms and in many cases could not ascertain whether symptoms preceded job changes. Despite this limitation, we could still see an obvious relationship between worsening symptoms and work ability during the study, and significantly higher number of job changes in symptomatic versus asymptomatic workers, whose job changes could not have been due to symptoms. The timing of job changes with relation to symptoms should be explored in greater detail to inform early intervention efforts.

The major strength of the study is the longitudinal design, which followed a large cohort of workers over a long period of time. We collected repeated measures on several important factors that affect work ability and performance. Our follow-up rates were very good with 93% of subjects completing at least one follow-up survey, and 75% with complete follow-up data for all four surveys during the study period. Limiting study subjects to only those with complete data sets may have eliminated some workers with a high risk of symptoms and work limitations, as those with missing data were less educated and more likely to work in service oriented jobs such as housekeeping and food service. However, there was no difference in study subjects and those with missing data on baseline health indicators such as presence of UE symptoms, prior MSD diagnosis, or comorbid health conditions.

Upper extremity symptoms and work limitations due to symptoms are common and dynamic in their course. Our findings suggest that a significant proportion of working adults experience musculoskeletal symptoms at any given time and have difficulty performing their regular work activities. As shown in recent studies, the cost of absenteeism to employers is exceeded by the costs of presenteeism, due to workers who continue working but at decreased capacity^{10, 11}. Most research and social programs target the relatively smaller number of workers with lost time injuries due to the higher individual costs. Although the proportion of workers with symptoms and work limitations appears to be relatively stable over time, a sizeable proportion of workers fluctuate in and out of symptoms and corresponding changes in work ability. Our study population included a range of low and high physical exposure jobs, so the findings are likely to be generalizable across industries. In order to improve injury and disability prevention programs, it is important to gain a better understanding of the natural course of symptoms to identify better targets for intervention. Subsequent studies should identify the temporal sequence of work limitations and whether there are differences in risk factors for early or later stages of disease and disability and differences in age and social position.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study was supported by CDC/NIOSH (grant # R01OH008017-01), the National Center for Research Resources (NCRR) (grant # UL1 RR024992), a component of the National Institutes of Health (NIH), NIH Roadmap for Medical Research.

References

1. Sluiter JK, Rest KM, Frings-Dresen MH. Criteria document for evaluating the work-relatedness of upper-extremity musculoskeletal disorders. *Scand J Work Environ Health*. 2001; 27 (Suppl 1):1–102. [PubMed: 11401243]
2. Miranda H, Viikari-Juntura E, Heistaro S, Heliovaara M, Riihimaki H. A population study on differences in the determinants of a specific shoulder disorder versus nonspecific shoulder pain without clinical findings. *American Journal of Epidemiology*. 2005; 161:847–855. [PubMed: 15840617]
3. Gold JE, d'Errico A, Katz JN, Gore R, Punnett L. Specific and non-specific upper extremity musculoskeletal disorder syndromes in automobile manufacturing workers. *Am J Ind Med*. 2009; 52:124–132. [PubMed: 19016265]
4. Roquelaure YGY, Gillant JC, Delalieux P, Ferrari C, Mea M, Fanello S, Penneau-Fontbonne D. Transient hand paresthesias in champagne vineyard workers. *Am J Ind Med*. 2001; 40:639–645. [PubMed: 11757040]
5. Evanoff BA, Dale AM, Descatha A. A Conceptual Model On Musculoskeletal Disorders For Occupational Health Practitioners. *International Journal of Occupational Medicine and Environmental Health*. 2014 Feb 8. Epub ahead of print.
6. Gardner BT, Dale AM, Vandillen L, Franzblau A, Evanoff BA. Predictors of upper extremity symptoms and functional impairment among workers employed for 6 months in a new job. *Am J Ind Med*. 2008; 51:932–940. [PubMed: 18651568]
7. Amick BC, Lerner D, Rogers WH, Rooney T, Katz JN. A review of health-related work outcome measures and their uses, and recommended measures. *Spine*. 2000; 25:3152–3160. [PubMed: 11124731]
8. Beaton DE, Kennedy CA. Beyond return to work: testing a measure of at-work disability in workers with musculoskeletal pain. *Qual Life Res*. 2005; 14:1869–1879. [PubMed: 16155774]
9. Lederer V, Loisel P, Rivard M, Champagne F. Exploring the diversity of conceptualizations of work (dis)ability: a scoping review of published definitions. *J Occup Rehabil*. 2013 Jul. Epub ahead of print.
10. Stewart WF, Ricci JA, Chee E, Morganstein D, Lipton R. Lost productive time and cost due to common pain conditions in the US workforce. *JAMA*. 2003; 290:2443–2454. [PubMed: 14612481]
11. Stewart W, Ricci J, Leotta C. Health-related lost productive time (LPT): recall interval and bias in LPT estimates. *J Occup Environ Med*. 2004; 46:S12–S22. [PubMed: 15194892]
12. Aronsson G, Gustafsson K, Dallner M. Sick but yet at work. An empirical study of sickness presenteeism. *J Epidemiol Community Health*. 2000; 54:502–509. [PubMed: 10846192]
13. Hemp P. Presenteeism: at work but out of it. *Harvard Business Review*. 2004; 82:49–58. [PubMed: 15559575]
14. Dew K, Keefe V, Small K. Choosing to work when sick: workplace presenteeism. *Soc Sci Med*. 2005; 60:2273–2282. [PubMed: 15748675]
15. Schulz A, Edington D. Employee health and presenteeism: a systematic review. *J Occup Rehabil*. 2007; 17:547–579. [PubMed: 17653835]
16. Hu X, Markson L, Lipton R, Stewart W, Berger M. Burden of migraine in the United States. *Arch Intern Med*. 1999; 159:813–818. [PubMed: 10219926]
17. Von Korff M, Stewart W, Simon D, Lipton R. Migraine and reduced work performance. A population-based diary study. *Neurology*. 1998; 50:1741–1745. [PubMed: 9633720]
18. Burton W, Conti D, Chen C, Schultz A, Edington D. The impact of allergies and allergy treatment on worker productivity. *J Occup Environ Med*. 2001:64–71. [PubMed: 11201771]
19. Crystal-Peters J, Crown W, Goetzel R, Schutt D. The cost of productivity losses associated with allergic rhinitis. *Am J Manag Care*. 2000; 6:373–378. [PubMed: 10977437]
20. Henke C, Levin T, Henning J, Potter L. Work loss costs due to peptic ulcer disease and gastroesophageal reflux disease in a health maintenance organization. *Am J Gastroenterology*. 2000; 95:788–792.

21. Dean B, Aguilar D, Barghout V, Kahler K, Frech F, Groves D, Ofman J. Impairment in work productivity and health-related quality of life in patients with IBS. *Am J Manag Care*. 2005; 11:S17–S26. [PubMed: 15926760]
22. Sandler R, Everhart J, Donowitz M, Adams E, Cronin K, Goodman C, Gemmen E, Shah S, Avdic A, Rubin R. The burden of selected digestive diseases in the United States. *Gastroenterology*. 2002; 122:1500–1511. [PubMed: 11984534]
23. Lerner D, Reed J, Massarotti E, Wester L, Burke T. The work limitations questionnaire's validity and reliability among patients with osteoarthritis. *J Clin Epidemiol*. 2002; 55:197–208. [PubMed: 11809359]
24. Burton W, Morrison A, Maclean R, Ruderman E. Systematic review of studies of productivity loss due to rheumatoid arthritis. *Occup Med*. 2006; 56:18–27.
25. Adler D, McLaughlin T, Rogers W, Chang H, Lapitsky L, Lerner D. Job performance deficits due to depression. *Am J Psychiatry*. 2006; 163:1569–1576. [PubMed: 16946182]
26. Lerner D, Adler D, Chang H, Lapitsky L, Hood M, Perissinotto C, Reed J, McLaughlin T, Berndt E, Rogers W. Unemployment, job retention, and productivity loss among employees with depression. *Psychiatric Services*. 2004; 55:1371–1378. [PubMed: 15572564]
27. Collins J, Baase C, Sharda C, Ozminkowski R, Nicholson S, Billotti G, Turpin R, Olson M, Berger M. The assessment of chronic health conditions on work performance, absence, and total economic impact for employers. *J Occup Environ Med*. 2005; 47:547–557. [PubMed: 15951714]
28. Bunn W, Pikelny D, Paralkar S, Slavin T, Borden S, Allen H. The burden of allergies-and the capacity of medications to reduce this burden- in a heavy manufacturing environment. *J Occup Environ Med*. 2003; 45:941–955. [PubMed: 14506338]
29. Hagberg M, Tornquist E, Toomingas A. Self-reported reduced productivity due to musculoskeletal symptoms: associations with workplace and individual factors among white-collar computer users. *J Occup Rehabil*. 2002; 12:151–162. [PubMed: 12228946]
30. Campo M, Darragh A. Work-related musculoskeletal disorders are associated with impaired presenteeism in allied health care professionals. *J Occup Environ Med*. 2012; 54:64–70. [PubMed: 22157700]
31. Roy J, MacDermid J, Amick B III, Shannon H, McMurty R, Roth J, Grewal R, Tang K, Beaton D. Validity and responsiveness of presenteeism scales in chronic work-related upper extremity disorders. *Phys Ther*. 2011; 91:254–266. [PubMed: 21212376]
32. Tang K, Pitts S, Solway S, Beaton D. Comparison of the psychometric properties of four at-work disability measures in workers with shoulder or elbow disorders. *J Occup Rehabil*. 2009; 19:142–154. [PubMed: 19301108]
33. Johns G. Presenteeism in the workplace: a review and research agenda. *J Organiz Behav*. 2010; 31:519–542.
34. Franzblau A, Salerno DF, Armstrong TJ, Werner RA. Test-retest reliability of an upper-extremity discomfort questionnaire in an industrial population. *Scand J Work Environ Health*. 1997; 23:299–307. [PubMed: 9322821]
35. Salerno DF, Franzblau A, Armstrong TJ, Werner RA, Becker MP. Test-retest reliability of the Upper Extremity Questionnaire among keyboard operators. *Am J Ind Med*. 2001; 40:655–666. [PubMed: 11757042]
36. SPSS. SPSS Base 20.0. Chicago, IL: SPSS Inc; 2011.
37. Bern SV, Brauer C, Moller KL, Koblauch H, Thygesen LC, Simonsen EB, Alkjaer T, Bonde JP, Mikkelsen S. Baggage handler seniority and musculoskeletal symptoms: is heavy lifting in awkward positions associated with risk of pain? *BMJ Open*. 2013; 3:e004055.
38. Punnett L, Wegman DH. Work-related musculoskeletal disorders: the epidemiologic evidence and the debate. *J Electromyogr Kinesiol*. 2004; 14:13–23. [PubMed: 14759746]
39. Silverstein BA, Viikari-Juntura E, Fan ZJ, Bonauto DK, Bao S, Smith C. Natural course of nontraumatic rotator cuff tendinitis and shoulder symptoms in a working population. *Scandinavian Journal of Work Environment & Health*. 2006; 32:99–108.
40. Descatha A, Dale AM, Franzblau A, Evanoff B. Natural history and predictors of long-term pain and function among workers with hand symptoms. *Arch Phys Med Rehabil*. 2013; 94:1293–1299. [PubMed: 23416766]

41. Riihimäki H. Hands up or back to work: future challenges in epidemiological research on musculoskeletal disorders. *Scand J Work Environ Health*. 1995; 21:401–403. [PubMed: 8824744]
42. National Academy of Sciences and National Research Council. Executive Summary: Musculoskeletal Disorders and the Workplace: Low Back and Upper Extremities. 2001.
43. Franzblau A, Werner R, Valle J, Johnston E. Workplace surveillance for carpal tunnel syndrome: a comparison of methods. *Journal of Occupational Rehabilitation*. 1993; 3:1–14. [PubMed: 24243148]
44. d’Errico A, Caputo P, Falcone U, Fubini L, Gilardi L, Mamo C, Migliardi A, Quarta D, Coffano E. Risk factors for upper extremity musculoskeletal symptoms among call center employees. *J Occup Health*. 2010; 52:115–124. [PubMed: 20179379]
45. Douphrate DI, Gimeno D, Nonnenmann MW, Hagevoort R, Rosas-Goulart C, Rosecrance JC. Prevalence of work-related musculoskeletal symptoms among U.S. large-herd dairy parlor workers. *Am J Ind Med*. 2014; 57:370–379. [PubMed: 24338602]




Figure 1.
Pyramid of Disability.




Figure 2. Study design and survey response at each follow-up.
Note: Excluded subjects represented in grey circles due to missing surveys.




Figure 3.
 Natural course of symptoms and work limitations in the study population.

Table 1

Demographic characteristics of the study population at baseline.

Characteristic	All Participants at baseline (n=1107)		Participants with complete surveys at all time-points (n=827)		Participants missing at least 1 survey (n=280)		p value*
	Mean (SD)	n (%)	Mean (SD)	n (%)	Mean (SD)	n (%)	
Age (years)	30.3 (10.3)		30.4 (10.4)		30.1 (9.9)		0.664
Body mass index (kg/m ²)	28.5 (6.5)		28.3 (6.2)		29.0 (7.4)		0.116
Male Gender		719 (65)		530 (64)		189 (68)	0.301
Job category							0.000 [§]
Construction		450 (41)		334 (40)		116 (42)	
Service		301 (27)		186 (23)		115 (41)	
Technical		146 (13)		123 (15)		23 (8)	
Office/Clerical		210 (19)		184 (22)		26 (9)	
Upper extremity symptoms at baseline		334 (30)		259 (31)		75 (27)	0.150
With at least high school education		1022 (92)		779 (94)		243 (87)	0.000 [§]
Comorbid health condition [†]		58 (5)		46 (6)		12 (4)	0.407
Prior musculoskeletal disorder diagnosis [‡]		114 (10)		91 (11)		23 (8)	0.205

SD- standard deviation, kg- kilograms, m-meters.

* Comparing participants with complete surveys at all time-points and those missing at least 1 survey.

[†] Diabetes mellitus, osteoarthritis, rheumatoid arthritis, or hypertension.[‡] Tendonitis in the fingers, hands, wrists, forearms, or elbows, rotator cuff disorder, ganglion cyst, tendonitis in the shoulders, carpal tunnel syndrome, or ulnar neuropathy.[§] Statistically significant, p<0.05.

Table 2

Comparison of employment changes among workers by upper extremity symptom and functional status during the study period (n=827).

Symptom status (baseline through 36 months)	%	Job change ever[†] (%)	<i>p</i>[*]
Remained Asymptomatic throughout study	28	47	---
Symptoms			
Persistent Symptoms at all time-points	12	47	0.805
Improved from baseline to 36 months	10	43	0.352
Worsened from baseline to 36 months	22	57	0.010 [‡]
Fluctuated	27	45	0.231
Missing	1	---	---
Total	100		

Functional Status (6 to 36 months)	%	Job change ever[†] (%)	<i>P</i>[*]
No work limitations at any time-point	68	44	---
Work limitations			
Persistent at all time-points	3	61	0.181
Progressively improved	10	49	0.855
Progressively worsened	10	62	0.007 [‡]
Fluctuated	8	57	0.117
Missing	1	---	---
Total	100		

* Compared to the proportion of job changes among workers who remained asymptomatic or had no work limitations at any time-point during the study.

[†] Defined as a change in job title or company.

[‡] Statistically significant, $p < 0.05$.