



# HHS Public Access

Author manuscript

*J Autism Dev Disord.* Author manuscript; available in PMC 2015 June 30.

Published in final edited form as:

*J Autism Dev Disord.* 2015 May ; 45(5): 1271–1280. doi:10.1007/s10803-014-2287-3.

## Using Standardized Diagnostic Instruments to Classify Children with Autism in the Study to Explore Early Development

**Lisa D. Wiggins,**

National Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention, 1600 Clifton Road MS E-86, Atlanta, GA 30333, USA

**Ann Reynolds,**

University of Colorado, School of Medicine, Aurora, CO, USA

**Catherine E. Rice,**

National Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention, 1600 Clifton Road MS E-86, Atlanta, GA 30333, USA

**Eric J. Moody,**

University of Colorado, School of Medicine, Aurora, CO, USA

**Pilar Bernal,**

Autism Spectrum Disorders Center, San Jose Medical Center, Kaiser Permanente Northern California, San Jose, CA, USA

**Lisa Blaskey,**

Center for Autism Research, Children's Hospital of Philadelphia, Perelman School of Medicine at University of Pennsylvania, Philadelphia, PA, USA

**Steven A. Rosenberg,**

University of Colorado, School of Medicine, Aurora, CO, USA

**Li-Ching Lee,** and

Department of Epidemiology, Johns Hopkins University, Bloomberg School of Public Health, Baltimore, MD, USA

**Susan E. Levy**

Center for Autism Research, Children's Hospital of Philadelphia, Perelman School of Medicine at University of Pennsylvania, Philadelphia, PA, USA

### Abstract

The Study to Explore Early Development (SEED) is a multi-site case-control study designed to explore the relationship between autism spectrum disorder (ASD) phenotypes and etiologies. The goals of this paper are to (1) describe the SEED algorithm that uses the Autism Diagnostic Interview-Revised (ADI-R) and Autism Diagnostic Observation Schedule (ADOS) to classify children with ASD, (2) examine psychometric properties of different ASD classification methods, including the SEED method that incorporates rules for resolving ADI-R and ADOS discordance,

and (3) determine whether restricted interests and repetitive behaviors were noted for children who had instrument discordance resolved using ADI-R social and communication scores. Results support the utility of SEED criteria when well-defined groups of children are an important clinical or research outcome.

### Keywords

ADI-R; ADOS; Autism; Classification; Phenotypes; Study methods

---

### Introduction

Autism spectrum disorder (ASD) is a developmental disorder that affects social, communication, and behavioral development and is sometimes associated with co-occurring features such as intellectual disability (ID), attention and activity deficits, and abnormal sensory response [American Psychiatric Association (APA) 2006, 2013; Levy et al. 2010], among others. The number of children identified with ASD has increased substantially over the past decade, rising from about 6.4 per 1,000 children in 2002 to about 11.3 per 1,000 children in 2008 [Centers for Disease Control and Prevention (CDC) 2007, 2012]. This increase in the number of children identified with ASD has fueled interest in understanding the complex genetic and environmental factors that may affect ASD phenotypes. Yet defining ASD phenotypes has been challenging given the heterogeneity in symptom presentation and severity and the reliance on behavioral criteria to define the disorder. As such, developing standardized criteria that can both classify and characterize children with ASD is important for clinical practice and epidemiologic research.

It is now well established that best estimate clinical judgment by experienced and reliable clinicians informed by standardized diagnostic instruments is the best predictor of stable ASD diagnoses (Lord et al. 2006). For instance, Lord et al. (2006) found that clinical judgment at 2 years of age predicted ASD diagnoses at 9 years of age more than results of standardized diagnostic instruments—the Autism Diagnostic Interview Revised (ADI-R) and Autism Diagnostic Observation Schedule (ADOS)—when these instruments were used alone (Gotham et al. 2007; Lord et al. 1994, 1999, 2000, 2012; Rutter et al. 2003b). However, scores from the ADI-R and ADOS are both sensitive and specific in detecting children with ASD when used in combination (Risi et al. 2006), and offer several advantages to classify children with ASD in clinical practice and research studies. First, ADI-R and ADOS scores are assigned by experienced and reliable clinicians and offer a uniform method of characterizing ASD symptoms in large cohorts of children that can be replicated in other studies. Second, symptom profiles gleaned from the ADI-R and ADOS allow the opportunity to create ASD sub-groups based on observed and/or reported symptoms that could represent a range of behavioral trajectories and phenotypes. Consequently, using the ADI-R and ADOS to classify children with ASD may be advantageous when well-defined groups of children are an important clinical or research outcome (Schendel et al. 2012).

One of the primary challenges of using ADI-R and ADOS scores to classify children with ASD is that results of the ADI-R and ADOS are not always consistent with each other and there is not a standard method to resolve discordance when one instrument suggests the presence of an ASD and the other instrument does not suggest presence of an ASD (de Bildt et al. 2004; Le Couter et al. 2008). There have been numerous proposals to resolve ADI-R and ADOS discordance or relax ADI-R criteria to include the broader spectrum of ASD in addition to the formerly defined Autistic Disorder (IMGSAC 2001; Risi et al. 2006; Sung et al. 2005). Some of these proposals overlap with one another (Risi et al. 2006; Sung et al. 2005) and define relaxed ADI-R criteria for ASD as meeting the cutoff score on the social deficits domain or communication deficits domain and being within two points of the cutoff score on the social or communication domain not met. These relaxed criteria have been shown to be both sensitive and efficient in detecting children with ASD in a variety of populations (de Bildt et al. 2013; Risi et al. 2006; Sung et al. 2005). However, these criteria have not been evaluated to determine whether they capture children with restricted interests and repetitive behaviors (RRB) noted on the ADOS or ADI-R, which is now a required diagnostic criteria in the Diagnostic and Statistical Manual of Mental Disorders [American Psychiatric Association (APA) 2013].

The Study to Explore Early Development (SEED) is a multi-site case–control study of children 30–68 months of age designed to explore the relationship between ASD phenotypes and various genetic and environmental risk factors (Schendel et al. 2012). A SEED workgroup consisting of ASD diagnostic experts convened to develop a final classification algorithm based on results of the ADI-R and ADOS and to address the complexities of classifying children with ASD based on ADI-R and ADOS scores. The objectives of this paper are to (1) describe the final classification algorithm based on results of the ADI-R and ADOS adopted in SEED, (2) examine the psychometric properties of different ASD classification methods, including the SEED method that incorporates rules for resolving ADI-R and ADOS discordance, and (3) determine whether RRB were noted for children defined as ASD in SEED who had instrument discordance resolved using ADI-R social and communication scores. We hypothesized that SEED ASD criteria would have a good balance of sensitivity and specificity using informed clinical judgment as the referent standard and classify more well-defined children than other ASD classification schemes.

## Methods

### Participant Ascertainment

SEED is a case–control study conducted in six sites across the United States: California, Colorado, Georgia, Maryland, North Carolina, and Pennsylvania. Children born between September 1, 2003 and August 31, 2006 were eligible if they were born in and still resided in one of the six study catchment areas, and lived with a knowledgeable caregiver who was competent to communicate orally in English (or at the California and Colorado sites, in English or Spanish). SEED participants were recruited from one of three ascertainment groups: (1) the general population (POP); (2) children with a broad array of developmental delays or disorders (DD); and (3) children with ASD. Children in the POP group were identified by a sample of state vital records of children born in the target years. Children

with DDs and ASD were identified from multiple education and health providers in each study area who diagnose and serve children with a range of developmental disabilities, including ASD or any ASD-related diagnosis (e.g., ID or language disorder). Children with an ASD or DD diagnosis from a community provider could self-refer into SEED; children without a diagnosis could not self-refer for enrollment in SEED.

### **ASD Screening and Study Data Collection Procedures**

Families in all three recruitment groups were sent a written invitation to participate in the study and if they expressed interest they received a follow-up invitation telephone call; some sites initially contacted participants with both invitation letters and telephone calls. Those families that expressed interest in SEED during an invitation telephone call were given further information about the study. During this call, families were also asked to complete the Social Communication Questionnaire (SCQ) (Rutter et al. 2003a) to screen for ASD. The SCQ recommends a score of 15 points or greater as an indicator of risk for an ASD. However, based on past research that indicates an SCQ score of 11 maximizes sensitivity and specificity in young children, SEED investigators defined an SCQ score of 11 points or higher as an indicator of risk for an ASD (Allen et al. 2007; Lee et al. 2007; Wiggins et al. 2007). Analyses that support a SCQ cutoff score of 11 points in young children were subsequently replicated in the SEED sample (unpublished data).

Children with an SCQ score less than 11 points and without a previous ASD diagnosis were asked to complete a clinic visit consisting of the Mullen Scales of Early Learning (MSEL) (Mullen 1995). If children scored a standard score less than 78 standard points on the MSEL, the Vineland Adaptive Behavior Scales-Second Edition (VABS-II) (Sparrow et al. 2005) was also administered to the parent. Children and parents of children who obtained a score of 11 or higher on the SCQ or had a previous ASD diagnosis were asked to complete a more comprehensive developmental evaluation that consisted of the ADOS, ADI-R, MSEL, and VABS-II. A small number of children who scored less than 11 points on the SCQ and did not have a previously documented ASD diagnosis were also administered the comprehensive evaluation if the study clinician suspected ASD during the clinic visit. Details on these instruments, including average SEED administration time and cut-off scores used in these analyses, are provided in Table 1.

SEED clinicians have advanced degrees in developmental pediatrics, developmental psychology, clinical psychology and related fields and experience with the assessment and diagnosis of children with ASD. Clinicians who administered the ADI-R and ADOS participated in pre-data collection exercises to establish reliability, at least quarterly exercises to maintain reliability, and at least yearly exercises to verify administration fidelity. Overall, quarterly inter-site reliability among SEED clinicians was 99 % on first-pass ADI-R and ADOS exercises and 100 % on second-pass ADI-R and ADOS exercises, for those who did not achieve reliability on the first pass. Quarterly intrasite reliability among SEED clinicians was 87 % on first-pass ADI-R exercises, 99 % on first-pass ADOS exercises, and 100 % on second-pass ADI-R and ADOS exercises for those who did not achieve reliability on the first pass.

### **Clinical Judgment of ASD**

For children who completed a comprehensive developmental evaluation, including the ADI-R and ADOS, a study clinician was asked to complete the Ohio State University (OSU) Autism Rating Scale (OSU Research Unit 2005) adapted for SEED (OARS-adapted). The original OARS has clinicians note the presence of each of the diagnostic criteria outlined in the Diagnostic and Statistical Manual of Mental Disorders-Fourth Edition Text Revision (DSM4; APA 2006) as well as global impression of ASD symptom severity (including whether ASD symptoms can be better accounted for by another disorder). SEED adapted the OARS by having the clinician note his or her degree of certainty that the child had an ASD, which represents the SEED best estimate clinical judgment and was obtained after from considering all information available on the child. Clinical judgment was dichotomized into ASD (scores that indicated clinician was certain the child had an ASD) and non-ASD (scores that indicated the clinician was certain the child did not have an ASD).

### **Final Study Classification**

Based on results from the ADOS and ADI-R, the SEED final classification algorithm was used to classify each child into one of four study groups: (1) ASD, (2) Suspected ASD but incomplete data (i.e., Incomplete Classification), (3) DD, and (4) POP. Some children classified as DD and POP were further divided into one of several sub-classifications to characterize the child by the presence of observed or reported ASD symptoms noted on the SCQ, ADOS, and/or ADI-R if they received a comprehensive evaluation for ASD.

The SEED final classification algorithm was based on best practice guidelines, review of the literature, and clinical experience. The SEED method for resolving discordance between results of the ADI-R and ADOS was originally designed in consultation with the instrument authors and required the child to meet ASD criteria on the revised ADOS algorithms and one of three relaxed ADI-R criteria. The ADI-R criteria were relaxed since the ADI-R was developed to detect individuals with Autistic Disorder rather than the broader range of ASD sought in SEED. The relaxed ADI-R criteria adopted in SEED include the overlapping criteria proposed by Risi et al. (2006), Sung et al. (2005), which are (1) the child met the cutoff score on the ADI-R social deficits domain and was within two points of the cutoff score of the communication deficits domain and (2) the child met the cutoff score on the ADI-R communication deficits domain and was within two points of the cutoff score of the social deficits domain (Risi et al. 2006; Sung et al. 2005) and an additional criteria suggested by an instrument author (3) the child met the cutoff score on the ADI-R social deficits domain and had at least two points noted on the behavioral domain (C. Lord, personal communication).

### **SEED Criteria for ASD**

SEED ASD criteria for children with a mental age of at least 24 months were (1) met the ASD cutoff score on the revised ADOS diagnostic algorithms (Gotham et al. 2007) and the autism cutoff score on the ADI-R diagnostic algorithm or (2) met the ASD cutoff score on the revised ADOS diagnostic algorithms and any one of the three relaxed criteria on the ADI-R diagnostic algorithm noted above.

SEED ASD criteria for children with a mental age less than 24 months were (1) met the ASD cutoff score on the revised ADOS diagnostic algorithms and the autism cutoff score on the ADI-R diagnostic algorithm and the clinician defined the child as ASD on the OARS-adapted or (2) met the ASD cutoff score on the revised ADOS diagnostic algorithms and any one of the three relaxed criteria on the ADI-R diagnostic algorithm noted above and the clinician defined the child as ASD on the OARS-adapted. Clinical judgment was considered for children with a mental age of less than 24 months since the ADI-R is appropriate only for children with a mental age of at least 24 months. However, SEED investigators recognized that many children who have ID also have co-occurring ASD, despite invalid ADI-R scores, and wanted to classify a representative sample of children with ASD. Children with a mental age less than 24 months were excluded from psychometric analyses since clinical judgment was used as the referent standard and to define ASD status for these children.

### **SEED Criteria for Study Classifications Other Than ASD**

Children defined as DD were those who were recruited from an education or clinic source and completed a limited developmental evaluation because ASD risk was not indicated on the SCQ or they completed the comprehensive developmental evaluation but did not meet study criteria for a child with ASD. Children defined as POP were those who were recruited from state vital records and completed a limited developmental evaluation because ASD risk was not indicated on the SCQ or they completed the comprehensive developmental evaluation but did not meet study criteria for a child with ASD. Children defined as Incomplete Classification were those who were asked to complete a comprehensive developmental evaluation but did not complete the evaluation for any reason or completed the evaluation, had a mental age less than 24 months, and the study clinician did not define the child as ASD on the OARS-adapted.

### **Statistical Analyses**

Statistical analyses were conducted with SPSS version 19.0. Descriptive statistics are provided on the number of children defined as ASD, DD, and POP and age, race, sex, and ID. Descriptive statistics are also provided to note whether ASD characteristics were observed, reported, or neither for children who completed a comprehensive evaluation but did not meet SEED ASD criteria. Frequencies of children defined as ASD who had ADI-R and ADOS discordance resolved are provided for the three ADI-R discordance criteria, as well as the ADI-R and ADOS RRB scores for these children.

Sensitivity, specificity, PPV and NPV were assessed for the following ASD classification schemes: (1) ADOS alone, (2) ADI-R alone, (3) concordant ADOS and ADI-R results (i.e., child met the ASD cutoff score on the ADOS and autism cutoff score on the ADI-R), and (4) SEED ASD criteria (noted above; only including children with a mental age of at least 24 months). ASD versus non-ASD, defined by clinician certainty the child had an ASD noted on the OARS-adapted, was used to define best estimate clinical judgment. In our analyses, sensitivity gauged the number of true positives (e.g., ASD based on both the SEED algorithm and clinical judgment) divided by the number of children defined as ASD by the clinician (ASD based on clinical judgment). Specificity gauged the number of true negatives

(e.g., non-ASD based on both the SEED algorithm and clinical judgment) divided by the number of children defined as non-ASD based on clinical judgment. PPV gauged the number of true positives divided by the number of children who were classified as ASD by a particular classification scheme and NPV gauged the number of true negatives divided by the number of children who were not classified as ASD by a particular classification scheme. It is important to note that these statistics represent psychometric properties of a clinical sample that had ASD risk noted on an ASD screen rather than population sensitivity, specificity, PPV, and NPV.

## Results

A total of 3,769 index children were enrolled in SEED, 2,722 (72.2 %) completed a clinic visit, and 2,600 (68.9 %) completed enough of the clinic visit to be classified as ASD or non-ASD. The sample described in this paper is those 2,600 children who were classified as ASD or non-ASD. The SEED final classifications for these children were ASD ( $n = 707$ ), DD ( $n = 995$ ), and POP ( $n = 898$ ). Children defined as DD had a variety of diagnosed conditions reported by parents on the caregiver interview: speech delay (59.3 %), developmental delay (36.2 %), non-specific developmental or learning problem (38.0 %), movement or coordination problem (14.6 %), sensory integration problem (10.5 %), hearing problem (9.0 %), behavior problem (8.5 %), birth defect (7.5 %), attention deficit hyperactivity disorder (7.4 %), Down syndrome (3.9 %), vision impairment (2.8 %), sleep problem (2.6 %), seizure disorder or epilepsy (2.9 %), cerebral palsy (2.4 %), self-injurious behavior (2.0 %), reading disorder (1.6 %), obsessive compulsive disorder (1.1 %), bipolar disorder (0.4 %), Fragile  $\times$  syndrome (0.3 %), neurofibromatosis (0.3 %), reactive attachment disorder (0.2 %), Tourette syndrome (0.1 %), and tuberous sclerosis (0.1 %). These conditions were not mutually exclusive so more than one condition could be noted for each child.

The mean child age at time of clinic visit was 59.4 months ( $SD = 7.1$  months). The racial distribution of the child sample was White (63.6 %), Black (15.9 %), multi-racial (11.7 %), Asian (4.1 %), not stated (4.3 %), and other (0.4 %). Hispanic ethnicity was reported in at least one parent in 15.2 % of the sample. Sixty-six percent of the sample was male; children defined as having ASD had a higher proportion of male children (82.0 % of 707) than children defined as DD (66.7 % of 995) or POP (53.1 % of 898),  $p < .001$ . Children defined as ASD had a higher proportion of ID as defined by the MSEL (62.4 % of 694) than children defined as DD (24.2 % of 988) or POP (2.8 % of 890),  $p < .001$ . The mean MSEL early learning composite for children in our sample were 66.9 points for children classified as ASD, 86.3 points for children classified as DD, and 102.3 points for children classified as POP; MSEL scores between 85 and 115 indicate average learning abilities.

Figure 1 shows final classification details for children defined as ASD, DD, and POP in SEED. There were 1,064 children who received a comprehensive evaluation for ASD and 707 (66.4 %) of these children were defined as ASD. Of the 707 children defined as ASD, 600 (84.9 %) had a previous ASD diagnosis, 97 (13.7 %) had a previous DD but not ASD diagnosis, and 10 (1.4 %) were identified from state vital records. Only 372 children were reported to have a known previous ASD diagnosis at the time of recruitment; the other 228

children with a previous ASD diagnosis had the diagnosis reported by a parent on the invitation call or during the caregiver interview.

There were 305 children defined as DD who received a comprehensive evaluation for ASD but did not meet SEED ASD criteria: 122 (40.0 %) of these children had a previous ASD diagnosis with ASD characteristics observed on the ADOS (n = 38), reported on the ADI-R (n = 26), or neither observed on the ADOS or reported on the ADI-R (n = 58) and 183 (60.0 %) of these children had a previous DD but not ASD diagnosis with ASD characteristics observed on the ADOS (n = 44), reported on the ADI-R (n = 19), or neither observed on the ADOS or reported on the ADI-R (n = 120). There were 52 children defined as POP who received a comprehensive evaluation for ASD but did not meet SEED ASD criteria: 10 had ASD characteristics observed on the ADOS, 3 had ASD characteristics reported on the ADI-R, and 39 did not have ASD characteristics observed on the ADOS or reported on the ADI-R.

A total of 1,017 children who completed a comprehensive evaluation for ASD also had a study clinician complete the OARS-adapted to gauge his or her clinical judgment of whether the child met criteria for ASD. Psychometric properties of the ADOS alone, ADI-R alone, concordant ADI-R and ADOS and SEED ASD criteria were conducted on the sample of children who completed a comprehensive evaluation for ASD, had a study clinician complete the OARS-adapted, and had a mental age of at least 24 months (n = 922). For these children, the ADOS alone yielded the highest sensitivity (.92) but the lowest specificity (.61) compared to other classification schemes (Table 2). The sensitivity and specificity of other classification schemes were relatively comparable, with SEED ASD criteria showing higher sensitivity than the ADI-R alone or concordant positive ADOS and ADI-R results (0.86, 0.77, and 0.75, respectively) and concordant positive ADOS and ADI-R results showing higher specificity than ADI-R alone or SEED ASD criteria (0.82, 0.73, and 0.74, respectively; Table 2). Concordant positive ADOS and ADI-R missed 62 more children defined as ASD by the clinician than SEED ASD criteria. Likewise, SEED ASD criteria classified 27 more children defined as non-ASD by the clinician than concordant positive ADOS and ADI-R.

Figure 2 displays how many of the 1,017 children who completed a comprehensive evaluation for ASD and had clinical judgment noted on the study form had concordant positive ADI-R and ADOS results, concordant negative ADI-R and ADOS results, and discordant ADI-R and ADOS results. Figure 2 also demonstrates the SEED criteria used to resolve ADI-R and ADOS discordance and the mean ADI-R and ADOS RRB scores for children who met each of the discordance criteria. It is important to note that only one of the 96 children who met SEED ADI-R and ADOS discordance criteria did not have any RRB noted on the ADI-R or ADOS. The endorsement rate (i.e., a score of other than 0 or none reported) of RRB noted on the ADI-R for children who had instrument discordance resolved was 58.9 % for repetitive use of objects, 57.9 % for sensitivity to noise, 49.4 % for unusual sensory interests, 40.0 % for circumscribed interests, 32.6 % resistance to changes in own routine or environment, 32.6 % for complex mannerisms, 30.9 % for hand and finger mannerisms, 22.1 % for unusual reactions to specific sensory stimuli, 21.1 % for unusual preoccupations, 20.9 % for unusual attachment to objects, 17.9 % for compulsions or rituals,

5.3 % for midline hand movements, and 3.2 % for resistance to changes in general environment. Children who had an ADI-R and ADOS discordance resolved were more likely to be defined as having ASD by the study clinician (71.9 %) than children who did not have an ADI-R and ADOS discordance resolved (43.4 %),  $p < .001$ .

## Discussion

Diagnostic instruments alone cannot replace informed clinical judgment when diagnosing children with ASD. However, use of diagnostic instruments to classify children with ASD may be advantageous when well-defined groups of children are an important clinical or research outcome. We found that SEED ASD classification based on results of both the ADI-R and ADOS had a good balance of sensitivity and specificity, which supports the utility of these standardized instruments to classify well-defined groups of children in clinical practice and research studies. Use of the SEED-specific ASD classification algorithm offers two major advantages over use of other instrument-based ASD classification schemes because it provides: (1) a method for resolving discordance between the ADI-R and ADOS that classifies more children defined as ASD by an experienced and reliable clinician than concordant positive ADOS and ADI-R results and (2) detailed classifications and sub-classifications that can be replicated in other studies. The SEED final classification algorithm may therefore help guide other clinical and research decisions when ASD risk is indicated on the ADOS but not the ADI-R or detailed sub-classifications are needed to explore ASD behavioral presentations and/or risks.

Our results support previous claims that the ADI-R and ADOS should be used together rather than alone since they offer additive contributions to ASD classification (Risi et al. 2006). For instance, we found high sensitivity but low specificity of the ADOS; sensitivity decreased but specificity improved when the ADOS and ADI-R were used together (concordant positive ADOS and ADI-R). Moreover, sensitivity continued to improve when SEED ASD criteria were applied. Specifically, SEED ASD criteria classified 62 more children with a mental age of at least 24 months than concordant positive ADOS and ADI-R, although there was a slight compromise in specificity (there was no compromise in PPV and an increase in NPV). SEED criteria also classified 87 children with ASD also classified as ASD by the study clinician but had a mental age less than 24 months (data on these children were not presented because clinical judgment was used to classify these children and also used as the referent standard for psychometric analyses). These findings are significant for shaping the future of population-based studies that seek an adequate balance of sensitivity and specificity and rely on sample size to yield the statistical power needed to conduct etiologic and other analyses.

As mentioned previously, the ADOS alone yielded the lowest specificity and highest false positive rate compared to any other classification scheme, which could be an artifact of the young mean age and cognitive abilities of our sample. Previous studies have found the probability of ADOS agreement with diagnostic criteria is inversely related to level of functioning for the child (de Bildt et al. 2004) and the mean cognitive standard score for children classified as ASD in SEED was 66.9 points. Yet the mean cognitive standard score for children in our sample is similar to other studies with similar samples of young children

with ASD (Risi et al. 2006). Specificity of the ADOS and ADI-R improved in our sample when the instruments were used in conjunction rather than when they were used alone. Thus, our findings continue to support use of both the ADI-R and ADOS for evaluation of current developmental status and improvement of ASD classification.

Another advantage of the SEED final classification algorithm is the introduction of detailed classifications and sub-classifications that allow exploration of different pathways of development and the range of behavioral phenotypes. These classifications and sub-classifications are not available when children with ASD are classified using clinician judgment, diagnostic criteria, or one diagnostic instrument alone. We found that 16.9 % of children with a previous ASD diagnosis did not meet SEED criteria for an ASD, which could suggest improvement in symptoms or challenges with ASD assessment and diagnosis at young ages (Lord et al. 2006; Kleinman et al. 2008; Sutera et al. 2007; Turner et al. 2006; Wiggins et al. 2012). Conversely, we found many children met SEED criteria for an ASD or had ASD symptoms noted on the SCQ, ADOS, and/or ADI-R despite the fact they had not been recognized as having an ASD in community settings. Specifically, of the children defined as ASD in SEED, 84.9 % were identified with a previous ASD diagnosis (52.6 % were identified with a previous ASD diagnosis at the time of recruitment), 13.7 % were identified from community providers as a child with DD (but not ASD) and 1.4 % were identified from state vital records. Moreover, many children defined as DD and POP had ASD symptoms noted on the SCQ, ADOS, and/or ADI-R. These results highlight the heterogeneous nature of ASD and could indicate problems with differential diagnosis, continuous distribution of ASD symptoms in child samples, or selection bias among families who agreed to participate in SEED. Further research is needed to describe these different pathways of development, and the detailed nature of the SEED final classification algorithm allows further exploration of characteristics that distinguish children in one sub-classification from children in other sub-classifications. These findings also indicate caution is needed when classifying children for research based solely on existing diagnoses at the time of recruitment or lack of diagnoses at these young ages.

One challenge to using ADI-R and ADOS results to classify children with ASD is that results of these instruments do not always agree with one another. We found discordant ADI-R and ADOS results in 17.6 % of children administered a comprehensive developmental evaluation in SEED, which supports previous findings (de Bildt et al. 2004; Le Couter et al. 2008). Discordant ADI-R and ADOS results thus present a challenge to both clinicians and researchers who rely on these instruments for ASD classification. The discordance criteria adopted in SEED required the child to meet either the cutoff score in the social deficits domain or communication domain but not the RRB domain, even though RRB are now a necessary component of the ASD diagnosis (APA 2013). However, our analyses found that children who had ADI-R and ADOS discordance resolved had some RRB noted on both instruments; only one child who had ADI-R and ADOS discordance resolved did not have any RRB noted on either instrument. More than half of these children had repetitive use of objects and sensitivity to noise noted on the ADI-R. Additionally, these children were more likely to be defined as ASD by a study clinician than children who did not have an ADI-R and ADOS discordance resolved. These results could indicate that young children who meet ADOS ASD criteria and relaxed ADI-R criteria show social-

communication and behavioral deficits reflective of ASD, even if clinically significant RRB have not fully manifested in the child or are not recognized and reported by a parent. These children should therefore still be considered for an ASD classification given the importance of early recognition of and intervention for all children with ASD.

A few limitations to our analyses warrant mention. First, clinical judgment determined by the OARS-adapted and results of the ADI-R and ADOS were not independent of one another, although clinical judgment took into account all information available to the clinician which is comparable to best estimate clinical diagnosis. The clinician who completed the OARS-adapted was often, but not always, the clinician who administered the ADOS and/or had access to ADI-R results. This clinician also had access to SCQ screen results, medical and psychiatric histories, and results of self-administered questionnaires that assessed the behavioral and social functioning of the child. Thus, clinical judgment was based on all available information instead of one or a few diagnostic instruments. Second, clinical judgment was used to define ASD in children with a mental age less than 24 months since low mental age deems the ADI-R invalid. Thus, the psychometric properties of SEED ASD criteria were not presented for these children. Future research is needed to examine the utility of ASD classification schemes for children with very low cognitive scores since ASD and ID often co-occur and may represent an important phenotype to guide etiologic analyses.

Despite these limitations, our findings support the use of ADI-R and ADOS scores in classifying children with ASD in clinical practice and research studies when creating well-defined groups of children is important to the diagnostic or treatment process or research design. SEED ASD criteria may detect more well-defined groups of children than other ASD classification schemes. Future research should explore whether SEED classifications yield children with different behavioral, developmental, and medical profiles so they can be used as important outcomes to explore the relationship between ASD phenotypes and etiologies.

## Acknowledgments


We would like to thank Lisa Croen, Julie Daniels, Ellen Giarelli, Rebecca Landa, Cordelia Robinson, Diana Schendel, Amy Sims, and Patrick Thompson for their contributions to the development of the SEED final classification algorithm and/or comments on previous versions of this paper. We would also like to thank Aimee Alexander, Rebecca Cantrell, and Laura Schieve for their assistance with data cleaning and the SEED principal investigators, co-principal investigators, project coordinators, project staff, and children and families who participated in this research. This publication was supported by six cooperative agreements from the Centers for Disease Control and Prevention: Cooperative Agreement Number U10DD000180, Colorado Department of Public Health; Cooperative Agreement Number U10DD000181, Kaiser Foundation Research Institute (CA); Cooperative Agreement Number U10DD000182, University of Pennsylvania; Cooperative Agreement Number U10DD000183, Johns Hopkins University; Cooperative Agreement Number U10DD000184, University of North Carolina at Chapel Hill; and Cooperative Agreement Number U10DD000498, Michigan State University. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

## References


Allen CW, Silove N, Williams K, Hutchins P. Validity of the Social Communication Questionnaire in assessing risk of autism in preschool children with developmental problems. *Journal of Autism and Developmental Disorders*. 2007; 37:1272–1278. [PubMed: 17080270]

- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4th ed.. American Psychiatric Association; Washington, DC: 2006.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed.. American Psychiatric Publishing; Arlington, VA: 2013.
- Centers for Disease Control and Prevention; Autism and Developmental Disabilities Monitoring Network Surveillance Year 2002 Principal Investigators. Prevalence of autism spectrum disorders—Autism and Developmental Disabilities Monitoring Network, 14 sites, United States, 2002. *Morbidity and Mortality Weekly Report*. 2007; 56(No. SS-1):12–28.
- Centers for Disease Control and Prevention; Autism and Developmental Disabilities Monitoring Network Surveillance Year 2008 Principal Investigators. Prevalence of autism spectrum disorders—Autism and Developmental Disabilities Monitoring Network, 14 sites, United States, 2008. *Morbidity and Mortality Weekly Report*. 2012; 61:1–19.
- De Bildt A, Oosterling I, van Lang N, Kuijper S, Dekker V, Sytema S, et al. How to use the ADI-R for classifying autism spectrum disorders? Psychometric properties of criteria from the literature in 1,204 Dutch children. *Journal of Autism and Developmental Disorders*. 2013; 43:2280–2294. [PubMed: 23397166]
- De Bildt A, Sytema S, Ketelaars C, Kraijer D, Mulder E, Volmar F, et al. Inter-relationship between Autism Diagnostic Observation Schedule—Generic (ADOS-G), Autism Diagnostic Interview—Revised (ADI-R), and the Diagnostic and Statistical Manual of Mental Disorders (DSM4-TR) classification in children and adolescents with mental retardation. *Journal of Autism and Developmental Disorders*. 2004; 34:129–137. [PubMed: 15162932]
- Gotham K, Risi S, Pickles A, Lord C. The Autism Diagnostic Observation Schedule: Revised algorithms for improved diagnostic validity. *Journal of Autism and Developmental Disorders*. 2007; 37:613–627. [PubMed: 17180459]
- International Molecular genetic Study of Autism Consortium (IMG-SAC). A genomewide screen for autism: Strong evidence for linkage to chromosomes 2q, 7q, and 16p. *American Society of Human Genetics*. 2001; 69:570–581.
- Kleinman JM, Ventola P, Pandey J, Verbalis AD, Barton M, Hodgson S, et al. Diagnostic stability in very young children with an autism spectrum disorder. *Journal of Autism and Developmental Disorders*. 2008; 38:606–615. [PubMed: 17924183]
- Le Couter A, Haden G, Hammal D, McConachie H. Diagnosing autism spectrum disorders in pre-school children using two standardized assessment instruments: The ADI-R and ADOS. *Journal of Autism and Developmental Disorders*. 2008; 38:362–372. [PubMed: 17605097]
- Lee L-C, David AB, Rusyniak J, Landa R, Newschaffer CJ. Performance of the Social Communication Questionnaire in children receiving preschool special education services. *Research in Autism Spectrum Disorders*. 2007; 1:126–138.
- Levy SE, Giarelli E, Lee L-C, Schieve LA, Kirby RS, Cunniff C, et al. Autism spectrum disorder and co-occurring developmental, psychiatric, and medical conditions among children in multiple populations of the United States. *Journal of Developmental and Behavioral Pediatrics*. 2010; 31:267–275. [PubMed: 20431403]
- Lord C, Risi S, DiLavore P, Shulman C, Thurm A, Pickles A. Autism from 2 to 9 years of age. *Archives of General Psychiatry*. 2006; 63:694–701. [PubMed: 16754843]
- Lord C, Risi S, Lambrecht L, Cook EH Jr, Leventhal BL, DiLavore PC, et al. The Autism Diagnostic Observation Schedule-generic: A standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders*. 2000; 30(3):205–223. [PubMed: 11055457]
- Lord, C.; Rutter, M.; DiLavore, PC.; Risi, S. Autism Diagnostic Observation Schedule. Western Psychological Services; Los Angeles, CA: 1999.
- Lord, C.; Rutter, M.; DiLavore, PC.; Risi, S. Autism Diagnostic Observation Schedule. 2nd ed.. Western Psychological Services; Los Angeles, CA: 2012.
- Lord C, Rutter M, Le Couteur AL. Autism Diagnostic Interview-Revised: A revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism and Developmental Disorders*. 1994; 24(5):659–685. [PubMed: 7814313]
- Mullen, E. Mullen scales of early learning. Pearson; San Antonio, TX: 1995.

- Risi S, Lord C, Gotham K, Corsello C, Chrysler C, Szatmari P, et al. Combining information from multiple sources in the diagnosis of autism spectrum disorders. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2006; 45:1094–1103. [PubMed: 16926617]
- Rutter, M.; Bailey, A.; Lord, C. SCQ: Social Communication Questionnaire. Western Psychological Services; Los Angeles, CA: 2003a.
- Rutter, M.; Le Couteur, A.; Lord, C. ADI-R: The Autism Diagnostic Interview-Revised. Western Psychological Services; Los Angeles, CA: 2003b.
- Schendel DE, DiGiuseppi C, Croen LA, Fallin MD, Reed PL, Schieve LA, et al. The Study to Explore Early Development (SEED): A multisite epidemiologic study of autism by the Centers for Autism and Developmental Disabilities Research and Epidemiology (CADDRE) Network. *Journal of Autism and Developmental Disabilities*. 2012 epub ahead of print.
- Sparrow, S.; Balla, D.; Cicchetti, D. Vineland adaptive behavior scales. 2nd ed.. Pearson; San Antonio, TX: 2005.
- Sung Y, Dawson G, Munson J, Estes A, Schellenberg GD, Wijsman EM. Genetic investigation of quantitative traits related to autism: Use of multivariate polygenic models with ascertainment adjustment. *American Journal of Human Genetics*. 2005; 76:68–81. [PubMed: 15547804]
- Sutera S, Pandey J, Esser E, Rosenthal MA, Wilson LB, Barton M, et al. Predictors of optimal outcome in toddlers diagnosed with autism spectrum disorders. *Journal of Autism and Developmental Disorders*. 2007; 37:98–107. [PubMed: 17206522]
- The Ohio State University (OSU) Research Unit on Pediatric Psychopharmacology. OSU Autism Rating Scale (OARS; adapted for SEED) and Clinical Global Impression (CGI; adapted for SEED). 2005. Retrieved August 30, 2010 from, <http://psychmed.osu.edu/resources.htm>
- Turner L, Stone W, Pozdol S, Coonrod E. Follow-up of children with autism spectrum disorders from age 2 to age 9. *Autism*. 2006; 10:243–265. [PubMed: 16682397]
- Wiggins LD, Baio J, Lee L-C, Nicholas J, Rice C. Retention of autism spectrum disorder diagnoses by community professionals: Findings from the Autism and Developmental Disabilities Monitoring Network, 2000 & 2006. *Journal of Developmental and Behavioral Pediatrics*. 2012; 33:387–395. [PubMed: 22580734]
- Wiggins LD, Bakeman R, Adamson LB, Robins DL. The utility of the Social Communication Questionnaire in screening for autism in children referred for early intervention. *Focus on Autism and Other Developmental Disabilities*. 2007; 22:33–38.



**Fig. 1.** Final study classifications for children enrolled in the Study to Explore Early Development. All children defined as ASD received a comprehensive evaluation for autism spectrum disorder (ASD) and 11 of these children scored less than 11 points on the Social Communication Questionnaire (SCQ) and did not have a previous ASD diagnosis; Children were not tested for ASD if they did not have a previous ASD diagnosis, scored less than 11 points on the SCQ, and a study clinician did not suspect ASD during a limited evaluation.



**Fig. 2.** Performance on the Autism Diagnostic Interview-Revised (ADI-R) and Autism Diagnostic Observation Schedule (ADOS) among Children in the Study to Explore Early Development (SEED) tested for Autism Spectrum Disorder (ASD). Note that SEED ADI-R and ADOS discordance criteria were not mutually exclusive so children could meet more than one criteria.

**Table 1**  
Instruments administered during Study to Explore Early Development (SEED) developmental evaluations

Data collection and average administration time <sup>a</sup>	Reference; version	Domains	Cut-off scores required for SEED ASD classification	Scores used in SEED ASD classification
Autism Diagnostic Interview-Revised (ADI-R); 150 min	Lord et al. (1994, Rutter et al. 2003b) WPS Version	Social Communication Restricted interests and repetitive behaviors (RRB) Developmental delays or deficits	Standard autism cut-off scores: Social = 10 Communication = 8 for verbal children and 7 for nonverbal children RRB = 3 Relaxed ASD criteria when the child meets ADOS criteria for ASD but not ADI-R criteria for autism: Met the social cutoff and was within two points of the communication cutoff (5 or more points for nonverbal children and 6 or more points for verbal children) Met the communication cutoff and was within two points of the social cutoff (8 or more points) Met the social cutoff and had at least two points noted on the behavioral domain.	Meets standard autism criteria Meets one of three relaxed ASD criteria
Autism Diagnostic Observation Scale-2 (ADOS); 60 min	Gotham et al. (2007), Lord et al. (2012) WPS version	Social affect RRB	Module 1 with no words = 11 Module 1 with some words = 8 Module 2 less than 59 months = 7 Module 2 more than 59 months = 8 Module 3 = 7	Meets ASD criteria
Mullen Scales of Early Learning; 60 min	Mullen (1995); 1995 AGS/Pearson version	Early learning composite Expressive language Receptive language Visual reception Fine motor skills	N/A	Clinic visit age Visual reception age equivalent dichotomized into less than 24 months and 24 month or more to determine mental age
Ohio State University (OSU) Autism Rating Scale (OARS)—adapted for SEED; 10 min	OSU Research Unit, 2005; adapted for SEED	Presence and degree of severity of ASD diagnostic symptoms Degree of impairment associated with ASD Clinician degree of certainty the child has an ASD	N/A	Five-point Likert rating of clinician degree of certainty the child has an ASD dichotomized into “uncertain” (scores of 1–3 or note that ASD symptoms better accounted for by another disorder) and “certain” (scores of 4–5)
Vineland Adaptive Behavior Scales, Second Edition; 60 min	Sparrow et al. (2005); 2005 AGS/Pearson Version	Adaptive behavior composite Social Communication Daily living skills Motor skills	N/A	N/A

Average administration time based on feedback from SEED supervising site clinicians

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 2**

Psychometric properties of autism spectrum disorder (ASD) classification schemes for children in the Study to Explore Early Development (SEED)

	ASD	Non-ASD	Sensitivity	Specificity	Positive predictive value	Negative predictive value
ADOS met	536	133	0.92	0.61	0.80	0.81
ADOS not met	48	205				
ADI-R met	450	90	0.77	0.73	0.83	0.65
ADI-R not met	134	248				
Concordant ADOS + ADI-R met	438	60	0.75	0.82	0.88	0.66
Concordant ADOS + ADI-R not met	146	278				
SEED ASD criteria met	500	87	0.86	0.74	0.85	0.75
SEED ASD criteria not met	84	251				

The sample of children in these analyses were those who had a comprehensive evaluation for ASD and clinical judgment noted on the study form (n = 1,017) and a mental age of at least 24 months (n = 922 of 1,017)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript