



# HHS Public Access

Author manuscript

*J Occup Environ Med.* Author manuscript; available in PMC 2016 July 01.

Published in final edited form as:

*J Occup Environ Med.* 2015 July ; 57(7): 743–750. doi:10.1097/JOM.0000000000000484.

## Predictors Associated with Changes of Weight and Total Cholesterol among two Occupational Cohorts over 10 Years

**Ulrike Ott, PhD, MSPH,**

Rocky Mountain Center for Occupational and Environmental Health, Department of Family and Preventive Medicine, University of Utah School of Medicine, 391 Chipeta Way, Suite C, Salt Lake City, Utah 84108, Fax: 801-581-1224, Phone: 801-581-4800

**Joseph B. Stanford, MD, MSPH,**

Department of Family and Preventive Medicine, University of Utah School of Medicine, Salt Lake City, Utah

**Maureen A. Murtaugh, PhD, RD,**

Division of Epidemiology, University of Utah School of Medicine, Salt Lake City, Utah

**Jessica L.J. Greenwood, MD, MSPH,**

Department of Family and Preventive Medicine, University of Utah School of Medicine, Salt Lake City, Utah

**Lisa H. Gren, PhD, MSPH,**

Division of Family and Preventive Medicine, Department of Public Health, University of Utah School of Medicine, Salt Lake City, Utah

**Kurt T. Hegmann, MD, MPH, and**

Rocky Mountain Center for Occupational and Environmental Health, Department of Family and Preventive Medicine, University of Utah School of Medicine, Salt Lake City, Utah

**Matthew S. Thiese, PhD, MSPH**

Rocky Mountain Center for Occupational and Environmental Health, Department of Family and Preventive Medicine, University of Utah School of Medicine, Salt Lake City, Utah

Ulrike Ott: [Ulrike.ott@hsc.utah.edu](mailto:Ulrike.ott@hsc.utah.edu)

### Abstract

**Objective**—To ascertain worker health characteristics and psychosocial factors associated with changes in body weight and total cholesterol (TC) among two production operation populations.

**Methods**—We performed descriptive and predictive analysis of questionnaire data and biomedical measurements from two prospective cohort studies. Our key outcomes were changes in weight, and TC over 5–10 years between baseline and exit assessments.

---

Correspondence to: Ulrike Ott, [Ulrike.ott@hsc.utah.edu](mailto:Ulrike.ott@hsc.utah.edu).

The authors have nothing to disclose.

**Results**—146 subjects were analyzed. Increases in weight were associated with belief in being overweight and baseline overweight and obesity. Increases in TC levels were associated with female gender, belief that TC levels were “not good,” and feeling depressed.

**Conclusion**—Most of the reported associations with increases in weight and TC levels are amenable to interventions and may be a target for workplace intervention programs.

## Introduction

One in every six adults (16.3%) has high total cholesterol (TC)  $\geq 240$ mg/dl and 35.5% of US adults are obese (Body Mass Index (BMI)  $\geq 30.0$  kg/m<sup>2</sup>).<sup>1, 2</sup> Both elevated cholesterol and BMI have been associated with workplace absenteeism and health care costs.<sup>3</sup> Henke et al estimated that cholesterol, weight, blood pressure and glucose, had the greatest impact on total health care costs among workers in a manufacturing plant.<sup>4</sup>

The CDC estimates that overall medical costs related to obesity for U.S. adults were \$147 billion in 2008.<sup>5</sup> The US economic productivity losses due to obesity are projected to be between \$48 billion to \$66 billion per year by 2030.<sup>6</sup> In addition to cardiovascular health concerns, obesity has also been associated with musculoskeletal or joint-related pain in the feet<sup>7</sup>, knees<sup>8–12</sup>, back<sup>13–17</sup>, shoulders<sup>18–23</sup>, and hands.<sup>24, 25</sup> Additionally, obesity has been associated with an increased risk of occupational injuries.<sup>26–28</sup>

An individual’s perceived risk of developing a certain health condition is likely essential in motivating behavior.<sup>29, 30</sup> Adults with elevated TC or BMI may be more motivated to alter their lifestyle because of health concerns. Therefore, it appears necessary that adults have an understanding of key health indicators as well as recommended target levels. A study to assess whether better knowledge improves adherence to lifestyle changes in patients with coronary heart disease concluded that “patient education must be formalized and acknowledged as an official part of the health care system.”<sup>31</sup> This suggests that people who are aware of their TC levels may be more likely to reduce their blood cholesterol levels.

To the best of our knowledge, no prior research has been conducted assessing the changes in TC and weight over time among production workers. Therefore, the goal of this study was to ascertain characteristics associated with changes in weight and TC from baseline enrollments to study completion. We were particularly interested in assessing whether knowledge of TC and BMI levels were associated with changes.

## Methods

This research study was nested within the Utah populations of two prospective cohorts (the WISTAH Distal Upper Extremity (DUE) cohort and the BackWorks Low Back Pain (LBP) cohort).<sup>32, 33</sup> Both cohorts were approved by the University of Utah’s Institutional Review Board (#s 00010930 and 00011889). Baseline data for these cohorts were collected during worksite enrollments conducted between 2002 and 2007. Additional data were collected during study completion visits in the Spring of 2012 and analyzed in 2013. The parent cohort studies have detailed methods papers published.<sup>32, 33</sup> Thus, a brief summary of the methods follows.

## Subjects

Subjects were at least 18 years of age at enrollment and employed at one of eight participating companies in Utah. Participants were excluded if they could not give informed consent, did not speak either English or Spanish and were planning to retire within 4 years of study enrollment. Subjects for this nested study were recruited from five different employment settings in Utah which included: airbag manufacturing, sewing facility, office work, red meat processing, and printing operations.<sup>32, 33</sup> Only a subset of both cohorts had TC measured at baseline and this subset was eligible for the present analysis.

## Baseline Measures

At baseline enrollments, workers completed a laptop administered questionnaire under the supervision of a research assistant. Data quantified at baseline included demographics (age, gender, race, marital status, and education level), leisure-time physical activity, tobacco use, psychosocial factors (e.g., depression, job satisfaction, family problems), and health status (e.g. “Have you even been told by a physician that you have high cholesterol (Laboratory test result over 200 mg/dl)”).

Questions addressed 21 leisure-time physical activities (e.g., walking, baseball, basketball) and could include additional activities beyond those 21. Each of those activities was further queried for the number of months per year, the average number of times per week, and the average number of minutes each activity was performed. A composite of all these activities was calculated and the total reported leisure-time physical activity in minutes per week was determined.

Height and weight were measured in stocking feet to calculate BMI. Height was assumed to have not changed appreciably during the study. If weight exceeded 200kg, two scales were used simultaneously and the sum was recorded. BMI  $<18.5\text{kg/m}^2$  was classified as underweight, between  $18.5\text{--}24.9\text{kg/m}^2$  was classified as normal weight,  $25\text{--}29.9\text{kg/m}^2$  was classified as overweight and  $>30\text{kg/m}^2$  was classified as obese.

Serum non-fasting TC, low density lipoprotein (LDL), high density lipoprotein (HDL), triglycerides, C - reactive protein, and hemoglobin A1c levels were measured in blood collected by venipuncture and analyzed at ARUP Labs in Salt Lake City or by fingerstick and analyzed using Alere Cholestech LDX system (Alere Inc., Waltham, MA).

Blood pressure was measured in a seated position after a minimum of 5 minutes of rest using automated cuffs (Omron HEM-780).

Study participants were informed of their weight and blood pressure results upon completion of baseline enrollments by a trained researcher. Immediate feedback regarding those results was provided in writing indicating desired ranges for each of those measures. A handout was given to each participant which listed the measured systolic and diastolic blood pressure. Recommendations for normal, pre-hypertension, stage 1 hypertension, and stage 2 hypertension were also listed indicating whether a lifestyle modification is encouraged or not. Blood test results were mailed to the participants upon receipt of the blood results from the laboratory. The mailer contained the current classifications and recommendations for

adult TC, low density lipoprotein, high density lipoprotein, triglycerides, C - reactive protein, and hemoglobin A1c. Participants were advised to consult with either their healthcare provider if any of the reported values were out of range.

### Study Completion Measures

Age at study completion was recalculated. Participants completed another laptop-administered questionnaire. Survey items quantified leisure-time physical activity outside work, and psychosocial factors (e.g., depression, job satisfaction, and family problems) with the same questions as in the baseline questionnaire. It also included items regarding knowledge of BMI, knowledge of TC levels, and fruit and vegetable intake.

Dietary intake questions were included that have been previously developed.<sup>34</sup> Fruit and vegetable intake was assessed by asking “*How many times do you typically eat a serving of fruit in one day?*” and “*How many times do you typically eat a serving of vegetables in one day?*” Binary dummy variables were created for both of these variables (<5 vs. 5).

Breakfast and fast food consumptions were assessed by asking “*How many times do you eat restaurant or fast food in a typical week?*” and “*How many times do you typically eat breakfast in one week (7 days)?*” Dummy variables reporting tertiles were created for both breakfast and fast food consumption.

Participants were asked whether they could recall their current TC and BMI (*Yes/No*). Questions on how study participants perceive their weight and TC were also included. Subjects were asked “*Do you think your Total Cholesterol is: Good, Not Good, Unsure?*” and “*Do you believe you are Underweight, Normal Weight, Overweight, Obese, Unsure?*” Intake of cholesterol-lowering medication was also assessed.

Having received any education (e.g. doctor, the internet, magazines) in weight management, diet, nutrition or physical fitness throughout the study duration was also assessed.

### Outcome Variables

Changes in weight were determined by comparing the measured weight at study completion visit with the weight at the baseline visit.

Blood samples were drawn to measure at study completion via finger stick method using the Alere Cholestech LDX system (Alere Inc., Waltham, MA). These data enabled assessment of changes in TC levels throughout the course of the study.

### Data Analysis

All analyses were performed using SAS 9.3 (SAS Institute, Cary, North Carolina). Outliers and missing data were verified by pulling individual charts for each participant. Imputation using the study population mean was used when missing data could not be verified. Less than 0.3% of all data were imputed.

Variables were analyzed for normality and skewness. Mean differences for weight and TC changes between baseline enrollments and study completion visits were determined by using

a paired *t*-tests (normal distribution) and Wilcoxon signed rank sum test (not normally distributed). Statistical significance was determined using  $\alpha$  level of 0.05. Frequencies, means and standard deviations were used to describe the population.

We assessed the data for attrition bias since a large proportion of our population exited the study. We aimed to determine whether those workers who exited the study have different characteristics than those who completed the study and would have therefore introduced attrition bias. Differences were assessed for demographics (age, gender, race, education, and marital status) by using chi square test analyses. In addition, we assessed differences for our main continuous health outcomes (weight and TC) by using the Wilcoxon-Mann-Whitney test.

Multivariate linear regression was conducted to identify which factors were independently associated with the main outcomes. Factors for the multivariate model were selected based on evidence published in other research articles and biological plausibility. Stepwise backwards regression analyses were performed separately for each outcome.

To investigate whether the relationships between participant characteristics, BMI, and TC would differ between the cohorts, we ran linear regression models including interaction terms between the cohort status (DUE or LBP) and the predictor variables. We used an a-priori p-value of 0.1 for significance of each interaction term.

## Results

A total of 366 subjects met the baseline inclusion criteria. More than half exited the study for various reasons unrelated to the study, which are detailed in Figure 1 (e.g. leaving employment to take another job, retirement, termination, etc.). Another 38 were lost to follow up at study completion visits. A total of 146 subjects remained in the cohort through study completion and participated in the end of study visits.

At study completion, participants were between ages 35–55 years ( $n=86$ , 58.9%) with a mean age of 49.6 (SD=10.6) years. The majority were female ( $n=74$ , 50.7%), and white ( $n=98$ , 67.1%). Most ( $n=106$ , 72.6%) were married and 46.6% ( $n=68$ ) had a high school degree or GED. More than half of the population was obese ( $n=84$ , 57.5%) with a mean BMI of 31.7kg/m<sup>2</sup> (SD=7.4). Only 4.8% ( $n=7$ ) of the subjects at exit reported knowing their BMI, although 58.9% ( $n=86$ ) believed themselves to be overweight. Less than 10% reported knowing their TC. Only 57.5% ( $N=84$ ) of workers consumed 5 servings of fruits and vegetables combined per day. (Table 1)

A plurality of the participants ( $n=65$ , 45%) had abnormal TC levels ( $> 200$ mg/dl) at baseline enrollments. However, 57% ( $n=37$ ) with abnormal baseline TC levels reported not having been informed about those abnormal levels by a health professional.

Table 2 shows the mean weight and TC comparing baseline measures and study completion measures. The mean TC became significantly lower ( $194.4\pm 36.3$  vs.  $182.1\pm 37.8$ ;  $P<0.0001$ ). In contrast, mean weight was significantly higher at study completion ( $86.9\pm 23.6$  vs.  $90.5\pm 24.5$ ;  $P=0.003$ ).

Attrition bias analyses indicated that only age significantly differed between those who exited the study and those who completed it ( $p < 0.0001$ ). Statistical differences between groups were not found for gender ( $p = 0.3$ ), race ( $p = 0.2$ ), marital status ( $p = 0.2$ ), education ( $p = 0.3$ ), weight ( $p = 0.1$ ) and TC levels ( $p = 0.6$ ).

Associations with weight changes among production workers are shown in Table 3. Workers who believed they were overweight or obese at study completion gained significantly more weight (6.7kg increase, 14.4g increase) compared to those who believed they were of normal weight.

Those who consumed breakfast  $< 6$  times per week (the lowest tertile of breakfast consumption) lost 4.4kg ( $p = 0.058$ ), as compared to those eating breakfast daily. However that association was only borderline significant. No significant associations were found between fast food consumption and weight changes. Reported physical activity at baseline was not associated with weight changes.

Associations with TC changes are reported in Table 4. TC level changes for workers who indicated “always” feeling depressed were significantly higher (112.4 mg/dl,  $p = 0.009$ ) than those who indicated “never.” However, only two workers indicated “always” feeling depressed. Workers who indicated “often” having family problems experienced lower TC levels ( $-36.5$  mg/dl,  $p = 0.002$ ) as compared to those who indicated “never.” Workers who believed their TC levels to be “no good” increased their TC levels (17.6 mg/dl,  $p = 0.02$ ) than those who indicated “good.” TC levels among female workers increased more so than among males throughout the course of the study (16.0 mg/dl,  $p = 0.007$ ). Eating breakfast  $< 6$  times per week (the lowest tertile of breakfast consumption) was associated with reduced TC levels as compared to eating breakfast every day ( $-16.9$  mg/dl,  $p = 0.005$ ).

## Discussion

This study found the sole characteristic associated with weight gain over the study duration is a belief they are overweight or obese. A characteristic associated with weight reduction was being overweight or obese at baseline. Characteristics associated with increases in TC levels over the study duration included: female gender, belief their TC levels were “not good,” and feeling depressed. Characteristics associated with TC reductions included: having family problems and consuming breakfast  $< 6$  times per week. To our surprise, fast food consumption was not associated with weight or TC level changes, except for breakfast consumption.

Surprisingly, baseline leisure-time physical activity was not associated with TC changes. Workers reported baseline leisure-time physical activity that met exercise guidelines<sup>35</sup> (mean  $282.6 \pm 314.3$ ), but not at study completion ( $119.6 \pm 173.1$ ). A meta-analysis of 95 studies assessing exercise effects on serum lipid and lipoprotein levels found exercise lowered cholesterol levels by 7 to 13 mg/dl compared with controls,<sup>36, 37</sup> with larger reductions among those losing weight. However, questionnaires assessing physical activity levels are subject to recall errors and biases. We suspect that most workers over-reported their leisure-time physical activity levels.

We were also surprised that fat intake and fast food consumption were not significantly associated with TC or weight changes. Reductions in saturated fat, dietary cholesterol, and weight are considered to offer the most effective dietary strategies for reducing total cholesterol.<sup>38, 39</sup> However, controlled studies have reported only modest long-term reductions in TC.<sup>37</sup> Short-term decreases in TC of 10% to 20% have resulted from a controlled low-fat diet.<sup>37</sup> This study's findings may be partially due to a single assessment of dietary intake at exit among the workers.

We found significant associations between depression and increasing TC levels. These findings are somewhat concurrent with other research studies. Symptoms of depression and anxiety have been associated with decreased levels of HDL cholesterol and increased abdominal obesity.<sup>40</sup> Researchers found that anxiety is a proxy risk factor for depression severity in aggravating dyslipidemia.<sup>40</sup>

Less than 10% of workers reported knowing their TC levels at study exit, although all had received results from this study many years previously. Most workers (53.9%) who believed their TC was "not good" actually had normal TC levels (<200mg/dl). Yet, considering only 57% of workers with abnormal baseline TC levels recalled having been informed of their TC levels by a physician, suggests a need for more intensive interventions.

These results also showed no significant associations between health education and changes in TC and weight. It is widely believed that awareness and knowledge of factors associated with negative health outcomes is necessary before health promotion programs can be successfully implemented.<sup>31, 41-44</sup>

The prevalence of obesity in this working population (57.5%) was greater than the U.S. adult population (35.7%).<sup>1</sup> This is also greater than the reported obesity prevalence among the general population in Utah (22.5%)<sup>45</sup>, and is twice as high as a recently reported prevalence for U.S. workers (27.7%).<sup>46</sup>

Given the impacts of BMI and TC on healthcare and occupational costs,<sup>3-7</sup> this research analyzed a target population that should be disposed to improvements in weight, TC and other weight-related comorbidities.

Strengths of this study include 1) anthropometric measures, 2) ability to collect data from the same population up to 9 years apart, 3) recruitment from a wide array of employment settings to improve generalizability of the results, and 4) computerized data collection methods of questionnaires.

Several factors limit these findings. Measurements were taken at two time points, which were mostly 5 to 9 years apart. We also cannot address temporal relationships for measures taken only at the exit, particularly knowledge of BMI, TC and dietary recall. Recall error and bias may particularly affect reporting of physical activity levels. Prior evidence suggests that overweight individuals tend to over-estimate their activity level.<sup>47, 48</sup>

## Conclusion

Results suggest that most of the reported associations with increases in weight and TC levels among production workers are amenable to interventions and may be a target for workplace intervention programs. The need for these programs is also warranted by the high prevalence of obesity among workers. Given that the state of Utah ranks among the lowest obesity prevalence rates in the U.S., we were able to fill a gap in the literature but also identify a target population that is in need of weight loss interventions.

## Acknowledgments

The authors wish to acknowledge the contributions of Arun Garg, PhD and Jay Kappelusch, PhD on the WISTAH Distal Upper Extremity cohort and the BackWorks Low Back Pain cohort.

This research was supported in part by Centers for Disease Control and Prevention/National Institute of Occupational Safety and Health (NIOSH) Grant Number 1R01OH009155-01 and NIOSH Training Grant Number T42/CCT810426.


## References

1. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity in the United States, 2009–2010. NCHS data brief. 2012;1–8.
2. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA*. 2012; 307:491–497. [PubMed: 22253363]
3. Bertera RL. The effects of behavioral risks on absenteeism and health-care costs in the workplace. *J Occup Med*. 1991; 33:1119–1124. [PubMed: 1765851]
4. Henke RM, Carls GS, Short ME, et al. The relationship between health risks and health and productivity costs among employees at Pepsi Bottling Group. *J Occup Environ Med*. 2010; 52:519–527. [PubMed: 20431407]
5. [Accessed: February 4, 2013] Chronic Disease Prevention and Health Promotion: Obesity. Available at: <http://www.cdc.gov/chronicdisease/resources/publications/aag/obesity.htm>
6. Trust for America's Health and the Robert Wood Johnson Foundation. F as in Fat: How Obesity Threatens America's Future. [serial online]. 2012;2013(September). Available from: <http://healthyamericans.org/report/100/>
7. Irving DB, Cook JL, Young MA, Menz HB. Obesity and pronated foot type may increase the risk of chronic plantar heel pain: a matched case-control study. *BMC musculoskeletal disorders*. 2007; 8:41. [PubMed: 17506905]
8. Blagojevic M, Jinks C, Jeffery A, Jordan KP. Risk factors for onset of osteoarthritis of the knee in older adults: a systematic review and meta-analysis. *Osteoarthritis and cartilage/OARS, Osteoarthritis Research Society*. 2010; 18:24–33.
9. Wendelboe AM, Hegmann KT, Biggs JJ, et al. Relationships between body mass indices and surgical replacements of knee and hip joints. *Am J Prev Med*. 2003; 25:290–295. [PubMed: 14580629]
10. Kohatsu ND, Schurman DJ. Risk factors for the development of osteoarthrosis of the knee. *Clinical orthopaedics and related research*. 1990:242–246. [PubMed: 2245553]
11. Bagge E, Bjelle A, Eden S, Svanborg A. Factors associated with radiographic osteoarthritis: results from the population study 70-year-old people in Goteborg. *The Journal of rheumatology*. 1991; 18:1218–1222. [PubMed: 1941827]
12. Felson DT. The epidemiology of knee osteoarthritis: results from the Framingham Osteoarthritis Study. *Seminars in arthritis and rheumatism*. 1990; 20:42–50. [PubMed: 2287948]
13. Heuch I, Hagen K, Zwart JA. Body mass index as a risk factor for developing chronic low back pain: a follow-up in the Nord-Trondelag Health Study. *Spine*. 2013; 38:133–139. [PubMed: 22718225]



14. Leino-Arjas P, Solovieva S, Kirjonen J, Reunanen A, Riihimaki H. Cardiovascular risk factors and low-back pain in a long-term follow-up of industrial employees. *Scandinavian journal of work, environment & health*. 2006; 32:12–19.
15. Miranda H, Viikari-Juntura E, Punnett L, Riihimaki H. Occupational loading, health behavior and sleep disturbance as predictors of low-back pain. *Scandinavian journal of work, environment & health*. 2008; 34:411–419.
16. Tubach F, Leclerc A, Landre MF, Pietri-Taleb F. Risk factors for sick leave due to low back pain: a prospective study. *J Occup Environ Med*. 2002; 44:451–458. [PubMed: 12024690]
17. Van Nieuwenhuysse A, Crombez G, Burdorf A, et al. Physical characteristics of the back are not predictive of low back pain in healthy workers: a prospective study. *BMC musculoskeletal disorders*. 2009; 10:2. [PubMed: 19123931]
18. Morken T, Moen B, Riise T, et al. Prevalence of musculoskeletal symptoms among aluminium workers. *Occup Med (Lond)*. 2000; 50:414–421. [PubMed: 10994244]
19. Silverstein BA, Bao SS, Fan ZJ, et al. Rotator cuff syndrome: personal, work-related psychosocial and physical load factors. *J Occup Environ Med*. 2008; 50:1062–1076. [PubMed: 18784555]
20. Miranda H, Viikari-Juntura E, Martikainen R, Takala EP, Riihimaki H. A prospective study of work related factors and physical exercise as predictors of shoulder pain. *Occupational and environmental medicine*. 2001; 58:528–534. [PubMed: 11452048]
21. Luime JJ, Verhagen AP, Miedema HS, et al. Does this patient have an instability of the shoulder or a labrum lesion? *Jama*. 2004; 292:1989–1999. [PubMed: 15507585]
22. Wendelboe AM, Hegmann KT, Gren LH, Alder SC, White GL Jr, Lyon JL. Associations between body-mass index and surgery for rotator cuff tendinitis. *The Journal of bone and joint surgery American volume*. 2004; 86-A:743–747. [PubMed: 15069138]
23. Roquelaure Y, Ha C, Rouillon C, et al. Risk factors for upper-extremity musculoskeletal disorders in the working population. *Arthritis and rheumatism*. 2009; 61:1425–1434. [PubMed: 19790112]
24. Nathan PA, Keniston RC, Myers LD, Meadows KD. Obesity as a risk factor for slowing of sensory conduction of the median nerve in industry. A cross-sectional and longitudinal study involving 429 workers. *J Occup Med*. 1992; 34:379–383. [PubMed: 1564575]
25. Andersen RE, Crespo CJ, Bartlett SJ, Bathon JM, Fontaine KR. Relationship between body weight gain and significant knee, hip, and back pain in older Americans. *Obesity research*. 2003; 11:1159–1162. [PubMed: 14569039]
26. Janssen I, Bacon E, Pickett W. Obesity and its relationship with occupational injury in the canadian workforce. *Journal of obesity*. 2011; 2011:531403. [PubMed: 21773008]
27. Pollack KM, Sorock GS, Slade MD, et al. Association between body mass index and acute traumatic workplace injury in hourly manufacturing employees. *American journal of epidemiology*. 2007; 166:204–211. [PubMed: 17485732]
28. Garg A, Kapellusch J, Hegmann K, et al. The Strain Index (SI) and Threshold Limit Value (TLV) for Hand Activity Level (HAL): risk of carpal tunnel syndrome (CTS) in a prospective cohort. *Ergonomics*. 2012; 55:396–414. [PubMed: 22397385]
29. Marcus BH, Rakowski W, Rossi JS. Assessing motivational readiness and decision making for exercise. *Health psychology: official journal of the Division of Health Psychology, American Psychological Association*. 1992; 11:257–261.
30. Marcus BH, Simkin LR. The stages of exercise behavior. *The Journal of sports medicine and physical fitness*. 1993; 33:83–88. [PubMed: 8350613]
31. Alm-Roijer C, Stagmo M, Uden G, Erhardt L. Better knowledge improves adherence to lifestyle changes and medication in patients with coronary heart disease. *Eur J Cardiovasc Nurs*. 2004; 3:321–330. [PubMed: 15572021]
32. Garg A, Hegmann KT, Moore JS, et al. Study protocol title: a prospective cohort study of low back pain. *BMC musculoskeletal disorders*. 2013; 14:84. [PubMed: 23497211]
33. Garg A, Hegmann KT, Wertsch JJ, et al. The WISTAH hand study: a prospective cohort study of distal upper extremity musculoskeletal disorders. *BMC musculoskeletal disorders*. 2012; 13:90. [PubMed: 22672216]

34. Greenwood JL, Murtaugh MA, Omura EM, Alder SC, Stanford JB. Creating a clinical screening questionnaire for eating behaviors associated with overweight and obesity. *J Am Board Fam Med.* 2008; 21:539–548. [PubMed: 18988721]
35. [Accessed: March 25, 2014] 2008 Physical Activity Guidelines for Americans. Available at: <http://www.health.gov/paguidelines/pdf/paguide.pdf>
36. Tran ZV, Weltman A. Differential effects of exercise on serum lipid and lipoprotein levels seen with changes in body weight. A meta-analysis. *JAMA: the journal of the American Medical Association.* 1985; 254:919–924.
37. [Accessed: August 20, 2012] Screening for Lipid Disorders. Available at: <http://www.ahrq.gov/downloads/pub/prevent/pdfser/lipidser.pdf>
38. Lichtenstein AH, Van Horn L. Very low fat diets. *Circulation.* 1998; 98:935–939. [PubMed: 9738652]
39. Fletcher B, Berra K, Ades P, et al. AHA Scientific Statement. Managing abnormal blood lipids: a collaborative approach. *Circulation.* 2005; 112:3184–3209. [PubMed: 16286609]
40. van Reedt Dortland AK, Giltay EJ, van Veen T, Zitman FG, Penninx BW. Longitudinal relationship of depressive and anxiety symptoms with dyslipidemia and abdominal obesity. *Psychosomatic medicine.* 2013; 75:83–89. [PubMed: 23197842]
41. Han W, Piterman L, Morris N, McCall L. The Knowledge Of And Attitudes Towards Serum Cholesterol In The Chinese Patients In A Family Practice In Hong Kong. *Hong Kong Practitioner.* 1998; 20:307–318.
42. Tanaka T, Okamura T, Yamagata Z, et al. Awareness and treatment of hypertension and hypercholesterolemia in Japanese workers: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. *Hypertens Res.* 2007; 30:921–928. [PubMed: 18049023]
43. Celentano A, Panico S, Palmieri V, et al. Citizens and family doctors facing awareness and management of traditional cardiovascular risk factors: results from the Global Cardiovascular Risk Reduction Project (Help Your Heart Stay Young Study). *Nutr Metab Cardiovasc Dis.* 2003; 13:211–217. [PubMed: 14650353]
44. Celentano A, Palmieri V, Arezzi E, et al. Cardiovascular secondary prevention: patients' knowledge of cardiovascular risk factors and their attitude to reduce the risk burden, and the practice of family doctors. The "Help Your Heart Stay Young" study. *Ital Heart J.* 2004; 5:767–773. [PubMed: 15626274]
45. [Accessed: April 20, 2014] Overweight and Obesity: Utah State Nutrition, Physical Activity, and Obesity Profile. Available at: <http://www.cdc.gov/obesity/stateprograms/fundedstates/pdf/utah-state-profile.pdf>
46. Luckhaupt SE, Cohen MA, Li J, Calvert GM. Prevalence of obesity among U.S. Workers and associations with occupational factors. *Am J Prev Med.* 2014; 46:237–248. [PubMed: 24512862]
47. Jakicic JM, Polley BA, Wing RR. Accuracy of self-reported exercise and the relationship with weight loss in overweight women. *Medicine and science in sports and exercise.* 1998; 30:634–638. [PubMed: 9565948]
48. Lichtman SW, Pisarska K, Berman ER, et al. Discrepancy between self-reported and actual caloric intake and exercise in obese subjects. *The New England journal of medicine.* 1992; 327:1893–1898. [PubMed: 1454084]



**Figure 1. Flowchart of study participants from baseline enrollments to study completion measures**

\*Participants were missed for the following reasons: vacation, sick, all-day meetings, worked graveyard shift

**Table 1**

Population Demographic Characteristics with Self-reported Health Indicators at Study Completion (N=146)

	N	%
<b>Age</b>		
<35	29	19.9
35– 55	86	58.9
>55	31	21.2
<b>Gender*</b>		
Female	74	50.7
Male	72	49.3
<b>Race*</b>		
White	98	67.1
African American or Black	2	1.4
Asian	2	1.4
Pacific Islander or Native Hawaiian	15	10.2
Hispanic or Latino	29	19.9
<b>Marital Status*</b>		
Never Married (Single)	23	15.8
Married	106	72.6
Divorced	13	8.9
Separated	4	2.7
<b>Education Level*</b>		
Some high school	7	4.8
High school graduated or GED	68	46.6
Some college	57	39.0
College graduate (Bachelor's degree or higher)	14	9.6
<b>Measured BMI Status (kg/m<sup>2</sup>)</b>		
Underweight (BMI < 18.5)	2	1.4
Normal (BMI 18.5–24.9)	19	13.0
Overweight (BMI 25–29.9)	41	28.1
Obese (BMI ≥ 30)	84	57.5
<b>Do you know what your Body Mass Index is?</b>		
Yes	7	4.8
No	139	95.2
<b>Do you believe you are:</b>		
Underweight	9	6.2
Normal Weight	31	21.2
Overweight	86	58.9

	N	%
Obese	18	12.3
Unsure	2	1.4
<b>Do you know what your current TOTAL Blood Cholesterol is?</b>		
Yes	12	8.2
No	134	91.8
<b>Do you think your Total Cholesterol is:</b>		
Good	67	45.9
Not good	26	17.8
Unsure	53	36.3
<b>Do you currently take Cholesterol –Lowering Medication?</b>		
Yes	29	19.9
No	117	80.1
<b>Restaurant or Fast Food Consumption (per week)</b>		
1	92	63.0
2	23	15.8
3	31	21.2
<b>Breakfast Consumption (per week)</b>		
<6	51	34.9
6	10	6.9
7	85	58.2
<b>Fruit and Vegetables Combined (Servings/Day)</b>		
<5	62	42.5
5	84	57.5
<b>How Often do you have Family Problems that Irritate or Bother You</b>		
Never	28	19.2
Sometimes	101	69.2
Often	14	9.6
Always	3	2.0
<b>How Often do you feel Down, Blue or Depressed</b>		
Never	54	37.0
Sometimes	77	52.7
Often	13	8.9
Always	2	1.4
<b>How Often do you Feel Nervous</b>		
Never	37	25.4
Sometimes	90	61.6
Often	18	12.3
Always	1	0.7

	N	%
<b>How satisfied are you with your job?</b>		
Very satisfied	72	49.3
Somewhat satisfied	58	39.7
A little satisfied	13	8.9
Not at all satisfied	3	2.1
<b>Mean±SD</b>		
Age (Years)	49.6±10.6	
BMI (kg/m <sup>2</sup> )	31.6±7.4	
Leisure-Time Physical Activity (min/week)*	282.6±314.3	
Leisure-Time Physical Activity (min/week)	119.6±173.1	

\* Assessed at Baseline

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 2**

Comparison of Weight and Total Cholesterol at baseline and study completion (N=146)

<b>Mean ± SD</b>			
	<b>Baseline Measures</b>	<b>Study Completion Measures</b>	<b>P-Value</b>
Weight (kg)	86.9±23.6	90.5±24.5	0.003
Total Cholesterol (mg/dl)	194.4±36.3	182.1±37.8	<0.0001
<200mg/dl N (%) <sup>*</sup>	81 (55.5)	102 (69.9)	
≥200mg/dl N (%) <sup>*</sup>	65 (44.5)	44 (30.1)	

<sup>\*</sup>We categorized TC based on the ATP III Classification: Desirable (<200mg/dl) and Borderline High/High (≥200mg/dl)<sup>35</sup>

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 3**

Multivariate linear regression analyses of associations with weight change (kg)\* from baseline to study completion.

	Change in weight (kg) from baseline	Standard Error	P-Value
<b>Baseline Age (Years)</b>	-0.1	0.1	0.56
<b>Baseline BMI (kg/m<sup>2</sup>)</b>			
Underweight	1.1	7.8	0.89
Normal Weight		Reference	
Overweight	-7.5	3.0	0.01
Obese	-11.2	3.1	0.0003
<b>Baseline Leisure-Time Total Physical Activity (min/week)</b>			
0-240		Reference	
240-480	3.9	2.9	0.16
>480	0.9	2.9	0.75
<b>Gender</b>			
Female	-1.4	2.3	0.53
Male		Reference	
<b>Breakfast Consumed (Times/Week)</b>			
<6	-4.4	2.3	0.06
6	-5.3	4.3	0.21
7		Reference	
<b>Nutrition/Diet Education**</b>			
Yes		Reference	
No	-0.3	2.9	0.92
<b>Physical Fitness Education**</b>			
Yes		Reference	
No	-0.5	2.5	0.83
<b>Weight Management Education**</b>			
Yes		Reference	
No	4.1	2.7	0.13
<b>Do you know what your Body Mass Index (BMI) is?</b>			
Yes		Reference	
No	5.7	4.7	0.23
<b>Do you believe you are:</b>			
Underweight	-4.0	5.1	0.43
Normal Weight		Reference	
Overweight	6.7	2.7	0.02



	Change in weight (kg) from baseline	Standard Error	P-Value
Obese	14.4	3.9	0.0004
Not sure or don't know	-2.6	9.0	0.77
<b>Do you think your Total Cholesterol is:</b>			
Good		Reference	
Not Good	-0.4	2.8	0.9
Unsure	1.2	2.4	0.6
<b>Fast Food Consumed/Week***</b>			
1		Reference	
2	0.7	2.9	0.8
3	1.3	2.6	0.6
<b>Cohort</b>			
DUE		Reference	
LBP	-0.4	2.3	0.9

\* Weight Changes=Weight Study Completion (kg) – Weight Baseline (kg)

\*\* Throughout the duration of this study, have you received any education (your doctor, the internet, magazines etc) in:

\*\*\* How many times do you eat restaurant or fast food in a typical week? (For example Chili's™, McDonald's™, Burger King™, or your local diner. Take home leftovers count twice)

**Table 4**

Multivariate linear regression analyses of associations with TC changes (mg/dl)\* from baseline to study completion.

	Change in TC (mg/dl) from baseline	Standard Error	P-Value
<b>Baseline Age (Years)</b>	-0.7	0.3	0.04
<b>Baseline BMI (kg/m<sup>2</sup>)</b>			
Underweight	12.6	19.1	0.51
Normal Weight		Reference	
Overweight	-7.3	7.2	0.31
Obese	-8.3	6.9	0.23
<b>Baseline Leisure-Time Total Physical Activity (min/week)</b>			
0-240		Reference	
240-480	5.9	7.8	0.45
>480	9.7	7.5	0.20
<b>Baseline Cholesterol (md/dl)</b>			
<200		Reference	
200	-15.0	5.5	0.007
<b>Gender</b>			
Female	16.0	5.8	0.007
Male		Reference	
<b>Nutrition/Diet Education</b> ***			
Yes		Reference	
No	10.0	7.7	0.19
<b>Physical Fitness Education</b> ***			
Yes		Reference	
No	-3.5	6.6	0.60
<b>Weight Management Education</b> ***			
Yes		Reference	
No	-7.6	6.8	0.27
<b>Do you think your Total Cholesterol is:</b>			
Good		Reference	
Not Good	17.6	7.4	0.02
Unsure	-2.3	6.3	0.72
<b>Do you know what your current TOTAL Blood Cholesterol is?</b>			
Yes		Reference	
No	-3.1	10.3	0.76
<b>Do you currently take Cholesterol – Lowering Medication?</b>			

	Change in TC (mg/dl) from baseline	Standard Error	P-Value
Yes		Reference	
No	29.0	6.8	<0.0001
<b>Fast Food Consumed/Week **</b>			
1		Reference	
2	7.9	7.6	0.3
3	3.8	6.7	0.6
<b>Fruit and Vegetables Combined (Servings/Day)</b>			
<5	-3.9	5.5	0.48
5		Reference	
<b>Breakfast Consumed (Times/Week)</b>			
<6	-16.9	5.9	0.005
6	-11.3	11.2	0.31
7		Reference	
<b>Feeling Down, Blue or Depressed</b>			
Never		Reference	
Sometimes	5.2	5.6	0.35
Often	14.7	11.0	0.18
Always	112.4	42.3	0.009
<b>Family Problems that Irritate or Bother You</b>			
Never		Reference	
Sometimes	-3.4	6.8	0.62
Often	-36.5	11.6	0.002
Always	-35.5	35.7	0.32
<b>Cohort</b>			
DUE		Reference	
LBP	4.6	6.3	0.46

\* TC Changes=Study Completion TC – Baseline TC

\*\* How many times do you eat restaurant or fast food in a typical week? (For example Chili's™, McDonald's™, Burger King™, or your local diner. Take home leftovers count twice)

\*\*\* Throughout the duration of this study, have you received any education (your doctor, the internet, magazines etc) in: