

HHS Public Access

Author manuscript *Mol Cancer Ther.* Author manuscript; available in PMC 2016 March 01.

Published in final edited form as:

Mol Cancer Ther. 2015 March ; 14(3): 740-746. doi:10.1158/1535-7163.MCT-14-0972.

CD30 is a potential therapeutic target in malignant mesothelioma

Snehal Dabir¹, Adam Kresak², Michael Yang², Pingfu Fu³, Gary Wildey¹, and Afshin Dowlati^{1,*}

¹Division of Hematology and Oncology, Case Comprehensive Cancer Center and University Hospitals Case Medical Center, Cleveland, Ohio, 44106

²Pathology, Case Comprehensive Cancer Center and University Hospitals Case Medical Center, Cleveland, Ohio, 44106

³Biostatistics, Case Comprehensive Cancer Center and University Hospitals Case Medical Center, Cleveland, Ohio, 44106

Abstract

CD30 is a cytokine receptor belonging to the tumor necrosis factor superfamily (TNFRSF8) that acts as a regulator of apoptosis. The presence of CD30 antigen is important in the diagnosis of Hodgkin's disease and anaplastic large cell lymphoma. There have been sporadic reports of CD30 expression in non-lymphoid tumors, including malignant mesothelioma. Given the remarkable success of brentuximab vedotin, an antibody-drug conjugate directed against CD30 antigen, in lymphoid malignancies, we undertook a study to examine the incidence of CD30 in mesothelioma and to investigate the ability to target CD30 antigen in mesothelioma. Mesothelioma tumor specimens (N = 83) were examined for CD30 expression by immunohistochemistry. Positive CD30 expression was noted in 13 mesothelioma specimens, primarily those of epithelial histology. There was no significant correlation of CD30 positivity with either tumor grade, stage or survival. Examination of four mesothelioma cell lines (H28, H2052, H2452, and 211H) for CD30 expression by both FACS analysis and confocal microscopy showed that CD30 antigen localized to the cell membrane. Brentuximab vedotin treatment of cultured mesothelioma cells produced a dose-dependent decrease in cell growth and viability at clinically relevant concentrations. Our studies validate the presence of CD30 antigen in a subgroup of epithelial-type mesothelioma tumors and indicate that selected mesothelioma patients may derive benefit from brentuximab vedotin treatment.

Keywords

Mesothelioma; CD30; brentuximab vedotin; immunohistochemistry; predictive biomarkers

Conflict of interest: The authors have no conflict of interest to declare

^{*}Corresponding Author: Afshin Dowlati, MD, Department of Medicine, Division of Hematology and Oncology, Case Comprehensive Cancer Center and University Hospitals Case Medical Center, 11100 Euclid Avenue, Mail Stop LKS 5079, Cleveland, Ohio, 44016. Phone: 216-286-6741; FAX: 216-844-5234; afshin.dowlati@case.edu.

Introduction

Malignant mesothelioma is a locally aggressive tumor with the most common form being malignant pleural mesothelioma, representing approximately 70% of all mesothelioma diagnoses. Diagnosis at an early stage of the disease is often difficult because of its nonspecific signs and symptoms, and consequently a large majority of the patients are initially diagnosed at an advanced stage. Mesothelioma is associated with occupational asbestos exposure in most industrialized countries (1). Many emerging countries have yet to prohibit asbestos usage, which predicts that a great number of the patients will suffer from this disease in the future. In addition, there is also growing concern that the widespread use of nano-sized particles for medical and industrial purposes may create new risks for developing mesothelioma. Treatment for advanced malignant mesolthelioma is comprised of combination chemotherapy containing platinum and pemetrexed (2). Since approval of this disease. Thus, there is an urgent need to develop new therapeutic strategies against this disease.

Antibody-based therapies and immunotoxins have shown remarkable therapeutic activities in various tumors and might represent a promising approach against mesothelioma (2, 3). This approach targets antigens specifically expressed on cancerous cells, resulting in an increased therapeutic efficacy with minimal systemic toxicity. CD30 is a member of the tumor necrosis factor receptor super family (TNFRSF8) and is highly expressed in lymphoma with limited expression in healthy tissues, making it an ideal target for antibodybased therapies (4-8). Recently, in clinical trials for relapsed/refractory Hodgkin lymphoma (HL) and anaplastic large cell lymphoma (ALCL) (9, 10), brentuximab vedotin demonstrated high response rates as a single agent leading to its accelerated approval by the Food and Drug Administration (FDA) for treatment of these lymphomas under the brand name Adcetris®. Brentuximab vedotin is an antibody-drug conjugate, which couples an anti-CD30 antibody (SGN-30) to the synthetic anti-tubulin agent MMAE (11).

While several reports have identified CD30 antigen in other types of solid tumors, including malignant mesothelioma (4, 5, 12), these findings have not been systematically followed up, which is important for broadening the family of CD30-positive cancers that are potential targets for brentuximab vedotin. Here we validate the presence of CD30 in a panel of mesothelioma tumors by immunohistochemistry and demonstrate its cell surface location in mesothelioma cell lines by confocal microscopy and flow cytometry. Importantly, the efficacy of brentuximab vedotin on mesothelioma cell lines is demonstrated using clinically relevant drug concentrations.

Materials and Methods

Cell Lines

The human small cell lung cancer cell line H526 and mesothelioma cell lines H2052, H2452, 211H and H28 were purchased from the American Type Culture Collection (Manassas, VA) within the last three years and used at a passage of 7. These thoracic cell lines were not authenticated because of their direct purchase from the ATCC and low

passage number. Karpas cells were a gift from Dr. H. Koon and while not authenticated, served as a positive control and responded to drug treatment in accordance with published reports (9, 10, 13, 14) Cell lines were maintained in Dulbecco's Modified Eagle Medium (DMEM)/Ham's F-12 medium supplemented with 10% (v/v) fetal bovine serum (FBS, Hyclone, ThermoFisher Scientific, Waltham, MA) in a 5% CO₂ humidified incubator at 37° C.

Confocal Microscopy

Mesothelioma cells were seeded onto a glass-bottomed culture dish (MatTek Corp.). Cells were fixed with 4% formaldehyde diluted in PBS. After 1 h blocking with 1% bovine serum albumin in PBS, cells were incubated with a rabbit polyclonal anti-CD30 antibody (Novus Biologicals, NBP1-72175; 1:100) at room temperature for 2 h and washed three to four times with PBS. Incubation with Alexa Fluor 488–conjugated goat anti-rabbit IgG (Molecular Probes; 1:500) was done for 1 h at room temperature, followed by repeated washes using PBS. After subsequent staining of nuclei with 1 μ M DRAQ5 in PBS for 30 min, confocal images were obtained on a Zeiss LSM510 NLO laser-scanning microscope using single-line (488 nm) or multitrack sequential excitation (488 and 633 nm). Images were acquired and processed with Zeiss LSM Image Browser software.

Flow cytometry analysis of CD30 expression

Expression of CD30 on the surface of mesothelioma cell lines was confirmed by FACS analysis, using a fluorescent dye conjugated secondary antibody Alexa Fluor 488– conjugated anti-rabbit IgG (Molecular Probes; 1:500). Mesothelioma cells were collected by centrifugation, fixed with ethanol by incubating at least 30 min at room temperature. Centrifuged again and supernatant decanted off. Cells were permeablized or non-permeabilized with or without Triton X-100 (0.1% in PBS) for 10 min, briefly centrifuged and the resulting supernatant removed. The cells were suspended in FACS buffer with or without diluted primary (rabbit polyclonal anti-CD30 antibody, NBP1-72175, 1:100) or secondary (Alexa Fluor 488–conjugated anti-rabbit IgG, 1:500) antibodies followed by 30 min incubation on ice. The stained cells were analyzed on a LSR II flow cytometer (BD Biosciences; San Jose, CA). FACS data were analyzed by using FlowJo software (Tree Star, Ashland, OR).

Cell proliferation analysis

Cell growth and viability were assessed in manually counted cells by either trypan blue dye exclusion or the MTS assay, as described previously (15).

Cell cycle analysis

Karpas 299, H2052, H2452 and H526 cells were seeded in 100 mm dishes and the next day treated with 0.05 μ g to 60 μ g /mL of brentuximab vedotin for 48 h. All attached and floating cells were washed with PBS and fixed in 0.125% formaldehyde followed by methanol. After incubation at -20° C, cells were washed with PBS and incubated for 45 min at 4°C in 50 μ g/ml propidium iodide, 0.1% Nonidet P-40, 20 μ g/ml RNase A and 0.1% sodium azide in

PBS. Propidium iodide fluorescence was measured on an EPICS XL-MCL cytometer (Beckman Coulter, Pasadena, CA).

Immunohistochemistry

The TMA slides were obtained from the National Mesothelioma Virtual Bank (NMVB) (http://mesotissue.org/). The slides were deparaffinized with xylene then transferred through graded ethanol to H2O. Endogenous peroxidase activity was blocked by a 10-min incubation in a 3% hydrogen peroxide solution. Antigen retrieval was performed by boiling the slides in a pressure cooker filled with a sodium citrate buffer (pH 6.0). After antigen retrieval, the slides were blocked using Background Sniper (Biocare, BS966M) for 20 min. The tissues were incubated with a mouse monoclonal anti-human CD30 antibody (Cell Marque/Clone Ber-H2) overnight at 4°C. Bound antibody was detected as described previously (16). The intensity of immunostaining for CD30 was scored visually by a thoracic pathologist and stratified into three staining groups based on staining intensity and percentage of tumor cells involved. Multiple cores for a single tumor sample were viewed to determine the score.

Statistical Analysis

All the described experiments were done more than three times, and the data are presented as mean values \pm SEM. *P* values were determined by *t* test using either Prism or SAS software (SAS Institute, Cary, NC). *P* < 0.05 was considered statistically significant.

Results

CD30 mRNA expression is high in mesothelioma tissues

We analyzed the gene expression patterns of CD 30 in all types of adult solid tumor cancer cell lines in the Cancer Cell Line Encyclopedia (CCLE) database (http:// www.broadinstitute.org/ccle/home) (17). When we compared cell lines derived from solid tumors we found the highest CD30 expression levels in mesothelioma, excluding lymphomas (Figure 1A). Based on these observations, we searched the Oncomine database (https://www.oncomine.org/resource/login.html) for expression in tumor tissues. A study by Gordon et al (18) performed expression profiling in thoracic cancers. When we specifically looked at CD30 expression we found that mesothelioma showed significantly higher expression of CD30 (p<0.0001) compared to adenocarcinoma (Figure 1B). A study by Lopez-Rios et al (19) performed expression profiling in mesothelioma subtypes. When we specifically looked at CD30 expression in this dataset we observed significantly higher CD30 expression in epithelioid compared to biphasic (p<0.006) and sarcomatoid (p<0.001) subtypes of mesothelioma tumors. Biphasic expression was not statistically different from sarcomatoid (p<0.506) (Figure 1C).

CD30 protein is expressed in mesothelioma tumors

We obtained tissue microarrays from the National Mesothelioma Virtual Bank (NMVB) to determine the level of CD30 staining in mesothelioma tumors by immuno-histochemistry. Descriptive statistics for the entire cohort of analyzed samples, as well as for CD30+ samples, is given in Table 1. The cohort demonstrated a typical distribution of histologic subtypes. Overall, CD30+ expression was noted in 13 out of 83 total mesothelioma

specimens and 12 of the CD30+ tumors demonstrated epithelial histology. The incidence of CD30 positivity was 4.6% in the biphasic and sarcomatoid group vs. 19.7% for the epithelioid group, approaching significance (p < 0.094). Membrane-associated staining was evident in the highest scoring tumors, similar to that observed in the Hodgkin's lymphoma positive control. Representative results of CD30 positivity are shown in Figure 2 at three different magnifications. Diffuse cytoplasmic staining of variable intensity was observed in lower scoring mesothelioma tumors. The percentage of tumor cells stained positive varied greatly, from nearly the whole tumor core down to only 5-10% of the core, as shown in the low power images in Figure 2, and generally decreased with scoring grade. One epithelioid core demonstrated 3+ and three demonstrated 2+ staining and the rest exhibited 1+ staining. The remaining CD30+ tumor was a high grade, biphasic metastatic tumor that demonstrated 1+ staining. We also looked for CD30 expression in three mesothelioma specimens obtained from our own hospital and found only one positive sample, although it demonstrated strong (3+) CD30 expression (data not shown). The mesothelioma cohort contained survival data on 63 patients, however univariate analyses using both Kaplan-Meier (p<0.935) and continuous measurement Cox regression models (p<0.82) predicted no statistical significance of CD30 positivity on overall survival.

CD30 protein is expressed on the cell surface of mesothelioma cells

The immunohistochemistry of CD30 in tumors clearly demonstrated a pattern of membraneassociated staining in some samples, which may represent an important feature in predicting sensitivity to brentuximab vedotin. We therefore sought to validate this finding in mesothelioma cell lines. We obtained all the available mesothelioma cell lines from the ATCC and determined the subcellular localization of CD30 by confocal microscopy. Data from the CCLE indicated that all four mesothelioma cells expressed CD30 mRNA. Figure 3A shows that CD30 staining outlines the shape of two mesothelioma cell lines, H2052 and H2452, consistent with CD30 residing on the cell membrane. Further examination of other focal planes confirmed this finding and detected little or no staining of cytoplasm. Similar results were obtained for 211H and H28 cells (data not shown). These results indicated that in mesothelioma cell lines that CD30 resides on, or near, the cellular membrane.

In order to determine if CD30 is expressed on the cell surface, H2052 and H2452 cells were fixed with ethanol, prior to being stained with an anti-CD30 antibody, and then analyzed by flow cytometry (Figure 3B). Both cell lines were found to be positive for CD30 when intact (non-permeabilized). However, when cells were permeabilized with Triton X-100 prior to staining to reveal intracellular CD30, variable amounts of additional CD30 were detected. Taken together, these results indicate that a significant amount of cellular CD30 antigen is located on the cell-surface of mesothelioma cells.

Brentuximab vedotin decreases cell growth and viability in mesothelioma cells

We hypothesize that the expression of CD30 on the cell surface should sensitize mesothelioma cells to brentuximab vedotin. We therefore investigated the effect of this antibody-drug conjugate on cell growth in mesothelioma cell lines. We selected Karpas 299, a CD30+ lymphoma cell line, as a positive control and H526, a small cell lung cancer cell line, as a negative control for these experiments. Cell growth determination by viable cell

count demonstrated that Karpas cells were sensitive to brentuximab vedotin in a dosedependent manner after 48 h of treatment, as expected, whereas H526 cells were largely resistant, indicating that most of the toxic drug remained conjugated to the CD30 antibody (Figure 4A). The two mesothelioma cell lines tested, H2052 and H2452, were both sensitive to the antibody-drug conjugate, with H2052 cells demonstrating more sensitivity than H2452 cells, although both were less sensitive than Karpas cells. We further investigated the effect of brentuximab vedotin on mesothelioma cell viability using the MTS assay. Treatment of H2052 and H2452 cells for 48 h induced a dose-dependent decrease in cell viability compared to untreated control cells (Figure 4B). As expected, the Karpas cells demonstrated greater sensitivity than mesothelioma cells whereas the H526 cells were

To confirm that brentuximab vedotin induced cell death in mesothelioma cells, we performed cell cycle analysis by flow cytometry. As shown in Figure 5, brentuximab vedotin greatly increased the proportion of Karpas cells in the G0 (sub G1) phase at the lowest dose after 48 h of incubation. This effect was not observed with H526 cells at any dose. Importantly, the two mesothelioma cell lines tested, H2052 and H2452, also demonstrated a dose-dependent increase in the proportion of cells in the sub G1 phase in response to brentuximab vedotin treatment, with H2052 cells again being more sensitive to drug than H2452 cells. The increased proportion of mesothelioma cells in sub G1 phase was accompanied by a parallel increase in cells in the G2/M phase. This is consistent with the anti-microtubule activity of the conjugated drug in brentuximab vedotin. MMAE, to produce mitotic arrest and G2/M blockade. Taken together, these results demonstrate that brentuximab vedotin treatment has an inhibitory effect on cell growth and survival in mesothelioma cells.

Discussion

resistant.

The goal of this study was to determine the therapeutic potential of targeting CD30 with brentuximab vedotin in mesothelioma. We looked for CD30 expression in the largest cohort of mesothelioma tumors studied to date and found that only 15.6% (13 of 83 tumors) were positive for CD30 expression using an antisera routinely used in the diagnosis of Hodgkin's lymphoma. There was a trend toward CD30 expression in mesothelioma tumors with epithelioid histology and high grade. Previous studies of CD30 expression in mesothelioma had noted expression in a single index case (4), two of five tumors (20) and two of eight tumors and 16 of 28 pleural and peritoneal effusions (5)

Although CD30 expression was not prognostic for overall survival in our study, of greater importance was whether or not CD30 expression could predict sensitivity to the antibodydrug conjugate targeting CD30, brentuximab vedotin. We used mesothelioma cell lines as a model system to test this hypothesis. Our results demonstrated that CD30 was expressed on the cell surface of mesothelioma cells using two complementary methods, confocal microscopy and FACS analysis. We then demonstrated that brentuximab vedotin decreased cell growth and viability in CD30 positive mesothelioma cells by three different methods. It is important to highlight three crucial points in these experiments. First, the doses of brentuximab vedotin used in our study, 0.5, 15 and 60 µg/ml, were within the range of drug

clinically achievable. That is, the typical C_{max} and AUC for brentuximab vedotin after a single 1.8 mg/kg dose was approximately 32 µg/ml and 80 µg /ml/day, respectively, in a phase I study (Acedris® insert). Second, the loss of mesothelioma cell viability after brentuximab vedotin treatment was associated with an increase in the proportion of cells in G2/M phase, as expected since the antibody conjugated toxin, MMAE, is a microtubule inhibitor. Third, there was no effect of brentuximab vedotin on a small cell lung cancer cell line, H526, suggesting therapeutic specificity.

It is unclear, however, to what extent our demonstration of brentuximab vedotin sensitivity in vitro using cultured mesothelioma cells is translatable to clinical drug sensitivity in mesothelioma patients. In this regard a recent interim report from a phase II study by Jalal et al on brentuximab vedotin in patients with malignant mesothelioma reported that 34% were positive for CD30 expression and that five positive responses were obtained to drug treatment out of 21 evaluable CD30+ patients (21). Although we obtained a lower percentage of CD30+ tumors (15.6%), likely attributable to scoring methods, the interim results of this phase II study confirm our hypothesis that a subgroup of malignant mesothelioma patients will respond to brentuximab.

It is clear from our study that mesothelioma tumor specimens demonstrated a range of CD30 expression, both in terms of their staining intensity and percentage of tumor cells involved. However, there was clearly a subset of high-scoring tumors that demonstrated obvious membrane-associated staining. It is intriguing to think that these tumors may demonstrate the greatest sensitivity to brentuximab vedotin. It has been suggested, however, that the MMAE toxin is freely diffusible into the surrounding microenvironment once it is internalized and cleaved from CD30 antibody in target CD30+ lymphoma cells, as evidenced by the sometimes-low occupancy (3%) of CD30-binding sites in lymphoma cells from highly responsive patients treated with brentuximab vedotin (22). Thus, the efficacy of brentuximab vedotin in mesothelioma tumors may not simply correlate with CD30 positivity. The answer to these questions awaits more detailed analysis of CD30 expression in clinical trials of brentuximab vedotin in malignant mesothelioma.

Acknowledgement

We are thankful to the Mesothelioma Research Bank [CDC NIOSH 1-U19-OH009077-01 National Mesothelioma Virtual Bank for Translational Research] for providing us with the tissue microarrays of mesothelioma. We are thankful to Dr. H. Koon at the Case Comprehensive Cancer Center for providing the Karpas cell line. We are also very thankful to Karen McCall for her support in the project.

Financial Support:

Supported by funds from the University Hospitals Seidman Cancer Center (A. Dowlati) as well as the Tissue Resources Core Facility and the Biostatistics & Bioinformatics Core Facility of the Case Comprehensive Cancer Center (National Cancer Institute P30 CA043703).

References

 Robinson BW, Lake RA. Advances in malignant mesothelioma. N Engl J Med. 2005; 353:1591– 603. [PubMed: 16221782]

- Hassan R, Cohen SJ, Phillips M, Pastan I, Sharon E, Kelly RJ, et al. Phase I clinical trial of the chimeric anti-mesothelin monoclonal antibody MORAb-009 in patients with mesothelin-expressing cancers. Clin Cancer Res. 16:6132–8. [PubMed: 21037025]
- Weldon JE, Xiang L, Zhang J, Beers R, Walker DA, Onda M, et al. A recombinant immunotoxin against the tumor-associated antigen mesothelin reengineered for high activity, low off-target toxicity, and reduced antigenicity. Mol Cancer Ther. 12:48–57. [PubMed: 23136186]
- 4. Dunphy CH, Gardner LJ, Bee CS. Malignant mesothelioma with CD30-positivity. A case report and review of the literature. Arch Pathol Lab Med. 2000; 124:1077–9. [PubMed: 10888787]
- Durkop H, Foss HD, Eitelbach F, Anagnostopoulos I, Latza U, Pileri S, et al. Expression of the CD30 antigen in non-lymphoid tissues and cells. J Pathol. 2000; 190:613–8. [PubMed: 10727988]
- Kadin ME. Regulation of CD30 antigen expression and its potential significance for human disease. Am J Pathol. 2000; 156:1479–84. [PubMed: 10793058]
- Vaklavas C, Forero-Torres A. Safety and efficacy of brentuximab vedotin in patients with Hodgkin lymphoma or systemic anaplastic large cell lymphoma. Ther Adv Hematol. 3:209–25. [PubMed: 23606932]
- Hu S, Xu-Monette ZY, Balasubramanyam A, Manyam GC, Visco C, Tzankov A, et al. CD30 expression defines a novel subgroup of diffuse large B-cell lymphoma with favorable prognosis and distinct gene expression signature: a report from the International DLBCL Rituximab-CHOP Consortium Program Study. Blood. 121:2715–24. [PubMed: 23343832]
- Hamblett KJ, Senter PD, Chace DF, Sun MM, Lenox J, Cerveny CG, et al. Effects of drug loading on the antitumor activity of a monoclonal antibody drug conjugate. Clin Cancer Res. 2004; 10:7063–70. [PubMed: 15501986]
- Doronina SO, Toki BE, Torgov MY, Mendelsohn BA, Cerveny CG, Chace DF, et al. Development of potent monoclonal antibody auristatin conjugates for cancer therapy. Nat Biotechnol. 2003; 21:778–84. [PubMed: 12778055]
- Deng C, Pan B, O'Connor OA. Brentuximab vedotin. Clin Cancer Res. 19:22–7. [PubMed: 23155186]
- Garcia-Prats MD, Ballestin C, Sotelo T, Lopez-Encuentra A, Mayordomo JI. A comparative evaluation of immunohistochemical markers for the differential diagnosis of malignant pleural tumours. Histopathology. 1998; 32:462–72. [PubMed: 9639123]
- Francisco JA, Cerveny CG, Meyer DL, Mixan BJ, Klussman K, Chace DF, et al. cAC10vcMMAE, an anti-CD30-monomethyl auristatin E conjugate with potent and selective antitumor activity. Blood. 2003; 102:1458–65. [PubMed: 12714494]
- Kim KM, McDonagh CF, Westendorf L, Brown LL, Sussman D, Feist T, et al. Anti-CD30 diabody-drug conjugates with potent antitumor activity. Mol Cancer Ther. 2008; 7:2486–97. [PubMed: 18723494]
- Dabir S, Kluge A, Dowlati A. The association and nuclear translocation of the PIAS3-STAT3 complex is ligand and time dependent. Molecular cancer research : MCR. 2009; 7:1854–60. [PubMed: 19903771]
- Dabir S, Kluge A, Kresak A, Yang M, Fu P, Groner B, et al. Low PIAS3 Expression in Malignant Mesothelioma Is Associated with Increased STAT3 Activation and Poor Patient Survival. Clin Cancer Res. 2014; 20:5124–32. [PubMed: 25124686]
- Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature. 2012; 483:603–7. [PubMed: 22460905]
- Gordon GJ, Jensen RV, Hsiao LL, Gullans SR, Blumenstock JE, Ramaswamy S, et al. Translation of microarray data into clinically relevant cancer diagnostic tests using gene expression ratios in lung cancer and mesothelioma. Cancer research. 2002; 62:4963–7. [PubMed: 12208747]
- Lopez-Rios F, Chuai S, Flores R, Shimizu S, Ohno T, Wakahara K, et al. Global gene expression profiling of pleural mesotheliomas: overexpression of aurora kinases and P16/CDKN2A deletion as prognostic factors and critical evaluation of microarray-based prognostic prediction. Cancer research. 2006; 66:2970–9. [PubMed: 16540645]
- 20. Sharman JP, Goldschmidt JH, Burke JM, Hellerstedt BA, McIntyre K, Yasenchak CA, et al. CD30 expression in nonlymphomatous malignancies. J Clin Oncol. 2012; 30(suppl) abstr 3069.

- 21. Jalal SI, Einhorn L, Shapiro GI, Hilton J, Wheler J, Dowlati A, Josephson N, Sharman J, Burke JM. Brentuximab Vedotin in Patients with CD30+ Mesothelioma. Mesothelioma, Thymic Malignancies, and Other Thoracic Malignancies. Int J Rad Oncol. 2014; 90(5):S8.
- 22. Fromm JR, McEarchern JA, Kennedy D, Thomas A, Shustov AR, Gopal AK. Clinical binding properties, internalization kinetics, and clinicopathologic activity of brentuximab vedotin: an antibody-drug conjugate for CD30-positive lymphoid neoplasms. Clin Lymphoma Myeloma Leuk. 12:280–3. [PubMed: 22542449]



Figure 1. Mesothelioma shows high CD30 expression in public datasets

(A) Gene expression pattern for CD30 across solid tumor cancer cell lines in the Cancer Cell Line Encyclopedia (CCLE) data base (B) Whisker plots showing the Log2 median centered intensity of CD30 expression in Gordon lung dataset (18). Compared to adenocarcinoma, CD30 expression in mesothelioma tissue samples is significantly higher. (C) Whisker plots of the Log2 median centered intensity of CD30 expression in Lopez-Rios dataset (19) shows higher CD30 expression in epithelioid compared to biphasic and sarcomatoid.



Figure 2. Mesothelioma tumor specimens demonstrate high CD30 expression

CD30 expression was examined by immuno-histochemistry (IHC) with an anti-BerH2 antibody used for the clinical diagnosis of Hodgkin disease. The positive control is Hodgkin lymphoma. Rectangles show area magnified in image below. The 3+ core was scored higher than the 2+ core due to a greater area of positivity, not because of intensity.





(A) Confocal analysis of mesothelioma cell lines. The DAPI stain identifies the nucleus. (B) Mesothelioma CD30 expression by FACS analysis. In these experiments, control is staining in the absence of primary antibody.



Figure 4. Effect of brentuximab vedotin treatment on cell growth and viability

(A) Mesothelioma cell lines H2052, H2452, 211H and positive control Karpas 299 treated with brentuximab vedotin and cells were counted on day 1 and 2. (B) MTS assay of cell proliferation.



Figure 5. Effect of brentuximab vedotin treatment on cell cycle

Results demonstrate a dose response of mesothelioma cells to brentuximab vedotin treatment after 48 h. The percentage of cells in the G0, G1, S and M phase is shown in the figure.

Table 1

CD30 positivity in mesothelioma cohort

	Whole cohort	CD30+	P-value
Histologic type			
Biphasic, Sarcomatoid	22	1	0.094
Epithelioid	61	12	
Stage ^a			
1, 2 and 3	52	6	0.465
4	10	2	
Tumor grade ^{<i>a</i>} (epitheliod only)			
High	24	7	0.163
Intermediate and low	35		

 $a_{\rm incomplete \ staging \ and \ grading \ data \ available}$