



Published in final edited form as:

Gastroenterology. 2011 May ; 140(6): 1776–1784.e5. doi:10.1053/j.gastro.2011.02.015.

Blocking Lymphocyte Localization to the Gastrointestinal Mucosa as a Therapeutic Strategy for IBD

Eduardo J. Villablanca^{1,3}, Barbara Cassani^{1,3}, Ulrich H. von Andrian^{2,*}, and J. Rodrigo Mora^{1,*}

¹ Gastrointestinal Unit, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114

² Immune Disease Institute, Harvard Medical School, Boston, MA 02115

Abstract

Lymphocyte migration (homing) to specific tissues has an important role during protective and pathological immune responses, including inflammatory bowel diseases (IBDs). Lymphocytes use integrin $\alpha 4\beta 7$ and the chemokine receptor CCR9 to localize to the gastrointestinal (GI) mucosa; their respective ligands, MAdCAM-1 and CCL25, are displayed on endothelial cells in intestinal post-capillary venules. Whereas GI-homing receptors are required for lymphocyte migration to the intestine in the non-inflamed steady state, their role during inflammation is a matter of debate. Reagents designed to block interactions between these receptors and their ligands have had variable degrees of success in animal models of IBD and patients. We discuss the mechanisms involved in lymphocyte localization to the intestinal mucosa and how they can be applied to therapy for IBD.

Introduction

Lymphocytes localize to specific tissues during the protective immune response and in inflammatory disorders. Learning how these cells localize to different organs is important for understanding basic immunology as well as disease pathogenesis.

Circulating lymphocytes are exposed to extreme shear forces so they do not randomly adhere to endothelial cells; instead, they express adhesion receptors for ligands expressed on endothelial cells. Adhesion usually takes place in post-capillary venules via a multistep process. First, lymphocytes are captured and loosely adhere to the endothelial cells (tethering and rolling, respectively), a step that usually requires selectins and their ligands, although the integrins $\alpha 4\beta 7$ and $\alpha 4\beta 1$ can also contribute to this step in some tissues. While lymphocytes are rolling they can be stimulated, generally via chemokine receptors (activation), which increases integrins' binding affinity and avidity. Integrin activation

© 2011 The American Gastroenterological Association. Published by Elsevier Inc. All rights reserved.

Corresponding authors: Ulrich H. von Andrian (uva@hms.harvard.edu) or J. Rodrigo Mora (j_rodrigo_mora@hms.harvard.edu).

³Authors with equal contribution

Author's contributions: EJV, BC, UHvA and JRM contributed equally in writing this review article.

Disclosures: The authors declare that no conflict of interest exists and have no other disclosures to make.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

causes the lymphocytes to adhere to the endothelium (sticking) and then extravasation into non-inflamed or inflamed tissues.

Lymphocyte migration and adhesion to specific tissues are determined by the combination of receptors involved in each step, rather than a single receptor and adhesive molecule. The diversity of receptors use in each step of the adhesion process allows for versatile and tissue-specific localization of lymphocytes, making lymphocyte adhesion amenable to modulation for therapeutic purposes.

The mechanisms that regulate lymphocyte homing to different tissues have been reviewed;²⁻⁴ we focus on lymphocyte migration to the gastrointestinal (GI) mucosa and discuss how this process might be modulated in patients, to reduce GI inflammation.

Compartmentalized homing to the intestine

Naïve T and B cells constantly transit between the blood and secondary lymphoid organs (SLO), such as spleen, lymph nodes and Peyer's patches (PP). Upon activation in SLO, naïve lymphocytes become effector and/or memory T and B cells and express receptors that control their migration to extra-lymphoid tissues such as the skin, GI lamina propria, central nervous system (CNS), liver, and lungs⁵.

Whereas migration to SLO occurs through the mechanism described above, lymphocyte migration to some extra-lymphoid tissues requires expression of specific receptors. T-cell localization the GI mucosa and the skin—the largest surfaces in the body that are exposed to the external environment—has been well characterized. T-cell migration to the skin requires ligands for P- and E-selectins, CCR4, and the integrin lymphocyte function antigen (LFA)-1⁶.

In contrast to the skin, migration of T and B cells to the small intestine requires the integrin $\alpha 4\beta 7$ and CCR9, whose induction depends on the vitamin A metabolite *all-trans* retinoic acid (RA)³ (Figure 1). Localization to colon partially requires $\alpha 4\beta 7$, but not CCR9;⁷ the chemokine receptor(s) required for leukocyte migration to the colon have not been identified.

The ligand for CCR9, CCL25/TECK, is differentially distributed in a proximal-to-distal gradient in the small bowel; CD8+ T cells localize to the ileum partially via CCR9-independent mechanisms (Figure 1)⁷. Alternative candidates for T-cell migration to the small bowel include CXCR3 and CXCR4, whose ligands (CXCL10 and CXCL12, respectively), are expressed in the GI mucosa⁸. Consistent with an *in vivo* role for these alternative chemokine pathways, *CXCR3*^{-/-} mice have lower numbers of CD8+ intestinal epithelial cells in the lamina propria;⁹ blocking the interaction between CXCR4 and CXCL12 inhibits entry of T cells to the small intestine in steady-state and inflammatory conditions¹⁰.

Localization of lymphocytes to the colon differs in some ways from migration to the small bowel—it requires either $\alpha 4\beta 7$ or $\alpha 4\beta 1$, but not CCR9^{6, 11}. The ligand for $\alpha 4\beta 7$, MAdCAM-1, is expressed in small bowel and colon, whereas CCL25 is expressed in only the small bowel^{12, 13}. Moreover, whereas migration to the small intestine was impaired in *CCR9*^{-/-} and $\beta 7$ integrin chain-deficient (*$\beta 7$* ^{-/-}) T-helper (Th)17, homing to the colon was reduced in only the *$\beta 7$* ^{-/-} Th17 cells¹⁴. Transfer of *$\beta 7$* ^{-/-} Th17 cells into SCID mice induced less inflammation in the small and large bowel than transfer of wild-type Th17 cells (Table 1), whereas transfer of *CCR9*^{-/-} Th17 induced less inflammation than wild-type cells in only the small bowel¹⁴. Together, these data indicate that CCR9 is required for T-cell migration

and pathogenicity primarily in the small intestine, whereas $\alpha 4\beta 7$ is required for T-cell migration and pathogenicity in the small bowel and colon.

Migration of T cells to the intestinal mucosa also depends on their specific subset and phenotype. Recently activated CD8⁺ T cells require CCR9 for migration to the small bowel, whereas effector CD4⁺ T cells are less dependent on CCR9 for homing into this GI compartment¹⁵. Moreover, homing of *CCR6*^{-/-} Th17 cells to PP and small bowel was significantly reduced, compared to wild-type Th17 cells, whereas homing of Th1 or Foxp3⁺ regulatory T cells (T_{REG}) to these compartments did not require CCR6¹⁶. RA induces $\alpha 4\beta 7$ and CCR9 on T_{REG}^{17,18}; however, mice given diets that did not contain vitamin A (and therefore lack RA synthesis) did not have decreased numbers of T_{REG} in the small bowel, although Th17 cells were markedly reduced in this compartment¹⁹. Moreover, T_{REG} isolated from mice depleted of dietary vitamin A were equally efficient in suppressing ileitis as T_{REG} from mice on a vitamin A sufficient diet (or that received extra vitamin A)²⁰. However, these studies did not discriminate between thymus-derived or inducible T_{REG}. Further studies are needed to determine the in vivo roles of RA in localization of T_{REG} in the GI mucosa and their immunoregulatory functions there.

Another example of differential gut-homing requirements is cells that secrete immunoglobulin (Ig)A (IgA-secreting cells). CCR10 is expressed primarily by IgA-secreting cells, whereas cells that secrete antibodies against IgG or IgM do not express this receptor²¹. Moreover, IgA-secreting cells require CCR10 to localize to the intestine, although CCR10 is not expressed on gut-associated T cells^{4,21}. In addition to CCR10, IgA-secreting cells also express CCR9; mice deficient in this receptor have reduced numbers of these cells in the small intestine²². Interestingly, blocking CCR9 prevented localization of IgA-secreting cells to the small bowel, whereas blocking either CCR10 or its mucosal ligand, CCL28, impaired their localization to small and large bowel²³. Similar to T cells, RA was sufficient to induce CCR9 and $\alpha 4\beta 7$ (but not CCR10) on activated B cells and mice depleted of vitamin A had very low numbers of small bowel IgA-secreting cells.^{19, 24, 25}

Aberrant recruitment of lymphocyte in IBDs

IBDs, which include Crohn's disease (CD) and ulcerative colitis (UC), are associated with a massive influx of immune cells into the GI tract. During disease development, altered production of pro-inflammatory cytokines induces expression of alternative adhesion receptors and chemokines on intestinal endothelial cells, which might allow lymphocytes to migrate to the intestine without expression of the receptors that normally regulate their localization in that compartment¹¹. These alternative pathways of lymphocyte recruitment might have important implications for IBD therapy.

Studies of animal models and human tissues have indicated the role for gut-homing effector T cells in IBD pathogenesis. In experimental models of IBD, MAdCAM-1 is upregulated in the intestinal lamina propria.^{26-28,29} Similar MAdCAM-1 upregulation is observed in active inflamed tissues from patients with CD or UC, which is associated with increased numbers of $\alpha 4\beta 7$ ⁺ T cells, compared with normal tissues^{30,31}. Deficiency of $\beta 7$ integrin subunit inhibits inflammation in a mouse model of CD³² and antibodies against $\beta 7$ or MAdCAM-1 reduced inflammation in mice with TNBS-induced or cell transfer-induced colitis^{33,34} (Table 1). Blocking $\alpha 4$ or $\alpha 4\beta 7$ reduced colitis in a non-human primate model of IBD^{35,36}. Additional mechanistic insights have come from studies of SAMP1/YitFc mice, which spontaneously develop CD-like ileitis. Although SAMP1/YitFc mice deficient in $\beta 7$ have reduced intestinal inflammation³⁷, antibodies that block $\alpha 4\beta 7$ or MAdCAM-1 did not reduce the inflammation; only combined blockade of VCAM-1 and MAdCAM-1 significantly improved ileitis²⁹.

Blocking $\alpha 1\beta 1$, a collagen-binding integrin that is upregulated in inflamed tissues, reduced TNBS-induced colitis in mice³⁸. Results from mice with dextran sodium sulphate (DSS)-induced colitis have varied results—some studies reported that blockers of MAdCAM-1 reduced inflammation,^{33, 39, 40} whereas others showed that development of colitis required $\alpha 4\beta 1$ -VCAM-1 and $\beta 2$ -ICAM-1 interactions, but not $\alpha 4\beta 7$ -MAdCAM-1^{41, 42}. Pathogenic effector T cells might not require only interaction between lymphocyte $\alpha 4\beta 7$ and epithelial cell MAdCAM-1 to promote chronic inflammation, but other integrins that mediate immune cell localization during general inflammation might fulfill redundant roles in intestinal pathology.

SAMP1/YitFc mice have increased numbers of IgA-secreting cells in the MLN and LP⁴³. Adoptive transfer of B cells and T cells from SAMP1/YitFc into SCID mice increased ileitis, compared with transfer of only T cells⁴³. Moreover, B cells required $\alpha 4\beta 7$ to exacerbate ileitis³⁷, indicating that B-cell localization to the GI tract might also be involved in IBD pathogenesis.

T_{REG} cells are believed to prevent or even cure intestinal inflammation, based on studies from different models of IBD. However, the precise role of $\alpha 4\beta 7$ and CCR9 in trafficking and function of T_{REG} cells during GI inflammation is unclear. T_{REG} cells seem to require $\beta 7$ -independent pathways—rather, those that involve CCR7 and CCR4—for their immune suppressive functions and to prevent colitis⁴⁴⁻⁴⁶. These alternative migratory pathways might allow T_{REG} cells function in lymphoid compartments, such as prophylactic suppression before the onset of inflammation in MLN or PP. However, T_{REG} cells might need GI homing receptors to suppress immune activity in the lamina propria during active inflammation⁴⁷, which is probably most relevant for development of therapeutics.

CCL5 and CCR5 are also up-regulated in experimental models of ileitis and mediate the specific recruitment of T_{REG} cells and some subsets of effector CD4+ T cells⁴⁸. These alternative chemokine pathways could account for the observation that blocking CCL25 or CCR9 is only effective at early stages of disease, even though expression of CCL25 increases in the small bowel of patients with CD.⁴⁹ Other, perhaps non GI-specific, chemokine signals might mediate lymphocyte homing at later stages during inflammation⁵⁰. Human lamina propria and intraepithelial lymphocytes also express CXCR3, CX3CR1, and CCR2, and levels of their ligands are increased in tissues of patients CD⁵¹. Moreover, recruitment of T cells, monocytes, and DC to the inflamed mucosa might also involve CX3CL1 and its receptor CX3CR1, which contributes to pathogenesis of IBD⁵²⁻⁵⁴.

Some of the extra-intestinal pathologies associated with IBD might arise from aberrant homing of immune cells. For instance, MAdCAM-1 and CCL25 are upregulated in the liver during primary sclerosing colangitis (PSC), a chronic disease characterized by progressive inflammation and scarring of the bile ducts. PSC has been associated with UC in epidemiologic studies⁵⁵.

Therapies for IBD

Patients with IBD usually require life-long therapy with corticosteroids and other immunosuppressive drugs. Choice of therapy depends on the primary clinical goal (induction or maintenance of remission), the extent and severity of disease, the response to current or prior treatments, and the occurrence of side effects (summarized in Supplementary Table 1A). Many drugs for IBD can have serious adverse effects, and some patients become refractory to treatment during disease progression and require surgery. Therefore, new therapeutic approaches, that target specific inflammatory mediators, are needed.

Although the primary causes of IBD are not clear, many molecules that are involved in disease pathogenesis have been identified as targets for therapy. Therapeutics that have been developed include inhibitors of T-cell activation, co-stimulatory pathways, pro-inflammatory cytokine receptors, Th1 polarization, cytokines and their regulatory proteins, growth hormone, and growth factors (Supplementary Table 1B, C).

Although many of these biologic agents showed efficacy in preclinical studies, most of them have not shown efficacy in clinical trials, or have caused significant side effects⁵⁶. Antibodies to the cytokine TNF- α have been the most effective, and are currently used to treat patients with refractory moderate-to-severe active CD or UC. Infliximab binds the soluble bioactive and membrane-bound forms of human TNF- α and reduces its toxicity. Although Infliximab is effective in reducing the symptoms of IBDs, its immunosuppressive effects predispose patients to infections and increase risk of malignancies, such as lymphomas^{57, 58}.

Blocking adhesion receptors

Molecules that mediate lymphocyte localization to the GI mucosa, in the steady state or during development of inflammatory diseases such as IBDs, are attractive targets for drug development. Antibodies or compounds that selectively block homing receptors,¹¹ or reagents that sequester lymphocytes in secondary lymphoid organs (to prevent their migration to sites of inflammation),⁵⁹ have shown efficacy in animal models and in clinical trials for psoriasis^{60, 61}, asthma⁶², graft-versus-host disease,⁶³ and multiple sclerosis⁵⁹.

Because $\alpha 4\beta 7$ and CCR9 are the primary mediators of lymphocyte migration to the intestine, reagents that block their function should reduce inflammation in the intestinal mucosa, yet cause low levels of systemic immunosuppression. Agents developed for treatment of IBD disrupt interactions between LFA-1 and ICAM-1, $\alpha 4\beta 1$ and VCAM-1, as well as $\alpha 4\beta 7$ and MAdCAM-1 (Figure 2, Table 2).

The first successful clinical use of anti-ICAM-1 was in treatment of rheumatoid arthritis⁶⁴. Antibodies against ICAM-1 or antisense oligonucleotides that disrupt expression of ICAM-1 showed efficacy in mouse models of IBD, including DSS-induced colitis and SAMP-1/Yit mice ileitis^{65 66}. Interestingly, in SAMP-1/Yit mice, anti-ICAM-1 was only effective when administered in combination with anti-VCAM-1 or anti- $\alpha 4$ integrins, indicating redundancy between LFA-1/ICAM-1 and $\alpha 4\beta 1$ /VCAM-1 pathways during inflammation in mice. Clinical trials that investigated the effects of reagents against ICAM-1 in patients with IBD included investigation of Alicaforsen (ISIS 2302), an antisense oligonucleotide that prevents expression of ICAM-1. Whereas an early-stage clinical trial suggested a therapeutic potential for Alicaforsen in patients with mild-to-moderate active CD⁶⁷, two subsequent, larger, multicenter trials failed to demonstrate significant efficacy^{68 69}. Despite this setback, patients treated with Alicaforsen, in an enema formulation, had significant improvements in distal UC in a randomized, placebo-controlled trial.⁷⁰ However, given the important role of LFA-1 interaction with ICAM-1 in leukocyte localization to many lymphoid and non-lymphoid tissues, as well as in T-cell activation,¹¹ it is likely that blockers of ICAM-1 will cause significant systemic immunosuppression.

Natalizumab is a recombinant, humanized, monoclonal IgG4 against $\alpha 4$; it inhibits MAdCAM-1 signaling through integrin $\alpha 4\beta 7$ and VCAM-1 signaling through integrin $\alpha 4\beta 1$ ⁷¹. In placebo-controlled, randomized trials, 40% patients with moderate-to-severe CD responded to Natalizumab and went to remission, compared to 8% in the group that received placebo⁷².

However, phase III clinical trials that included clinical response, remission, and maintenance as endpoints showed that the drug was more effective when given in combination with other immunosuppressants or before therapy with an anti-TNF- α reagent^{73,74}. Since blockers of $\alpha 4$ probably do not affect T cells that have already localized to intestinal tissues, Natalizumab might not be sufficient to reduce ongoing inflammation—its combination with other immunosuppressant drugs might be required. Natalizumab has also shown potential for treatment of UC,⁷⁵ probably due to its ability to block $\alpha 4\beta 7$ and $\alpha 4\beta 1$ (integrins involved in localization of lymphocytes to the colon).

Natalizumab has also been used to treat patients with multiple sclerosis,⁷⁶ based on the role of $\alpha 4\beta 1$ interaction with VCAM-1 in leukocyte homing to the CNS and experimental allergic encephalomyelitis⁷⁷. However, cases of progressive multifocal leukoencephalopathy (PML)—a rare and often fatal opportunistic infection of the CNS—have developed in some patients given Natalizumab (approximate incidence 1/1000), raising concerns about its safety^{78,79}. Sporadic cases of melanoma have also been reported in patients treated with Natalizumab, which might be associated with impaired immunosurveillance of the skin following $\alpha 4\beta 1$ blockade^{80,81}. However, larger cohorts of patients that have received Natalizumab need to be studied to determine more precisely the incidence of these rare side effects. Similar safety concerns might also apply to an orally bioavailable inhibitor of $\alpha 4$ (AJM300), which has also shown to be effective in patients with active CD⁸². Some cases of PML also occurred in patients with psoriasis who were treated with Efalizumab, a monoclonal antibody against LFA-1⁸³. Reagents that selectively block homing of lymphocytes to the GI without affecting immunosurveillance in other tissues (including the CNS), are urgently required for IBD.

Vedolizumab (MLN-02) is a humanized monoclonal IgG1 against integrin $\alpha 4\beta 7$. In a phase II trial in 181 patients with UC, remission rates were significantly higher among subjects treated with Vedolizumab than those given placebo⁸⁴. Another placebo-controlled trial of 185 patients with mild-to-moderate active CD showed that Vedolizumab was significantly more effective than placebo at inducing remission in patients with CD⁸⁵.

A monoclonal antibody against MAdCAM-1 (Pf-0054,659, human IgG2) is being tested in a phase I/II clinical trial of patients with UC. Although the study includes a small number of patients, endoscopic examinations identified improvements among in patients treated with anti-MAdCAM-1, without major side effects⁸⁶.

Nevertheless, because $\alpha 4\beta 1$ might also have a role in chronic intestinal inflammation^{29,41,42}, it is possible that selective targeting of $\alpha 4\beta 7$ or MAdCAM-1 might not be as effective as reagents designed to block both $\alpha 4$ integrins.

Chemokines as therapeutic targets

The effects of blocking the interaction between CCL25 and CCR9 were demonstrated in SAMP-1/Yit mice with ileitis. Blocking either CCL25 or CCR9 during early stages of ileitis reduced inflammation, whereas no effect was observed when mice were given the reagents at later stages of disease progression⁵⁰. Moreover, administration of CCX282 (Traficet-EN), an orally bioavailable antagonist of CCR9, reduced the inflammatory response in mice when given before or after gut inflammation induced by TNF- α overexpression⁸⁷.

CCX282 is being tested in trials of patients with CD and refractory celiac disease (Figure 2, Table 2); preliminary efficacy and safety evaluations look promising⁸⁸. In a phase II trial, 74 patients with CD were given either CCX282 or placebo for 28 days. Fifty-eight percent of patients had a significant reduction in CD scores, compared with 31% in the placebo group; in the CCX282 group, the effect that was associated with reduced levels of pro-

inflammatory cytokines and C-reactive protein⁸⁸. A subsequent phase II/III trial showed significant improvement in the CCX282 group; disease scores were reduced in 81% of patients and 41% experienced clinical remission—effects that were maintained even upon withdrawal of corticosteroids⁸⁸.

Results from larger phase III trials of CCX282 for CD and UC are pending. Although the drug was generally well tolerated and not associated with an increased risk of infection, a large cohort of patients must be followed for long time period to exclude risk for rare diseases such as PML. The fact that CCX282 is orally bioavailable offers a clear advantage to parenteral therapies, decreasing the cost of the treatment, eliminating morbidity associated with parenteral administration, and potentially increasing compliance.

Interestingly, tolerogenic plasmacytoid DC also express CCR9,⁸⁹ which seems to be required for localization of these cells to the small bowel⁹⁰. Inhibitors of CCR9 might affect migration of plasmacytoid DC and their tolerogenic functions in the intestine, with potential effects that should be considered and explored in models of IBD pathogenesis.

Conclusion

Although the exact cellular and molecular mechanisms of IBD pathogenesis are undefined, lymphocyte homing has an important role. Improved understanding of lymphocyte localization to the non-inflamed and inflamed intestinal mucosa has led to specific and effective therapies for IBD and improved the benefit–risk profile for patients.

Nevertheless, results from studies of animal models of IBD have identified alternative homing receptors that, in addition to $\alpha 4\beta 7$ and/or CCR9, have roles in lymphocyte migration to the GI tract and might contribute to inflammation; these redundant homing pathways could account for the varying degrees of effectiveness of reagents that target GI-specific homing receptors in clinical trials. It is important to better define which receptors, adhesion molecules, and chemokine pathways contribute to chronic intestinal inflammation in humans. It is also important to determine the precise role of T_{REG} cells in intestinal inflammation and whether GI-specific reagents that interfere with lymphocyte adhesion affect T_{REG} functions. Combination therapies, which target more than one step in adhesion of lymphocytes to the intestinal epithelium, might be the most effective strategy for IBD. Combinations such as anti- $\alpha 4\beta 7$ and antagonists of CCR9 could have additive effects to reduce inflammation in patients with IBD.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding: We are grateful to Allison McNulty for editorial assistance. JRM is indebted to Ingrid Ramos for constant support. EJv was supported by grants from Crohn's & Colitis Foundation of America (CCFA). UHvA was supported by grants from NIH. JRM was supported by grants from CCFA, Cancer Research Institute (CRI), Howard H. Goodman (MGH), Massachusetts Life Science Center (MLSC) and NIH DP2 2009A054301.

References


1. von Andrian UH, Mackay CR. T-cell function and migration. Two sides of the same coin. *N Engl J Med*. 2000; 343:1020–1034. [PubMed: 11018170]
2. Mora JR, von Andrian UH. Differentiation and homing of IgA-secreting cells. *Mucosal Immunology*. 2008; 1:96–109. [PubMed: 19079167]

3. Mora JR, Iwata M, von Andrian UH. Vitamin effects on the immune system: vitamins A and D take centre stage. *Nat Rev Immunol*. 2008
4. Mora JR. Homing imprinting and immunomodulation in the gut: role of dendritic cells and retinoids. *Inflamm Bowel Dis*. 2008; 14:275–89. [PubMed: 17924560]
5. Masopust D, Vezyz V, Marzo AL, Lefrancois L. Preferential localization of effector memory cells in nonlymphoid tissue. *Science*. 2001; 291:2413–7. [PubMed: 11264538]
6. Mora JR, von Andrian UH. T-cell homing specificity and plasticity: new concepts and future challenges. *Trends Immunol*. 2006; 27:235–43. [PubMed: 16580261]
7. Stenstad H, Svensson M, Cucak H, Kotarsky K, Agace WW. Differential homing mechanisms regulate regionalized effector CD8 α T cell accumulation within the small intestine. *Proc Natl Acad Sci U S A*. 2007; 104:10122–7. [PubMed: 17551016]
8. Wright N, Hidalgo A, Rodriguez-Frade JM, Soriano SF, Mellado M, Parmo-Cabanas M, Briskin MJ, Teixido J. The chemokine stromal cell-derived factor-1 α modulates α 4 β 7 integrin-mediated lymphocyte adhesion to mucosal addressin cell adhesion molecule-1 and fibronectin. *J Immunol*. 2002; 168:5268–5277. [PubMed: 11994484]
9. Annunziato F, Cosmi L, Liotta F, Lazzeri E, Romagnani P, Angeli R, Lasagni L, Manetti R, Marra F, Gerard C, Petrai I, Dello Sbarba P, Tonelli F, Maggi E, Romagnani S. CXCR3 and α 4 β 7 integrin identify a subset of CD8 $^{+}$ mature thymocytes that share phenotypic and functional properties with CD8 $^{+}$ gut intraepithelial lymphocytes. *Gut*. 2006; 55:961–8. [PubMed: 16368781]
10. Oyama T, Miura S, Watanabe C, Hokari R, Fujiyama Y, Komoto S, Tsuzuki Y, Hosoe N, Nagata H, Hibi T. CXCL12 and CCL20 play a significant role in mucosal T-lymphocyte adherence to intestinal microvessels in mice. *Microcirculation*. 2007; 14:753–66. [PubMed: 17885999]
11. Rodrigo Mora J, Von Andrian UH. Specificity and plasticity of memory lymphocyte migration. *Curr Top Microbiol Immunol*. 2006; 308:83–116. [PubMed: 16922087]
12. Kunkel EJ, Campbell JJ, Haraldsen G, Pan J, Boisvert J, Roberts AI, Ebert EC, Vierra MA, Goodman SB, Genovese MC, Wardlaw AJ, Greenberg HB, Parker CM, Butcher EC, Andrew DP, Agace WW. Lymphocyte CC chemokine receptor 9 and epithelial thymus-expressed chemokine (TECK) expression distinguish the small intestinal immune compartment: Epithelial expression of tissue-specific chemokines as an organizing principle in regional immunity. *J Exp Med*. 2000; 192:761–8. [PubMed: 10974041]
13. Papadakis KA, Prehn J, Nelson V, Cheng L, Binder SW, Ponath PD, Andrew DP, Targan SR. The role of thymus-expressed chemokine and its receptor CCR9 on lymphocytes in the regional specialization of the mucosal immune system. *J Immunol*. 2000; 165:5069–5076. [PubMed: 11046037]
14. Wang C, Kang SG, HogenEsch H, Love PE, Kim CH. Retinoic acid determines the precise tissue tropism of inflammatory Th17 cells in the intestine. *J Immunol*. 2010; 184:5519–26. [PubMed: 20400707]
15. Stenstad H, Ericsson A, Johansson-Lindbom B, Svensson M, Marsal J, Mack M, Picarella D, Soler D, Marquez G, Briskin M, Agace WW. Gut associated lymphoid tissue primed CD4 $^{+}$ T cells display CCR9 dependent and independent homing to the small intestine. *Blood*. 2006
16. Wang C, Kang SG, Lee J, Sun Z, Kim CH. The roles of CCR6 in migration of Th17 cells and regulation of effector T-cell balance in the gut. *Mucosal Immunol*. 2009; 2:173–83. [PubMed: 19129757]
17. Kang SG, Lim HW, Andrisani OM, Broxmeyer HE, Kim CH. Vitamin A metabolites induce gut-homing FoxP3 $^{+}$ regulatory T cells. *J Immunol*. 2007; 179:3724–33. [PubMed: 17785809]
18. Benson MJ, Pino-Lagos K, Roseblatt M, Noelle RJ. All-trans retinoic acid mediates enhanced T reg cell growth, differentiation, and gut homing in the face of high levels of co-stimulation. *J Exp Med*. 2007; 199:1775–1774. [PubMed: 17620362]
19. Cha HR, Chang SY, Chang JH, Kim JO, Yang JY, Kim CH, Kweon MN. Downregulation of Th17 cells in the small intestine by disruption of gut flora in the absence of retinoic acid. *J Immunol*. 2010; 184:6799–806. [PubMed: 20488794]
20. Kang SG, Wang C, Matsumoto S, Kim CH. High and low vitamin A therapies induce distinct FoxP3 $^{+}$ T-cell subsets and effectively control intestinal inflammation. *Gastroenterology*. 2009; 137:1391–402 e1-6. [PubMed: 19632226]

21. Kunkel EJ, Butcher EC. Plasma-cell homing. *Nat Rev Immunol*. 2003; 3:822–9. [PubMed: 14523388]
22. Pabst O, Ohl L, Wendland M, Wurbel MA, Kremmer E, Malissen B, Forster R. Chemokine receptor CCR9 contributes to the localization of plasma cells to the small intestine. *J Exp Med*. 2004; 199:411–6. [PubMed: 14744993]
23. Hieshima K, Kawasaki Y, Hanamoto H, Nakayama T, Nagakubo D, Kanamaru A, Yoshie O. CC chemokine ligands 25 and 28 play essential roles in intestinal extravasation of IgA antibody-secreting cells. *J Immunol*. 2004; 173:3668–75. [PubMed: 15356112]
24. Mora JR, Iwata M, Eksteen B, Song SY, Junt T, Senman B, Otipoby KL, Yokota A, Takeuchi H, Ricciardi-Castagnoli P, Rajewsky K, Adams DH, von Andrian UH. Generation of gut-homing IgA-secreting B cells by intestinal dendritic cells. *Science*. 2006; 314:1157–60. [PubMed: 17110582]
25. Uematsu S, Fujimoto K, Jang MH, Yang BG, Jung YJ, Nishiyama M, Sato S, Tsujimura T, Yamamoto M, Yokota Y, Kiyono H, Miyasaka M, Ishii KJ, Akira S. Regulation of humoral and cellular gut immunity by lamina propria dendritic cells expressing Toll-like receptor 5. *Nat Immunol*. 2008; 9:769–76. [PubMed: 18516037]
26. Shigematsu T, Specian RD, Wolf RE, Grisham MB, Granger DN. MAdCAM mediates lymphocyte-endothelial cell adhesion in a murine model of chronic colitis. *Am J Physiol Gastrointest Liver Physiol*. 2001; 281:G1309–15. [PubMed: 11668040]
27. Kawachi S, Morise Z, Jennings SR, Conner E, Cockrell A, Laroux FS, Chervenak RP, Wolcott M, van der Heyde H, Gray L, Feng L, Granger DN, Specian RA, Grisham MB. Cytokine and adhesion molecule expression in SCID mice reconstituted with CD4+ T cells. *Inflamm Bowel Dis*. 2000; 6:171–80. [PubMed: 10961589]
28. Matsuzaki K, Tsuzuki Y, Matsunaga H, Inoue T, Miyazaki J, Hokari R, Okada Y, Kawaguchi A, Nagao S, Itoh K, Matsumoto S, Miura S. In vivo demonstration of T lymphocyte migration and amelioration of ileitis in intestinal mucosa of SAMP1/Yit mice by the inhibition of MAdCAM-1. *Clin Exp Immunol*. 2005; 140:22–31. [PubMed: 15762871]
29. Rivera-Nieves J, Olson T, Bamias G, Bruce A, Solga M, Knight RF, Hoang S, Cominelli F, Ley K. L-selectin, alpha4beta1, and alpha4beta7 integrins participate in CD4+ T cell recruitment to chronically inflamed small intestine. *J Immunol*. 2005; 174:2343–52. [PubMed: 15699171]
30. Souza HS, Elia CC, Spencer J, MacDonald TT. Expression of lymphocyte-endothelial receptor-ligand pairs, alpha4beta7/MAdCAM-1 and OX40/OX40 ligand in the colon and jejunum of patients with inflammatory bowel disease. *Gut*. 1999; 45:856–63. [PubMed: 10562584]
31. Arihiro S, Ohtani H, Suzuki M, Murata M, Ejima C, Oki M, Kinouchi Y, Fukushima K, Sasaki I, Nakamura S, Matsumoto T, Torii A, Toda G, Nagura H. Differential expression of mucosal addressin cell adhesion molecule-1 (MAdCAM-1) in ulcerative colitis and Crohn's disease. *Pathol Int*. 2002; 52:367–74. [PubMed: 12100519]
32. Apostolaki M, Manoloukos M, Roulis M, Wurbel MA, Muller W, Papadakis KA, Kontoyiannis DL, Malissen B, Kollias G. Role of beta7 integrin and the chemokine/chemokine receptor pair CCL25/CCR9 in modeled TNF-dependent Crohn's disease. *Gastroenterology*. 2008; 134:2025–35. [PubMed: 18439426]
33. Goto A, Arimura Y, Shinomura Y, Imai K, Hinoda Y. Antisense therapy of MAdCAM-1 for trinitrobenzenesulfonic acid-induced murine colitis. *Inflamm Bowel Dis*. 2006; 12:758–65. [PubMed: 16917232]
34. Picarella D, Hurlbut P, Rottman J, Shi X, Butcher E, Ringler DJ. Monoclonal antibodies specific for beta 7 integrin and mucosal addressin cell adhesion molecule-1 (MAdCAM-1) reduce inflammation in the colon of scid mice reconstituted with CD45RBhigh CD4+ T cells. *J Immunol*. 1997; 158:2099–106. [PubMed: 9036954]
35. Hesterberg PE, Winsor-Hines D, Briskin MJ, Soler-Ferran D, Merrill C, Mackay CR, Newman W, Ringler DJ. Rapid resolution of chronic colitis in the cotton-top tamarin with an antibody to a gut-homing integrin alpha 4 beta 7. *Gastroenterology*. 1996; 111:1373–80. [PubMed: 8898653]
36. Podolsky DK, Lobb R, King N, Benjamin CD, Pepinsky B, Sehgal P, deBeaumont M. Attenuation of colitis in the cotton top tamarin by anti- α 4 integrin monoclonal antibody. *J Clin Invest*. 1993; 92:372–380. [PubMed: 7686922]

37. Gorfu G, Rivera-Nieves J, Hoang S, Abbott DW, Arbenz-Smith K, Azar DW, Pizarro TT, Cominelli F, McDuffie M, Ley K. Beta7 integrin deficiency suppresses B cell homing and attenuates chronic ileitis in SAMP1/YitFc mice. *J Immunol.* 2010; 185:5561–8. [PubMed: 20926792]
38. Fiorucci S, Mencarelli A, Palazzetti B, Sprague AG, Distrutti E, Morelli A, Novobrantseva TI, Cirino G, Kotliansky VE, de Fougères AR. Importance of innate immunity and collagen binding integrin alpha1beta1 in TNBS-induced colitis. *Immunity.* 2002; 17:769–80. [PubMed: 12479823]
39. Farkas S, Hornung M, Sattler C, Edtinger K, Steinbauer M, Anthuber M, Schlitt HJ, Herfarth H, Geissler EK. Blocking MAdCAM-1 in vivo reduces leukocyte extravasation and reverses chronic inflammation in experimental colitis. *Int J Colorectal Dis.* 2006; 21:71–8. [PubMed: 15856265]
40. Kato S, Hokari R, Matsuzaki K, Iwai A, Kawaguchi A, Nagao S, Miyahara T, Itoh K, Ishii H, Miura S. Amelioration of murine experimental colitis by inhibition of mucosal addressin cell adhesion molecule-1. *J Pharmacol Exp Ther.* 2000; 295:183–9. [PubMed: 10991977]
41. Taniguchi T, Tsukada H, Nakamura H, Kodama M, Fukuda K, Saito T, Miyasaka M, Seino Y. Effects of the anti-ICAM-1 monoclonal antibody on dextran sodium sulphate-induced colitis in rats. *J Gastroenterol Hepatol.* 1998; 13:945–9. [PubMed: 9794195]
42. Soriano A, Salas A, Sans M, Gironella M, Elena M, Anderson DC, Pique JM, Panes J. VCAM-1, but not ICAM-1 or MAdCAM-1, immunoblockade ameliorates DSS-induced colitis in mice. *Lab Invest.* 2000; 80:1541–51. [PubMed: 11045571]
43. Olson TS, Bamias G, Naganuma M, Rivera-Nieves J, Burcin TL, Ross W, Morris MA, Pizarro TT, Ernst PB, Cominelli F, Ley K. Expanded B cell population blocks regulatory T cells and exacerbates ileitis in a murine model of Crohn disease. *J Clin Invest.* 2004; 114:389–98. [PubMed: 15286805]
44. Denning TL, Kim G, Kronenberg M. Cutting Edge: CD4+CD25+ Regulatory T Cells Impaired for Intestinal Homing Can Prevent Colitis. *J Immunol.* 2005; 174:7487–91. [PubMed: 15944246]
45. Schneider MA, Meingassner JG, Lipp M, Moore HD, Rot A. CCR7 is required for the in vivo function of CD4+ CD25+ regulatory T cells. *J Exp Med.* 2007; 204:735–45. [PubMed: 17371928]
46. Yuan Q, Bromley SK, Means TK, Jones KJ, Hayashi F, Bhan AK, Luster AD. CCR4-dependent regulatory T cell function in inflammatory bowel disease. *J Exp Med.* 2007; 204:1327–34. [PubMed: 17548518]
47. Mottet C, Uhlig HH, Powrie F. Cutting edge: cure of colitis by CD4+CD25+ regulatory T cells. *J Immunol.* 2003; 170:3939–43. [PubMed: 12682220]
48. Kang SG, Piniacki RJ, Hogenesch H, Lim HW, Wiebke E, Braun SE, Matsumoto S, Kim CH. Identification of a chemokine network that recruits FoxP3(+) regulatory T cells into chronically inflamed intestine. *Gastroenterology.* 2007; 132:966–81. [PubMed: 17324406]
49. Papadakis KA, Prehn J, Moreno ST, Cheng L, Kouroumalis EA, Deem R, Breaverman T, Ponath PD, Andrew DP, Green PH, Hodge MR, Binder SW, Targan SR. CCR9-positive lymphocytes and thymus-expressed chemokine distinguish small bowel from colonic Crohn's disease. *Gastroenterology.* 2001; 121:246–54. [PubMed: 11487533]
50. Rivera-Nieves J, Ho J, Bamias G, Ivashkina N, Ley K, Oppermann M, Cominelli F. Antibody blockade of CCL25/CCR9 ameliorates early but not late chronic murine ileitis. *Gastroenterology.* 2006; 131:1518–29. [PubMed: 17101325]

Biographies



Mora headshot



Andrian headshot



Villablanca headshot



Cassani headshot




Figure 1. Different Lymphocyte Subsets Use Distinct Homing Receptors and Ligands to Localize to Specific Regions of the Intestine

A) Effector CD8⁺ T cells use CCR9 and $\alpha 4\beta 7$, and possibly CXCR4 and/or CXCR3, to localize to the GI mucosa. Th17 cells might also use CCR6 to localize to small bowel and IgA-secreting cells use CCR10 to localize to GI and other mucosal tissue compartments. **B)** Expression of addressins varies throughout the intestine, even in the steady state. MAdCAM-1 is expressed along the whole intestine (small and large bowel) and it is upregulated during inflammation. CCL25, a ligand for CCR9, is expressed in a proximal-to-distal gradient in the small bowel but absent from the colon. CCL28, a ligand for CCR10, is expressed mostly in colon and other mucosal sites; it regulates localization of IgA-secreting cells, but not T cells. CCL20, a ligand for CCR6, is most highly expressed in Peyer's patches and the small bowel, but also it is upregulated in inflamed colon.




Figure 2. Interfering with Homing Receptors as Therapy for IBD

Natalizumab is a monoclonal antibody (mAb) that blocks the integrins $\alpha 4\beta 7$ and $\alpha 4\beta 1$, preventing their binding to MAdCAM-1 and VCAM-1, respectively. Similarly, AJM300 is an orally bioavailable antagonist of the integrin $\alpha 4$ subunit. The mAbs Vedolizumab and Pf-00547,659 bind specifically to $\alpha 4\beta 7$ and MAdCAM-1, respectively, blocking their interaction. Traficet-EN (CCX282-B) is an orally bioavailable selective antagonist of CCR9 that blocks its functional interaction with CCL25. Alicaforsen (ISIS 2302) is an antisense oligodeoxynucleotide that binds to the 3' UTR portion of the *ICAM1* mRNA and prevents its translation.

Table 1

Role of gut-homing receptors in experimental IBD models

Model	Gut Segment	Pathogenic cells	Advantages	Limitations	Blocking homing receptors	Ref.
Mouse DSS colitis	Colon	Innate immunity T/B cell independent (Occurs in RAG1 ^{-/-} mice)	Easy to set up, fast readout	Mostly colon inflammation, acute disease (can also be made chronic), T/B cell independent, little resemblance to human IBD?	α4β7/MAdCAM-1 Not determined (n.d.)	33, 39-42
Mouse TNBS colitis	Colon	Th1	Easy to set up, fast readout	Mostly colon, acute disease, little resemblance to human IBD?	Decreases colitis	33
Naive CD4 T cell transfer into RAG1 ^{-/-} or SCTD mice	Colon	Th1	Chronic disease, easy to set up, reproducibility	Mostly colon inflammation	Decreases colitis	34
Gut-tropic Th1 7 cell transfer into RAG1 ^{-/-} mice	Ileum and Colon	<i>Ex vivo</i> differentiated gut-homing Th17	Involvement of both small bowel (ileitis) and colon	Need transfer of <i>ex vivo</i> differentiated gut-homing Th17 cells	Decreases ileitis and colitis	14
Cotton-top tamarin	Colon	Th1?	Non-human primate	Cost, logistical limitations	Decreases colitis	35, 36
TNF [±] are mice (TNF-α overproduction)	Ileum	Th1	Affects small bowel (ileitis), chronic disease, model for human CD	Logistical (mice are non-commercially available)	Suppresses ileitis	32
Samp 1/Yit mice	Ileum	Th2, Th17, B cells	Affects small bowel (ileitis), chronic disease, model for human CD	Logistical (mice are non-commercially available)	Reduced ileitis in SAMP1/Yit β7 ^{-/-} mice Decreased ileitis when blocking both MAdCAM-1 and VCAM-1	29, 37, 50

Table 2

Targeting homing receptors in IBD

Target Adhesion molecules	Name	Indication	Mechanism	Clinical Phase	Advantages	Side effects	Ref.
Human ICAM-1 antisense oligodeoxynucleotide	Alicaforsen (ISIS 2302)	CD, UC	Reduction of ICAM-1 protein expression	Phase II/III	Clinical improvements and well tolerated	Injection/infusion site reactions; Systemic immunosuppression	64, 65, 66, 69, 70
Humanized IgG4 mAb anti- $\alpha 4$ integrin	Natalizumab	CD, UC	Inhibition of $\alpha 4\beta 7$ /MAdCAM-1 interaction and $\alpha 4\beta 1$ /VCAM-1 binding.	Phase IV FDA-approved	Long-term clinical response and/or remission following the withdrawal of concomitant corticosteroids; well tolerated.	Rare cases of progressive multifocal leukoencephalopathy (PML); Rare cases of melanoma?	71-79
Orally bioavailable $\alpha 4$ integrin inhibitor	AJM300	CD (UC?)	Inhibition of $\alpha 4\beta 7$ /MAdCAM-1 interaction and $\alpha 4\beta 1$ /VCAM-1 binding.	Phase II	Orally bioavailable	Same as Natalizumab?	82
Humanized IgG1 mAb anti- $\alpha 4\beta 7$ integrin	Vedolizumab (MLN-02)	CD, UC	Inhibition of MAdCAM-1-mediated leukocyte adhesion	Phase II	Good clinical response and well-tolerated	Infusion reaction with angioedema, nausea and nasopharyngitis; Efficacy limited by development of anti-drug antibody	84, 85
Human IgG2 mAb anti-MAdCAM	PF-00547,659	UC (CD?)	Inhibition of $\alpha 4\beta 7$ /MAdCAM-1 interaction	Phase I/II	Clinical improvements and well tolerated	No major side effects reported	86
Chemokine receptors							
Orally bioavailable antagonist for CCR9	Traflet-EN (CCX282-B)	CD GVHD? Cellac Disease? Intestinal transplant?	Inhibition of leukocyte chemotaxis to the ligand CCL25	Phase III	Clinical remission; well tolerated; orally bioactive; no risk of DTH reactions; no increased risk of systemic infection; no formation of anti-human Ab	No major side effects reported	50, 87, 88, 91