



Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2011 July ; 20(7): 1341–1349. doi:10.1158/1055-9965.EPI-11-0204.

VALIDATION OF A LIGHT QUESTIONNAIRE WITH REAL-LIFE PHOTOPIC ILLUMINANCE MEASUREMENTS: THE HARVARD LIGHT EXPOSURE ASSESSMENT QUESTIONNAIRE

Archna Bajaj¹, Bernard Rosner², Steven Lockley^{3,4}, and Eva S. Schernhammer^{2,3,5}

¹Case Western Reserve University School of Medicine, Cleveland, OH

²Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA

³Division of Sleep Medicine, Harvard Medical School, Boston, MA

⁴Circadian Physiology Program, Division of Sleep Medicine, Brigham and Women's Hospital, Boston, MA

⁵Department of Epidemiology, Harvard School of Public Health, Boston, MA

Abstract

Background—Light exposure at night is now considered a probable carcinogen. To study the effects of light on chronic diseases like cancer, methods to measure light exposure in large observational studies are needed. We aimed to investigate the validity of self-reported current light exposure.

Methods—We developed a self-administered semiquantitative light questionnaire, the Harvard Light Exposure Assessment (H-LEA) questionnaire, and compared photopic scores derived from this questionnaire with actual photopic and circadian measures obtained from a real-life 7-day light meter application among 132 women (85 rotating night shift workers and 47 day workers) participating in the Nurses' Health Study II.

Results—After adjustment for age, BMI, collection day, and night work status, the overall partial Spearman correlation between self-report of light exposure and actual photopic light measurements was 0.72 ($P < 0.001$; Kendall $\tau = 0.57$) and 0.73 ($P < 0.0001$; Kendall $\tau = 0.58$) when correlating circadian light measurements. There were only minimal differences in accuracy of self-report of light exposure and photopic or circadian light measurement between day ($r = 0.77$ and 0.78 , respectively) and rotating night shift workers ($r = 0.68$ and 0.69 , respectively).

Conclusions—The results of this study provide evidence of the criterion validity of self-reported light exposure using the H-LEA questionnaire. Impact: This questionnaire is a practical method of assessing light exposure in large scale epidemiologic studies.

Keywords

light; melatonin; validation; questionnaire

Correspondence to Eva S. Schernhammer, MD, DrPH, Channing Laboratory, 181 Longwood Avenue, Boston, MA 02115. Tel: (617) 525-4648; Fax: (617) 525-2008; eva.schernhammer@channing.harvard.edu.

AUTHOR CONTRIBUTIONS A.B. data collection, analysis, interpretation and manuscript preparation, B.R. data analysis and manuscript preparation. S.L. manuscript preparation, E.S. funding, data collection, analysis, interpretation, and manuscript preparation.

All of the authors declare no relevant conflict of interest.

Introduction

Environmental lighting powerfully influences the circadian system in humans (1). In 2007, the International Agency for Research on Cancer concluded that shift-work involving circadian disruption is “probably carcinogenic” to humans (2). Light exposure at night may increase cancer risk through suppression of melatonin, a hormone intimately linked to the circadian system that demonstrates cancer-protective effects (3). Observational studies have consistently associated night work with an increase primarily in breast cancer risk, but more recent work also suggests increased risk of colorectal, endometrial, and prostate cancer (4–13), potentially mediated through the melatonin pathway. Further explorations of the physiologic potential of the association between circadian disruption and disease risk are imperative.

Even though all measurements are prone to error, a light meter is likely to have few correlated errors and thus can be considered the criterion standard for light measurements. However, while standard light meters can be used, paper-based questionnaires are a more practical method of assessing light exposure in large-scale studies. Most prior studies of the reliability and validity of paper-based questionnaires investigated the correlation between self-reported sun exposure and ultraviolet radiation exposure measured by personal dosimeter to help understand the etiology of sun-related cancers (14–20). Studies were often limited to children and adolescents (15, 17, 20). Study designs described various methods of self-recording with different levels of success, such as recall of sun exposure time over the prior four days (15), and recording exposure and activity for periodic intervals (30 min. or 60 min.) during daytime hours (14, 16, 18).

To the best of our knowledge, ours is the first study to attempt to validate a paper-based questionnaire measuring current exposure to both sunlight and artificial light sources. We investigated in the Nurses' Health Study II the criterion validity of self-reported current light exposure by comparing light measures based on our paper-based tool, the Harvard Light Exposure Assessment (H-LEA) questionnaire, and actual light exposure as measured by the Daysimeter (Lighting Research Center, Rensselaer Polytechnic Institute, Troy, NY, USA) (21). Goals of our study were to establish the validity of the H-LEA questionnaire in ranking high versus low levels of light throughout a 24-hour day, and to evaluate for differences in validity by rotating night work status.

Materials and Methods

Study Population

Eligible study participants included women from the Nurses' Health Study II (NHS2) cohort, which was established in 1989 when 116,609 female nurses between 25 and 42 years old completed a mailed questionnaire. The women have since been followed by mailed questionnaires every two years to update exposure and disease information. In 2005, 90,051 participants were still alive and being followed. Volunteer participants for the light measurement validation study were recruited between November 2006 and April 2008, based on predefined eligibility criteria from the overall cohort, as outlined below.

Eligibility and Recruitment

Women who, at enrollment, presently worked or had worked rotating night shifts as well as women who never worked night shifts were identified from the NHS2 cohort. Roughly 60% of the nurses in the NHS2 cohort worked on rotating night shifts (alternating among day-, afternoon-, evening-, and night shifts; from this point forward the term “night workers” will be used when referring to these rotating night shift workers). Eligible participants were randomly selected from all women within the NHS2 cohort who met the following

inclusion/exclusion criteria: returned the 2005 questionnaire; answered the night work questions in the NHS2 questionnaires, from 1989 (baseline) onwards; were cancer-free in 2005 (with the exception of nonmelanoma skin cancer); did not report having cardiovascular disease up to 2005; presently (in 2005), did not take any hormonal preparations (oral contraceptives or postmenopausal hormones); were not pregnant or lactating; and were presently (in 2005) in the work force. After these exclusions, 18,044 women remained eligible, and of these, participants were randomly chosen, in two waves, using a computer-based random number generator. They were contacted initially by mail, with follow-up by email and telephone (see Figure 1). The first wave of these invitations was mailed in September 2006 to 720 participants. Because the majority of participants from this first wave were day workers, and the remaining nurses from the first wave who were not contacted were also day workers, a second wave of invitation letters was mailed in November 2007 in order to collect more night workers who would be willing to participate in the study. These invitation letters were mailed to NHS2 cohort members who met the following inclusion/exclusion criteria: answered the 2007 online questionnaire to determine current night work status; answered the night work questions in the NHS2 questionnaires, from 1989 (baseline) onwards; were cancer-free in 2005 (with the exception of nonmelanoma skin cancer); did not report having cardiovascular disease up to 2005; presently (in 2005) did not take any hormonal preparations (oral contraceptives or postmenopausal hormones); were not pregnant or lactating; answered the question about rotating night work in the 2007 online questionnaire (“Since June 2007, how many months have you worked rotating night shifts – none, 1–4 months, ..., 20+ months) and did not answer with “none”; and were not in the prior selection for the first wave of having a history of night work. To maximize enrollment of night workers, we selected all 242 current night workers who passed all exclusion criteria, were not part of the first wave, and worked at least 15 months of night work between 2005 and 2007. Further details regarding enrollment from these two recruitment waves are offered in Figure 1. Based on our initial target enrollment of 150 women, combining waves one and two, we accrued a total of 148 study participants until the end of the study.

Horne Ostberg Morningness-Eveningness Questionnaire

Morningness-eveningness preference was assessed with the Horne-Ostberg Morningness-Eveningness Questionnaire (MEQ) (22), a 19-item questionnaire with a total score ranging from 16 to 86, previously used extensively in adults and adolescents (23–26). A higher score is indicative of morning types, a lower score of evening types. We added the MEQ only halfway through our study, hence we obtained MEQ scores to define morningness-eveningness preference only in a subgroup of 88 participants.

Circadian Light Meter

The Daysimeter was developed by the Lighting Research Center (LRC) at Rensselaer Polytechnic Institute (RPI), Troy, NY (USA), as a personal circadian light exposure and activity meter to measure circadian light-dark and activity-rest patterns (21). It is a self-contained, battery-operated data logger worn as a lightweight headset. It has two optical sensors placed near the plane of one cornea; the first (photopic) sensor detects optical radiation and closely matches the standard photopic luminous efficiency function; the second (blue) sensor has an intrinsic long-wavelength response cut-off at approximately 580 nm together with a UV blocking filter creating a spectral response peaking at approximately 460 nm. From both the photopic and blue sensors, the Daysimeter stores a light value proportional to the logarithm of the short circuit current of the cell. An accelerometer within the Daysimeter is used to detect the subject's activity by measuring acceleration in both horizontal and vertical directions. The Daysimeter stores the root mean square value of the acceleration data for each axis over 30 seconds. Current through a reference diode in the Daysimeter is used to approximate the temperature. When data logging, the Daysimeter

takes a sequence of ten readings once a second, spread out over one cycle of 60 Hz. The maximum run time of the Daysimeter is 72 days, but in practice this is limited by battery life to 30 days of active logging. Further technical details about Daysimeter have been described elsewhere (27).

Measures obtained from the Daysimeter were summarized by calculating hourly averages of the 30-second data points of light exposure for each day. Mean photopic illuminance values (lux) were derived from the photopic sensor on the Daysimeter, and mean spectrally weighted illuminance values as a measure for circadian light were obtained from the blue sensor. These circadian light levels were approximated using post-processing algorithms based upon the model of circadian phototransduction by Rea et al.(28) and reported in lux, as outlined in more detail in Miller et al. (27). Mean corneal illuminance was approximated from the light questionnaire (H-LEA questionnaire) using estimated average illuminance values (lux) for each light source offered on the questionnaire. Values for these approximations were obtained from the Lighting Research Center (LRC) at RPI (Table 1). In the H-LEA questionnaire, we assigned “other” light sources a value of 10 lux based on our observation that nurses tended to report in their logs almost exclusively low light levels such as TV in a dark room, or PC, or candle light, as “other” light sources. If a woman noted more than one light source for an individual hour on the H-LEA questionnaire, the average illuminance for these light sources was calculated; for example, an “F” (representing fluorescent lamp) and “H” (representing halogen lamp) would amount to $100 \text{ lux} + 20 \text{ lux} = 120 \text{ lux} / 2 = 60 \text{ lux}$.

Harvard Light Exposure Assessment (H-LEA) Questionnaire

The initial version of the H-LEA questionnaire used in the present study was developed by investigators from the Division of Sleep Medicine at Harvard Medical School and subsequently modified for our validation study (see Figure 2). Light exposure was grouped into 7 types of light sources: halogen lamp, fluorescent lamp, incandescent light, other artificial light source, indoor natural light, sunlight/outdoor natural light, and darkness. Participants were asked to record the light source(s) they were exposed to hourly for seven consecutive days.

Study Design and Study Procedures

All study materials were mailed to participants. The study was approved by the Harvard School of Public Health and the Brigham and Women's Hospital Human Subjects Committee, and written informed consent was obtained from subjects. Each participant completed the self-administered H-LEA while simultaneously wearing the Daysimeter for seven consecutive days and received \$100 compensation upon successful completion of the study. Efforts to increase compliance rates included maintaining ample contact and communication with all participants. Investigator-initiated telephone contact was established prior to enrollment. During the seven day study period, study investigators supplied email and phone number information in case subjects had technical issues or questions related to the rather complex study procedures and materials. If a problem with the Daysimeter arose, a replacement Daysimeter was shipped to the participant as soon as possible, and the study dates were changed in accordance with the study protocol to a new seven day study period. Study material was accompanied by detailed instructions and thorough study information, including a precaution that study investigators were going to assess the recorded Daysimeter data to check for adherence to study guidelines, and if the participant was determined to be noncompliant, she would not receive the full \$100 compensation.

Participants were instructed to wear the Daysimeter for a seven-day period, during all waking hours, except when in water. We did not record whether participants wore

sunglasses when outdoors, but assumed that outdoor light levels that required them to wear sunglasses would still accurately be reflected by Daysimeter recordings much higher in relation to any measurement obtained from indoor light exposure. To protect the light meter while in, for example, the shower or a swimming pool, they were asked to place the Daysimeter nearby, on a bathroom counter or any surface away from the water, but in the same light environment. They were instructed to keep it next to their bed while sleeping. The Daysimeter is suited for use in working environments, including sterile operating rooms.

Exclusions Based on Faulty Data

Of 148 participants, 16 (10.8%) were excluded because of either technical issues or gross noncompliance with wearing the Daysimeter, leaving a total of 132 women for our analyses. Six of the women that were excluded used a particular Daysimeter for which it was detected only in hindsight that after its use in May 2007, a technical problem with the photopic sensor rendered the recorded data less useful. For four other women, the Daysimeter was either started incorrectly or it malfunctioned such that no data were recorded. It was possible to identify nurses who did not comply with the protocol by using data from the Daysimeter's on-board temperature sensor and accelerometer; room-temperature readings, as opposed to elevated temperatures when the Daysimeter is in closer contact with the body, and extended periods of inactivity were certain signs that the nurse did not wear the device when required. Based on these criteria, three women were excluded because they wore the Daysimeter for less than three days, and another three women were excluded because their recorded data showed flagrant noncompliance with wearing the Daysimeter over the course of the seven days, for example, only wearing the Daysimeter for a couple of hours each day. Of the remaining 132 women, ten others, whose recorded data indicated they were compliant less than 100% of the study time, but was not sufficient to conclude they 'barely' (i.e., less than 50% of the time) wore the Daysimeter, were still included for the analysis. A sensitivity analysis was performed excluding these ten subjects.

Statistical Methods

To assess criterion validity of our paper-based questionnaire (i.e., the degree to which measures obtained from this questionnaire correlate with an established external criterion (29)), we calculated partial Spearman correlation coefficients (with p-values) between light values from the H-LEA questionnaire, which were converted into illuminance values (lux) based on estimates provided by the Lighting Research Center, Rensselaer Polytechnic Institute, Troy, NY (Table 1) and the two light values obtained from the Daysimeter, - photopic illuminance (lux) and circadian light (lux), adjusting for nurses' ID. In secondary analyses, we additionally adjusted for day of the 7-day collection and factors that have previously been linked to the circadian system including age, BMI (which is associated with rotating night shift work in this cohort (30)), and whether or not participants were current night workers. We also conducted correlation analyses stratified by night work and day work as well as by night time (defined 7:00pm – 6:59am) and day time (defined as 7:00am – 6:59pm). In addition, we quantified the agreement between the three light measures using the Kendall tau coefficient, a measure of the degree of correspondence between two rankings (31). In subanalyses, we excluded 10 women whose recorded data indicated they were compliant less than 100% of the study over the course of the seven days, but not so much as to conclude they 'barely' wore the Daysimeter. Their exclusion did not change correlations appreciably, and we therefore kept them in our main validation subgroup.

Results

After all exclusions, 132 women were left in our validation sample for a total of 924 days (22,276 hours) of light collection. Characteristics of the 132 women in the validation study subgroup from within the NHS2 cohort, stratified by day and night work status, are shown in Table 2, along with respective characteristics of the full NHS2 cohort. Overall, 47 of the women in the validation sample were day workers and 85 were night workers. Participants came from 24 states. The 132 women who participated in this study were, for the most part, representative of women participating in the NHS2 cohort in 2005 (90,051 women). However, women in the validation group were on average slightly younger and more likely to be premenopausal, reflecting our selection criteria for entry (i.e., they still had to be in the work force). There were also fewer current smokers among the women who participated in our validation study, and they tended to exercise more. On average they had worked more years of night work than women from the NHS2 cohort overall. Similarly, when stratifying by day and night work status, night workers tended to be heavier, exercise less, consume less alcohol, and were more likely to currently smoke than were day workers.

Among the subset of 88 women from whom we queried chronotype, night workers were more likely to be definite evening types than were day workers (17.7% versus 10.6%), but there was no statistically significant difference in mean Horne-Osterberg scores between the two groups (mean scores 53.2 and 55.2 for day and night workers, respectively).

Light measures obtained from the Daysimeter and H-LEA questionnaire were averaged over each hour of a 24-hour day for all participants combined as well as stratified by night work status. These averages of the absolute values ranged from 0.31 lux to 1,148 lux of photopic illuminance, from 0.24 lux to 1,375 lux of circadian light, and from 2.2 lux to 504 lux for corneal illuminance as derived from the H-LEA questionnaire. When graphing these data (Figure 3), relative photopic illuminance and circadian light values closely resembled each other and appeared to be well-approximated by the estimates derived from the H-LEA questionnaire (Figure 2).

The close approximation of the Daysimeter values by the H-LEA questionnaire in Figure 3 was reflected in an overall partial Spearman correlation coefficient of $r = 0.72$ after adjusting for age, BMI, day of the 7-day collection, and night work status ($P < 0.001$; Kendall $\tau = 0.57$) for the correlation between H-LEA questionnaire estimates and photopic illuminance, and an $r = 0.73$ ($P < 0.0001$; Kendall $\tau = 0.58$) for the correlation between the H-LEA questionnaire and circadian light if taking the average across all seven days (Table 3). Crude partial correlation coefficients were very similar to those additionally adjusted for age, BMI, day of collection, and night work status. The overall positive correlations were only slightly weaker among night workers than among day workers (e.g., for photopic illuminance, $r = 0.77$ for day workers and $r = 0.68$ for night workers; all $P < 0.0001$). Moreover, strength of correlations between the individual days of the 7-day validation study did not vary appreciably (Table 3).

When defining daytime as the time between 7:00am and 6:59pm, and nighttime as the time between 7:00pm and 6:59am, Spearman rank correlations between light measures obtained from the H-LEA and circadian light measures from the Daysimeter among all women combined were 0.42 ($P < 0.001$) during daytime hours and 0.78 ($P < 0.001$) during nighttime hours. Among day workers, these correlations were 0.38 and 0.74; and among night workers, 0.44 and 0.80, respectively (all P -values < 0.001).

Discussion

In this, to our knowledge the first attempt to validate a paper questionnaire in relation to standard light measures of both sun light and artificial light sources, we provide evidence of the criterion validity of current light exposure by self report. We found strong correlations of 0.68 and above for light measures calculated from our paper-based H-LEA questionnaire and photopic illuminance as well as circadian light, both in women who worked day shifts and night shifts.

In validating a paper-based light questionnaire against light measures obtained from a standard light meter, it is important to use light meters that measure light as accurately as possible. The Daysimeter in our study had to be worn close to the eye to truly reflect light falling onto the retina – this is particularly important when considering the effects of light on the circadian system, because it is light reaching the retina that affects the circadian system, and illuminance on the cornea is a close surrogate to this value.

There is scant literature on the validity of using paper-based light exposure questionnaires to estimate actual light exposure. Most prior studies looked only at sun exposure as measured by ultraviolet radiation (UVR) dosimeters. O'Riordan, et al. (15) compared UVR exposure of mothers and infants less than one year old measured by the dosimeter worn on the wrist with a measure of estimated exposure using a diary based on recall by mothers of the prior four days. Study authors concluded that the association between estimated exposure and dosimeter readings was poor and needed improvement, citing the possible bias resulting from recall error. Thieden et al. (16) asked 44 Danish adult indoor workers to record outdoor activity between 07:00 and 19:00 in 30-minute intervals for a mean of 13 days during holiday periods and a mean of 17 days during work periods (June – September). Participants were asked to record their time outdoors, the location and clothing. Authors calculated skin area exposure hours from the diaries and found a highly significant ($P < 0.001$) correlation between the calculated exposure hours with UVR exposure as measured by dosimeter worn on the wrist. They concluded that questions other than exposure hours (e.g. type of clothing) were not crucial to obtain valid information about UV exposure, and by reducing the number of questions in the diary, participant compliance would likely increase. In 2006, Thieden et al. (19) used a shortened version of the diary and achieved better compliance amongst 407 participants, including children, adolescents, and adults, who were asked to answer “yes” or “no” to questions about sun exposure behavior on a daily basis for a total of 54,943 participation days. Diary completion rate was 95%, and subject compliance rate for wearing the dosimeter on days the diary was completed was 86.4%. Thieden et al. found a tendency to wear the dosimeter less during weekends than on workdays. Chodick et al. (14) similarly asked 124 volunteers from a cohort of radiologic technologists to wear a dosimeter on the shoulder and complete a daily activity diary for seven days, listing all activities undertaken for each 30-minute interval between 09:00 and 17:00. Results showed a significant correlation between the amount of recorded time spent outdoors and personal UVR dose measurement. In a pilot study, O'Riordan et al. (18) asked 27 participants to wear a personal dosimeter and complete a sun habits diary for four consecutive days, including recording their primary activity for each hour between 10:00 and 16:00. Results indicated a fair agreement between personal UVR exposure as measured by the dosimeter and time spent outside as reported in the diaries ($r_s = 0.32$, $P = 0.03$).

A paper-based log of light exposure is superior to other methods of assessing light exposure in larger-scale cohort studies because it is easy to administer. Strengths of the H-LEA questionnaire developed for this study include that it asks participants in the form of a structured diary to record light exposure on an hourly basis, limiting the potential for measurement error. The H-LEA questionnaire could possibly be further simplified by

including only three major light categories – indoors, outdoors, and darkness. Other strengths of our study are high rates of participation and success with participants enrolled from a well-documented cohort with longitudinal information on a number of important covariates such as BMI, age, and shift-work status.

Limitations of our study include the potential for noncompliance with wearing the Daysimeter, i.e., our criterion standard. A likely reason for noncompliance is that the Daysimeter is a conspicuous object worn on the head, and participants may have found this uncomfortable or felt awkward wearing it in public places. Note, however, that this is not a limitation for the paper questionnaire. Compliance would likely be much higher if study participants were asked only to fill out the paper questionnaire, as opposed to also wearing a light meter for seven days. Furthermore, participants in our study were asked to complete several other questionnaires, including keeping a diary of daily activity; participants are likely to be more compliant with less paperwork. Another limitation of our findings is that they may not be reflective of self-reported past light exposure. Future studies should assess the validity of the H-LEA questionnaire when used to assess average light exposure during a past period of time, as opposed to current light exposure akin to a structured diary, like in our study. Our estimate for the amount of light derived from sun exposure was conservatively based on 2,000 lux and may have underestimated actual light levels. However, correlation coefficients are based on the relative ranking of light levels, hence our results should not have been impacted by this. Whether the validity of our questionnaire, which is based on a group of middle-aged, highly motivated and healthcare educated Caucasian women can be generalized to other population groups including men or other age and ethnic groups needs to be studied. Similarly, women who volunteered to participate in our study appeared more health-conscious, as reflected in the fact that night workers in the validation group had much lower rates of smoking than night workers in the overall NHS2 cohort (see Table 2), and were thus perhaps more compliant.

In sum, our study provides evidence that adults are able to report exposure to both artificial and natural light using the H-LEA questionnaire with shown validity. This tool could prove useful in large-scale studies examining associations between light exposure and various disease endpoints.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This article was supported by the Cooperative Agreement Number R01 OH04192 from Centers for Disease Control and Prevention (CDC) and the U.S. National Institute of Health (R01 OH008171 “Effects of Light at Night on the Circadian System in Nurses”, PI E. Schemhammer and R01 CA50385, PI W. Willett). We are indebted to all the participants of the NHS2 cohorts for their dedication and we are particularly grateful to those nurses who agreed to additionally participate in our validation substudy. We would also like to express our thanks to Mark Rea, PhD (Lighting Research Center, RPI) and Walter Willett, MD, DrPH (Harvard School of Public Health, PI of the NHS2 cohort) for their insightful comments. The content of this article is solely the responsibility of the authors and does not necessarily represent the official views of the Department of Health and Human Services/CDC/NIOSH/NIH. Funding agencies did not have any role in the design of the study; the collection, analysis, or interpretation of the data; the decision to submit the manuscript for publication; or the writing of the manuscript.

References

1. Czeisler, C.; Dijk, D. Chapter 20: Human circadian physiology and sleep-wake regulation. In: Takahashi, J.; Turek, F.; Moore, R., editors. *Handbook of Behavioral Neurobiology: Circadian Clocks*. Plenum Publishing Co.; New York: 2001. p. 521-69.

2. Straif K, Baan R, Grosse Y, et al. Carcinogenicity of shift-work, painting, and fire-fighting. *Lancet Oncol.* 2007; 8:1065–6. [PubMed: 19271347]
3. Vijayalaxmi, Thomas CR Jr, Reiter RJ, Herman TS. Melatonin: from basic research to cancer treatment clinics. *J Clin Oncol.* 2002; 20:2575–601. [PubMed: 12011138]
4. Davis S, Mirick DK, Stevens RG. Night shift work, light at night, and risk of breast cancer. *J Natl Cancer Inst.* 2001; 93:1557–62. [PubMed: 11604479]
5. Hansen J. Increased breast cancer risk among women who work predominantly at night. *Epidemiology.* 2001; 12:74–7. [PubMed: 11138824]
6. Pukkala E, Auvinen A, Wahlberg G. Incidence of cancer among Finnish airline cabin attendants, 1967–92. *BMJ.* 1995; 311:649–52. [PubMed: 7549630]
7. Rafnsson V, Tulinius H, Jonasson JG, Hrafnkelsson J. Risk of breast cancer in female flight attendants: a population-based study (Iceland). *Cancer Causes Control.* 2001; 12:95–101. [PubMed: 11246849]
8. Schernhammer ES, Laden F, Speizer FE, et al. Rotating night shifts and risk of breast cancer in women participating in the nurses' health study. *J Natl Cancer Inst.* 2001; 93:1563–8. [PubMed: 11604480]
9. Schernhammer ES, Laden F, Speizer FE, et al. Night-shift work and risk of colorectal cancer in the nurses' health study. *J Natl Cancer Inst.* 2003; 95:825–8. [PubMed: 12783938]
10. Tynes T, Hannevik M, Andersen A, Vistnes AI, Haldorsen T. Incidence of breast cancer in Norwegian female radio and telegraph operators. *Cancer Causes Control.* 1996; 7:197–204. [PubMed: 8740732]
11. Conlon M, Lightfoot N, Kreiger N. Rotating shift work and risk of prostate cancer. *Epidemiology.* 2007; 18:182–3. [PubMed: 17179764]
12. Kubo T, Ozasa K, Mikami K, et al. Prospective cohort study of the risk of prostate cancer among rotating-shift workers: findings from the Japan collaborative cohort study. *Am J Epidemiol.* 2006; 164:549–55. [PubMed: 16829554]
13. Viswanathan AN, Hankinson SE, Schernhammer ES. Night shift work and the risk of endometrial cancer. *Cancer Res.* 2007; 67:10618–22. [PubMed: 17975006]
14. Chodick G, Kleinerman RA, Linet MS, et al. Agreement between diary records of time spent outdoors and personal ultraviolet radiation dose measurements. *Photochem Photobiol.* 2008; 84:713–8. [PubMed: 18435619]
15. O'Riordan DL, Stanton WR, Eyeson-Annan M, Gies P, Roy C. Correlations between reported and measured ultraviolet radiation exposure of mothers and young children. *Photochem Photobiol.* 2000; 71:60–4. [PubMed: 10649890]
16. Thieden E, Agren MS, Wulf HC. Solar UVR exposures of indoor workers in a Working and a Holiday Period assessed by personal dosimeters and sun exposure diaries. *Photodermatol Photoimmunol Photomed.* 2001; 17:249–55. [PubMed: 11722749]
17. Dwyer T, Blizzard L, Gies PH, Ashbolt R, Roy C. Assessment of habitual sun exposure in adolescents via questionnaire--a comparison with objective measurement using polysulphone badges. *Melanoma Res.* 1996; 6:231–9. [PubMed: 8819126]
18. O'Riordan DL, Glanz K, Gies P, Elliott T. A pilot study of the validity of self-reported ultraviolet radiation exposure and sun protection practices among lifeguards, parents and children. *Photochem Photobiol.* 2008; 84:774–8. [PubMed: 18179624]
19. Thieden E, Philipsen PA, Wulf HC. Compliance and data reliability in sun exposure studies with diaries and personal, electronic UV dosimeters. *Photodermatol Photoimmunol Photomed.* 2006; 22:93–9. [PubMed: 16606414]
20. Gies P, Roy C, Toomey S, MacLennan R, Watson M. Solar UVR exposures of primary school children at three locations in Queensland. *Photochem Photobiol.* 1998; 68:78–83. [PubMed: 9679453]
21. Bierman A, Klein TR, Rea MS. The Daysimeter: A device for measuring optical radiation as a stimulus for the human circadian system. *Measurement Science and Technology.* 2005; 16:2292–2299.
22. Horne JA, Ostberg O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int J Chronobiol.* 1976; 4:97–110. [PubMed: 1027738]

23. Park YM, Matsumoto K, Seo YJ, Shinkoda H. Sleep and chronotype for children in Japan. *Percept Mot Skills*. 1999; 88:1315–1329. [PubMed: 10485118]
24. Andrade MM, Benedito-Silva AA, Menna-Barreto L. Correlations between morningness-eveningness character, sleep habits and temperature rhythm in adolescents. *Braz J Med Biol Res*. 1992; 25:835–839. [PubMed: 1342618]
25. Chung MH, Chang FM, Yang CC, Kuo TB, Hsu N. Sleep quality and morningness-eveningness of shift nurses. *J Clin Nurs*. 2009; 18:279–84. [PubMed: 19120754]
26. Taillard J, Philip P, Chastang JF, Bioulac B. Validation of Horne and Ostberg morningness-eveningness questionnaire in a middle-aged population of French workers. *J Biol Rhythms*. 2004; 19:76–86. [PubMed: 14964706]
27. Miller D, Bierman A, Figueiro MG, Schernhammer E, Rea MS. Ecological measurements of light exposure, activity, and circadian disruption. *Lighting Res technol*. 2010; 42:271–284.
28. Rea MS, Figueiro MG, Bullough JD, Bierman A. A model of phototransduction by the human circadian system. *Brain Res Brain Res Rev*. 2005; 50:213–28. [PubMed: 16216333]
29. Tulsy DS. An introduction to test theory. *Oncology (Williston Park)*. 1990; 4:43–8. discussion 69–70. [PubMed: 2143409]
30. Schernhammer ES, Kroenke CH, Laden F, Hankinson SE. Night work and risk of breast cancer. *Epidemiology*. 2006; 17:108–11. [PubMed: 16357603]
31. Kendall M. A New Measure of Rank Correlation. *Biometrika*. 1938; 30:81–89.




Figure 1. Flow diagram: Subject recruitment

*Of these, only 132 participants were included in the final analysis, see explanation in Methods section.

During a typical day, describe your exposure to the below specified light sources. Please fill in as applies, for each single day of your 7-day trial, and circle the hours at which you had a meal, like indicated in the example below.*

H...Halogen Lamp F...Fluorescent Lamp I...Incandescent light O...Other Artificial Light Source
 N...Natural Light (Indoors) S...Sunlight, Natural Light (Outdoors) D...Darkness

EXAMPLE:

DATE	<input type="checkbox"/> NIGHT SHIFT WORKHOURS FROM												TO		OR <input type="checkbox"/> DAYSHIFT WORKHOURS FROM												TO		OR <input type="checkbox"/> OFF WORK	
01/01/2007	1am	2am	3am	4am	5am	6am	7am	8am	9am	10am	11am	noon	1pm	2pm	3pm	4pm	5pm	6pm	7pm	8pm	9pm	10pm	11pm	12am						
	D	D	D	D	D	D	I	N	F	F	F	S	F	F	F	F	H	F	F	F	F	I	I	H	H					

Figure 2. Harvard Light Exposure Assessment (H-LEA) Questionnaire

*Mealtime data was collected for other research purposes but not used in this study.




Figure 3. Mean hourly light values over 24-hour period (averaged over seven days) as measured by the Daysimeter and the H-LEA questionnaire (means, and standard deviations as indicated in graph by error bars), overall and stratified by night work status in validation subgroup of 132 participants from the NHS2.

Table 1

Estimated corneal illuminance provided by the Lighting Research Center, Rensselaer Polytechnic Institute (Troy, NY).

Assumed application	Estimated corneal illuminance
H...Halogen Lamp residence	20 lux
F...Fluorescent Lamp office	100 lux
I...Incandescent light residence	20 lux
O...Other Artificial Light Source *	10 lux
N...Natural Light (Indoors)	200 lux
S...Sunlight, Natural Light (Outdoors)	2000 lux
D...Darkness, suburban night sky	0.2 lux

* Mostly low light levels like TV in otherwise dark room, or candle light, or PC (information from diaries)

Table 2

Age and age-standardized characteristics of 132 participants from the Nurses' Health Study 2, stratified by work schedule and compared to full cohort of n=90,051 women participating in NHS2 (2005).

Characteristics/Lifestyle covariates	Validation Subgroup (n=132)		Full NHS2 cohort (n=90,051)	
	Day worker (n=47)	Night worker (n=85)	Never or past night worker (n=68,350)	Current night worker (n=21,701)
Age (yrs; mean, SD)	48.3 (4.4)	47.0 (5.0) [‡]	48.5 (4.7)	48.2 (4.7) [‡]
Body mass index (kg/m ² ; mean, SD)	25.8 (5.3)	28.2 (7.1) [‡]	26.3 (7.0)	27.5 (7.2) [‡]
Alcohol consumption (grams; mean, SD)	6.6 (11.2)	3.3 (5.6) [‡]	5.7 (9.6)	4.7 (8.5) [‡]
Current smokers (%)	1.8	2.5 [‡]	7.3	15.9 [‡]
Premenopausal (%)	54.5	60.3 [‡]	45.3	43.0
Number of years worked rotating night shifts (mean, SD)	6.9 (6.7) [‡]	10.7(6.0) ^{‡, †}	4.1 (3.7)	5.8 (5.0) [‡]
Parity [*] (mean, SD)	2.0 (0.9)	2.0 (0.9)	2.1 (0.9)	2.1 (0.9)
Exercise (METs [*] ; mean, SD)	23.9 (20.9)	25.7 (30.6)	23.2 (29.5)	22.7 (30.5)
Chronotype:[§]				
Definite morning type (%)	21.3	21.2	NA	NA
Probable morning type	46.8	32.9	NA	NA
Probable evening type (%)	21.3	27.1	NA	NA
Definite evening type (%)	10.6	17.7	NA	NA
HO scores (mean, sd)	53.2 (12.8)	55.2 (9.7)	NA	NA

* Among parous women only. METs = metabolic equivalents

[‡] Difference in frequency (chi square test) or mean (t-test) statistically significant between day and night worker groups, P<0.05.

[†] Difference in frequency (chi square test) or mean (t-test) statistically significant between day or night workers, respectively, comparing the validation subgroup with the full cohort, P<0.05.

[§] Home-Osterberg (HO) scores only available from a subgroup of women (11 day workers, 71 night workers)

Table 3

Kendall tau and Spearman correlation coefficient (95% CI) for correlation between corneal illuminance approximated from H-LEA questionnaire and photopic illuminance and circadian irradiance from Daysimeter measurements (criterion standard).*

	Kendall tau τ	Crude Spearman correlation r^*	Adjusted Spearman correlation r^*
PHOTOPIC ILLUMINANCE			
Overall (n=132)	0.57	0.72	0.72
Day workers (n=47)	0.63	0.77	0.77
Night workers (n=85)	0.54	0.68	0.68
Day 1	0.57	0.71	0.71
Day 2	0.61	0.76	0.76
Day 3	0.59	0.74	0.74
Day 4	0.55	0.69	0.70
Day 5	0.55	0.69	0.69
Day 6	0.54	0.68	0.69
Day 7	0.59	0.73	0.73
CIRCADIAN LIGHT			
Overall (n=132)	0.58	0.72	0.73
Day workers (n=47)	0.63	0.78	0.78
Night workers (n=85)	0.55	0.69	0.69
Day 1	0.57	0.72	0.72
Day 2	0.61	0.76	0.77
Day 3	0.59	0.74	0.74
Day 4	0.56	0.70	0.70
Day 5	0.55	0.70	0.70
Day 6	0.55	0.69	0.69
Day 7	0.59	0.74	0.74

* All correlation and Kendall τ coefficients are statistically significant ($P < 0.0001$). Crude values are adjusted for nurses' ID only; adjusted values are adjusted for nurses' ID, age, body mass index (BMI), day of the 7-day collection, and whether or not they were current night workers.