



Published in final edited form as:

Environ Res. 2012 August ; 117: 132–137. doi:10.1016/j.envres.2012.04.012.

Exposure to Polycyclic Aromatic Hydrocarbons and Serum Inflammatory Markers of Cardiovascular Disease

John D. Clark III^a, Berrin Serdar^b, David J. Lee^a, Kristopher Arheart^a, James D. Wilkinson^{a,c}, and Lora E. Fleming^{a,d}

^aDepartment of Epidemiology and Public Health, Miller School of Medicine, University of Miami, 1120 NW 14th Street, Room 1049, CRB Building (R669) Miami, FL 33136

^bDepartment of Environmental and Occupational Health, Robert Stempel College of Public Health and Social Work, Florida International University, University Park, HLS-II 591, 11200 SW 8th Street, Miami, FL, 33199

^cDepartment of Pediatrics, Miller School of Medicine, University of Miami, Batchelor Children's Research Institute, Room 521, P.O. Box 016820 (D-820) Miami, FL 33101

^dEuropean Centre for Environment and Human Health (ECEHH), Peninsula College of Medicine and Dentistry, Truro, Cornwall TR1 3HD UK

Abstract

Polycyclic aromatic hydrocarbons (PAHs) are environmental and occupational carcinogens produced by the incomplete combustion of organic materials, such as coal and petroleum product combustion, tobacco smoking, and food cooking, that may be significant contributors to the burden of cardiovascular disease in human populations. The purpose of this study was to investigate associations between ten monohydroxy urinary metabolites of four PAHs and three serum biomarkers of cardiovascular disease (fibrinogen, homocysteine, and white blood cell count). Using data on 3,219 participants aged 20 years and older from the National Health and Nutrition Examination Survey (NHANES) 2001–2004 dataset, the associations between PAH metabolites and serum inflammatory markers were analyzed using Spearman correlations and multiple linear regression modeling. The PAH metabolites of naphthalene, fluorene, phenanthrene, and pyrene each showed both positive and negative correlations with homocysteine, fibrinogen, and white blood cell count (correlation coefficient range: -0.077 to 0.143) in nonsmoking participants. Using multiple linear regression models adjusted for age, gender, race/ethnicity, and body mass index, estimates of weighted geometric means of inflammatory marker levels were not significantly different between high and low levels (75th vs. 25th percentiles) for all PAH metabolites in nonsmoking subjects. The results of this study do not provide evidence for a relationship between PAH exposure (as measured by urinary levels of PAH metabolites) and serum biomarkers of cardiovascular disease after controlling for tobacco use.

© 2012 Elsevier Inc. All rights reserved.

Corresponding Author: John D. Clark III, PhD, c/o Dept of Epidemiology & Public Health, University of Miami Miller School of Medicine, 1120 NW 14th Street, Room 1049, CRB Building (R669), Miami, FL 33136, tel: 323 382 2113, fax: 305 421 4833, jclark2@med.miami.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Polycyclic aromatic hydrocarbons; cardiovascular disease; homocysteine; fibrinogen; white blood cells

1. Introduction

Polycyclic aromatic hydrocarbons (PAH) are environmental and occupational carcinogens that are produced by the incomplete combustion of organic materials, such as coal and petroleum product combustion, tobacco smoking, and food cooking (ATSDR, 1995; Bartle, 1991; U.S. Department of Health and Human Services, 2005). In addition to causing cancers in different organs, exposure to high levels of PAHs has been suggested to contribute to the development of atherosclerosis and to increased rates of cardiovascular disease in humans (Burstyn et al., 2005; Tuchsén et al., 1996). Considering the contribution of tobacco smoke exposure, ambient air pollution, and occupational hazards to cardiovascular morbidity and mortality, PAHs may be significant contributors to the burden of cardiovascular disease in human populations.

Evidence from both basic science and public health research has suggested associations between PAH exposure and cardiovascular disease. Animal studies in various species have demonstrated that injections of PAHs significantly increase aortic plaque volumes in a dose dependent relationship, and that chronic PAH exposure significantly increases the number of inflammatory cells in atherosclerotic plaques (Albert et al., 1977; Curfs et al., 2005; Penn and Snyder, 1988). Studies investigating the public health impact of exposure to tobacco smoke and to fine particulate matter air pollution, both of which are known to be sources of PAHs in human populations, have shown both exposures to be associated with adverse cardiovascular events (Dominici et al., 2006; Mensah et al., 2005; Peel et al., 2007; Taylor et al., 1992; Walker et al., 2008). Additionally, employees in occupations with known exposures to high levels of PAHs have been reported to have an increased risk of ischemic heart disease and cardiovascular mortality due to cardiovascular disease (Burstyn et al., 2005; Evanoff et al., 1993; Hansen and Hansen, 1983; Ronneberg and Ronneberg, 1995; Tuchsén et al., 1996).

One mechanism by which PAHs may play a role in the development of atherosclerosis and cardiovascular disease is through inflammation. A number of inflammatory factors are important predictors and possible promoters of atherosclerosis (Wilson et al., 2004). Serum biomarkers of cardiovascular disease (such as white blood cell count, homocysteine, fibrinogen, and c-reactive protein) have been investigated in human populations as possible proximate mechanisms of adverse vascular outcomes and cardiovascular disease. Studies have shown that an elevated white blood cell count and serum c-reactive protein are associated with the development of cardiovascular disease (Bassuk et al., 2004; Danesh et al., 1998; Ridker et al., 2003). Elevated levels of homocysteine are associated with increased risk of cardiovascular disease while the mechanism is not yet completely elucidated, increased levels of homocysteine have been associated with impaired fibrinolysis (Tofler et al., 2002). A large body of evidence has established fibrinogen as an independent risk factor for cardiovascular disease and as an important mediator through which traditional risk factors, such as low-density lipoprotein, exert their effects (Kannel and Kannel, 2005). C-reactive protein has since been found to be associated with high levels of urinary monohydroxy PAH metabolites, suggesting that PAH exposure may play a role in the progression of atherosclerosis through inflammation (Everett et al., 2010).

The purpose of this study was to determine if associations existed between the serum biomarkers of cardiovascular disease fibrinogen, homocysteine, and white blood cell count and ten urinary metabolites of four PAHs using data from a study involving a nationwide probability sample representative of the non-institutionalized US population.

2. Materials and Methods

Using data from the National Health and Nutrition Examination Surveys (NHANES) 2001–2004, this study investigated associations between the urinary metabolites of PAHs and serum biomarkers of cardiovascular disease in a representative sample of the US civilian population. Conducted by the National Center for Health Statistics, the 2001–2004 NHANES used a complex sampling strategy to obtain a representative sample of the non-institutionalized U.S. civilian population aged two months or older from 15 US communities per survey cycle (NHANES, 2008). Participants completed detailed questionnaires, received physical examinations, and received laboratory analysis of biologic samples.

Measurements of urinary metabolites of PAHs were assessed in participants aged 6 years and over on a one-third sample of all NHANES participants. Within the database of NHANES participants from the 2001–2002 and 2003–2004 survey cycles, some participants did not have complete demographic data to allow for sampling weight calculation needed for analyses to be representative of the U.S. population. Therefore, for this study, a subsample of participants were identified who had a) valid sampling weights for use in analyses adjusting for the complex sampling design of the study and b) were age 20 years or older since children are believed to have different PAH exposure and metabolism compared to adults (Cohen Hubal et al., 2000).

White blood cell counts were performed on samples using the Beckman Coulter MAXM instrument in the Mobile Examination Centers. Homocysteine was measured using fluorescence polarization immunoassay on the Abbott IMX analyzer. Fibrinogen was measured by the clot detection technique to determine sample values. All variables (except for fibrinogen) were available for analysis in all two 2-year survey cycles; fibrinogen was available only for analysis for the first NHANES cycle (2001–2002).

At the time of this study, data on ten monohydroxy PAH metabolites were available from the 2001–2004 NHANES survey cycles for analysis. Urinary concentrations of PAH metabolites were measured by means of capillary gas chromatography combined with high-resolution mass spectroscopy methods that have been described previously (NHANES, 2001). Urinary concentrations of PAH metabolites were adjusted for urinary creatinine concentration as was done in the Center for Disease Control's Third National Report on Human Exposure to Environmental Chemicals (CDC, 2005).

Additional variables shown to be associated with urinary PAH metabolite levels in past studies were investigated for inclusion in adjusted linear regression models. These included age, gender, race/ethnicity, body mass index (BMI) (equal to the weight in pounds divided by the square of the height in inches), and tobacco smoke exposure (either being a current smoker or having either home or workplace exposure to secondhand smoke) (Suwan-ampai et al., 2009). Age, gender, and race were attained by self-report. Race/ethnicity was classified as Non-Hispanic White, Non-Hispanic Black, Hispanic, and other/mixed race. BMI was calculated using measured weight and height attained by physical examination.

All statistical analyses were conducted with SAS version 9.2 (SAS institute, Inc., Cary, NC) using survey procedures that adjusted for the complex survey design and included appropriate sampling weights developed by the National Center for Health Statistics for this random subsample to obtain accurate estimates representative of the non-institutionalized

civilian US population (CDC, 2005). Relationships between demographic characteristics and inflammatory markers were performed to identify covariates to be used in later analyses. Due to urinary PAH metabolites having a detection limit, analyses using ranks and percentiles were performed to properly handle left-censored PAH data. Spearman correlations were performed to investigate associations between levels of urinary PAH metabolites and inflammatory markers for current smokers and nonsmokers. Lastly, associations between urinary PAH metabolites and serum biomarkers of cardiovascular disease were analyzed using simple linear regression with adjustment for sampling weights. To investigate possible relationships between PAH metabolite and serum cardiovascular disease biomarkers in nonsmoking participants geometric mean differences were calculated for inflammatory markers by the interquartile range (75th vs 25th percentile) of each urinary PAH metabolite in nonsmoking subjects a multiple linear regression models adjusting the variables previously shown to be associated with PAH metabolites (age, gender, race, and BMI).

3. Results

Data on 21,161 NHANES participants were available from the 2001–2004 survey cycles. A random one-sixth subsample of NHANES participants received laboratory examination of urine samples for PAHs, for a total of 5,598 subjects. A total of 3,219 participants (57.5% of PAH subjects; 15.2% of all NHANES subjects) with PAH measurements were age 20 years or older and had valid sampling weights for inclusion in this analysis. Fibrinogen measurements were only available for subjects aged 40 years and older and in the 2001–2002 survey cycle, for a total of 1,008 participants. Sample size and weighted percentages for demographic characteristics of the NHANES 2001–2004 sample population included in this analysis are shown in Table 1.

To investigate associations between demographic covariables and serum inflammatory markers, weighted geometric mean differences of individual inflammatory markers were calculated by covariate category (Table 2). Mean homocysteine levels were significantly increased for participants aged 40–59 or 60+ years compared to age 20–39 years; mean fibrinogen level was significantly increased for age being 60+ years compared to those aged 20–39 years; mean white blood cell count was significantly decreased for age being 40–50 and 60+ years. Female gender was found to be associated with significantly decreased levels of homocysteine and significantly increased levels of fibrinogen and white blood cell count. Significant differences in mean levels of all inflammatory markers were seen in at least one racial category compared to White non-Hispanic subjects. Self-report of being a current smoker was significantly associated with increased levels of all inflammatory markers. Lastly, having a BMI defined as either overweight or obese was associated with significantly increased mean levels of fibrinogen and white blood cell count compared to normal and underweight subjects.

Spearman correlations of urinary metabolites of naphthalene, fluorene, phenanthrene, and pyrene with serum cardiovascular disease biomarkers were performed with stratification by smoking status (see online appendix, Table A). For homocysteine, current smokers were found to have weakly yet positive correlations with all urinary PAH metabolites (range: 0.133 to 0.259), while nonsmokers were found to have both positive and negative weak correlations with PAH metabolites (range: –0.077 to 0.143). For fibrinogen, current smokers and nonsmokers were found to have both positive and negative weak correlations with urinary PAH metabolites (smokers: –0.034 to 0.144, nonsmokers: –0.061 to 0.023). For white blood cell count, current smokers were found to have weak yet positive correlations with all urinary PAH metabolites (range: 0.024 to 0.118), while nonsmokers were found to

have both positive and negative weak correlations with PAH metabolites (range: -0.036 to 0.076).

Table 3 shows associations between estimates of adjusted, weighted geometric mean levels of inflammatory markers by the interquartile range (75th vs. 25th percentile) of monohydroxy PAH metabolites and serum biomarkers of cardiovascular disease in nonsmokers using age-adjusted and multiple linear regression models after adjusting demographic characteristics. For both homocysteine and white blood cell count, elevated levels of all urinary PAH metabolites were associated with small increases on inflammatory marker levels that were not statistically significant in nonsmoking subjects in both age-adjusted and multiple linear regression models (homocysteine range: 0.00 to 0.08 , white blood cell count range: 0.00 to 0.06). For fibrinogen, elevated levels of urinary PAH metabolites were associated with small increases and decreases in inflammatory marker levels that were not statistically significant in nonsmoking subjects in both age-adjusted and multiple linear regression models (range: -0.45 to 2.88).

4. Discussion

The results of this study do not provide evidence for a relationship between PAH exposure (as measured by urinary levels of PAH metabolites) and serum biomarkers of cardiovascular disease after controlling for tobacco use. While current smokers were found to have consistently small yet positive correlations between urinary PAH metabolites with levels of homocysteine and white blood cell count, this trend in correlations was not seen in nonsmokers. In the case of fibrinogen, no consistent trend was seen, with both small positive and negative correlations with urinary PAH metabolites for both smokers and nonsmokers. In our analysis of nonsmoking subjects using multiple linear regression models, no statistically significant differences existed in the estimated mean levels of inflammatory markers between high and low levels (75th vs. 25th percentiles) of each PAH metabolite. Thus, the results of our analyses do not support the tentative hypothesis that PAH exposure contributes to cardiovascular disease development.

The results of this study also do not support the hypothesis that environmental exposure to PAHs, such as with exposure to secondhand tobacco smoke and/or environmental air pollution, are a significant contributor to the burden of cardiovascular disease via inflammation in the U.S. population. Past research has shown that exposure to tobacco smoke, which contains mixtures of PAHs, is associated with elevated levels of serum biomarkers of cardiovascular disease (Bazzano et al., 2003; Wilkinson et al., 2007). Similarly, past studies have found that people exposed to high levels of ambient air particulate matter, which also contains mixtures of PAHs, may have increased levels of c-reactive protein, fibrinogen, and white blood cells (Ruckerl et al., 2007; Ruckerl et al., 2006; Schwartz, 2001). While a recent study that used data from the NHANES 2003–2004 found associations between urinary PAH metabolites and c-reactive protein, believed to be the most reliable of the inflammatory biomarkers, the results of this study do not provide additional evidence of such an association (Everett et al., 2010; Pearson et al., 2003). The lack of associations between urinary metabolites of the PAHs naphthalene, fluorene, phenanthrene, and pyrene with serum biomarkers of inflammation, however, suggests that PAH exposure may not be a significant contributor to the development of cardiovascular disease through an inflammatory mechanism in the US population.

One important strength of this study is that our analysis incorporated 4 years of collected data (2001–2004) with more than 3,000 adult subjects of a nationally representative sample of the entire US civilian population whose PAH metabolites were assessed under controlled conditions. In addition, this study used 3 biomarkers of inflammation as surrogate measures

of inflammation involved in cardiovascular disease development. However, the cross-sectional nature of NHANES limits the conclusions that can be made based on the results of this analysis. Additionally, many statistical tests were performed in our analysis, which may result in some statistically significant results that may not accurately represent the true state of tested relationships.

5. Conclusions

In conclusion, the results of this study do not provide evidence for a relationship between PAH exposure (as measured by urinary monohydroxy PAH metabolites) and serum biomarkers of cardiovascular disease independent of tobacco smoke exposure.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was funded in part by the National Institute of Environmental Health Science (NIEHS F30 ES015969) and the National Institute of Occupational Safety and Health (NIOSH R01 OH03915).

References

- Albert RE, Vanderlaan M, Burns FJ, Nishizumi M, Albert RE, Vanderlaan M, Burns FJ, Nishizumi M. Effect of carcinogens on chicken atherosclerosis. *Cancer Research*. 1977; 37:2232–2235. [PubMed: 405095]
- ATSDR. Toxicological Profile for Polycyclic Aromatic Hydrocarbons (PAHs). Atlanta: Agency for Toxic Substances and Disease Registry; 1995. p. 485
- Bartle, K. Analysis and occurrence of polycyclic aromatic hydrocarbons in food. In: Creaser, C.; Purchase, R., editors. *Food Contaminants: Sources and Surveillance*. Cambridge: The Royal Society of Chemistry; 1991.
- Bassuk SS, Rifai N, Ridker PM, Bassuk SS, Rifai N, Ridker PM. High-sensitivity C-reactive protein: clinical importance. *Current Problems in Cardiology*. 2004; 29:439–493. [PubMed: 15258556]
- Bazzano LA, He J, Muntner P, Vupputuri S, Whelton PK. Relationship between cigarette smoking and novel risk factors for cardiovascular disease in the United States. *Annals of Internal Medicine*. 2003; 138:891–897. [summary for patients in *Ann Intern Med*. 2003 Jun 3;138(11):145; PMID: 12779312]. [PubMed: 12779299]
- Burstyn I, Kromhout H, Partanen T, Svane O, Langard S, Ahrens W, Kauppinen T, Stucker I, Shaham J, Heederik D, Ferro G, Heikkila P, Hooiveld M, Johansen C, Randem BG, Boffetta P, Burstyn I, Kromhout H, Partanen T, Svane O, Langard S, Ahrens W, Kauppinen T, Stucker I, Shaham J, Heederik D, Ferro G, Heikkila P, Hooiveld M, Johansen C, Randem BG, Boffetta P. Polycyclic aromatic hydrocarbons and fatal ischemic heart disease. *Epidemiology*. 2005; 16:744–750. [PubMed: 16222163]
- CDC. Centers for Disease Control and Prevention (CDC), Department of Health and Human Services (DHHS); 2005. Third National Report on Human Exposure to Environmental Chemicals.
- Cohen Hubal EA, Sheldon LS, Burke JM, McCurdy TR, Berry MR, Rigas ML, Zartarian VG, Freeman NC. Children's exposure assessment: a review of factors influencing Children's exposure, and the data available to characterize and assess that exposure. *Environ Health Perspect*. 2000; 108:475–486. [PubMed: 10856019]
- Curfs DMJ, Knaapen AM, Pachen DMFA, Gijbels MJJ, Lutgens E, Smook MLF, Kockx MM, Daemen MJAP, van Schooten FJ. Polycyclic aromatic hydrocarbons induce an inflammatory atherosclerotic plaque phenotype irrespective of their DNA binding properties. *FASEB J*. 2005; 19:1290–1292. [PubMed: 15939734]

- Danesh J, Collins R, Appleby P, Peto R. Association of fibrinogen, C-reactive protein, albumin, or leukocyte count with coronary heart disease: meta-analyses of prospective studies. *JAMA*. 1998; 279:1477–1482. [PubMed: 9600484]
- Dominici F, Peng RD, Bell ML, Pham L, McDermott A, Zeger SL, Samet JM, Dominici F, Peng RD, Bell ML, Pham L, McDermott A, Zeger SL, Samet JM. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA*. 1998; 295:1127–1134. [see comment]. [PubMed: 16522832]
- Evanoff BA, Gustavsson P, Hogstedt C, Evanoff BA, Gustavsson P, Hogstedt C. Mortality and incidence of cancer in a cohort of Swedish chimney sweeps: an extended follow up study. *British Journal of Industrial Medicine*. 1993; 50:450–459. [PubMed: 8507598]
- Everett CJ, King DE, Player MS, Matheson EM, Post RE, Mainous AG 3rd. Association of urinary polycyclic aromatic hydrocarbons and serum C-reactive protein. *Environmental Research*. 2010; 110:79–82. [PubMed: 19836015]
- Hansen ES, Hansen ES. Mortality from cancer and ischemic heart disease in Danish chimney sweeps: a five-year follow-up. *American Journal of Epidemiology*. 1983; 117:160–164. [PubMed: 6829545]
- Kannel WB, Kannel WB. Overview of hemostatic factors involved in atherosclerotic cardiovascular disease. *Lipids*. 2005; 40:1215–1220. [PubMed: 16477805]
- Mensah GA, Brown DW, Croft JB, Greenlund KJ, Mensah GA, Brown DW, Croft JB, Greenlund KJ. Major coronary risk factors and death from coronary heart disease: baseline and follow-up mortality data from the Second National Health and Nutrition Examination Survey (NHANES II). *American Journal of Preventive Medicine*. 2005; 29:68–74. [PubMed: 16389129]
- NHANES. Laboratory Procedures Manual. 2001; Vol. 2008
- NHANES. Key Concepts About Survey History. 2008
- Pearson TA, Mensah GA, Alexander RW, Anderson JL, Cannon RO 3rd, Criqui M, Fadl YY, Fortmann SP, Hong Y, Myers GL, Rifai N, Smith SC Jr, Taubert K, Tracy RP, Vinicor F. Markers of inflammation and cardiovascular disease: application to clinical and public health practice: A statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation*. 2003; 107:499–511. [PubMed: 12551878]
- Peel JL, Metzger KB, Klein M, Flanders WD, Mulholland JA, Tolbert PE, Peel JL, Metzger KB, Klein M, Flanders WD, Mulholland JA, Tolbert PE. Ambient air pollution and cardiovascular emergency department visits in potentially sensitive groups. *American Journal of Epidemiology*. 2007; 165:625–633. [PubMed: 17194748]
- Penn A, Snyder C. Arteriosclerotic plaque development is 'promoted' by polynuclear aromatic hydrocarbons. *Carcinogenesis*. 1988; 9:2185–2189. [PubMed: 3142695]
- Ridker PM, Bassuk SS, Toth PP, Ridker PM, Bassuk SS, Toth PP. C-reactive protein and risk of cardiovascular disease: evidence and clinical application. *Current Atherosclerosis Reports*. 2003; 5:341–349. [PubMed: 12911843]
- Ronneberg A, Ronneberg A. Mortality and cancer morbidity in workers from an aluminium smelter with prebaked carbon anodes--Part III: Mortality from circulatory and respiratory diseases. *Occupational & Environmental Medicine*. 1995; 52:255–261. [PubMed: 7795741]
- Ruckerl R, Greven S, Ljungman P, Aalto P, Antoniadis C, Bellander T, Berglund N, Chrysohoou C, Forastiere F, Jacquemin B, von Klot S, Koenig W, Kuchenhoff H, Lanki T, Pekkanen J, Perucci CA, Schneider A, Sunyer J, Peters A, Group AS, Ruckerl R, Greven S, Ljungman P, Aalto P, Antoniadis C, Bellander T, Berglund N, Chrysohoou C, Forastiere F, Jacquemin B, von Klot S, Koenig W, Kuchenhoff H, Lanki T, Pekkanen J, Perucci CA, Schneider A, Sunyer J, Peters A. Air pollution and inflammation (interleukin6, C-reactive protein, fibrinogen) in myocardial infarction survivors. *Environmental Health Perspectives*. 2007; 115:1072–1080. [PubMed: 17637925]
- Ruckerl R, Ibald-Mulli A, Koenig W, Schneider A, Woelke G, Cyrus J, Heinrich J, Marder V, Frampton M, Wichmann HE, Peters A, Ruckerl R, Ibald-Mulli A, Koenig W, Schneider A, Woelke G, Cyrus J, Heinrich J, Marder V, Frampton M, Wichmann HE, Peters A. Air pollution and markers of inflammation and coagulation in patients with coronary heart disease. *American Journal of Respiratory & Critical Care Medicine*. 2006; 173:432–441. [PubMed: 16293802]

- Schwartz J. Air pollution and blood markers of cardiovascular risk. *Environmental Health Perspectives*. 2001; 109(Suppl 3):405–409. [PubMed: 11427390]
- Taylor AE, Johnson DC, Kazemi H, Taylor AE, Johnson DC, Kazemi H. Environmental tobacco smoke and cardiovascular disease. A position paper from the Council on Cardiopulmonary and Critical Care, American Heart Association. *Circulation*. 1992; 86:699–702. [PubMed: 1638735]
- Tofler GH, D'Agostino RB, Jacques PF, Bostom AG, Wilson PW, Lipinska I, Mittleman MA, Selhub J, Tofler GH, D'Agostino RB, Jacques PF, Bostom AG, Wilson PWF, Lipinska I, Mittleman MA, Selhub J. Association between increased homocysteine levels and impaired fibrinolytic potential: potential mechanism for cardiovascular risk. *Thrombosis & Haemostasis*. 2002; 88:799–804. [PubMed: 12428097]
- Tuchsen F, Andersen O, Costa G, Filakti H, Marmot MG, Tuchsen F, Andersen O, Costa G, Filakti H, Marmot MG. Occupation and ischemic heart disease in the European Community: a comparative study of occupations at potential high risk. *American Journal of Industrial Medicine*. 1996; 30:407–414. [PubMed: 8892545]
- U.S. Department of Health and Human Services. P. H. S., National Toxicology Program., Report on Carcinogens. Eleventh Edition 2005.
- Walker B Jr, Mouton CP, Walker B Jr, Mouton CP. Environmental influences on cardiovascular health. *Journal of the National Medical Association*. 2008; 100:98–102. [PubMed: 18277816]
- Wilkinson JD, Lee DJ, Arheart KL. Secondhand smoke exposure and C-reactive protein levels in youth. *Nicotine Tob Res*. 2007; 9:305–307. [PubMed: 17365762]
- Wilson PW, Cdc Aha, Wilson PWF. CDC/AHA Workshop on Markers of Inflammation and Cardiovascular Disease: Application to Clinical and Public Health Practice: ability of inflammatory markers to predict disease in asymptomatic patients: a background paper. *Circulation*. 2004; 110:e568–e571. [PubMed: 15611383]

Polycyclic aromatic hydrocarbon metabolites are not associated with cardiovascular biomarkers.

No consistent relationships between exposure and outcome were found independent of tobacco use.

Polycyclic aromatic hydrocarbon exposure is not a major contributor to cardiovascular disease.

Table 1

Demographic characteristics of sample populations from NHANES 2001–2004.

	n^a	Weighted %	95% Confidence Interval	
Age (years)				
20–39	1,172	39.90%	37.10%	- 42.69%
40–59	989	38.56%	36.10%	- 41.01%
60+	1,058	21.55%	19.66%	- 23.44%
Gender				
Male	1,547	47.87%	45.85%	- 49.89%
Female	1,672	52.13%	50.11%	- 54.15%
Race/ethnicity				
White, non-Hispanic	1,701	72.42%	68.16%	- 76.68%
Black, non-Hispanic	632	11.21%	8.87%	- 13.56%
Mexican American	662	7.48%	5.18%	- 9.77%
Other Hispanic	118	4.82%	2.52%	- 7.12%
Other	106	4.07%	2.92%	- 5.22%
Tobacco Smoke Exposure^b				
Secondhand Smoke Unexposed Nonsmoker	2,122	63.10%	60.06%	- 66.15%
Secondhand Smoke Exposed Nonsmoker	363	12.11%	10.55%	- 13.67%
Current Smoker	731	24.79%	22.33%	- 27.25%
Body Mass Index				
Normal/Underweight	1,004	34.35%	31.80%	- 36.91%
Overweight	1,111	33.54%	31.40%	- 35.68%
Obese	1,000	32.10%	29.43%	- 34.77%

^a – Sample size varies due to item non-response^b – Self-reported exposure to secondhand smoke in either the home or the workplace

Table 2

Weighted geometric mean differences and 95% confidence intervals of inflammatory markers by demographic characteristics.

	Homocysteine (mg/dL)				Fibrinogen (mg/dL)				White Blood Cell count (SI)				
	N ^a	Weighted Geometric Mean	95% Confidence Interval	N ^b	Weighted Geometric Mean	95% Confidence Interval	N ^a	Weighted Geometric Mean	95% Confidence Interval	Weighted Geometric Mean	95% Confidence Interval	Weighted Geometric Mean Difference	95% Confidence Interval
Age (years)													
20–39	1,110	7.35	reference	-			1,117	7.18	reference			reference	
40–59	954	8.42	0.85 – 1.30	508	355.18	reference	957	6.94	reference			-0.25	-0.41 – -0.09
60+	1,022	10.21	2.55 – 3.17	500	395.92	40.74	1,025	6.74	32.12 – 49.55			-0.44	-0.60 – -0.29
Gender													
Male	1,481	9.16	reference	515	352.18	reference	1,483	6.89	reference			reference	
Female	1,605	7.61	-1.55	493	376.14	23.96	1,616	7.08	10.95 – 37.43			0.19	0.04 – 0.35
Race/ethnicity													
White, non-Hispanic	1,651	8.48	reference	572	364.13	reference	1,657	7.06	reference			reference	
Black, non-Hispanic	584	8.42	-0.06	193	389.61	25.48	588	6.30	12.10 – 39.35			-0.76	-0.99 – -0.52
Mexican American	635	7.12	-1.36	174	351.63	-12.50	638	7.17	-27.91 – 3.62			0.11	-0.12 – 0.35
Other Hispanic	116	8.03	-0.45	25	349.03	-15.10	116	7.57	-36.45 – 7.64			0.51	0.15 – 0.88
Other	100	8.01	-0.46	44	350.33	-13.80	100	6.69	-43.14 – 18.24			-0.37	-0.69 – -0.02
Tobacco Smoke Exposure													
Secondhand													
Smoke Unexposed	2,027	8.12	reference	712	362.10	reference	2,039	6.75	reference			reference	
Nonsmoker													
Secondhand													
Smoke Exposed	357	8.19	0.07	104	355.09	-7.02	357	6.67	-34.88 – 23.23			-0.08	-0.34 – 0.19
Nonsmoker													
Current Smoker	699	8.92	0.80	191	378.47	16.37	700	7.80	2.48 – 30.78			1.04	0.84 – 1.25
Body Mass Index													
Normal/Underweight	966	8.19	reference	266	340.21	reference	969	6.69	reference			reference	
Overweight	1,073	8.40	0.21	367	355.53	15.32	1,076	6.91	1.32 – 29.90			0.22	0.07 – 0.38
Obese	958	8.35	0.16	325	391.65	51.44	965	7.39	39.55 – 63.71			0.70	0.49 – 0.93

^a – Sample size due to inflammatory marker availability for subjects aged 20 years and older from NHANES 2001–2004.

^b – Sample size due to inflammatory marker availability from subjects aged 40 years and older from NHANES 2001–2002.

Table 3

Estimates of weighted, adjusted geometric mean differences of inflammatory markers by the interquartile range (75th vs. 25th percentile) of urinary polycyclic aromatic hydrocarbon metabolite levels in nonsmoking subjects of NHANES 2001–2004 using multiple linear regression models.

Urinary PAH metabolite	Homocysteine (mg/dL)				Fibrinogen (mg/dL)				White Blood Cell count (SI)			
	N	Age-Adjusted Geometric Mean Difference	95% Confidence Intervals	N	Age-Adjusted Geometric Mean Difference	95% Confidence Intervals	N	Age-Adjusted Geometric Mean Difference	95% Confidence Intervals	N	Age-Adjusted Geometric Mean Difference	95% Confidence Intervals
1-Hydroxynaphthalene	2,272	0.02	-0.14 – 0.18	767	0.44	-13.09 – 14.48	2,284	0.01	-0.08 – 0.10	2,284	0.01	-0.08 – 0.10
2-Hydroxynaphthalene	2,261	0.07	-0.08 – 0.23	767	2.88	-9.89 – 16.11	2,273	0.06	-0.03 – 0.14	2,273	0.06	-0.03 – 0.14
2-Hydroxyfluorene	2,230	0.06	-0.10 – 0.23	767	0.18	-13.26 – 14.14	2,242	0.02	-0.07 – 0.11	2,242	0.02	-0.07 – 0.11
3-Hydroxyfluorene	2,223	0.11	-0.05 – 0.27	767	0.44	-13.01 – 14.40	2,295	0.03	-0.06 – 0.12	2,295	0.03	-0.06 – 0.12
9-Hydroxyfluorene ^a	1,051	0.07	-0.13 – 0.27		--- <i>b</i>	--- <i>b</i>	1,055	0.05	-0.11 – 0.20	1,055	0.05	-0.11 – 0.20
1-Hydroxyphenanthrene	2,227	0.03	-0.13 – 0.20	765	0.04	-13.33 – 13.92	2,239	0.02	-0.07 – 0.12	2,239	0.02	-0.07 – 0.12
2-Hydroxyphenanthrene	2,221	0.03	-0.14 – 0.19	766	0.29	-13.03 – 14.12	2,233	0.03	-0.06 – 0.12	2,233	0.03	-0.06 – 0.12
3-Hydroxyphenanthrene	2,186	0.00	-0.16 – 0.17	765	0.26	-13.06 – 14.07	2,198	0.00	-0.09 – 0.09	2,198	0.00	-0.09 – 0.09
4-Hydroxyphenanthrene ^a	1,027	0.08	-0.13 – 0.29		--- <i>b</i>	--- <i>b</i>	1,091	0.05	-0.09 – 0.19	1,091	0.05	-0.09 – 0.19
1-Hydroxypyrene	2,231	0.07	-0.09 – 0.23	767	0.62	-12.49 – 14.22	2,243	0.02	-0.07 – 0.12	2,243	0.02	-0.07 – 0.12
Urinary PAH metabolite	N	Multivariable Adjusted Geometric Mean Difference ^c	95% Confidence Interval	N	Multivariable Adjusted Geometric Mean Difference ^d	95% Confidence Interval	N	Multivariable Adjusted Geometric Mean Difference ^d	95% Confidence Interval	N	Multivariable Adjusted Geometric Mean Difference ^d	95% Confidence Interval
1-Hydroxynaphthalene	2,272	0.03	-0.14 – 0.20	767	0.45	-11.10 – 12.02	2,284	0.01	-0.08 – 0.08	2,284	0.01	-0.08 – 0.08
2-Hydroxynaphthalene	2,261	0.02	-0.14 – 0.19	767	1.51	-9.50 – 13.12	2,273	0.06	-0.04 – 0.13	2,273	0.06	-0.04 – 0.13
2-Hydroxyfluorene	2,230	0.02	-0.13 – 0.17	767	0.01	-11.01 – 11.37	2,242	0.03	-0.04 – 0.11	2,242	0.03	-0.04 – 0.11
3-Hydroxyfluorene	2,223	0.02	-0.14 – 0.18	767	0.45	-10.51 – 10.79	2,295	0.05	-0.02 – 0.12	2,295	0.05	-0.02 – 0.12
9-Hydroxyfluorene ^a	1,051	0.03	-0.21 – 0.28		--- <i>b</i>	--- <i>b</i>	1,055	0.06	-0.04 – 0.14	1,055	0.06	-0.04 – 0.14
1-Hydroxyphenanthrene	2,227	0.02	-0.15 – 0.20	765	-0.25	-11.45 – 10.77	2,239	0.02	-0.08 – 0.11	2,239	0.02	-0.08 – 0.11
2-Hydroxyphenanthrene	2,221	0.02	-0.17 – 0.20	766	-0.05	-11.02 – 11.29	2,233	0.02	-0.08 – 0.13	2,233	0.02	-0.08 – 0.13
3-Hydroxyphenanthrene	2,186	0.01	-0.14 – 0.16	765	0.25	-10.11 – 11.82	2,198	0.01	-0.07 – 0.08	2,198	0.01	-0.07 – 0.08
4-Hydroxyphenanthrene ^a	1,027	0.03	-0.19 – 0.25		--- <i>b</i>	--- <i>b</i>	1,091	0.04	-0.04 – 0.14	1,091	0.04	-0.04 – 0.14
1-Hydroxypyrene	2,231	0.01	-0.16 – 0.18	767	-0.45	-11.06 – 9.98	2,243	0.04	-0.04 – 0.11	2,243	0.04	-0.04 – 0.11

^a – PAH metabolites were not available for the 2001–2002 survey cycle.

b – Sample size due to inflammatory marker availability for subjects aged 20 years and older from NHANES 2001–2004.

c – adjusting for age, gender, race and body mass index.

d – adjusting for age, gender, race, and body mass index