



Published in final edited form as:

*Infect Control Hosp Epidemiol.* 2011 November ; 32(11): 1140–1142. doi:10.1086/662587.

## Not all nosocomial *Escherichia coli* bacteriurias are catheter-associated

Jonas Marschall, MD<sup>1</sup>, Kyle N Ota, MPH<sup>1</sup>, Jeffrey P Henderson, MD, PhD<sup>1</sup>, and David K Warren, MD, MPH<sup>1</sup> for the CDC Prevention Epicenters Program

<sup>1</sup>Division of Infectious Diseases, Washington University School of Medicine, St. Louis, MO

### Abstract

We prospectively determined what proportion of nosocomial *E. coli* bacteriurias are associated with urinary catheters. Only 46% (95% CI 37–56%) of nosocomial *E. coli* bacteriurias were catheter-associated. Compared to bacteriuric patients with catheters, non-catheterized patients were less likely to be male and have renal insufficiency or a recent urogenital procedure.

### Keywords

*Escherichia coli*; Bacteriuria; Urinary Tract; Catheter; Nosocomial

---

Urinary tract infections (UTIs) are the most common hospital-acquired infections and are thought to be primarily a consequence of urinary catheterization.<sup>1</sup> Strategies to prevent hospital-acquired UTIs focus almost exclusively on urinary catheter management. However, data to support the assumption that hospital-acquired UTIs can be equated with catheter-associated UTIs are very limited. In a recent editorial for a nationwide survey of practices to prevent hospital-acquired UTIs the author stated that 80% of these infections were catheter-associated but did not provide a reference.<sup>2</sup>

### METHODS

We performed a 12-month prospective cohort study of inpatients with *E. coli* bacteriuria (defined as  $5 \times 10^4$  colony-forming units/ml in a clean-voided urine culture, or  $5 \times 10^3$  in urine culture from a catheterized patient) starting August 1<sup>st</sup>, 2009, at Barnes-Jewish Hospital, a 1250-bed teaching hospital in Missouri. Adult patients with a first positive urine culture 48 hours after admission were included. Urine cultures were performed at the treating physician's discretion. We excluded patients with polymicrobial bacteriuria and/or concurrent, non-*E. coli* bacteremia. Using medical records, the patients' clinical presentation, vital signs, laboratory, radiological, and pharmacy data were recorded prospectively.

A bacteriuria episode was considered to be catheter-associated if a catheter had been in place within 48 hours before urine cultures were obtained. Asymptomatic bacteriuria (ASB) was defined as absence of urinary symptoms;<sup>3</sup> cystitis as presence of dysuria, frequency, or urinary retention; pyelonephritis as flank pain or tenderness and/or fever ( $\pm$  cystitis

---

Corresponding author and reprint requests to: Jonas Marschall, MD, Division of Infectious Diseases, Washington University School of Medicine, 660 S. Euclid, St. Louis, 63110, MO, USA, Phone: 314 454-8225, Fax: 314 454-5392, jmarscha@dom.wustl.edu.

*Potential conflicts of interest.* JM, KNO, and JPH: no conflicts of interest. DKW is a consultant for 3M Healthcare, C. R. Bard, Inc., and Cardinal Health, and receives research funding from Sage Products, Inc., Cubist Pharmaceuticals and bioMérieux.

symptoms). Sepsis was defined using established criteria. We reviewed blood cultures that were drawn within  $\pm 1$  day of the bacteriuria.

We used SPSS 18 (SPSS Inc., Chicago, IL) for data analysis. Univariate comparisons of categorical variables were performed with  $\chi^2$  test or Fisher's exact test as appropriate. Continuous independent variables were analyzed using Student's t test or Mann-Whitney U test. A two-sided  $p$  value of  $<0.05$  was considered significant. We entered variables with a  $p < 0.1$  in univariate testing into a multivariate logistic regression model. The study was approved by the Washington University Human Research Protection Office.

## RESULTS

One hundred eighty-three patients had hospital-acquired *E. coli* bacteriuria during the study period and met study criteria. Patients were diagnosed with ASB (77; 42%), cystitis (28; 15%), pyelonephritis (55; 30%), or unclassifiable bacteriuria (e.g., intubated patients) (23; 13%). Among asymptomatic patients, 65 (84%) were female.

Eighty-five [46% (95% confidence interval, 37–56%)] of 183 episodes were catheter-associated. Patients with catheter-associated bacteriuria were more likely to be male ( $p=0.003$ ), have renal insufficiency ( $p=0.02$ ) and a recent urological procedure ( $p=0.03$ ) (Table 1). There was no difference in frequency of ASB ( $p=0.6$ ).

One hundred fifty-one bacteriuric patients (83%) received antibiotic treatment, including 64 (83%) of patients with ASB. The presence of a catheter did not determine whether antibiotics were given [70/71 (99%) with vs. 77/80 (96%) without catheter;  $p=0.6$ ]. Among patients tested for it there was no difference in frequency of bacteremia ( $p=0.5$ ). In-hospital mortality was similar in catheter-associated vs. non-catheter-associated bacteriuria ( $p=1.0$ ) as was length of hospital stay after bacteriuria ( $p=0.08$ ). Independent factors predisposing to catheter-association in bacteriuric patients are shown in Table 1.

## DISCUSSION

It is widely assumed that the terms “hospital-acquired bacteriuria” and “catheter-associated bacteriuria” are synonymous. However, few data actually quantify urinary catheterization as a precursor of bacteriuria. The 1983 CDC guideline for prevention of catheter-associated UTIs states that 66–86% episodes of hospital-acquired bacteriuria are secondary to urinary instrumentation.<sup>4</sup> The corresponding reference does not explicitly provide this information.<sup>5</sup> Also, to our knowledge, this statement has not been reevaluated over the past three decades. We found that only 46% of hospital-acquired bacteriurias in a tertiary-care hospital were catheter-associated, lower than previously suspected. Why there was such a high proportion of non-catheterized bacteriuric patients is unclear. Changes in genitourinary hygiene during hospitalization may play a role as could medications that alter the bladder function. Possibly, hospital policies to reduce unnecessary device use resulted in a lower proportion of catheter-associated bacteriuria. The development of targeted preventive measures clearly depends on a better understanding of the pathogenesis of non-catheter-associated nosocomial bacteriuria.

In non-catheterized patients ASB may have been present before admission but remained undetected until later in the hospital course, leading to patients being mislabeled as having nosocomial bacteriuria. Testing this hypothesis would require admission urinary cultures to be obtained on patients. Antibiotic treatment of ASB was common (83%), independent of catheter status. Although ASB-related antibiotic overuse in long-term care facilities has stimulated interventions,<sup>6</sup> comparable data for acute-care hospitals are lacking. ASB may be

a major driver of antibiotic use (and antimicrobial resistance) in hospitals and therefore represents a target for antimicrobial stewardship.<sup>7</sup>

We identified a number of plausible independent predictors of catheter-associated bacteriuria. The lower level of catheter-association among bacteriuric women could be in line with their predisposition to ASB.<sup>8</sup> The need for monitoring fluid intake and output may contribute to higher catheterization frequency among bacteriuric patients with renal insufficiency. Lastly, catheters have been shown to result in post-procedure UTIs.<sup>9</sup>

Our data was obtained from medical records, including both physician and nurses' notes. It is possible that some urinary catheterizations were unrecorded, leading us to underestimate the number of catheter-associated bacteriurias. The imperfect correlation between catheterization and its documentation has been addressed in a recent study.<sup>10</sup> Also, the proportion of catheter-associated episodes might be higher for nosocomial pathogens other than *E. coli*.

In summary, we found catheter-associated infection to be less common in nosocomial *E. coli* bacteriuria than previously reported. A better understanding of non-catheter-associated bacteriuria could lead to improved infection prevention strategies among hospitalized patients.

## Acknowledgments

We thank Cherie Hill and Dorothy Sinclair for the invaluable help with data management.

This publication was made possible by Grant Number UL1 RR024992, Sub-Award KL2 RR024994 from the NIH-National Center for Research Resources (NCRR), a component of the National Institutes of Health (NIH), and NIH Roadmap for Medical Research. Its contents are solely the responsibility of the authors and do not necessarily represent the official view of NCRR or NIH.

JM is the recipient of a KL2 Career Development Award (5KL2RR024994-03). JPH holds a Career Award for Medical Scientists from the Burroughs Wellcome Fund and is also supported by K12 HD001459-09. DKW received a CDC Prevention Epicenter Program grant (CDC 1U1CI000033 301).

## References

1. Hooton TM, Bradley SF, Cardenas DD, et al. Diagnosis, prevention, and treatment of catheter-associated urinary tract infection in adults: 2009 International Clinical Practice Guidelines from the Infectious Diseases Society of America. *Clin Infect Dis*. Mar 1; 50(5):625–663. [PubMed: 20175247]
2. Nicolle LE. The prevention of hospital-acquired urinary tract infection. *Clin Infect Dis*. Jan 15; 2008 46(2):251–253. [PubMed: 18171257]
3. Nicolle LE, Bradley S, Colgan R, Rice JC, Schaeffer A, Hooton TM. Infectious Diseases Society of America guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clin Infect Dis*. Mar 1; 2005 40(5):643–654. [PubMed: 15714408]
4. Wong ES. Guideline for prevention of catheter-associated urinary tract infections. *Am J Infect Control*. Feb; 1983 11(1):28–36. [PubMed: 6551151]
5. Martin CM, Bookrajian EN. Bacteriuria prevention after indwelling urinary catheterization. A controlled study. *Arch Intern Med*. 1962; 110:703–711.
6. Zabarsky TF, Sethi AK, Donskey CJ. Sustained reduction in inappropriate treatment of asymptomatic bacteriuria in a long-term care facility through an educational intervention. *Am J Infect Control*. Sep; 2008 36(7):476–480. [PubMed: 18786450]
7. Gross PA, Patel B. Reducing antibiotic overuse: a call for a national performance measure for not treating asymptomatic bacteriuria. *Clin Infect Dis*. Nov 15; 2007 45(10):1335–1337. [PubMed: 17968830]

8. Colgan R, Nicolle LE, McGlone A, Hooton TM. Asymptomatic bacteriuria in adults. *Am Fam Physician*. Sep 15; 2006 74(6):985–990. [PubMed: 17002033]
9. Wells TH, Steed H, Capstick V, Schepanksy A, Hiltz M, Faught W. Suprapubic or urethral catheter: what is the optimal method of bladder drainage after radical hysterectomy? *J Obstet Gynaecol Can*. Nov; 2008 30(11):1034–1038. [PubMed: 19126285]
10. Meddings J, Saint S, McMahon LF Jr. Hospital-acquired catheter-associated urinary tract infection: documentation and coding issues may reduce financial impact of Medicare's new payment policy. *Infect Control Hosp Epidemiol*. Jun; 31(6):627–633. [PubMed: 20426577]

**Table 1**  
Comparison of patients with hospital-acquired *Escherichia coli* bacteriuria depending on catheter association

Variable	Total n=183	Catheter-associated n=85	Not catheter-associated n=98	p value	Odds Ratio (95% CI)
Gender (male)	46 (25%)	30 (35%)	16 (16%)	0.003	2.8 (1.4–5.7)
Race (white)	124 (68%)	60 (71%)	64 (65%)	0.4	
Age (years, median, range)	70 (20–98)	68 (24–96)	71 (20–98)	0.9	
Body Mass Index (kg/m <sup>2</sup> , median, range)	27.1 (12.1–64.2)	27.0 (17.2–64.2)	27.3 (12.1–63.1)	0.7	
Charlson comorbidity score (median, range)	3 (0–13)	3 (0–13)	3 (0–11)	0.4	
Diabetes mellitus	64 (35%)	31 (37%)	33 (34%)	0.7	
Renal insufficiency (Cr>1.5 mg/dl)	42 (23%)	26 (31%)	16 (16%)	0.02	2.2 (1.0–4.6)
Any malignancy	50 (27%)	26 (31%)	24 (25%)	0.4	
Dementia	32 (18%)	10 (12%)	22 (22%)	0.06	0.5 (0.2–1.2)
Benign prostatic hyperplasia	12 (7%)	8 (9%)	4 (4%)	0.1	
Urological procedure this admission	8 (4%)	7 (8%)	1 (1%)	0.03	10.4 (1.2–88.8)
Fever	52 (28%)	23 (27%)	29 (30%)	0.7	
Confusion, altered mental status	47 (26%)	23 (27%)	24 (25%)	0.7	
Sepsis	98 (54%)	47 (55%)	51 (52%)	0.7	
Pyelonephritis	55 (30%)	25 (29%)	30 (31%)	0.9	
Asymptomatic bacteriuria	77 (42%)	34 (40%)	43 (44%)	0.6	
Urinalysis with >10 WBC	121 (66%)	61 (72%)	60 (61%)	0.1	
<b>Outcomes</b>					
Bacteremia (tested n=70)	9/70 (13%)	3/33 (9%)	6/37 (16%)	0.5	
Length of hospital stay after bacteriuria (days, median, range)	4.9 (0.1–66.1)	5.6 (0.2–36.5)	4.2 (0.1–66.1)	0.08	
In-hospital mortality	13 (7%)	6 (7%)	7 (7%)	1.0	

NOTES. All data expressed as n (%) unless otherwise specified.

CI=confidence interval. Cr=Creatinine. WBC=White blood cells. Hosmer-Lemeshow goodness-of-fit p=0.635 (for the multivariate logistic regression model).