



Published in final edited form as:

*Parasitology*. 2012 June ; 139(7): . doi:10.1017/S003118201200008X.

## The effect of spatial heterogeneity on the aggregation of ticks on white-footed mice

G. Devevey and D. Brisson\*

Department of Biology, 415 S. University Avenue, Philadelphia, PA 19104, USA

### SUMMARY

Parasites are often aggregated on a minority of the individuals in their host populations. Although host characteristics are commonly presumed to explain parasite aggregation on hosts, spatio-temporal aggregation of parasites during their host-seeking stages may have a dominant effect on the aggregation on hosts. We aimed to quantify, using mixed models, repeatability and autocorrelation analyses, the degree to which the aggregation of blacklegged ticks (*Ixodes scapularis*) on white-footed mice (*Peromyscus leucopus*) is influenced by spatio-temporal distributions of the host-seeking ticks and by heterogeneity among mice. Host-seeking ticks were spatially aggregated at both the larval and nymphal life-stages. However, this spatial aggregation accounted for little of the variation in larval and nymphal burdens observed on mice (3% and 0%, respectively). Conversely, mouse identity accounted for a substantial proportion of the variance in tick burdens. Mouse identity was a significant explanatory factor as the majority of ticks parasitized a consistent set of mice throughout the activity seasons. Of the characteristics associated with mouse identity investigated, only gender affected larval burdens, and body mass and home range sizes in males were correlated with nymphal burdens. These analyses suggest that aggregation of ticks on a minority of mice does not result from the distribution of host-seeking ticks but from characteristics of the hosts.

### Keywords

burden; host-seeking ticks; index of dispersion; *Ixodes scapularis*; Pennsylvania; *Peromyscus leucopus*; repeatability

### INTRODUCTION

In the natural world, many things are not evenly or randomly distributed among individuals in populations. This pattern is clearly demonstrated in many animal species where a few dominant males sire the majority of offspring (Clutton-Brock, 1990) as well as in human economics where the majority of material wealth is aggregated in an ever-shrinking minority of people (e.g. Davies *et al.* 2007). Thus, in many natural systems the majority of results are derived from a minority of inputs, a pattern referred to as the Pareto principle or the 20/80 rule (Juran *et al.* 1962). The degree of aggregation of organisms across their potential environments can have fundamental consequences on ecosystem dynamics. For example, the abundance of many parasites is often a function of their distribution in time, in space, and among hosts. Further, highly aggregated distributions of macroparasites that vector microbial pathogens can increase the probability of persistence of the pathogen (Woolhouse *et al.* 1997; Perkins *et al.* 2003). Identifying the factors that result in parasite aggregation is a

fundamental step towards understanding the natural causes of high parasite abundances (Woolhouse *et al.* 1997; Lloyd-Smith *et al.* 2005).

Highly aggregated distributions of parasites on their hosts have been observed in several natural systems (reviewed by Shaw *et al.* 1998; Krasnov *et al.* 2010; Sanchez *et al.* 2011; discussed by Poulin, 2007 in chapter 6). Although host characteristics have been investigated to explain this (e.g. Folstad and Karter, 1992; Ezenwa *et al.* 2006; Bize and Roulin, 2009), aggregation on hosts can result from other factors such as the intricacies of the biology of the parasite or interactions with a heterogeneous environment. Indeed, theoretical models suggest that aggregated spatial distributions of host-seeking parasites are sufficient to generate aggregated distributions of parasites among identical hosts (Leung, 1998; Hansen *et al.* 2004). Thus, the spatial distribution of host-seeking parasites may be a dominant factor determining the subsequent distribution of parasites on hosts yet has received surprisingly little empirical attention. The distribution of the host-seeking stages of many parasites is either virtually absent (e.g. fleas and lice) or is difficult to define (e.g. in the case of microscopic parasites). However, many tick species have extensive host-seeking stages and are ideally suited to simultaneously quantify the contribution of host-seeking parasite behaviours and host identity –defined here as the complex of host characteristics –to the aggregation of parasites on a minority of hosts. For example, it has been previously shown that ticks in the genus *Ixodes* aggregate on different vertebrate species due in part to the spatial distribution of host-seeking ticks (Randolph, 1975; Ostfeld *et al.* 1996a, b; Shaw *et al.* 2003; Perkins *et al.* 2003; Krasnov *et al.* 2007, 2010). Extensive studies have shown that white-footed mice (*Peromyscus leucopus*) feed more ticks than eastern chipmunks (*Tamias striatus*) (Schmidt *et al.* 1999; Shaw *et al.* 2003; Brunner and Ostfeld, 2008) because mice are much more likely to use tick-infested microhabitats (Shaw *et al.* 2003). The variation of parasite burdens among hosts within species is also striking (Randolph, 1975; Brunner and Ostfeld, 2008) and assessing the factors affecting aggregation on individuals within species is essential to understanding the natural causes of variation in parasite abundances.

In this study we examined the aggregation of immature *I. scapularis* ticks (larvae and nymphs) both while seeking a host and while attached to white-footed mice. These data were used to quantify the effects of the spatio-temporal aggregation of host-seeking ticks and of heterogeneity among mice on the aggregation of ticks on hosts. Aggregation of ticks on hosts driven by spatial heterogeneity in host-seeking ticks is expected to result in the aggregation of ticks on mice in specific locations with strong spatial autocorrelation. Alternatively, aggregation of ticks on hosts driven by host heterogeneity is expected to result in the aggregation of ticks on a consistent set of mice over time regardless of the spatial location. We conducted this study in 2 sites in southeastern Pennsylvania, a region endemic for several tick-borne diseases such as human babesiosis, Bartonellosis, Lyme disease, and human granulocytic anaplasmosis (Steiner *et al.* 2008; Yeagley *et al.* 2009; Dubey *et al.* 2009) which has been largely ignored in previous studies of tick-borne pathogens.

## MATERIALS AND METHODS

### Study site

We monitored the abundance of larval and nymphal *I. scapularis* both while host seeking and while attached to white-footed mice from mid-April to mid-September in 2009 and 2010 at 2 sites in Southeastern Pennsylvania, USA. The 2 sites, called Monocassy and Northside, are located in Crow's Nest Preserve (Natural Lands Trust), Chester County (40°11'N, 75°17'W), a deciduous mid-Atlantic forest with interspersed agricultural lands. The Monocassy and Northside sites have similar underlying geology and weather conditions but differ slightly in forest type: Monocassy is dominated by oaks (*Quercus* spp.) with limited

understory; Northside is dominated by maples (*Acer* spp.) and tulip trees (*Liriodendron tulipifera*) with a well-developed understory. At each site, 2 study plots (225 × 225 m each) were established and separated by 200 m. The Monocassy and Northside sites were separated by 500 m of agricultural lands, dwellings, roads, and a creek.

### Aggregation of host-seeking ticks in time and space

The abundance of host-seeking immature ticks was monitored weekly on four 20 m<sup>2</sup> transects per plot. Transects were randomly distributed on the margins of each study plot and sampled between 09:00 and 21:00 in dry conditions. A 1-m<sup>2</sup> panel of fabric was dragged at a slow, steady pace over bare ground and vegetation and was repeated forward and backward on each transect. The panel was checked for ticks every 10 m, and all ticks were identified visually and stored in 70% ethanol. To control for variance caused by surveys occurring at different times of day, operators simultaneously surveyed transects at each plot. To control for variance caused by operator technique, each operator sampled transects on each plot at each session. The study plots were thus simultaneously sampled, and the 4 transects in each plot were dragged by 4 different operators at each session. The spatial distribution of host-seeking ticks was assessed using the index of dispersion  $D = \sigma^2 / \mu$ , a common metric of the aggregation (Randolph and Steele, 1985; Ostfeld *et al.* 1996a; Poulin, 2007; Brunner and Ostfeld, 2008), where  $\mu$  and  $\sigma^2$  are respectively the mean and the variance of larvae and nymphs among transects of a site during a session. Values of  $D$  significantly greater than 1 indicate aggregation and significantly less than 1 indicate over-dispersion. The confidence intervals were computed by non-parametric bootstrapping in R 2.13.0 (R Development Core Team, 2010).

### Aggregation of feeding ticks on mice and in space

Trapping grids were established on each study plot to assess the distribution of ticks attached to white-footed mice. Two trapping grids were established in 2009 and an additional 2 were added in 2010 for 4 total. Each trapping grid contained 64 trapping stations spaced every 15 m with 1 Sherman live-trap per locality (Sherman Traps, Inc., Tallahassee, FL, USA). Traps were baited with rolled-oats, set at 17:00, and checked the following morning before 10:00. Each captured mouse was fitted with a unique 4-digit ear tag and the sex, age according to pelage (juvenile, subadult, adult), body mass, and the number of larval and nymphal ticks attached to the ears and the head were recorded once per week. Importantly, the entire cohort of ticks feeding on a mouse is replaced every 4 days (Brunner *et al.* 2011) suggesting that ticks are not counted multiple times. The number of ticks on the ears and head of mice was correlated with the total number of ticks per mouse using identical methods employed here (Brunner and Ostfeld, 2008). While ear tags may affect tick burdens on the affected ear, ear tags do not affect the relative tick burdens among mice (Ostfeld *et al.* 1993). The relative home range size of each mouse trapped on 3 or more occasions was estimated using Minimum Convex Polygons in Ranges VII (Anatrack Ltd, Wareham, Dorset, UK). The minimum convex polygon – the area enclosed by the localities where a mouse was caught – provides an estimate of the relative space used by each mouse which is correlated with, but are not a precise estimate of, the true home range size. However, the home range size estimates acquired by the minimum convex polygons in this study are very similar to estimations obtained previously by radiotelemetry (Wolff, 1985). All experiments were conducted in compliance with the Institutional Animal Care and Use Committee of the University of Pennsylvania.

The distribution of attached larvae and nymphs on mice or associated with each trap locality was assessed using the index of dispersion  $D$  and the dispersion parameter,  $k$ , of the negative binomial distribution (Randolph, 1975; Shaw *et al.* 1998; Brunner and Ostfeld, 2008). The best model describing the distribution of ticks on mice in each trapping session was

determined using the nls library for non-linear models under R 2.13.0 (R Development Core Team, 2010). This model was compared to a model assuming a random distribution of ticks on mice created by sampling from the empirical data (code in the supplemental file). The 95% confidence intervals of the model were determined by non-parametric bootstrapping.

### Factors affecting aggregation

We assessed whether larval and nymphal ticks were consistently aggregated at the same trap locations or on the same mice – indicating that factors unique to those locations or those mice results in greater tick burdens – by calculating the coefficient of repeatability,  $r$ , on standardized larval and nymphal burden data (see Lessells and Boag, 1987; Falconer and MacKay, 1996). The standardized tick burden was calculated as  $z_i = (x_i - \mu)/\sigma$  where  $x_i$  is the tick burden of the  $i^{th}$  mouse or trap station,  $\mu$  is the mean tick burden within each plot, and  $\sigma$  is the standard deviation in tick burden within each plot (Sokal and Rohlf, 1995). Standardizing tick burdens accounts for seasonal and among-plot heterogeneity of raw tick burdens (ANOVAs testing the effect of plot and season,  $P > 0.98$ ). Standardization was essential due to the aggregation of tick activity in both time and space (see results).

We quantified the effect of spatial heterogeneity and mouse identity on the variance in tick burdens among mice using mixed-effect models following the Residual Maximum Likelihood method for each tick life stage. The analyses modelled standardized tick burdens as a function of 2 random factors, trapping locality and mouse identity, and a random error term. The dataset analysed was restricted to mice trapped at multiple localities and to localities where several mice were observed in order to parse the effects of mouse identity and spatial location. This procedure partitions the proportion of the total variance explained by each random factor.

Variance that can be attributed to properties specific to a location are expected to be spatially and temporally correlated. Spatial autocorrelation among trapping localities was analysed in PASSaGE software (Rosenberg and Andersen, 2011) using *Moran's I*. A value of zero for *Moran's I* indicates a random spatial pattern, whereas values of 1 indicate a perfect spatial autocorrelation and  $-1$  perfect dispersion (Räty and Kangas, 2007). Temporal autocorrelation was tested by correlating average standardized tick burdens of localities visited in 2009 and in 2010.

The proportion of the variance that can be attributed to mouse identity was further investigated using mixed-effects models. The larval and nymphal burdens were modelled as a function of 4 mouse traits (sex, age, mass, home range size) using mouse ear tag number as a random factor. We followed a backward stepwise procedure starting with all second-degree interactions.

In all analyses, continuous variables were log-transformed when necessary to achieve normality and homoscedasticity. Values are presented as mean  $\pm$  s.e. unless otherwise noted, analyses were performed using JMP 9.0 (SAS institute Inc., Cary, NC, USA).

## RESULTS

### Aggregation of host-seeking ticks in time and space

The activity periods of larval and nymphal ticks were strongly aggregated in time and were restricted to the periods between July and early September (Fig. 1A) and early May and July (Fig. 1B), respectively. The larval and nymphal peak activity periods were not overlapping. We restricted all subsequent analyses to the time-frame that comprised 80% of the larvae or 80% of the nymphs observed as times when few ticks would likely have a strong and potentially misleading effect of analyses of distributions due to random sampling effects.

Host-seeking larval abundances were significantly different among sites. Interestingly the rank ordering of larval densities among sites varied among years such that sites did not retain consistently high or low larval densities across years (ANOVA,  $F_{3,248} = 4.9$ ,  $P = 0.003$ , interaction year\*site  $F_{1,248} = 7.51$ ,  $P = 0.007$ ). The densities of host-seeking larvae were lower in Monocassy than in Northside in 2009, whereas densities were lower in Northside than in Monocassy in 2010 (Fig. 1A). The density of host-seeking nymphs was much lower in 2010 than in 2009, without significant difference between sites (ANOVA,  $F_{2,305} = 71.7$ ,  $P < 0.001$ ; Year:  $F_{1,305} = 140.4$ ,  $P < 0.001$ ).

Within each plot, host-seeking larvae and host-seeking nymphs were significantly aggregated in space (larval average  $D = 201.4 \pm 31.0$ ,  $P < 0.05$ , nymphal average  $D = 2.89 \pm 0.6$ ,  $P < 0.05$ ) such that  $81.5 \pm 1.1\%$  of larvae and  $81.9 \pm 1.3\%$  of nymphs were collected on  $33.6 \pm 2.5\%$  and  $39.9 \pm 17.7\%$  of the sampled transects at each session.

### Aggregation of feeding ticks on mice and in space

The average larval burden on mice differed significantly among sites and between years ( $F_{2,20} = 8.98$ ,  $P = 0.002$ ; site:  $F_{1,20} = 7.33$ ,  $P = 0.014$ ; year:  $F_{1,20} = 8.93$ ,  $P = 0.007$ ) (Fig. 1A). Likewise, the average nymphal burden on mice differed significantly among sites and among years ( $F_{2,39} = 43.22$ ,  $P < 0.001$ ; site:  $F_{1,39} = 4.47$ ,  $P = 0.041$ ; year:  $F_{1,39} = 82.14$ ,  $P < 0.001$ ) (Fig. 1B). The seasonality of attached larvae and nymphs is consistent with the seasonality of the host-seeking activity patterns of each life stage.

Within each plot, the majority of larvae and the majority of nymphs were aggregated on a minority of mice at each session (larval average  $D = 11.0 \pm 2.0$ ,  $P < 0.05$ ; nymphal average  $D = 1.97 \pm 0.3$ ,  $P < 0.05$ ). The cumulative distributions of larvae on mice and nymphs on mice were best described by power functions (cumulative distribution of larvae =  $\text{mice}^{3.02}$ ,  $R^2 = 0.96$ ; cumulative distribution of nymphs =  $\text{mice}^{8.31}$ ,  $R^2 = 0.81$ ). These analyses demonstrate that most larvae ( $79.6 \pm 0.0\%$ ) and most nymphs ( $79.8 \pm 0.9\%$ ) were attached to a minority of mice ( $46.9 \pm 1.8\%$  and  $25.4 \pm 2.5\%$  for larval and nymphs, respectively). The distribution of ticks on mice is more aggregated than expected given a random distribution of ticks on mice (Fig. 2A and B). The distribution of ticks on mice follows a negative binomial distribution (larval distribution,  $k = 0.862$ ,  $X_{83}^2 = 70.33$ ,  $P = 0.84$ ; nymphal distribution,  $k = 0.299$ ,  $X_{14}^2 = 19.6$ ,  $P = 0.14$ ) that strongly supports a non-random distribution.

Importantly, the burdens of ticks on mice were significantly repeatable between captures in both the larval and nymphal datasets (Table 1A and B) such that mice with high tick burdens in one week had high tick burdens at all subsequent captures. That is, ticks were aggregated on a consistent subset of mice throughout the larval- or nymphal-activity seasons. The repeatabilities of nymphal burdens on mice were less distinct in 2010 due to the low numbers of observed nymphs (Table 1B).

The larval burden of mice in specific localities was also repeatable over the season (Table 1A), albeit weakly, suggesting that mice in some localities had more ticks than in others. On the contrary, we did not observe repeatability of the nymphal burden at localities (Table 1B).

### Mouse identity, not the distribution of host-seeking ticks, explains the observed aggregation on mice

A large proportion of the variance in standardized larval burden in each session was explained by mouse identity and only a small proportion by the trapping locality (GLM-REML:  $R^2 = 0.53$ ; Mouse: 35.1% of total variance; Locality: 3.7% of total variance) suggesting that mouse traits and not location determine larval burden. Likewise, variance in

standardized nymphal burdens could be attributed to mouse identity while none was due to trapping locality (GLM-REML:  $R^2 = 0.19$ ; Mouse: 10.9% of total variance; Locality: 0.0% of total variance).

Spatial auto-correlation in larval and nymphal burdens was not detected among the trapping locations. *Moran's I* was not significantly different from 0 for either the larval or the nymphal datasets (Fig. 3A and B). The absence of spatial autocorrelation in tick burdens despite the inherent territoriality of mice suggests that meso-scale habitat factors cannot account for the variance in tick burden. Furthermore, neither larval nor nymphal burdens at specific localities were repeatable between years (Larvae:  $F_{1,61} = 0.05$ ,  $P = 0.83$ ; Nymphs:  $F_{1,68} = 0.07$ ,  $P = 0.80$ ). That is, localities that had high tick burdens in 2009 were not more likely to have high tick burdens in 2010 (Fig. S1A and B-online version only).

Among the analysed mouse traits, only gender was significantly correlated with larval burdens (Males > Females) (Table 2A). Mouse age, mass, and home range size were not significantly correlated with larval burdens. Nymphal burdens were significantly and positively correlated with body mass and with the size of home range, but only in male mice (Table 2B). Mouse age was not included as a factor to explain nymphal burdens due to insufficient numbers of juveniles and subadults observed during the nymphal activity period.

## DISCUSSION

Immature blacklegged ticks are highly aggregated in time and in space while seeking a host and are also highly aggregated on a subset of mice. During their activity season, host-seeking larvae and nymphs are aggregated on a fraction of the sampled space at both large and local scales. Substantial inter-individual heterogeneity in burdens on mice was observed even after accounting for the strong seasonal and spatial heterogeneity in host-seeking ticks. Indeed, very little of the variation in burdens among mice could be explained by the areas where the mouse was captured, suggesting that the observed aggregation of host-seeking ticks in space has little effect on the observed aggregation of ticks on mice. The identity of the mouse, however, explained a substantial proportion of the variation in tick burdens suggesting that variation in individual mouse characteristics results in greater or fewer ticks per mouse. This result was further supported by the repeatability of ticks on mice over time such that highly parasitized mice remain highly parasitized throughout the season. This study suggests that the aggregation of ectoparasites on specific hosts, even parasites that are generalist with low dispersal rates, is not a byproduct of parasite ecologies but originate primarily from heterogeneity among individual hosts.

### Aggregation of host-seeking ticks in time and in space

As expected, season was the most important factor predicting the density of host-seeking ticks and variation of burdens between April and October. These data correspond in time with several other studies in the Northeastern US that describe the seasonal peaks in tick host-seeking activity and tick burdens on wildlife (Wilson and Spielman, 1985; Fish, 1993; Ostfeld *et al.* 1996a, b; Goodwin *et al.* 2001; Brunner and Ostfeld, 2008). The factors that affect host-seeking tick densities on a coarse scale are not permanent environmental characteristics. For example, the greatest larval densities in 2009 were reported from the Northside site, which had significantly lower densities than Monocassy in 2010. The factors that resulted in the observed differences in tick densities among sites are not discernable from the current data but are likely due to complex interactions between large- and local-scale factors (Wilson, 1998; Jones *et al.* 1998; Randolph, 2004; Ostfeld *et al.* 2006; Gern *et al.* 2008; Dobson *et al.* 2011).

The mechanisms that result in the fine-scale aggregation of host-seeking larvae are likely different than the mechanisms resulting in the observed aggregation of host-seeking nymphs. It is commonly postulated that larvae are highly aggregated as a natural consequence of the emergence of several thousand larvae from the single egg-mass produced by one fertilized female (for example Randolph and Steele, 1985; Daniels and Fish, 1990; Stafford, 1992). However, the spatial aggregation observed in the host-seeking nymphs must result from more complex mechanisms as the spatial location of nymphs is determined primarily by where they fall from a host after the larval bloodmeal (Ostfeld *et al.* 1996a). Potential hypotheses are that structural differences in microhabitats influence the spatial distribution of nymphs by causing nymphs to cluster in areas where they have better overwintering success or by engorged larvae dropping simultaneously from their hosts (Ostfeld *et al.* 1996a). The clustering of larvae and nymphs on a few of the 20 m<sup>2</sup> transects assessed in this study suggests that the phenomena concordant with these biological mechanisms occur at a very fine scale. Interactions among large- and local-scale environmental factors may be important to future research aimed at discovering characteristics that affect focal tick densities.

It is important to note that all measures of spatial aggregation are extremely sensitive to scale (size and number of transects; Bohan, 2000), which could result in the numerical differences in the degree of aggregation observed among studies (Randolph and Steele, 1985; Ostfeld *et al.* 1996a). Further, metrics of aggregation are sensitive to the mean number of organisms observed (Poulin, 2007) suggesting extreme caution when comparing the degree of aggregation among studies. For example, the degree of aggregation of nymphs attached to hosts in the current study is lower than that of attached larvae when assessed by the index of dispersion ( $D = 1.97$  vs  $11.0$  for nymphs and larvae, respectively, where greater numbers indicate greater aggregation) but greater when considering the aggregation parameter in the negative binomial ( $k = 0.299$  vs  $0.862$  where smaller numbers indicate greater aggregation) or the proportion of mice that host 80% of the ticks ( $25.4\%$  vs  $46.9\%$  for nymphs and larvae). Thus, metrics of aggregation can be used to statistically determine the presence of a non-random distribution but are of limited use in comparing the degree of aggregation among studies with different sampling. Finally, like all statistical tests, the power to detect aggregated distributions is correlated with the sample size of hosts, which often results in an underestimate of the true aggregation levels in the population (Poulin, 2007).

### Aggregation of attached ticks in space and on mice

Heterogeneity in the tick burdens among mice within plots was strongly apparent after accounting for spatial heterogeneity. At each trapping session, the number of ticks feeding on each mouse was highly variable (up to 85 larvae and 16 nymphs) with a minority of mice hosting the majority of ticks. These data are consistent with previous reports from the USA (Siegel *et al.* 1991; Schmidt *et al.* 1999; Brunner and Ostfeld, 2008) and from reports on other *Ixodes* species on a variety of mammals in Europe (Cotton and Watts, 1967; Randolph, 1975; Randolph and Steele, 1985; Randolph *et al.* 1999; Perkins *et al.* 2003; Bown *et al.* 2008; Kiffner *et al.* 2011a, b). These observations could result from 3 phenomena, each resulting in fundamentally different temporal patterns of tick burdens on mice and in localities. First, at each session a random subset of mice could utilize areas with ephemerally high host-seeking tick densities leading to an aggregation of ticks on different subsets of mice and at different subsets of locations at each trapping session. Second, spatial heterogeneity in the environment could lead to areas with permanently high host-seeking tick densities resulting in highly repeatable tick burdens within localities and strong autocorrelation among localities. Lastly, heterogeneity in mouse characteristics or behaviours would lead to highly repeatable tick burdens on mice without regard to locality.

Our data strongly support the last hypothesis as larvae and nymphs were highly aggregated on a consistent set of mice throughout the activity seasons.

Larval, but not nymphal, burdens were also significantly repeatable in localities within each year. However, the majority of the repeatability in localities was caused by resident mice visiting a consistent set of traps. That is, larval burdens of mice captured in the same localities were uncorrelated while the larval burdens of individual mice captured in different locations were strongly correlated. Further, we found no evidence of spatial autocorrelation of tick burdens among localities between years, nor evidence of temporal correlations of tick burdens at the same localities among years. These data strongly support the hypothesis that the spatial aggregation of host-seeking ticks is not responsible for the aggregation of ticks on individual mice.

Tick burdens on individual mice were repeatable throughout the seasons suggesting that ticks were consistently aggregated on the same subset of mice. For example, of the mice with the highest larval burden (top 20%) in July 2010, half remained in the group of mice with the greatest burden a month later while the remaining were more parasitized than the average. The repeatability of the tick burden on mice was significantly greater than random in all but 3 of the analyses likely due to the very small number of ticks observed in 1 larval and 2 nymphal datasets. Interestingly, these results are consistent with previous observations, suggesting that some mice with high nymphal burdens also have high larval burdens (Randolph *et al.* 1999; Randolph, 2004; Brunner and Ostfeld, 2008). The repeatability values of the tick burdens on mice measured in this study ( $r = 0.10-0.46$ ) are similar to repeatability values reported for heritable behavioural traits such as the exploratory behaviour of great tits ( $r = 0.27-0.48$ ; Dingemanse *et al.* 2002) or the risk-sensitivity of the three-spined stickle back ( $r = 0.05-0.7$ ; Dzielwczynski and Crovo, 2011). These data are especially intriguing as the entire cohort of ticks feeding on a mouse is replaced every 3–5 days (Brunner *et al.* 2011) suggesting that the variation among mice in tick burden results from perennial or slow-changing behavioural or physiological traits of individual mice.

Gender, age, mass, and home-range size are among the host traits that have been noted to affect ectoparasite burdens in other systems (for example Behnke and Wakelin, 1973; Brown *et al.* 1994; Perkins *et al.* 2003; Randolph, 2004; Sanchez *et al.* 2011). Of these traits, only gender significantly explained differences in larval burden with males more parasitized than females, a common trend in many ectoparasites (Randolph, 2004; Morand *et al.* 2006; Krasnov *et al.* 2005; Hillegeess *et al.* 2008; Kiffner *et al.* 2011a, b; Davidar *et al.* 1989; Ostfeld *et al.* 1996b; Brunner and Ostfeld, 2008). However, gender does not completely explain the variation among hosts as there was considerable inter-individual heterogeneity within each sex and there were no significant differences among the genders in nymphal burdens. Nymphal burdens were positively correlated with body mass and home range sizes in males but not in females. Unlike other rodents, the white-footed mouse is not sexually dimorphic for body mass ( $P > 0.6$ ) suggesting that size cannot account for the differences in tick burdens among the genders as it does in wood mice (*Apodemus sylvaticus*; Harrison *et al.* 2010). Similarly, the larger home-range size of males is often evoked as a cause of higher burden (Zuk and McKean, 1996; Robinson *et al.* 2009) but it is not consistent with the data from our study as the home ranges of males were not larger than those of females ( $P > 0.14$ ). However, differences among the sexes may become apparent using more accurate methods to estimate home range size such as radiotelemetry.

We expect that a complex combination of unmeasured factors such as ranging behaviour, timing and duration of activity, microhabitat use, grooming behaviours, hormones and immunological responses affect tick encounter rates and burdens. For example, testosterone,

which experimentally reduces resistance to tick feeding and increases tick burden in wood mice and bank voles (*Myodes glareolus*) parasitized by *I. ricinus* (Hughes and Randolph, 2001), may also affect space use behaviours (Ellis and Turek, 1983; Rosemitt, 1989; Lynn *et al.* 2000; Seivwright *et al.* 2005; Grear *et al.* 2009; but see also Minerly *et al.* 2008) which can affect encounter rates with ticks (Boyer *et al.* 2010). Future studies, which can exclude the spatial heterogeneity of host-seeking ticks as an explanatory factor, are needed to experimentally assess the potentially complex combination of behavioural, immunological, and physiological mechanisms underlying heterogeneity in tick burdens among individuals within populations. Understanding the mechanisms that lead to aggregation of ticks on specific hosts is particularly important for human disease risk as those hosts responsible for feeding many ticks are highly likely to be infected by pathogens, and to subsequently infect many naïve ticks. Therefore, a key element in understanding and controlling the transmission of these diseases is identifying the combination of characteristics of individuals that result in disproportionately large tick burdens.

## Acknowledgments

Authors want to acknowledge the field technicians, Maria Gomez-Solecki and Rick Ostfeld. Pierre Bize, Erica Foley, Chris Graves, Camillo Khatchikian, Jamie McKay and Maarten Voordouw gave very insightful comments. Finally, authors want to thank two anonymous referees who greatly helped to improve the manuscript.

### FINANCIAL SUPPORT

The study was supported by grant AI076342 from the National Institute of Health and grant CK000170 from the Center for Diseases Control and Prevention. G. D. was supported by the Swiss Science Foundation Fellowship for Prospective Researcher PBLAP3-127724/1.

## REFERENCES


- Behnke JM, Wakelin D. Survival of *Trichuris muris* in wild populations of its natural hosts. *Parasitology*. 1973; 67:157–164. [PubMed: 4748999]
- Bize P, Roulin A. Effects of common origin and common rearing environment on variance in ectoparasite load and phenotype of nestling Alpine Swifts. *Evolutionary Biology*. 2009; 36:301–310.
- Bohan DA. Spatial structuring and frequency distribution of the nematode *Steinernema feltiae* Filipjev. *Parasitology*. 2000; 121:417–425. [PubMed: 11072905]
- Bown KJ, Lambin X, Telford GR, Ogden NH, Telfer S, Woldehiwet Z, Birtles RJ. Relative importance of *Ixodes ricinus* and *Ixodes trianguliceps* as vectors for *Anaplasma phagocytophilum* and *Babesia microti* in Field Vole (*Microtus agrestis*) Populations. *Applied and Environmental Microbiology*. 2008; 74:7118–7125. [PubMed: 18820068]
- Boyer N, Reale D, Marmet J, Pisanu B, Chapuis J-L. Personality, space use and tick load in an introduced population of Siberian chipmunks *Tamias sibiricus*. *Journal of Animal Ecology*. 2010; 79:538–547. [PubMed: 20202009]
- Brown ED, Macdonald DW, Tew TE, Todd IA. *Apodemus sylvaticus* infected with *Heligmosomoides polygyrus* (Nematoda) in an arable ecosystem – Epidemiology and effects of infection on the movements of male mice. *Journal of Zoology*. 1994; 234:623–640.
- Brunner JL, Cheney L, Keesing F, Killilea M, Logiudice K, Previtali A, Ostfeld RS. Molting success of *Ixodes scapularis* varies among individual blood meal hosts and species. *Journal of Medical Entomology*. 2011; 48:860–866. [PubMed: 21845946]
- Brunner JL, Ostfeld RS. Multiple causes of variable tick burdens on small-mammal hosts. *Ecology*. 2008; 89:2259–2272. [PubMed: 18724736]
- Clayton, DH.; Moore, J. *Host-Parasite Evolution: General Principles and Avian Models*. Oxford, UK: Oxford University Press; 1997.
- Cotton MJ, Watts CHS. Ecology of tick *Ixodes trianguliceps* Birula (Arachnida Acarina Ixodoidea). *Parasitology*. 1967; 57:525–531. [PubMed: 6048570]

- Clutton-Brock, TH. Reproductive Success: Studies of Individual Variation in Contrasting Breeding Systems. Chicago, USA: University of Chicago Press; 1990.
- Daniels TJ, Fish D. Spatial-distribution and dispersal of unfed larval *Ixodes dammini* (ACARI, Ixodidae) in Southern New-York. Environmental Entomology. 1990; 19:1029–1033.
- Davidar P, Wilson M, Ribeiro JMC. Differential distribution of immature *Ixodes dammini* (ACARI, Ixodidae) on rodent hosts. Journal of Parasitology. 1989; 75:898–904. [PubMed: 2693676]
- Davies JB, Sandstroem S, Shorrocks A, Wolff EN. Estimating the level and distribution of global household wealth. \*\*\*UNU-WIDER, 2007/77. 2007
- Dingemanse NJ, Both C, Drent PJ, Van Oers K, Van Noordwijk AJ. Repeatability and heritability of exploratory behaviour in great tits from the wild. Animal Behaviour. 2002; 64:929–938.
- Dobson ADM, Taylor JL, Randolph SE. Tick (*Ixodes ricinus*) abundance and seasonality at recreational sites in the UK: Hazards in relation to fine-scale habitat types revealed by complementary sampling methods. Ticks and Tick-Borne Diseases. 2011; 2:67–74. [PubMed: 21771540]
- Dubey JP, Bhatia CR, Lappin MR, Ferreira LR, Thorn A, Kwok OCH. Seroprevalence of *Toxoplasma gondii* and *Bartonella spp.* antibodies in cats from Pennsylvania. Journal of Parasitology. 2009; 95:578–580. [PubMed: 19061304]
- Dziewieczynski TL, Crovo JA. Shyness and boldness differences across contexts in juvenile three-spined stickleback *Gasterosteus aculeatus* from an anadromous population. Journal of Fish Biology. 2011; 79:776–788. [PubMed: 21884112]
- Ellis GB, Turek FW. Testosterone and photoperio interact to regulate locomotor activity in male hamsters. Hormones and Behavior. 1983; 17:66–75. [PubMed: 6862394]
- Ezenwa VO, Price SA, Altizer S, Vitone ND, Cook KC. Host traits and parasite species richness in even and odd-toed hoofed mammals, Artiodactyla and Perissodactyla. Oikos. 2006; 115:526–536.
- Falconer, DS.; Mackay, TFC. Introduction to Quantitative Genetics. New-York, USA: Longman; 1996.
- Fish, D. Population ecology of *Ixodes dammini*. In: Ginsberg, HS., editor. Ecology and Environmental Management of Lyme Disease. New Brunswick, NJ, USA: Rutgers University Press; 1993. p. 25-42.
- Folstad I, Karter AJ. Parasites, bright males, and the immunocompetence handicap. American Naturalist. 1992; 139:603–622.
- Gern L, Cadenas FM, Burri C. Influence of some climatic factors on *Ixodes ricinus* ticks studied along altitudinal gradients in two geographic regions in Switzerland. International Journal of Medical Microbiology. 2008; 298:55–59.
- Goodwin BJ, Ostfeld RS, Schaub EM. Spatiotemporal variation in a Lyme Disease host and vector: black-legged ticks on white-footed mice. Vector borne and Zoonotic Diseases. 2001; 1:129–138. [PubMed: 12653143]
- Grear DA, Perkins SE, Hudson PJ. Does elevated testosterone result in increased exposure and transmission of parasites? Ecology Letters. 2009; 12:528–537. [PubMed: 19392718]
- Hansen F, Jeltsch F, Tackmann K, Staubach C, Thulke HH. Processes leading to a spatial aggregation of *Echinococcus multilocularis* in its natural intermediate host *Microtus arvalis*. International Journal for Parasitology. 2004; 34:37–44. [PubMed: 14711588]
- Harrison A, Scantlebury M, Montgomery WI. Body mass and sex-biased parasitism in wood mice *Apodemus sylvaticus*. Oikos. 2010; 119:1099–1104.
- Hillegass MA, Waterman JM, Roth JD. The influence of sex and sociality on parasite loads in an African ground squirrel. Behavioral Ecology. 2008; 19:1006–1011.
- Hughes VL, Randolph SE. Testosterone depresses innate and acquired resistance to ticks in natural rodent hosts: A force for aggregated distributions of parasites. Journal of Parasitology. 2001; 87:49–54. [PubMed: 11227902]
- Jones CG, Ostfeld RS, Richard MP, Schaub EM, Wolff JO. Chain reactions linking acorns to gypsy moth outbreaks and Lyme disease risk. Science. 1998; 279:1023–1026. [PubMed: 9461433]
- Juran, JM.; Seder, LA.; Gryna, FM. The Quality Control Handbook. New-York, New-York, USA: McGraw-Hill; 1962.


- Kiffner C, Loedige C, Alings M, Vor T, Ruehe F. Body-mass or sex-biased tick parasitism in roe deer (*Capreolus capreolus*)? A GAMLSS approach. *Medical and Veterinary Entomology*. 2011a; 25:39–45. [PubMed: 21118286]
- Kiffner C, Vor T, Hagedorn P, Niedrig M, Ruehe F. Factors affecting patterns of tick parasitism on forest rodents in tick-borne encephalitis risk areas, Germany. *Parasitology Research*. 2011b; 108:323–335. [PubMed: 20878183]
- Krasnov BR, Morand S, Hawlena H, Khokhlova IS, Shenbrot GI. Sex-biased parasitism, seasonality and sexual size dimorphism in desert rodents. *Oecologia*. 2005; 146:209–217. [PubMed: 16025350]
- Krasnov BR, Stanko M, Morand S. Host community structure and infestation by ixodid ticks: repeatability, dilution effect and ecological specialization. *Oecologia*. 2007; 154:185–194. [PubMed: 17684769]
- Krasnov BR, Stanko M, Morand S. Competition, facilitation or mediation via host? Patterns of infestation of small European mammals by two taxa of haematophagous arthropods. *Ecological Entomology*. 2010; 35:37–44.
- Lessells CM, Boag PT. Unrepeatable repeatabilities – a common mistake. *Auk*. 1987; 104:116–121.
- Leung B. Aggregated parasite distributions on hosts in a homogeneous environment: examining the Poisson null model. *International Journal for Parasitology*. 1998; 28:1709–1712. [PubMed: 9846607]
- Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on disease emergence. *Nature, London*. 2005; 438:355–359. [PubMed: 16292310]
- Lynn SE, Houtman AM, Weathers WW, Ketterson ED, Nolan V. Testosterone increases activity but not daily energy expenditure in captive male dark-eyed juncos, *Junco hyemalis*. *Animal Behaviour*. 2000; 60:581–587. [PubMed: 11082228]
- Minerly AE, Russo SJ, Kemen LM, Nazarian A, Wu HBK, Weierstall KM, Akhavan A, Jenab S, Quinones-Jenab V. Testosterone plays a limited role in cocaine-induced conditioned place preference and locomotor activity in male rats. *Ethnicity & Disease*. 2008; 18:S2.
- Morand, S.; Krasnov, BR.; Poulin, R. *Micromammals and macroparasites*. Tokyo: Springer-Verlag; 2006.
- Ostfeld RS, Canham CD, Oggenfuss K, Winchcombe RJ, Keesing F. Climate, deer, rodents, and acorns as determinants of variation in Lyme-disease risk. *Plos Biology*. 2006; 4:1058–1068.
- Ostfeld RS, Miller MC, Schnurr J. Ear tagging increases tick (*Ixodes dammini*) infestation rates of white-footed mice (*Peromyscus leucopus*). *Journal of Mammalogy*. 1993; 74:651–655.
- Ostfeld RS, Hazler KR, Cepeda OM. Temporal and spatial dynamics of *Ixodes scapularis* (Acari: Ixodidae) in a rural landscape. *Journal of Medical Entomology*. 1996a; 33:90–95. [PubMed: 8906910]
- Ostfeld RS, Miller MC, Hazler KR. Causes and consequences of tick (*Ixodes scapularis*) burdens on white-footed mice (*Peromyscus leucopus*). *Journal of Mammalogy*. 1996b; 77:266–273.
- Perkins SE, Cattadori IM, Tagliapietra V, Rizzoli AP, Hudson PJ. Empirical evidence for key hosts in persistence of a tick-borne disease. *International Journal for Parasitology*. 2003; 33:909–917. [PubMed: 12906875]
- Poulin, R. *Evolutionary Ecology of Parasites*. Princeton, NJ, USA: Princeton University Press; 2007.
- R Development Core Team. Vienna, Austria: R Foundation for Statistical Computing; 2010. R: a language and environment for statistical computing. <http://www.R-project.org>. ISBN3-900051-07-0.
- Randolph SE. Patterns of distribution of tick *Ixodes trianguliceps* Birula on its hosts. *Journal of Animal Ecology*. 1975; 44:451–474.
- Randolph SE. Tick ecology: processes and patterns behind the epidemiological risk posed by ixodid ticks as vectors. *Parasitology*. 2004; 129:S37–S65. [PubMed: 15938504]
- Randolph SE, Miklisova D, Lysy J, Rogers DJ, Labuda M. Incidence from coincidence: patterns of tick infestations on rodents facilitate transmission of tick-borne encephalitis virus. *Parasitology*. 1999; 118:177–186. [PubMed: 10028532]

- Randolph SE, Steele GM. An experimental evaluation of conventional control measures against the sheep tick, *Ixodes ricinus* (L) (Acari, Ixodidae). 2. The dynamics of the tick-host interaction. *Bulletin of Entomological Research*. 1985; 75:501–518.
- Raty M, Kangas A. Localizing general models based on local indices of spatial association. *European Journal of Forest Research*. 2007; 126:279–289.
- Robinson SA, Forbes MR, Hebert CE. Parasitism, mercury contamination, and stable isotopes in fish-eating double-crested cormorants: no support for the co-ingestion hypothesis. *Canadian Journal of Zoology-Revue Canadienne de Zoologie*. 2009; 87:740–747.
- Rosenberg MS, Anderson CD. PASSaGE: Pattern Analysis, Spatial Statistics and Geographic Exegesis. Version 2. *Methods in Ecology and Evolution*. 2011; 2:229–232.
- Rowsemitt CN. Activity of castrated male voles- Rhythms of responses to testosterone replacement. *Physiology & Behavior*. 1989; 45:7–13. [PubMed: 2657818]
- Sanchez A, Devevey G, Bize P. Female-biased infection and transmission of the gastrointestinal nematode *Trichuris arvicolae* infecting the common vole. *International Journal for Parasitology*. 2011; 41:1397–1402. [PubMed: 22056297]
- Schmidt KA, Ostfeld RS. Biodiversity and the dilution effect in disease ecology. *Ecology*. 2001; 82:609–619.
- Schmidt KA, Ostfeld RS, Schaub EM. Infestation of *Peromyscus leucopus* and *Tamias striatus* by *Ixodes scapularis* (Acari: Ixodidae) in relation to the abundance of hosts and parasites. *Journal of Medical Entomology*. 1999; 36:749–757. [PubMed: 10593076]
- Seivwright LJ, Redpath SM, Mougeot F, Leckie F, Hudson PJ. Interactions between intrinsic and extrinsic mechanisms in a cyclic species: testosterone increases parasite infection in red grouse. *Proceedings of the Royal Society of London, B*. 2005; 272:2299–2304.
- Shaw DJ, Grenfell BT, Dobson AP. Patterns of macroparasite aggregation in wildlife host populations. *Parasitology*. 1998; 117:597–610. [PubMed: 9881385]
- Shaw MT, Keesing F, McGrail R, Ostfeld RS. Factors influencing the distribution of larval blacklegged ticks on rodent hosts. *American Journal of Tropical Medicine and Hygiene*. 2003; 68:447–452. [PubMed: 12875294]
- Siegel JP, Kitron U, Bouseman JK. Spatial and temporal distribution of *Ixodes dammini* (Acari, Ixodidae) in a northwestern Illinois state park. *Journal of Medical Entomology*. 1991; 28:101–104. [PubMed: 2033600]
- Sokal, RR.; Rohlf, FJ. *Biometry*. New York, USA: W. H. Freeman and company; 1995.
- Steiner FE, Pinger RR, Vann CN, Grindle N, Civitello D, Clay K, Fuqua C. Infection and co-infection rates of *Anaplasma phagocytophilum* variants, *Babesia* spp, *Borrelia burgdorferi*, and the rickettsial endosymbiont in *Ixodes scapularis* (Acari : Ixodidae) from sites in Indiana, Maine, Pennsylvania, and Wisconsin. *Journal of Medical Entomology*. 2008; 45:289–297. [PubMed: 18402145]
- Stafford KC. Oviposition and larval dispersal of *Ixodes dammini* (ACARI, Ixodidae). *Journal of Medical Entomology*. 1992; 29:129–132. [PubMed: 1552522]
- Wilson K, Grenfell BT, Shaw DJ. Analysis of aggregated parasite distributions: A comparison of methods. *Functional Ecology*. 1996; 10:592–601.
- Wilson ML, Spielman A. Seasonal activity of immature *Ixodes dammini* (ACARI, Ixodidae). *Journal of Medical Entomology*. 1985; 22:408–414. [PubMed: 4045938]
- Wilson ML. Distribution and abundance of *Ixodes scapularis* (Acari : Ixodidae) in North America: Ecological processes and spatial analysis. *Journal of Medical Entomology*. 1998; 35:446–457. [PubMed: 9701926]
- Wolff JO. The effect of density, food, and interspecific interference on home range size in *Peromyscus leucopus* and *Peromyscus maniculatus*. *Canadian Journal of Zoology-Revue Canadienne de Zoologie*. 1985; 63:2657–2662.
- Woolhouse MEJ, Dye C, Etard JF, Smith T, Charlwood JD, Garnett GP, Hagan P, Hii JLK, Ndhlovu PD, Quinnell RJ, Watts CH, Chandiwana SK, Anderson RM. Heterogeneities in the transmission of infectious agents: Implications for the design of control programs. *Proceedings of the National Academy of Sciences, USA*. 1997; 94:338–342.

- Yeagley TJ, Reichard MV, Hempstead JE, Allen KE, Parsons LM, White MA, Little SE, Meinkoth JH. Detection of *Babesia gibsoni* and the canine small *Babesia* 'Spanish isolate' in blood samples obtained from dogs confiscated from dogfighting operations. *Javma-Journal of the American Veterinary Medical Association*. 2009; 235:535–539.
- Zuk M, McKean KA. Sex differences in parasite infections: Patterns and processes. *International Journal for Parasitology*. 1996; 26:1009–1023. [PubMed: 8982783]




**Fig. 1.** Blacklegged ticks have highly seasonal activity patterns with densities varying between sites and between years. The activity period of both (A) larval and (B) nymphal ticks were strongly aggregated in time. The average densities of host-seeking ticks (closed triangles  $\pm$  S.E.) and the average burdens on mice (open circles  $\pm$  S.E.) vary between sites and between years. The densities of host-seeking nymphs (closed triangles  $\pm$  S.E.) and the average nymphal burdens on mice (open circles  $\pm$  S.E.) were drastically lower in 2010 than in 2009. The black bars denote the peak period of host-seeking ticks (*ca.* 80% of collected ticks) and the grey bars figure the peak period of attached ticks (*ca.* 80% of counted ticks).



**Fig. 2.**

The majority of ticks parasitized a minority of mice. Data are plotted as the cumulative frequency distribution of attached larvae (A) and attached nymphs (B) on mice that are ranked from lowest to highest burden. The black curve represents the model that best explains the distribution of tick burdens on mice from all trapping sessions surrounded by the 95% confidence interval for this model (dashed curves). The dotted line represents the cumulative distribution of ticks on mice assuming ticks are randomly distributed among mice. The grey lines demonstrate that half of the mice host only 16% of the larvae, while the remaining mice host 84% of attached larvae. Similarly, 80% of the mice host only 17% of all attached nymphs while the remaining 20% of the mice host 83% of the nymphs.



**Fig. 3.** Tick burdens are not spatially auto-correlated. No spatial auto-correlation in standardized larval burdens (A) nor in standardized nymph burdens (B) were detected among the trapping locations as none of the *Moran's I* values were significantly different from 0 ( $P > 0.12$ ).

**Table 1**  
The tick burdens observed on mice and in specific localities are consistent across repeated observations

(A) Larvae		Mouse						Locality					
Year, site	Source	DF	SS	F ratio	P	r	Source	DF	SS	F ratio	P	r	
All	Mouse	167	278.2	2.77	<0.001	0.36	Locality	183	244.7	1.81	<0.001	0.22	
	Error	362	217.5				Error	330	244.5				
2009, Monocassy	Mouse	12	9.7	0.99	0.47	0.00	Locality	16	13.4	1.32	0.25	0.11	
	Error	40	32.6				Error	28	17.7				
2009, Northside	Mouse	14	18.5	2.45	0.019	0.32	Locality	14	23.2	2.28	0.043	0.35	
	Error	31	16.8				Error	21	15.3				
2010, Monocassy	Mouse	48	75.5	2.48	<0.001	0.31	Locality	55	74.3	1.67	0.013	0.20	
	Error	110	68.9				Error	99	79.9				
2010, Northside	Mouse	90	175.1	3.55	<0.001	0.46	Locality	95	133.5	1.94	<0.001	0.25	
	Error	181	99.2				Error	182	131.7				

(B) Nymphs		Mouse						Locality					
Year, site	Source	DF	SS	F ratio	P	r	Source	DF	SS	F ratio	P	r	
All	Mouse	227	64.8	1.34	0.004	0.10	Locality	236	55.7	0.94	0.72	0.00	
	Error	510	108.6				Error	616	155.3				
2009, Monocassy	Mouse	21	12.7	1.71	0.050	0.15	Locality	22	11.7	0.85	0.66	0.00	
	Error	69	24.4				Error	59	36.9				
2009, Northside	Mouse	39	21.1	2.45	<0.001	0.27	Locality	45	13.7	1.03	0.43	0.01	
	Error	119	26.2				Error	139	41.0				
2010, Monocassy	Mouse	50	9.7	0.81	0.800	0.00	Locality	54	10.4	0.88	0.88	0.00	
	Error	89	21.5	Error			Error	100	21.9				
2010, Northside	Mouse	114	20.8	1.17	0.164	0.05	Locality	112	18.3	0.94	0.65	0.00	
	Error	233	36.5				Error	318	55.5				

(The larval burdens (A) and nymphal burden (B) of mice were significantly repeatable across multiple capture events such that mice with high tick burdens in one observation had high tick burdens at subsequent captures. These data indicate that ticks were aggregated on a consistent subset of mice throughout the larval and nymphal activity seasons.)

**Table 2**

Different mouse-specific traits are correlated with larval and nymphal tick burdens

**(A) Larval burden**

Source	DF Num	DF Den	F ratio	<i>P</i>
Sex	1	117.4	20.82	<0.001
Age	2	362.6	0.00	0.99
Body mass	1	261	0.14	0.71
Home range size	1	74.5	0.07	0.79

**(B) Nymphal burden**

Source	DF Num	DF Den	F ratio	<i>P</i>
Sex	1	93.4	2.54	0.11
Body mass	1	264.4	5.09	0.025
Home range size	1	185.4	0.99	0.32
Sex * Body mass	1	271.1	8.41	0.004
Sex * Home range size	1	189.6	5.30	0.022

(The tables present the result of the mixed-model ANOVAs of traits on (A) larval burden and (B) nymphal burden, corrected for repeated measures on mice with the degrees of freedom approximated by the Welch-Satterthwaite equation.)