

Chapter 4: Hepatitis B

Lyn Finelli, DrPH and Beth P. Bell, MD, MPH

I. Disease Description

Hepatitis B is caused by infection with the hepatitis B virus (HBV), a double-stranded DNA virus of the family hepadnaviridae. HBV replicates in the liver and causes both acute and chronic hepatitis. Although the highest concentrations of virus are found in blood, other body fluids, such as semen and saliva, also have been demonstrated to contain HBV. Thus, HBV is predominantly a blood and sexually transmitted infection and is transmitted by percutaneous and mucosal exposure to infectious body fluids.

The incubation period for acute hepatitis B ranges from 45 to 160 days (average 120 days). The clinical manifestations of acute HBV infection are age dependent. Infants, young children (younger than 10 years of age), and immunosuppressed adults with newly acquired HBV infection are usually asymptomatic.¹ Older children and adults are symptomatic in 30%–50% of infections. When present, clinical symptoms and signs might include anorexia, malaise, nausea, vomiting, abdominal pain, jaundice, dark urine, and clay-colored or light stools. Occasionally, extrahepatic manifestations occur and include skin rashes, arthralgias, and arthritis. Fulminant hepatitis occurs with a case-fatality rate of 0.5%–1%.

From 2003-2008, there was an estimated 38,000–73,000 new HBV infections annually in the U.S.² Each year about 600,000 HBV-related deaths occur worldwide.³ Among adults with normal immune status, most (94%–98%) recover completely from newly acquired HBV infections, eliminating virus from the blood and producing neutralizing antibody that creates immunity from future infection. In infants, young children, and immunosuppressed persons, most newly acquired HBV infections result in chronic infection.⁴ Infants are at greatest risk, with a 90% chance of developing chronic infection if infected at birth. Although the consequences of acute hepatitis B can be severe, most of the serious sequelae occur in persons in whom chronic infection developed. Persons who acquire chronic HBV infection as infants or young children are often asymptomatic; however, chronic liver disease develops in two-thirds of these persons, and approximately 15%–25% die prematurely from cirrhosis or liver cancer. Persons with chronic HBV infection are often detected in screening programs, such as those for blood donors, pregnant women, and refugees. Persons with chronic HBV infection are a major reservoir for transmission of HBV infections. Any person testing positive for hepatitis B surface antigen (HBsAg) is potentially infectious to both household and sexual contacts.

II. Background

Each year during the 1970s and 1980s, an estimated 200,000–300,000 persons worldwide were newly infected with HBV. Until recently, hepatitis B was one of the most frequently reported vaccine-preventable diseases in the United States, with 15,000–20,000 cases reported annually to the National Notifiable Diseases Surveillance System (NNDSS). Since 1985, a steady decline has occurred in the number of cases of acute hepatitis B reported to the NNDSS. In 2007, approximately 4,500 cases of acute hepatitis B were reported, which after correcting for underreporting and asymptomatic infections, represented an estimated 43,000 new cases.² Based on testing from the National Health and Nutrition Examination Survey, the age-adjusted prevalence of hepatitis B core antibody, which indicates past or present infection, in the US population decreased from 5.4% in 1988-1994 to 4.7% during 1999-2006 among persons aged 6 years and older. The age adjusted prevalence of hepatitis B core antibody and hepatitis B surface antigen together, which indicates chronic active HBV infection, decreased from 0.38% to 0.27% among persons aged 6 years and older during the same time periods.³ An estimated 730,000 US adults had chronic active hepatitis B during 1999-2006.³

The extent to which children acquire HBV infection in the United States has not been appreciated, primarily because most infections in this age group are asymptomatic. In the United States, approximately 24,000 HBsAg-positive women gave birth in 2005. Without postexposure prophylaxis to prevent perinatal HBV infection, it is estimated that 12,000 infants

and children would be infected with HBV annually. Furthermore, before the implementation of universal infant hepatitis B immunization, an additional 16,000 children younger than 10 years old were infected annually in the United States through exposure to HBsAg-positive household members or community contacts. Populations with the highest rates of these early childhood infections included Alaska Natives, children of Pacific Islander parents, and children of first-generation immigrants from countries where HBV is of high or intermediate endemicity.⁵⁻⁸

Screening of all pregnant women for HBsAg to identify infants requiring postexposure prophylaxis has been recommended since 1988, universal childhood hepatitis B immunization since 1991, universal adolescent hepatitis B immunization since 1995,^{9,10} and universal hepatitis B birth dose administration since 2005. In the United States, without postexposure prophylaxis, HBV would annually infect 12,000 infants; without routine childhood immunization, 16,000 children would be infected.

Among persons who reported risk behaviors/exposures in 2007, the most frequently reported risk behavior/exposure for acute, symptomatic hepatitis B was multiple sex partners (38%), followed by men who have sex with men (MSM) (11%) and sexual contact with a person known to have hepatitis B (6%). Injection drug use (IDU) was reported for 15% of persons.¹¹ Although up to 25% of persons with newly acquired hepatitis B do not report a source for their infection, many of these persons have had a past history of high-risk sex or drug behaviors. Furthermore, more than half of persons with newly acquired hepatitis B were previously seen in medical settings where hepatitis B vaccine is routinely recommended, such as sexually transmitted disease (STD) treatment clinics. Thus, programs to vaccinate older adolescents and adults at increased risk for HBV infection need to be strengthened nationwide in order to have a significant impact on reducing HBV transmission in the next 2 decades.

III. Importance of Rapid Identification

Rapid identification and prompt reporting of cases of acute hepatitis B is important because measures such as postexposure prophylaxis can be taken to prevent transmission to other persons. Although outbreaks of hepatitis B are unusual, rapid recognition allows for identification of the source and prevention of further transmission. In addition, identification of risk behaviors/exposures for infection provides a means to assess the effectiveness of hepatitis B immunization activities in the community and identify missed opportunities for immunization.

In most states, HBsAg positivity is a laboratory reportable condition. Reporting of HBsAg-positive persons facilitates timely immunization of contacts. For HBsAg-positive pregnant women, reporting allows for initiation of case management to ensure prevention of perinatal HBV transmission (see “Postexposure prophylaxis” below). In 2003, chronic HBV infection became nationally notifiable and is reportable by state health departments to the NNDSS. All states are encouraged to report chronic hepatitis B infection. States should develop registries of persons with HBsAg-positive laboratory results to facilitate postexposure prophylaxis of contacts and reporting to NNDSS (see “Registries/databases for HBsAg-positive persons” below).

Postexposure prophylaxis

Hepatitis B immune globulin (HBIG) is prepared from human plasma known to contain a high titer of antibody to HBsAg (anti-HBs). The plasma from which HBIG is prepared is screened for HBsAg, hepatitis C virus (HCV), and human immunodeficiency virus, and since 1999, all products available in the United States have been manufactured by methods that inactivate HCV and other viruses. A regimen combining HBIG and hepatitis B vaccine is 85%–95% effective in preventing HBV infection when administered at birth to infants born to HBsAg-positive mothers. Regimens involving either multiple doses of HBIG alone or the hepatitis B vaccine series alone are 70%–75% effective in preventing HBV infection. It is recommended that HBIG should be given as soon as possible, after exposure to hepatitis B. Although the postexposure efficacy of the combination of HBIG and the hepatitis B vaccine series has not been evaluated for occupational or sexual exposures, it can be presumed that the increased efficacy of this regimen observed in the perinatal setting compared with HBIG alone would apply to these exposures.

Postexposure prophylaxis with HBIG and hepatitis B vaccine should be given to infants born to HBsAg-positive mothers, unvaccinated infants whose mothers or primary caregivers have acute hepatitis B, sexual contacts of persons with hepatitis B, and healthcare workers after occupational exposure to HBsAg-positive blood depending on their vaccination and vaccine response status. Household and sexual contacts of persons with chronic HBV infection should be vaccinated.

IV. Importance of Surveillance

Disease surveillance is used to 1) identify contacts of case-patients who require postexposure prophylaxis; 2) detect outbreaks; 3) identify infected persons who need counseling and referral for medical management; 4) monitor disease incidence and prevalence; and 5) determine the epidemiologic characteristics of infected persons, including the source of their infection, to assess and reduce missed opportunities for vaccination.

V. Disease Reduction Goals

The primary goal of hepatitis B vaccination is to prevent chronic HBV infection. However, because such a high proportion of persons with chronic HBV infection are asymptomatic and the consequences are not seen for many years, monitoring the direct impact of prevention programs on the prevalence of chronic infection is difficult. Consequently, the disease reduction goals that have been established for hepatitis B are a combination of process and disease outcome measures. Because most HBV infections among children younger than 10 years of age are asymptomatic, programs targeting infants and children are best evaluated by measuring vaccination coverage and not by measuring reduction in acute infection. In older age groups, monitoring the incidence of acute disease as well as measuring vaccine coverage levels provides data useful for measuring the effectiveness of prevention programs.

Healthy People 2020 disease reduction goals have been established for achieving the prevention of HBV transmission in the United States. Disease reduction goals for infants and children include reducing the estimated number of chronic HBV infections in infants and young children to 400 cases and the number of new hepatitis B cases reported among persons 2–18 years of age to zero cases. *Healthy People 2020* vaccination goals for infants and children include setting a target coverage level for infants age 0–3 days receiving the initial birth dose of hepatitis B vaccine to 85% and for children age 19–35 months completing the three-dose hepatitis B vaccination series to 90%.¹²

Disease reduction goals for adults include reducing the rate of acute hepatitis B to 1.5 cases per 100,000 in persons age 19 years and older. Among adults in high-risk groups, disease reduction goals include reducing the number of cases of acute hepatitis B to 215 cases in injection-drug users and to 45 new infections among men who have sex with men.¹²

VI. Case Definition

The following case definitions for acute hepatitis B, chronic hepatitis B virus infection and perinatal HBV infection have been adopted by the Council of State and Territorial Epidemiologists.¹³

Hepatitis B, acute (effective 2000)

Clinical case definition

An acute illness with

- a. a discrete onset of symptoms, and
- b. jaundice or elevated serum aminotransferase levels.

Laboratory criteria for diagnosis

- IgM antibody to hepatitis B core antigen (anti-HBc) positive or hepatitis B surface antigen (HBsAg) positive
- IgM anti-HAV negative (if done)

Case classification

Confirmed: a case that meets the clinical case definition and is laboratory confirmed

Chronic hepatitis B virus infection (effective 2007)**Clinical description**

Persons with chronic HBV infection may have no evidence of liver disease or may have a spectrum of disease ranging from chronic hepatitis to cirrhosis or liver cancer. Persons with chronic infection may be asymptomatic.

Laboratory criteria for diagnosis

- IgM anti-HBc negative AND a positive result on one of the following tests: HBsAg, HBeAg, or HBV DNA **OR**
- HBsAg positive or HBV DNA positive or HBeAg positive two times at least 6 months apart (Any combination of these tests performed 6 months apart is acceptable.)

Case classification

Confirmed: a case that meets either laboratory criterion for diagnosis

Probable: a case with a single HBsAg-positive or HBV DNA-positive or HBeAg-positive laboratory result when no IgM anti-HBc results are available

Comment: Multiple laboratory tests indicative of chronic HBV infection may be performed simultaneously on the same patient specimen as part of a “hepatitis panel.” Testing performed in this manner may lead to seemingly discordant results, e.g., HBsAg negative AND HBV DNA positive. For purposes of this case definition, any positive result among the three laboratory tests mentioned above is acceptable, regardless of other testing results. Negative HBeAg results and HBV DNA levels below positive cutoff level do not confirm the absence of HBV infection.

Perinatal hepatitis B virus infection (effective 1995)**Clinical description**

Perinatal hepatitis B in the newborn may range from asymptomatic to fulminant hepatitis.

Laboratory criteria for diagnosis

- Hepatitis B surface antigen (HBsAg) positive

Case classification

HBsAg positivity in any infant aged >1–24 months who was born in the United States or in U.S. territories to an HBsAg-positive mother

Comment: Infants born to HBsAg-positive mothers should receive HBIG and the first dose of hepatitis B vaccine within 12 hours of birth, followed by the second and third doses of vaccine at 1 and 6 months of age, respectively, if receiving the single-antigen vaccine. Dosage and scheduling will be different if the infant received a combination vaccine (see Table 1). Postvaccination testing for HBsAg and anti-HBs (antibody to HBsAg) is recommended from 3 to 6 months following completion of the vaccine series. If HBIG and the initial dose of vaccine are delayed for more than 1 month after birth, testing for HBsAg may determine if the infant is already infected.

Table 1. Hepatitis B combination vaccine schedule for newborn infants born to HBsAg positive mothers

Single antigen* combination vaccine	
Dose	Age
1	Birth (< 12 hrs)
HBIG	Birth (< 12 hrs)
2	2 mos
3	4 mos
4	6 mos (Pediatrix) or 12-15 mos (Comvax)

* *Recombivax HB or Engerix-B should be used for the birth dose. Comvax and Pediatrix cannot be administered at birth or before age 6 weeks.*

Source: CDC. *A comprehensive strategy to eliminate transmission of hepatitis B virus infection in the United States: Recommendations of the Advisory Committee on Immunization Practices (ACIP); Part 1: Immunization of infants, children and adolescents. MMWR 2005; 54(No. RR-16).*

VII. Laboratory Testing

Several specific and well-defined antigen–antibody systems are associated with HBV infection, including hepatitis B surface antigen (HBsAg) and antibody to HBsAg (anti-HBs); hepatitis B core antigen (HBcAg) and antibody to HBcAg (anti-HBc); and hepatitis B e antigen (HBeAg) and antibody to HBeAg (anti-HBe). Serologic assays are commercially available for all of these except HBcAg because no free HBcAg circulates in blood. One or more of these serologic markers are present in the blood during different phases of HBV infection (Table 2). Subtyping of HBsAg has occasionally been used to investigate outbreaks of hepatitis B, but this procedure is not routinely available in commercial laboratories.

Table 2. Interpretation of serologic test results for hepatitis B virus infection

Serologic Markers				Interpretation
HBsAg* Serologic Marker	Total Anti-HBc† Serologic Marker	IgM Anti-HBc§ Serologic Marker	Anti-HBs¶ Serologic Marker	
-	-	-	-	Susceptible, never infected
+	-	-	-	Early acute infection, transient (up to 18 days) after vaccination**
+	+	+	-	Acute infection
-	+	+	-	Acute resolving infection
-	+	-	+	Past infection, recovered and immune
+	+	-	-	Chronic infection
-	+	-	-	False positive (i.e., susceptible), past infection, or “low level” chronic infection
-	-	-	+	Immune (vaccination, prior infection) if titer is >10 mIU/mL

* Hepatitis B surface antigen

† Antibody to hepatitis B core antigen

§ Immunoglobulin M

¶ Antibody to hepatitis B surface antigen

** Transient HBsAg positivity (lasting <18 days) might be detected in some patients after vaccination.

The presence of HBsAg is an indication of ongoing HBV infection and the potential to spread the infection. In persons newly infected, HBsAg is present in serum 30–60 days after exposure to HBV and persists for variable periods. Anti-HBc develops in HBV infections, appearing at onset of symptoms or in liver test abnormalities in acute HBV infection, rising rapidly to high

levels, and persisting for life. The presence of the immunoglobulin M (IgM) class of anti-HBc, which persists for approximately 6 months, distinguishes acute or recently acquired infection. However, among infected infants, passively transferred maternal anti-HBc may persist beyond the age of 12 months, and IgM anti-HBc may not be present in newly infected children younger than 2 years of age, especially if they acquired their infection through perinatal transmission.

In persons who recover from HBV infection, HBsAg is eliminated from the blood system in 2–3 months, and anti-HBs develops during convalescence. The presence of anti-HBs indicates immunity from HBV infection. After recovery from natural infection, most persons will be positive for both anti-HBs and anti-HBc, whereas only anti-HBs develops in persons who are successfully vaccinated against hepatitis B. Additionally, anti-HBs can be present in persons who have received HBIG. Persons who do not recover from HBV infection and become chronically infected remain positive for HBsAg (and anti-HBc), although a small proportion (0.3% per year) eventually clear HBsAg and might develop anti-HBs.

Special laboratory studies

Molecular virologic methods such as polymerase chain reaction (PCR)-based assays are often used to amplify and sequence viral genomes. In conjunction with epidemiologic studies, these assays may be helpful for investigating common-source outbreaks of hepatitis B. In addition, these assays are essential for detecting the emergence of potential vaccine-resistant strains. Healthcare professionals with questions about molecular virologic methods or those who identify HBsAg-positive events among vaccinated persons should consult with their state health department or the Epidemiology and Surveillance Branch, Division of Viral Hepatitis, CDC, 404-718-8500.

For additional information on laboratory support for surveillance of vaccine-preventable diseases, see Chapter 22.

VIII. Reporting

In the United States, case reports of viral hepatitis are classified as hepatitis A, acute hepatitis B, acute hepatitis C, perinatal HBV infection, chronic HBV infection and hepatitis C, past or present. Serologic testing is necessary to determine the etiology of viral hepatitis, and case reports should be based on laboratory confirmation (see Section VII). Each state and territory has a list of reportable diseases and conditions of public health importance.¹⁴ These diseases and conditions that are to be reported and describe those persons or groups who are responsible for reporting, such as healthcare providers, hospitals, laboratories, and other institutions. Persons reporting these conditions should contact their state health department for state-specific reporting requirements.

Reporting to CDC

Case reports of acute hepatitis B, chronic HBV infection, perinatal HBV infection, and other reportable diseases are transmitted by the state health department weekly to CDC via the National Electronic Telecommunications System for Surveillance (NETSS). The NETSS core data elements include basic information (excluding personal identifiers)—age, race/ethnicity, sex, date of onset, date of report, county of residence. The NETSS extended data elements include clinical data, laboratory results, and exposure history. However, completeness of reporting of these extended data elements is often incomplete. In 2007, 52% of acute, symptomatic hepatitis B cases were reported in the United States with unavailable risk exposure/behavior information.¹¹ The Division of Viral Hepatitis has developed an extended Data Collection Worksheet to collect information about symptoms, risk behavior/exposures and serologic data (Appendix 6). This worksheet can be used for case investigation and data can be directly entered into the state's electronic reporting system. Sites reporting through the National Electronic Diseases Surveillance System (NEDSS)¹⁵ NEDSS-compatible, and Emerging Infections Program¹⁶ (Enhanced Hepatitis Surveillance) infrastructure collect data elements using this worksheet.

IX. Vaccination

Infants born to HBsAg-positive women should be administered by intramuscular injection hepatitis B immune globulin (HBIG; for hepatitis B postexposure prophylaxis) and the first dose of hepatitis B vaccine within 12 hours of birth. This combination also should be administered as soon as possible to unvaccinated infants whose primary caregivers have acute hepatitis B, unvaccinated healthcare personnel after occupational exposure, and sex partners of persons with acute hepatitis B. Studies are limited on the maximum interval after exposure during which postexposure prophylaxis is effective, but the interval is unlikely to exceed 7 days for percutaneous exposures and 14 days for sexual exposures. Exact treatment varies depending on source status and vaccination status. For infants born to women positive for HBsAg, the dose of HBIG is 0.5 mL. For all other applications, the dose is 0.06 mL/kg.¹⁷

Hepatitis B vaccine

Two single-antigen recombinant hepatitis B vaccines are commercially available, Recombivax HB® (Merck & Company, Inc.) and Engerix-B® (GlaxoSmithKline). Recombivax HB contains 5–40 µg of HBsAg protein per milliliter, depending on the formulation, whereas Engerix-B contains 20 µg/mL. Both vaccines are licensed for persons of all ages (Table 3).

Table 3. Recommended doses of currently licensed single-antigen hepatitis B vaccines

Group	Recombivax HB*		Engerix-B*	
	Dose (µg)	Volume (mL)	Dose (µg)	Volume (mL)
Infants, children and adolescents younger than 20 years of age	5	(0.5)		(0.5)
Adolescents 11–15 years† (adult formulation administered on 2-dose schedule)	10	(1.0)		
Adults 20 years of age or older	10	(1.0)	20	(1.0)
Hemodialysis patients and other immunocompromised persons	40	(1.0)§	40	(2.0)¶

* Both vaccines are routinely administered in three-dose series. Engerix-B also has been licensed for a four-dose series administered at 0, 1, 2, and 12 months.

† Two-dose schedule for adolescents using adult dose of Recombivax HB has been approved by ACIP, administered at 0, 4–6 months.

§ Special formulation administered on a 3-dose schedule at 0, 1, and 6 months.

¶ Two 1.0-mL doses administered at one site in a four-dose schedule at 0, 1, 2, and 6 months.

Twinrix® (GlaxoSmithKline), a combination of hepatitis A and B vaccine, is also available for use in persons aged 18 years and older. Twinrix consists of the antigenic components used in HAVRIX® (hepatitis A vaccine) and Engerix-B. In addition, there are two combination vaccines (Comvax® [Merck] and Pediarix® [GlaxoSmithKline]) that are used for vaccination of infants and young children. Comvax contains recombinant HBsAg and Haemophilus influenzae type b (Hib) polyribosylribitol phosphate conjugated vaccine. Pediarix contains recombinant HBsAg, diphtheria and tetanus toxoids and acellular pertussis adsorbed (DTaP), and inactivated poliovirus (IPV). However, these combination vaccines may not be administered to infants younger than 6 weeks of age (with exception of Twinrix®); only single-antigen hepatitis B vaccine may be used for the birth dose. Administration of four-dose hepatitis B vaccine schedules, including schedules with a birth dose followed by a combination vaccine series, is permissible (Table 4).

Table 4. Recommended doses of currently licensed combination hepatitis B vaccines*

Group	COMVAX		PEDIARIX		TWINRIX†	
	Dose (µg) ^{§, ¶}	Volume (mL)	Dose (µg) ^{§, **}	Volume (mL)	Dose (µg) ^{§, ††}	Volume (mL)
Infant (less than 1 year)— Mother HBsAg negative	5	0.5	10	0.5	NA	NA
Infant— Mother HBsAg positive	5	0.5	10	0.5	NA	NA
Children— (1–10 years)	5 ^{§§}	0.5	10	0.5	NA	NA
Adolescents 11–17 years	NA	NA	NA	NA	NA	NA
Adults— 18 years or older	NA	NA	NA	NA	20	1.0

* Hepatitis B vaccines are administered by intramuscular injection and may be given at the same time as other vaccines. Single-antigen vaccines may be administered with HBIG, but in a separate injection site.

† For persons ≥18 years of age at increased risk of both hepatitis B virus and hepatitis A virus infection

§ Recombinant HBsAg protein concentration

¶ Comvax also contains combined hepatitis B-Haemophilus influenzae type B conjugate vaccine. This vaccine cannot be administered at birth, before age 6 weeks, or after age 71 months.

** Pediarix also contains combined hepatitis B-25 Lf diphtheria toxoid, 10 Lf tetanus toxoid, 25 µg inactivated pertussis toxin, 25 µg filamentous hemagglutinin, 8 µg pertactin, 40 D-Wantigen Units (DU) Type 1 poliovirus, 8 DU Type 2 poliovirus, and 32 DU Type 3 poliovirus (cannot be administered at ages >7 years.

†† Twinrix also contains 720 ELISA Units (EL.U) inactivated hepatitis A virus.

§§ Maximum age at administration is 71 months.

An infant born to a HBsAg-positive woman who has not received HBIG and the first dose of hepatitis B vaccine by 12 hours of age or who has not received the third dose of hepatitis B vaccine by the age of 6 months is not adequately vaccinated.¹⁰ Infants born to HBsAg-positive mothers should be tested for HBsAg and antibody to HBsAg after completion of three or more doses in a licensed hepatitis B vaccination series, at age 9–18 months (generally at the next well-child visit). The testing should be done 1–2 months after the most recent hepatitis B vaccine dose to avoid a positive HBsAg result due to vaccine. Serologic testing can determine whether these infants are infected or have developed a protective antibody response after vaccination. Infants who do not respond to the primary vaccination series should be given three additional doses of hepatitis B vaccine on a 0, 1–2, 4–6-month schedule and re-tested 1–2 months after the final dose of vaccine.

The vaccination schedule for infants born to HBsAg-negative women includes three doses of vaccine in the first 18 months of life. The first dose should be given at birth with a minimum interval between doses 1 and 2 of 4 weeks; and a minimum interval of 8 weeks between doses 2 and 3.¹⁰ Dose 3 of hepatitis B vaccine should not be given before 24 weeks of age (164 days). Any infant born to an HBsAg-negative woman who has not received the third dose of hepatitis B vaccine by the age of 19 months is considered not up-to-date (Table 5).

Table 5. Hepatitis B vaccine schedules for newborn infants, by maternal hepatitis B surface antigen (HBsAg) status*

Maternal HBsAg Status	Single-Antigen Vaccine	Single-Antigen + Combination Vaccine
Positive	Dose: 1[†] Age: Birth (12 hours or younger)	Dose: 1[†] Age: Birth (12 hours or younger)
Positive	Dose: HBIG[§] Age: Birth (12 hours or younger)	Dose: HBIG[§] Age: Birth (12 hours or younger)
Positive	Dose: 2 Age: 1-2 months	Dose: 2 Age: 2 months
Positive	Dose: 3[¶] Age: 6 months	Dose: 3 Age: 4 months
Positive		Dose: 4[¶] Age: 6 months (Pediarix) or 12-15 months (Comvax)

Table 5. Hepatitis B vaccine schedules for newborn infants, by maternal hepatitis B surface antigen (HBsAg) status*

Maternal HBsAg Status	Single-Antigen Vaccine	Single-Antigen + Combination Vaccine
Unknown**	Dose: 1[†] Age: Birth (less than or equal 12 hours)	Dose: 1[†] Age: Birth (less than or equal 12 hours)
Unknown**	Dose: 2 Age: 1-2 months	Dose: 2 Age: 2 months
Unknown**	Dose: 3[¶] Age: 1-2 months	Dose: 3 Age: 4 months
Unknown**		Dose: 4[¶] Age: 6 months (Pediatrix) or 12-15 months (Comvax)
Negative	Dose: 1^{†, ††} Age: Birth (before discharge)	Dose: 1^{†, ††} Age: Birth (before discharge)
Negative	Dose: 2 Age: 1-2 months	Dose: 2 Age: 2 months
Negative	Dose: 3[¶] Age: 6 months	Dose: 3 Age: 4 months
Negative		Dose: 4[¶] Age: 6 months (Pediatrix) or 12-15 months (Comvax)

* Centers for Disease Control and Prevention. A comprehensive strategy to eliminate transmission of hepatitis B virus infection in the United States: Recommendations of the Advisory Committee on Immunization Practices (ACIP); Part 1: Immunization of Infants, Children and Adolescents. *MMWR* 2005;54(No. RR-16) p.9.

† Recombivax HB or Engerix-B should be used for the birth dose. Comvax and Pediatrix cannot be administered at birth or before age 6 weeks.

§ Hepatitis B immune globulin (0.5 mL) administered intramuscularly in a separate site from vaccine.

¶ The final dose in the vaccine series should not be administered before age 24 weeks (164 days).

** Mothers should have blood drawn and tested for HBsAg as soon as possible after admission for delivery; if the mother is found to be HBsAg positive, the infant should receive HBIG as soon as possible but no later than 7 days of age.

†† On a case-by-case basis and only in rare circumstances, the first dose may be delayed until after hospital discharge for an infant who weighs $\geq 2,000$ g and whose mother is HBsAg negative. When such a decision is made, a physician's order to withhold the birth dose and a copy of the original laboratory report indicating that the mother was HBsAg negative during this pregnancy should be placed in the infant's medical record.

Vaccination of pre-term infants should be delayed until they are 1 month old or at time of discharged from hospital, except for infants born to HBsAg-positive women and infants born to women with unknown HBsAg status. Infants born to HBsAg-positive women or women with unknown HBsAg status should be immunized with HBIG within 12 hours of birth regardless of birth weight.

Children and adolescents

Vaccination is routinely given as three-dose series at 0, 1, and 6 months. Acceptable alternative schedules include 0, 1, 2, 12 months and 0, 2, 4 months.

Adolescents 11–15 years of age

An alternative two-dose vaccination schedule has been developed for use in adolescents. The adult dose of Recombivax HB is administered to the adolescent, with the second dose given 4–6 months or 0, 1, 2, and 12 months after the first dose.

Adults (20 years of age or older)

Routinely given as three-dose series at 0, 1, and 6 months. Acceptable alternative schedules are 0, 1, 4 months and 0, 2, 4 months, and 0, 1, 2, and 12 months.

Dialysis patients and other immunocompromised persons

Either given as a three-dose series (0, 1, 6 months) or four-dose series (0, 1, 2, and 6 months), depending on formulation. Larger vaccine doses may be required to induce protective antibody levels in other immunocompromised persons (e.g., those taking immunosuppressive drugs, HIV infected), although few data are available concerning response to higher doses of vaccine in these patients and no data exist for children.

Combined hepatitis A and B vaccine

Primary vaccination of persons aged 18 years and older consists of three doses, administered on a 0, 1, and 6-month schedule.

X. Enhancing Surveillance

Establishing surveillance for acute hepatitis is difficult for several reasons. Five different viruses (A–E) cause viral hepatitis, and the clinical manifestations of the different types of acute hepatitis are similar. Infection with HBV, HCV and HDV can result in both acute and chronic infection. Therefore, serologic testing is necessary to establish an etiologic diagnosis for persons with symptoms of acute hepatitis and to evaluate case reports of persons who are reported with viral hepatitis. However, a lack of understanding about the epidemiology of these diseases and underutilization of serologic testing could result in significant misclassification in reporting of acute viral hepatitis.

Provider education

Providers should be educated about the importance of performing appropriate serologic tests to determine the etiology of viral hepatitis and reporting all cases of acute hepatitis B, chronic hepatitis B, and perinatal HBV. Case investigations of infected persons provide the best opportunity for postexposure prophylaxis of contacts and for reducing transmission.

Case investigation

Case investigation is essential for determining contacts who are eligible for prophylaxis and for collection of risk factor data. Analysis of risk factor data can identify populations where targeted interventions may be needed.

Laboratory reporting

Laboratories should be encouraged to report all persons with serologic markers of acute or chronic hepatitis to the state or local health department. All IgM anti-HBc, and HBsAg positive results should be reported. To facilitate reporting, these laboratory results could be included in the state's list of laboratory-reportable conditions.

Monitoring surveillance indicators

Regular monitoring of surveillance indicators, including date of report, timeliness, and completeness of reporting, may identify specific areas of the surveillance and reporting system that need improvement. Important program indicators that can be monitored through the surveillance, reporting and case investigation system include the following:

- Characteristics of cases of acute hepatitis B that occur in children and adolescents younger than 20 years of age and missed opportunities for vaccination.
- Characteristics of cases of acute hepatitis in which death has occurred.
- Characteristics of cases of acute hepatitis B in persons reporting a history of vaccination.
- Characteristics of cases of acute hepatitis B in persons over 70 years of age.
- Characteristics of cases of acute hepatitis B associated with healthcare transmission.
- Characteristics of cases of perinatal hepatitis B.

Registries/databases for HBsAg-positive persons

Reporting of HBsAg-positive test results and establishment of databases/registries for HBsAg-positive persons is encouraged. When any type of database is established, the confidentiality of individual identifying information needs to be ensured according to applicable laws and regulations. Computerized databases of persons with HBsAg-positive results can be used to:

- Distinguish newly reported cases of infection from previously identified cases and facilitate reporting of chronic hepatitis B;
- Facilitate case investigation and follow-up of persons with chronic HBV infection;
- Provide local, state, and national estimates of the proportion of persons with chronic HBV infection who have been identified.

Hospital-based reporting

Hospitals and infection control practitioners should be encouraged to report all persons with acute viral hepatitis (ICD-10 code B16), and all births to HBsAg-positive women.

XI. Case Investigation

Guidelines for investigating a suspected case of acute viral hepatitis include

1. determining a discrete onset of illness,
2. confirming evidence of acute liver disease (jaundice or elevated aminotransferase levels), and
3. obtaining serologic laboratory results.

The minimum recommended elements for investigating cases of chronic HBV infection and perinatal HBV infection include obtaining the serologic laboratory results needed to establish the case. Further investigation to determine the clinical characteristics of these cases may also be considered although it is not required to confirm the case.

Information to collect for acute hepatitis B infection

The following information is epidemiologically important to collect in a case investigation for acute hepatitis B.^{18, 19} Additional information may also be collected at the direction of the state health department.

- Demographic information
 - Clinical details
 - Date of illness onset
- Symptoms, including jaundice
- Laboratory results
- Vaccination status
- Risk behaviors/exposures
- Contact investigation and prophylaxis

Information to collect for chronic hepatitis B infection

The following information is epidemiologically important to collect in a case investigation for chronic hepatitis B virus infection. Additional information may also be collected at the direction of the state health department.¹⁹

- Demographic information
- Laboratory results
- Risk behaviors/exposures
- Pregnancy status. All HBsAg-positive pregnant women should be reported to the perinatal hepatitis B program manager so that they can be tracked and their infants can receive appropriate case management

The recommended elements of case investigation and follow-up of persons with chronic hepatitis B virus infection are detailed elsewhere.²⁰ The following should be included:

- Contact investigation and prophylaxis: Provision of hepatitis B vaccination for sexual, household, and other (needle-sharing) contacts of persons with hepatitis B, and counseling to prevent transmission to others
- Counseling and referral for medical management, including
 - assessing for biochemical evidence of chronic liver disease, and
 - evaluating eligibility for antiviral treatment.¹⁸

Information to collect for perinatal HBV infection

The following information is epidemiologically important to collect in a case investigation for perinatal HBV infection:

- Demographic information about the child and mother
- Laboratory results
- Immunization history of the child, including date and doses of HBIG and hepatitis B vaccine

Case investigation and follow-up of infants with hepatitis B virus infection should include the following:

- Referral for medical management, including
 - Assessing for biochemical evidence of chronic liver disease, and
 - Evaluating eligibility for antiviral treatment
- Identification of other susceptible infants and children in the household who require vaccination

References

1. McMahon BJ, Alward WL, Hall DB, et al. Acute hepatitis B virus infection: relation of age to the clinical expression of disease and subsequent development of the carrier state. *J Infect Dis* 1985; 151:599-603.
2. CDC. Division of Viral Hepatitis. Disease Burden from Viral Hepatitis A, B, and C in the United States. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. (Accessed February 10, 2011 at http://www.cdc.gov/hepatitis/PDFs/disease_burden.pdf).
3. Wasley A, Kruszon-Moran D, Kuhnert W, et al. The prevalence of hepatitis B virus infection in the United States in the era of vaccination. *J Infect Dis* 2010; 202(2):192-201.
4. Edmunds WJ, Medley GF, Nokes DJ, Hall AJ, Whittle HC. The influence of age on the development of the hepatitis B carrier state. *Lancet* 1993; 337:197-201.
5. Franks AL, Berg CJ, Kane MA, Browne BB, Sikes RK, Elsea WR, et al. Hepatitis B infection among children born in the United States to southeast Asian refugees. *N Engl J Med* 1989; 321:1301-5.
6. Hurie MB, Mast EE, Davis JP. Horizontal transmission of hepatitis B virus infection to United States-born children of Hmong refugees. *Pediatrics* 1992; 89:269-73.
7. Mahoney F, Lawrence M, Scott C, Le Q, Lambert S, Farley TA. Continuing risk for hepatitis B virus transmission among infants in Louisiana. *Pediatrics* 1995; 96:1113-6.
8. Pon EW, Ren H, Margolis H, Zhao Z, Schatz GC, Diwan A. Hepatitis B virus infection in Honolulu students. *Pediatrics* 1993; 92:574-8.
9. CDC. Hepatitis B virus: A comprehensive strategy for eliminating transmission in the United States through universal childhood vaccination. Recommendations of the Immunization Practices Advisory Committee *MMWR* 1991; 40(No. RR-13):1-25.
10. CDC. A comprehensive strategy to eliminate transmission of hepatitis B virus infection in the United States: Recommendations of the Advisory Committee on Immunization Practices (ACIP); Part 1: Immunization of infants, children and adolescents. *MMWR* 2005; 54(No. RR-16).

11. CDC. Surveillance for acute viral hepatitis – United States, 2007. *MMWR* 2009; 58 (No. SS-3).
12. Healthy People 2020. Healthy People 2020 Objectives. (Accessed March 9, 2011 at <http://www.healthypeople.gov/2020/topicsobjectives2020/default.aspx>).
13. CDC. Case definitions for infectious conditions under public health surveillance. Available at http://www.cdc.gov/osels/ph_surveillance/nndss/casedef/index.htm).
14. Council for State and Territorial Epidemiologists. State Reportable Conditions Assessment (2010), (Accessed March 10, 2011 at <http://www.cste.org/dnn/ProgramsandActivities/PublicHealthInformatics/StateReportableConditionsQueryResults/tabid/261/Default.aspx>).
15. CDC. National Electronic Disease Surveillance System (Accessed February 7, 2011 at http://www.cdc.gov/phin/library/phin_fact_sheets/111759_NEDSS.pdf).
16. Centers for Disease Control and Prevention. Emerging Infections Programs (Accessed February 7, 2011 at <http://www.cdc.gov/ncpdcid/deiss/eip/index.html>).
17. Dienstag JL. Hepatitis B virus infection. *N Engl J Med*. 2008; 359:1486-1500.
18. CDC. Division of Viral Hepatitis. Guidelines for viral hepatitis surveillance and case management. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, July 2005.
19. Sorrell MF, Belongia EA, Costa J, Gareen IF, Grem JL, Inadomi JM, Kern ER, McHugh JA, Petersen GM, Rein MF, Strader DB, Trotter HT. National Institutes of Health Consensus Development Conference Statement: Management of hepatitis B. *Ann Intern Med*. 2009;150(2):104-10.
20. CDC. A comprehensive strategy to eliminate transmission of hepatitis B virus infection in the United States: Recommendations of the Advisory Committee on Immunization Practices (ACIP); Part 2: Immunization of adults. *MMWR* 2006; 55(No. RR-16).