

MORBIDITY AND MORTALITY WEEKLY REPORT

September 28, 1990/Vol. 39/No. 38

- 653 Health Benefits of Smoking Cessation
- 653 Smokers' Beliefs about Smoking Cessation
- 656 Smoking-Related Mortality Decline among Physicians – Rhode Island
- 659 Smoking among Reproductive-Aged Women – Idaho, New York
- 662 Effects of Maternal Cigarette Smoking Ohio
- 665 Cigarette Brand Use U.S.
- **673** Employee Smoking Policy Colorado **676** Smoking Cessation U.S.
- 690 Smoking Cessation U.S
- 680 Smoking-Attributable Mortality Kentucky

Health Benefits of Smoking Cessation

This issue of MMWR focuses on the adverse effects of tobacco use on the public's health and on efforts to reduce and prevent those effects.

On September 25, 1990, The Health Benefits of Smoking Cessation: A Report of the Surgeon General, 1990, was released. The major conclusions of the report are: 1) smoking cessation has major and immediate health benefits for persons of all ages and provides benefits for persons with and without smoking-related disease; 2) former smokers live longer than continuing smokers; 3) smoking cessation decreases the risk for lung and other cancers, heart attack, stroke, and chronic lung disease; 4) women who stop smoking before pregnancy or during the first 3 to 4 months of pregnancy reduce their risk for having a lowbirth-weight infant to that of women who never smoked; and 5) the health benefits of smoking cessation substantially exceed any risks from the average 5-lb (2.3-kg) weight gain or any adverse psychologic effects that may follow quitting.

An executive summary of the report will be published as an MMWR Recommendations and Reports on October 5, 1990. Additional information is available from the Public Information Branch, Office on Smoking and Health, Center for Chronic Disease Prevention and Health Promotion, CDC, at (301) 443-5287.

Perspectives in Disease Prevention and Health Promotion

Smokers' Beliefs about the Health Benefits of Smoking Cessation – 20 U.S. Communities, 1989

The health risks associated with smoking and the reduction in risk associated with smoking cessation are well documented (1,2). Although public knowledge of the health hazards of smoking is high and has increased steadily since the 1950s (1), data are limited regarding public knowledge of the health benefits of smoking cessation. This report presents data on smokers' beliefs about their chances of avoiding disease by quitting smoking.

Smokers' Beliefs - Continued

Data were obtained from a telephone survey conducted from January through April 1989 of a random sample of 4351 smokers aged 25–64 years. The survey was conducted in 20 communities* in the United States as part of the National Cancer Institute's Community Intervention Trial for Smoking Cessation (3). Interviews were completed with 3669 (84%) eligible smokers regarding their knowledge, attitudes, and behavior relevant to cigarette smoking. For this report, responses to two items were analyzed: 1) "How likely do you think it is that you will avoid or decrease serious health problems from smoking if you quit?" (four response choices ranged from "very likely" to "very unlikely"); and 2) "If a person has smoked for more than 20 years, there is little health benefit to quitting" (four response choices ranged from "strongly agree" to "strongly disagree"). Responses were examined in relation to sex, age, level of education (high school graduate or less vs. some college or more), and daily cigarette consumption (<25 or \geq 25 cigarettes per day).

Overall, 83% of smokers responded that it was "very likely" or "likely" that by quitting they would avoid or decrease serious health problems from smoking. Eighty-five percent of smokers disagreed that little health benefit exists from quitting for a person who has smoked >20 years. For both items, beliefs about the benefits of quitting varied by age and education but not by sex. Within each age group, respondents who had attended college were more likely to both perceive benefits and disagree that there is little benefit from quitting than were those who had not (p<0.05, chi-square test) (Figure 1); this difference increased with age. For smokers with no college education, 87% of those aged 25–34 years and 67% of those aged 55–64 years believed they would avoid or decrease serious health problems by quitting (p<0.05). For college-educated smokers, age group differences did not vary significantly (Figure 1).

Reported by: KM Cummings, PhD, R Sciandra, Dept of Cancer Control and Epidemiology, Roswell Park Cancer Institute, Buffalo, New York, and TF Pechacek, PhD, WR Lynn, National Cancer Institute, National Institutes of Health, for the Community Intervention Trial for Smoking Cessation Research Group. Office on Smoking and Health, Center for Chronic Disease Prevention and Health Promotion, CDC.

Editorial Note: Former smokers most frequently cite concern about health as the reason for quitting smoking (4). Although most of the public is aware of the health risks associated with smoking and the health benefits of smoking cessation, smokers tend to be less aware of these risks and benefits, and sizable gaps in public knowledge persist in certain sociodemographic groups.

Educational level appears to be the best sociodemographic predictor of smoking behavior. Cessation rates are higher for college-educated than for noncollege-educated groups, a disparity that appears to be increasing (1,5). Educational status may be linked to attitudes and values that predispose a person to accept or reject warnings about tobacco use and may reflect exposure to antismoking messages (6). Future antismoking campaigns need to be more sensitive to educational status when defining messages and selecting communication channels.

Knowledge of the benefits of smoking cessation was lowest in smokers aged 55–64 years who had no college education. Thus, greater attention must be directed at informing this group about the health benefits of quitting smoking.

^{*}Bellingham and Longview/Kelso, Washington; Albany/Corvallis and Medford/Ashland, Oregon; Vallejo and Hayward, California; Santa Fe and Las Cruces, New Mexico; Cedar Rapids and Davenport, Iowa; Raleigh and Greensboro, North Carolina; Paterson and Trenton, New Jersey; Yonkers, New Rochelle, Utica, and Binghamton/Johnson City, New York; and Lowell and Fitchburg/Leominster, Massachusetts.

Smokers' Beliefs – Continued

CDC's Office on Smoking and Health, Center for Chronic Disease Prevention and Health Promotion (CCDPHP), is initiating a public information campaign on the health benefits of smoking cessation for older Americans based on the theme "It's never to late to quit smoking." The program is being conducted in collaboration with the National Institutes of Health, the Administration on Aging, the Department of Veterans Affairs, the Office of Disease Prevention and Health Promotion, the American Association of Retired Persons, and the Fox Chase Cancer Center. Information on this campaign and print materials are available from the Office on Smoking and Health, CCDPHP, CDC, 5600 Fishers Lane, Rockville, MD 20857; telephone (301) 443-5287.

References

- 1. CDC. Reducing the health consequences of smoking: 25 years of progress—a report of the Surgeon General, 1989. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1989; DHHS publication no. (CDC)89-8411.
- CDC. The health benefits of smoking cessation: a report of the Surgeon General, 1990. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1990; DHHS publication no. (CDC)90-8416.
- 3. Pechacek TF. A randomized trial for smoking cessation. In: Smoking and health, 1987-

FIGURE 1. Percentage of smokers who reported* that quitting reduces their risk for disease, by age and education level – 20 U.S. communities[†]



*Question number 1: Percentage who responded "very likely" or "likely" to the question "How likely do you think it is that you will avoid or decrease serious health problems from smoking if you quit?"

Question number 2: Percentage who responded "strongly disagree" or "disagree" to the statement "If a person has smoked for more than 20 years, there is little health benefit to quitting."

[†]Bellingham and Longview/Kelso, Washington; Albany/Corvallis and Medford/Ashland, Oregon; Vallejo and Hayward, California; Santa Fe and Las Cruces, New Mexico; Cedar Rapids and Davenport, Iowa; Raleigh and Greensboro, North Carolina; Paterson and Trenton, New Jersey; Yonkers, New Rochelle, Utica, and Binghamton/Johnson City, New York; and Lowell and Fitchburg/Leominster, Massachusetts.

Smokers' Beliefs - Continued

proceedings of the sixth World Conference on Smoking and Health. New York: Excerpta Medica, 1988:241–3.

- CDC. Tobacco use in 1986: methods and basic tabulations from Adult Use of Tobacco Survey. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1990.
- 5. Pierce JP, Fiore MC, Novotny TE, Hatziandrev EJ, Davis RM. Trends in cigarette smoking in the United States: educational differences are increasing. JAMA 1989;261:56–60.
- Warner KE. Selling smoke: cigarette advertising and public health. Washington, DC: American Public Health Association, 1986.

Current Trends

Smoking-Related Mortality Decline among Physicians - Rhode Island

Declines in smoking in the United States have contributed to declines in heart disease, stroke, and lung cancer among white men (1,2). In Rhode Island, where prevalence of smoking by physicians has been monitored since 1963, the proportion of physicians aged ≥ 25 years who smoke declined by 73% from 1963 to 1983 (Table 1). To characterize smoking-related mortality trends among white male physicians and other white males in Rhode Island, the Rhode Island Department of Health examined vital statistics data from that state. This report summarizes the findings from that study.

For 1968–1987, death certificate information for deaths of resident Rhode Island white men aged \geq 25 years was sorted by age, cause of death, and occupation. The eighth and ninth revisions of the *International Classification of Diseases* (ICD) were used to group deaths by the following categories: all causes, major smoking-related cancers (oral, larynx, pharynx, esophagus, trachea, bronchus, lung, pancreas, and bladder) and heart disease and stroke (*3,4*). Definitions from the 1970 U.S. Census were used to group deaths by occupational categories, including physicians, other professionals (professional, technical, and kindred workers), and others (*5*). ICD-8 and ICD-9 rubrics were used to aggregate deaths for 1968–1978 and 1979–1987, respectively.

	Percer	ntage who smoke cig	arettes
Location/Occupation	1965	1975	1985
United States*	51	42 [†]	31
Rhode Island		44	31
Physician ^s	33	19	9
Nonphysician		44	31
Professional		32	25
Other		46	33

TABLE 1. Percentage of white men aged \ge 25 years who smoke cigarettes, by occupation – Rhode Island and United States, circa 1965, 1975, 1985

*Source: NCHS. Health, United States, 1989. Hyattsville, Maryland: US Department of Health and Human Services, Public Health Service, CDC, 1990.

[†]U.S. population surveyed in 1974.

[§]Rhode Island physicians surveyed in 1963, 1973, and 1983.

Smoking-Related Mortality Decline – Continued

Census data for 1970 and 1980 were used to estimate the populations of physicians and "others"; the population of "other professionals" could not be estimated reliably from available census data. The 1970 U.S. population was used to standardize death rates by age. Rates were calculated for persons 25–64 years of age to ensure compatibility between the two sources of data; counts of deaths included retirees, and estimates of the populations at risk did not.

Proportionate mortality ratios (PMRs) (which do not require estimates of populations at risk) were used to compare the mortality of white male physicians aged \geq 25 years with that of white male nonphysicians aged \geq 25 years.

From 1968 through 1987, 89,593 white males died in Rhode Island, including 420 physicians and 10,640 other professionals. Smoking-related cancers accounted for 11% of deaths, and heart disease and stroke for 50%. Among persons aged 25–64 years, mortality from all causes declined substantially (among physicians, 38%; among nonphysicians, 19%) (Table 2). Among physicians, smoking-related cancer mortality decreased 38%, compared with a 3% decline among nonphysicians. Mortality from heart disease and stroke declined 57% among physicians and 32% among nonphysicians.

For both periods, PMRs for smoking-related cancers were <1.0 among physicians and other professionals and >1.0 among other white males (Table 3). PMRs for smoking-related cancers declined moderately among physicians and remained relatively constant among other professionals and other men. PMRs for heart disease and stroke in the earlier period were >1.0 among physicians and other professionals, decreasing over time among physicians but increasing over time among other professionals.

Reported by: HD Scott, MD, JP Fulton, PhD, JS Buechner, PhD, WJ Waters, PhD, JT Tierney, MSW, Rhode Island Dept of Health.

Editorial Note: These findings indicate that, for the two periods compared (1968– 1978 and 1979–1987), white male physicians in Rhode Island experienced greater declines in overall mortality, smoking-related cancers, and cardiovascular diseases than did white males in other occupations. However, these findings are based on

	196	68–1978	1979–1987			
Disease/Occupation	SDR	95% CI [†]	SDR	95% CI ⁺		
All causes						
Physician	536	414–658	331	228–434		
Nonphysician	755	744–766	611	600–623		
Smoking-related cancers						
Physician	74	28–120	46	9-83		
Nonphysician	98	94–102	95	9099		
Cardiovascular diseases						
Physician	246	164–328	105	47–163		
Nonphysician	352	344–359	241	234–249		

TABLE 2. Age-standardized death rates* (SDRs) from all causes, smoking-related
cancers, and cardiovascular diseases among resident white men aged 25-64 years,
by occupation – Rhode Island, 1968–1978 and 1979–1987

*Per 100,000 population at risk.

[†]Confidence interval (calculated in the manner of Keyfitz [6]).

1.00-1.00

Smoking-Related Mortality Decline – Continued

Rhode Island, 1968–1978 a	nd 1979–1987	white men ageu ≥ /	25 years, by c	leapution	
	190	68–1978	19	79–1987	
Disease/Occupation	PMR	95% CI*	PMR	95% CI*	
Smoking-related cancers					
Physician	0.83	0.51–1.36	0.70	0.46–1.06	
Nonphysician					
Professional	0.87	0.78-0.97	0.84	0.76-0.93	
Other	1.01	1.00–1.02	1.02	1.01–1.03	
Cardiovascular diseases					
Physician	1.04	0.95-1.13	0.98	0.01–1.46	
Nonphysician					
Professional	1.02	0.79–1.32	1.04	0.99–1.09	

TABLE 3. Proportionate mortality ratios (PMRs) for smoking-related cancers and many resident white man aged >25 years by occupation -

1.00 *Confidence interval (calculated from Mantel-Haenszel chi-square values [7]).

relatively small numbers of deaths and denominators and reflect moderate statistical variation. In addition, other risk factors for specific diseases are not considered in this analysis and may affect the results.

1.00-1.00

1.00

The Rhode Island data suggest a method for examining the population effects of smoking cessation on mortality trends among populations whose members have quit smoking in substantial numbers. Based on the study of physicians in Rhode Island, at least half the current cardiovascular and smoking-related cancer mortality of 25-64-vear-old nonphysician white men in that state may be preventable. The Rhode Island Department of Health will use these data to strengthen support for antismoking programs in the state.

References

Other

- 1. CDC. Reducing the health consequences of smoking: 25 years of progress -a report of the Surgeon General. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1989; DHHS publication no.(CDC)89-8411.
- 2. National Cancer Institute. Cancer statistics review 1973-1986. Bethesda, Maryland: US Department of Health and Human Services, Public Health Service, National Institutes of Health, 1989; NIH publication no. 89-2789.
- 3. Public Health Service. Eighth revision international classification of diseases, adapted for use in the United States. Washington, DC: US Department of Health, Education, and Welfare, Public Health Service, 1972; publication no. 1693.
- 4. World Health Organization. Manual of the international statistical classification of diseases, injuries, and causes of death. Geneva: World Health Organization, 1977.
- 5. Bureau of the Census. Census of population, 1970: detailed characteristics-final report. Washington, DC: US Department of Commerce, Bureau of the Census, 1972; PC(1)-D41.
- 6. Keyfitz N. Sampling variance of standardized mortality rates. Hum Biol 1966;38:309-17.
- 7. Kleinbaum DG, Kupper LL, Morgenstern H. Epidemiologic research. Belmont, California: Lifetime Learning Publications, 1982.

Cigarette Smoking among Reproductive-Aged Women – Idaho and New York

Smoking by mothers during pregnancy is associated with a range of serious adverse pregnancy outcomes. To identify strategies to reduce the prevalence of maternal smoking during pregnancy, state health departments should have current and specific information about smoking practices of these reproductive-aged women. This report presents findings from surveys conducted in Idaho and New York to determine family planning needs of reproductive-aged women; the surveys also gathered information on cigarette smoking practices of these women. The sampling methods and questionnaire were similar in both states (1,2).

During 1985, the Idaho Department of Health and Welfare conducted the first statewide Female Health Needs Assessment Telephone Survey. Clusters of residential telephone numbers were sampled to identify women aged 18–44 years; 2025 women were administered a standardized questionnaire regarding their smoking practices, their use of family planning methods, and other reproductive health topics (1). The New York Reproductive Health Survey was conducted during late 1988 and early 1989. Computer-assisted telephone interviews were used to collect data from 1910 women aged 15–44 years living in New York, excluding New York City (2). For this report, analysis of the New York data was restricted to 1809 women aged 18–44 years. In both surveys, current cigarette smoking was defined as responding "yes" to the question "Do you smoke cigarettes now?"

In Idaho and New York, 25.0% (95% confidence interval [CI] = 22.8–27.1) and 31.6% (95% CI = 29.0–34.1) of respondents, respectively, reported that they currently smoked cigarettes. Prevalence of current smoking did not vary substantially in either state by age group. In both states, however, unmarried women were more likely than married* women to be current smokers; 32.3% (95% CI = 26.8–37.7) and 36.7% (95% CI = 31.6–41.8) of unmarried women in Idaho and New York, respectively, were current smokers, compared with 23.1% (95% CI = 20.9–25.4) and 28.7% (95% CI = 25.8–31.7) of married women in Idaho and New York, respectively. Smoking prevalence also varied inversely with level of education in both states; in Idaho and New York, 55.2% (95% CI = 47.4–63.0) and 43.1% (95% CI = 38.8–47.5), respectively, of respondents with <12 years of education were current smokers, compared with 16.0% (95% CI = 13.5–18.4) and 18.6% (95% CI = 12.1–25.0) of respondents with >12 years of education in Idaho and New York, respectively.

In Idaho, where information was collected about religious affiliation, 11.4% of Mormons were current smokers, compared with 28.2% of Protestants, 31.9% of Roman Catholics, and 42.9% of women who reported no religious affiliation. In New York, women who reported an annual income <\$25,000 were more likely to smoke (40.4% [95% CI=34.4–46.4]) than those who reported an income \geq \$35,000 per year (26.3% [95% CI=22.8–29.9]). Among women who were current smokers, 20.0% (95% CI=16.4–23.8) in Idaho and 14.2% (95% CI=10.6–17.7) in New York reported smoking more than one pack of cigarettes per day.

In both states, women who reported having had a liveborn child were asked about their smoking practices during their most recent pregnancy. In Idaho and New York, 19.9% and 26.1% of women, respectively, smoked during their most recent pregnancy

^{*}Married women comprised those currently married and those living with a partner or boyfriend.

Smoking among Reproductive-Aged Women - Continued

(Table 1). In both states, women with less than a high school education were more likely to smoke during pregnancy, as were unmarried women. In Idaho, Mormon women were least likely to smoke during pregnancy (9.7%). In New York, white women and women with an annual income <\$25,000 were more likely to smoke during pregnancy. In Idaho and New York, nearly equal percentages of women smoked more than one pack of cigarettes per day during pregnancy (12.1% [95% CI=8.0–16.3] and 11.6% [95% CI=7.1–16.0], respectively).

In Idaho, 27.7% (95% CI = 22.1–33.2) of women taking oral contraceptives were current smokers; of oral contraceptive users aged 30–44 years, 30.4% (95% CI = 18.1–42.6) smoked. In New York, 33.3% (95% CI = 27.0–39.6) of women taking oral contraceptives also smoked; of oral contraceptive users 30–44 years of age, 20.3% (95% CI = 11.0–29.5) smoked.

	(n	Idaho = 1481)	New York (n = 1112)			
Characteristic	%*	95% CI [†]	%*	95% CI		
Age (yrs)						
18–24	21.2	14.5-28.0	25.6	15.2–36.0		
25–34	18.1	15.0-21.2	26.3	22.1–30.5		
35–44	21.4	17.8–24.9	25.9	21.5–30.4		
Education (yrs)						
<12	45.2	37.2-53.3	34.8	29.9–39.6		
12	22.7	18.8–26.6	18.7	14.8–22.7		
>12	10.9	8.3–13.4	15.5	8.0–23.0		
Marital status						
Married ^s	18.5	16.1–20.9	24.5	21.3-27.7		
Unmarried	30.7	23.4–38.0	33.7	25.8–41.6		
Religion						
Mormon	9.7	7.2-12.2	-	-		
Protestant	22.6	19.1–26.2	-	-		
Roman Catholic	23.3	16.5–30.1	_	-		
None	39.5	30.9–48.1	_	-		
Race						
White	_	_	28.1	24.8-31.3		
Other	-	-	15.2	8.4–21.9		
Annual income						
<\$25,000	21.3	18.4–24.2	35.3	28.4-42.1		
\$25,000-\$34,999	16.5	11.6–21.3	31.3	24.7–37.8		
≥\$35,000	17.7	12.5–23.0	21.7	17.6–25.9		
Total	19.9	17.6–22.2	26.1	23.2–29.0		

TABLE 1. Percentage of reproductive-aged women who smoked during most recent pregnancy, by selected characteristics – Idaho, 1985, and New York, 1988–89

*Percentages weighted to account for sampling.

[†]Confidence interval.

[§]Married women comprised those currently married and those living with a partner or boyfriend.

Smoking among Reproductive-Aged Women - Continued

Reported by: SE Ault, FR Dixon, MD, State Epidemiologist, Idaho Dept of Health and Welfare. ML Woelfel, MA, A Shuttleworth, DL Morse, MD, State Epidemiologist, New York State Dept of Health. Div of Reproductive Health, Center for Chronic Disease Prevention and Health Promotion, CDC.

Editorial Note: Maternal smoking during pregnancy is associated with a doubling in the risk for low birth weight and with an increased risk for placenta previa, abruptio placentae, bleeding during pregnancy, spontaneous abortion, and preterm rupture of membranes (3). The 1990 Health Objectives for the Nation recommended that the proportion of pregnant women who smoke should be no more than one half the proportion of all women who smoke (4); results from these surveys indicate this objective is unlikely to be met.

Based on the reported number of live births for 1987 in Idaho and New York (5) and on the prevalence of smoking during pregnancy (data from these surveys), each year approximately 3200 infants in Idaho and 71,000 infants in New York are exposed to the potentially harmful effects of maternal smoking during pregnancy.

In both states, a substantial proportion of women who used oral contraceptives also were current smokers. For women who use oral contraceptives and smoke cigarettes, the risk for both myocardial infarction and stroke is increased, especially for older women (6,7). Therefore, smoking cessation counseling is particularly important for women taking oral contraceptives (8).

Estimates of reproductive health needs within states are often based on national or regional estimates of such needs. However, data for local areas may not exist or may differ strikingly from national data – particularly for teenagers, unmarried women, and certain racial groups. For example, among women 15–17 years of age in New York, 29.3% were current smokers (2). National surveys may not adequately sample specific subpopulations important in particular states. In the Idaho study, for example, smoking practices among Mormon women, a religious group that advocates healthy behaviors, could be compared with that of women representing other religious groups in that state. These findings underscore the potential usefulness of data from state-specific surveys to program planners and administrators who must allocate and target available resources in local areas.

During the 1980s, the prevalence of smoking in the United States declined, although the decline occurred at a slower rate for women than for men (9). Therefore, smoking prevention and cessation efforts should be focused on women. Health-care personnel who provide family planning and prenatal care services should incorporate these efforts into their counseling of reproductive-aged women.

References

- 1. Idaho Department of Health and Welfare. Idaho Female Health Survey, 1985: final report. Atlanta: US Department of Health and Human Services, Public Health Service, CDC, 1987.
- New York State Department of Health. New York Reproductive Health Survey, 1989: final report. Atlanta: US Department of Health and Human Services, Public Health Service, CDC, 1990 (in press).
- 3. Office on Smoking and Health. Pregnancy and infant health. In: The health consequences of smoking for women: a report of the Surgeon General. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1980.
- Public Health Service. Promoting health/preventing disease: objectives for the nation. Washington, DC: US Department of Health and Human Services, Public Health Service, 1980.
- NCHS. Advance report of final natality statistics, 1987. Hyattsville, Maryland: US Department of Health and Human Services, Public Health Service, CDC, 1989. (Monthly vital statistics report; vol 38, no. 3S).
- 6. Petitti DB, Wingerd J. Use of oral contraceptives, cigarette smoking, and risk of subarachnoid hemorrhage. Lancet 1978;2:234–6.

Smoking among Reproductive-Aged Women - Continued

7. Shapiro S, Slone D, Rosenberg L, et al. Oral contraceptive use in relation to myocardial infarction. Lancet 1979;1:743-7.

MMWR

- 8. Goldbaum GM, Kendrick JS, Hogelin GC, Gentry EM. The behavioral risk factor surveys group: the relative impact of smoking and oral contraceptive use on women in the United States. JAMA 1987;258:1339–42.
- Fiore MC, Novotny TE, Pierce JP, Hatziandreu EJ, Patel KM, Davis RM. Trends in cigarette smoking in the United States. JAMA 1989;261:49–55.

Effects of Maternal Cigarette Smoking on Birth Weight and Preterm Birth – Ohio, 1989

In 1989, most states began using revised birth certificates that provide more detailed information about maternal behaviors during pregnancy and complications of pregnancy. The availability of information on cigarette smoking by mothers in Ohio permitted the Ohio Department of Health (ODH) to examine the proportion of low birth weight (LBW), very low birth weight (VLBW), and preterm births that were attributable to maternal cigarette smoking.

The ODH study included live infants born to Ohio resident mothers in Ohio hospitals from January 1 through June 30, 1989. The analysis was restricted to singleton infants of white (n = 62,732) and black (n = 11,407) mothers. Gestational age was imputed in the 12% of certificates for which a direct estimate from the date of the last menstrual period was not possible; calculations were based on both birth weight and months of completed gestation (1). An infant was classified as having LBW if the birth weight was <2500 g (<5 lbs 8 oz), having VLBW if the birth weight was <1500 g (<3 lbs 4 oz), and being born preterm if the gestational age was <37 weeks. The Ohio birth certificate includes these items: "Tobacco use during pregnancy" and "Average number of cigarettes per day."

Multiple logistic regression was used to control for factors that affect the risk for LBW and preterm delivery, including mother's educational attainment (a measure of socioeconomic status), age, race, prepregnancy weight, and weight gain and alcohol consumption during pregnancy; child's birth order; the month prenatal care began; and previous terminations of pregnancy.

Odds ratios (ORs) were estimated for LBW, VLBW, and preterm birth in relation to in utero exposure to maternal cigarette smoking; these ORs represent measures of the risk for these outcomes in women who smoked compared with nonsmoking women. These findings permitted estimation of the population-attributable risk percentage (PAR%) (i.e., the proportion of all LBW, VLBW, and preterm birth attributable to maternal smoking). The PAR% was approximated as (p × [OR–1]) × 100÷ (p × [OR–1] + 1), where p is the proportion of women in the total population who smoke and OR is estimated in the multivariate model.

Overall, 23% of Ohio mothers were reported to have smoked during pregnancy; this prevalence did not vary by race. Among smokers, white women were more likely than black women (8.8% and 4.7%, respectively) to smoke more than one pack of cigarettes per day during pregnancy. The overall rate of LBW was 5.7%: for whites it was 4.8%; for blacks, 12.1% (Table 1). Overall rates of VLBW and preterm birth were approximately 2–3 times higher for blacks than for whites. Among whites, all three outcomes were more prevalent among younger women; among black women, variation by age group was limited.

Maternal Cigarette Smoking – Continued

Infants born to smokers were more than twice as likely to have LBW as were infants born to nonsmokers (Table 2). In addition, among women who smoked, risk for LBW increased by level of exposure: adjusted ORs were 1.8, 2.2, and 2.4 for light (less than one half pack per day), moderate (one half pack to one pack per day), and heavy smokers (more than one pack per day), respectively. Consumption of even <10 cigarettes per day appeared to double the risk for LBW. For both blacks and whites, the risk was directly proportionate to levels of smoking.

Maternal cigarette smoking also increased the risk for VLBW and preterm birth (Table 3). However, these risks were similar for light and heavy smokers.

An estimated 20% of all LBW in the total Ohio population (i.e., smokers and nonsmokers) in the 6-month period was attributable to maternal smoking (Table 3). Similarly, more than 8% of all VLBW and more than 6% of all preterm deliveries were attributable to smoking. For each of the three outcomes, adjusted ORs and PAR% were slightly lower for blacks than for whites.

Reported by: RS Hopkins, MD, LE Tyler, MS, BK Mortensen, PhD, Div of Epidemiology and Toxicology, Bur of Preventive Medicine, Ohio Dept of Health. Pregnancy and Infant Health Br,

Race/Age (yrs)	% LBW	% VLBW	% Preterm
White			
<20	7.8	1.6	12.6
20–34	4.4	0.7	7.5
≥35	4.6	0.7	7.4
Total	4.8	0.8	8.1
Black			
<20	11.7	2.3	19.3
20–34	12.2	2.6	17.2
≥35	14.2	3.4	18.0
Total	12.1	2.6	17.8

TABLE 1. Percentage of low birth weight (LBW), very low birth weight (VLBW), and preterm birth, by mother's race and age – Ohio, January–June 1989*

*Data based on Ohio birth certificate information.

TABLE 2. Low birth weight (L	.BW) among sin	ngleton infants,	by mother's	cigarette
consumption and race - Ohic), January–June	e 1989*		

Packs per day		W	hite			0.1.1.					
	LBW	(%)	Total	R R [†]	LBW	(%)	Total	RR	ratio [§]		
None	1,744	(3.6)	48,427	1.0 ¶	871	(9.9)	8,780	1.0 ¶	1.0 *		
$<\frac{1}{2}$	223	(6.8)	3,303	1.9	167	(15.1)	1,103	1.5	1.8		
<u>1</u> –1	435	(8.0)	5,459	2.2	167	(16.8)	992	1.7	2.2		
² >1	497	(9.0)	5,543	2.5	125	(23.5)	532	2.4	2.4		
Total	2,899	(4.6)	62,732	_	1,330	(11.7)	11,407	-	_		

*Data based on Ohio birth certificate information.

[†]Relative risk.

[§]Adjusted for mother's educational attainment, age, race, prepregnancy weight, and weight gain and alcohol consumption during pregnancy; child's birth order; the month prenatal care began; and previous terminations of pregnancy. Birth certificates with unknowns for any of these variables were excluded.

[¶]Referent group.

Maternal Cigarette Smoking - Continued

TABLE 3. Crude and adjusted odds ratios and population-attributable risk percentage (PAR%)* for low birth weight (LBW), very low birth weight (VLBW), and preterm birth in relation to maternal cigarette smoking during pregnancy – Ohio, January–June 1989[†]

Measure	LBW	VLBW	Preterm birth
Crude odds ratio	2.2	1.6	1.5
Adjusted odds ratio ^s	2.1	1.4	1.3
PAR%	20.2%	8.4%	6.5%

*PAR% was approximated as ($p \times [OR-1]$) $\times 100 \div (p \times [OR-1] + 1$), where p is the proportion of persons in the total population exposed to the hazard and OR is the odds ratio estimated in the multivariate model.

[†]Data based on Ohio birth certificate information.

[§]Adjusted for mother's educational attainment, age, race, prepregnancy weight, and weight gain and alcohol consumption during pregnancy; child's birth order; the month prenatal care began; and previous terminations of pregnancy. Birth certificates with unknowns for any of these variables were excluded.

Div of Reproductive Health, Center for Chronic Disease Prevention and Health Promotion; National Center for Health Statistics, CDC.

Editorial Note: Smoking by mothers is an important preventable cause of adverse pregnancy outcome (2). In Ohio, the deleterious effects of cigarette smoking by mothers during pregnancy on the rates of LBW, VLBW, and preterm birth were substantial, even when adjusted for other risk factors identified from the birth certificates. The effect of smoking on fetal growth may be partially mediated through lower maternal weight gain. The adjustment for maternal weight gain in this multivariate model may have underestimated the ORs for LBW and VLBW and thus the PAR%. Conversely, the effects reported here could also partially reflect the impact of other factors (e.g., illegal drug use) that were not reported on the birth certificate but that are more common among smokers than nonsmokers (3). Under these circumstances, the PAR% may have been slightly overestimated.

This study relied on data collected during the first 6 months of use of the revised Ohio birth certificate; the reliability of the smoking-related and other data may be expected to improve over time as reporting of new information becomes routine. Nonetheless, the findings in Ohio are similar to those in other studies, some of which used different data sources (2,4-7).

Birth certificates are a useful surveillance tool for identifying subgroups of women who are likely to smoke during pregnancy. These subgroups can then be targeted for special prevention or cessation efforts. Birth certificate data can also be used to evaluate the success of a state's antismoking programs. In 1989, only seven states did not collect information about maternal smoking habits that was comparable to that collected in Ohio on birth certificates.

Smoking during pregnancy increases infant morbidity and mortality through effects on birth weight and preterm birth (5,6). In Ohio and other states, successful efforts to reduce or eliminate smoking during pregnancy could substantially reduce rates of LBW, VLBW, and preterm birth and, in turn, reduce infant morbidity and mortality and the cost of health care in the state (8).

References

 Taffel S, Johnson D, Heuser R, NCHS. A method of imputing length of gestation on birth certificates. Hyattsville, Maryland: US Department of Health and Human Services, Public

Maternal Cigarette Smoking - Continued

Health Service, CDC, 1982; DHHS publication no. (PHS)82-1367. (Vital and health statistics; series 2, no. 93).

- Office on Smoking and Health. The health consequences of smoking for women: a report of the Surgeon General. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1980.
- Zuckerman B, Frank DA, Hingson R, et al. Effects of maternal marijuana and cocaine use on fetal growth. N Engl J Med 1989;320:762–8.
- 4. Kleinman JC, Madans JH. The effects of maternal smoking, physical stature, and educational attainment on the incidence of low birth weight. Am J Epidemiol 1985;121:843–55.
- Kleinman JC, Pierre MB, Madans JH, Land GH, Schrämm WF. The effects of maternal smoking on fetal and infant mortality. Am J Epidemiol 1988;127:274–82.
- Malloy MH, Kleinman JC, Land GH, Schramm WF. The association of maternal smoking with age and cause of infant death. Am J Epidemiol 1988;128:46–55.
- 7. Alameda County Low Birth Weight Study Group. Cigarette smoking and the risk of low birth weight: a comparison in black and white women. Epidemiology 1990;1:201–5.
- 8. Oster G, Delea TE, Colditz GA. Maternal smoking during pregnancy and expenditures on neonatal health care. Am J Prev Med 1988;4:216–9.

Cigarette Brand Use among Adult Smokers – United States, 1986

Information about the use of cigarette brands is important to the development of smoking-prevention and smoking-cessation strategies. This report summarizes data from the 1986 Adult Use of Tobacco Survey (AUTS), which describe the brand of cigarettes smoked as reported by respondents; the data are presented by sex, race, age, and level of educational attainment.

The AUTS, conducted by CDC's Office on Smoking and Health, Center for Chronic Disease Prevention and Health Promotion, was designed to assess the knowledge, attitudes, and practices of adults regarding all forms of tobacco use. Data for this telephone survey, conducted primarily during October through December of 1986, were collected from a national probability sample of 13,031 respondents \geq 17 years of age and were weighted to represent the civilian, noninstitutionalized, adult U.S. population. According to the AUTS, an estimated 26.5% (approximately 46.8 million) of adults were smoking cigarettes in 1986 (1,2).

Data from the 4700 current cigarette smokers in 1986 who responded to the AUTS were used in this analysis. Current brand use was determined by responses to the question, "What brand of cigarettes do you usually smoke now?" (1). A series of follow-up questions were used to determine the specific variety of the brand used (e.g., mentholated vs. nonmentholated and "lights" vs. regular). In this report, however, data are presented only by overall brand categories. Market share data* are provided for comparison.

In 1986, the 12 most commonly named brands of cigarettes smoked were used by 74.7% of all current smokers and accounted for 72.6% of the cigarette market (3) (Table 1, page 671). Marlboro, Winston, Salem, Kool, and Newport-the top five brands smoked-were used by 52.0% of current smokers and accounted for 52.1% of the cigarette market. The percentage of smokers who reported using Marlboro (24.1%) was more than double the percentage who reported using Winston (9.6%), the next most commonly named brand (these findings were also consistent with known market share patterns [3]).

^{*}Percentage of all cigarettes sold in the United States, by brand. Market share data are collected quarterly by a tobacco industry analyst (3).

DISEASE DECREASE INCREASE CASES CURRENT 4 WEEKS Aseptic Meninaitis 1.394 Encephalitis, Primary 124 Hepatitis A 1.826 Hepatitis B 1.371 Hepatitis, Non-A, Non-B 159 Hepatitis, Unspecified 125 108 Legionellosis 103 Malaria 1.834 Measles, Total 124 Meningococcal Infections 215 Mumps 444 Pertussis Rabies, Animal 296 12 Rubella 0.25 8 0.5 2 1 4 Ratio(Log Scale)* Beyond Historical Limits Ratio

FIGURE I. Notifiable disease reports, comparison of 4-week totals ending September 22, 1990, with historical data – United States

*Ratio of current 4-week total to mean of 15 4-week totals (from comparable, previous, and subsequent 4-week periods for past 5 years).

TABLE I. Summary – cases of specified notifiable diseases, United States, cumulative, week ending September 22, 1990 (38th Week)

	Cum. 1990		Cum. 1990
AIDS Anthrax Botulism: Foodborne Infant Other Brucellosis Cholera Congenital rubella syndrome Diphtheria Encephalitis, post-infectious Gonorrhea: civilian military Leprosy Leptospirosis Measles: imported indigenous	30,875 14 47 56 4 3 2 73 481,068 6,421 156 34 1,038 20,227	Plague Poliomyelitis, Paralytic* Psittacosis Rabies, human Syphilis: civilian military Syphilis, congenital, age < 1 year Tetanus Toxic shock syndrome Trichinosis Tuberculosis Tuberculosis Tularemia Typhoid fever Typhus fever, tickborne (RMSF)	1 34,471 177 685 43 233 22 16,828 102 344 487

*Three cases of suspected poliomyelitis have been reported in 1990; five of 13 suspected cases in 1989 were confirmed and all were vaccine-associated.

	Г	Asentic	Encephalitis				Hepatitis (Viral), by type					
Reporting Area	AIDS	Menin- gitis	Primary	Post-in- fectious	Gond (Civi	orrhea ilian)	A	В	NA,NB	Unspeci- fied	Legionel- losis	Leprosy
	Cum. 1990	Cum. 1990	Cum. 1990	Cum. 1990	Cum. 1990	Cum. 1989	Cum. 1990	Cum. 1990	Cum. 1990	Cum. 1990	Cum. 1990	Cum. 1990
UNITED STATES	30,875	6,233	621	73	481,068	505,146	20,780	14,621	1,629	1,223	901	156
NEW ENGLAND	1,098	253	21	-	13,545	14,571	436	778	53	52	39	10
Maine	43	8	3	-	147	201	7	24	4	1	4	-
N.H. Vt	48	25	-	-	119	122	7	35	4	3	4	•
Mass.	638	25	10		5 710	5 650	296	38 484	31	46	5 19	- 9
R.I.	56	76	10	-	841	1.058	45	35	-	2	7	1
Conn.	300	30	5	-	6,685	7,493	76	162	10	-	-	-
MID. ATLANTIC	9,519	605	36	6	65,265	74,158	2,952	1,965	174	84	288	19
Upstate N.Y.	1,201	331	29	1	10,309	11,288	835	535	57	22	110	1
N.Y. City	5,592	114	3	2	27,232	30,256	442	531	23	43	75	13
N.J. Pa	1,823	160	1	-	10,848	11,131	342	434	35	- 10	43	4
	503	100	3	3	10,070	21,403	1,333	405	59	19	00	
E.N. CENTRAL	2,131	1,390	164	12	90,292	91,600	1,618	1,700	154	74	220	2
Ind	505	290	55	4	20,4/5	23,305	152	202	55	12	24	-
III.	846	205	51	2	28 666	30.038	784	335	33	15	15	1
Mich.	416	605	46	-	21,436	23,624	291	481	26	32	62	i
Wis.	187	59	6	-	5,589	7,826	278	283	30		33	-
W.N. CENTRAL	727	321	57	2	25 243	22 776	1 233	676	100	30	44	1
Minn.	145	41	23	1	3,185	2.535	175	88	21	-	1	-
lowa	25	59	5	-	1,843	1,963	232	47	8	4	4	-
Mo.	397	142	7	1	15,232	14,032	367	416	47	20	23	-
N. Dak.	2	16	3	-	76	108	14	5	2	1	1	-
5. Dak. Nebr	2	5	2	-	184	197	182	7	3	-	1	-
Kans.	43	25	10	-	3 417	2 9 1 9	190	20	4	5	9	
			10	-	3,417	2,313	150			5		_
5. ATLANTIC	6,354	1,242	144	21	138,193	136,280	2,475	2,800	236	178	137	5
Md	645	30	3		2,231	2,331	95	201	27	2	10	-
D.C.	521	8			9 558	8 287	13	31	4	-		-
Va.	571	214	40	1	12,902	11,482	219	178	33	128	11	-
W. Va.	55	40	38	-	851	1,026	17	62	4	6	3	-
N.C.	409	142	29	-	20,890	20,614	541	792	88	-	21	1
S.C.	258	15	1	-	11,054	12,502	33	451	14	8	15	-
Fla.	3 0/19	227	4	1	30,381	26,174	291	327	8	19	15	- 1
	3,040	405	12	18	33,905	38,005	430	497	42	10	3	'
E.S. CENTRAL	780	497	46	2	41,767	40,072	284	1,152	135	6	49	-
Tenn	130	117	19	-	4,399	3,911	69	406	39	4	20	-
Ala.	167	206	20	2	12,550	13,392	129	120	/8	1	13	
Miss.	216	84			10.377	9.942	1	4	2	1	-	-
W.S. CENTRAL	2 207	EAE	26	-	54 500	50.040	0.145	1 500	67	202	40	22
Ark.	168	045 13	30	/	51,585	53,048	2,145	1,509	70	203	40	- 32
La.	476	69	6	-	9 039	11.369	142	238	4	7	13	-
Okla.	158	55	3	6	4,434	4,622	409	113	19	26	13	-
lex.	2,495	408	26	1	31,639	31,041	1,207	1,098	37	151	7	32
MOUNTAIN	853	273	19	2	9.585	10.741	3.379	1,103	160	94	34	-
Mont.	11	4	-	-	132	142	132	53	6	4	3	-
Idano	19	7	-	-	105	137	, 78	62	8	-	3	-
Colo	2	1	1	-	117	75	48	13	5	1	1	-
N. Mex.	203	61	4	-	2,150	2,289	216	121	36	31	6	-
Ariz.	264	138		-	2 0 2 0	4 4 2 2	1 5 7 9	393	60	36	10	-
Utah	82	25	3	-	304	342	393	82	21	5	3	-
Nev.	137	24	4	2	1,924	2,336	254	231	15	10	5	-
PACIFIC	6 1 1 6	1 107	00	21	45 502	61 000	6 250	2 0 2 9	550	502	50	87
Wash.	436	1,107	6	21	40,092	4 776	1 046	2,530	92	28	11	5
Oreg.	230	-	-		1.872	2.260	649	306	43	7		-
Calif.	5,300	938	85	19	38,768	53,768	4,347	2,105	401	459	38	69
Alaska Hawaii	23	100	6	-	776	691	153	46	5	3	-	-
	127	69	1	1	359	405	63	52	9	5	1	13
Guam	2	2	-	-	162	123	11	2	-	10	-	-
r.n. V I	1,221	46	6	-	509	794	117	239	5	22	-	-
Amer Samoa	10	-	-	-	321	491	_1	10	-	-	-	-
C.N.M.I.	-	1	-	-	49	38	26	-	-	-	-	10
	-	-	-	-	150	73	10	9	-	15	-	4

TABLE II. Cases of specified notifiable diseases, United States, weeks ending September 22, 1990, and September 23, 1989 (38th Week)

N: Not notifiable

I

1

		Measles (Rubeola) Menin-			Buballa										
Reporting Area	Malaria	Indig	enous	Impo	orted*	Total	gococcal Infections	wumps			Pertussi	S		Rubella	1
	Cum. 1990	1990	Cum. 1990	1990	Cum. 1990	Cum. 1989	Cum. 1990	1990	Cum. 1990	1990	Cum. 1990	Cum. 1989	1990	Cum. 1990	Cum. 1989
UNITED STATES	856	197	20,227	3	1,038	12,645	1,836	50	4,060	110	2,662	2,608	-	796	304
NEW ENGLAND	73	-	256	-	25	322	139	1	37	14	308	289	-	8	6
Maine	1	-	27	-	2	1	12	-	-	:	10	17	-	1	;
N.H. Vt.	4		-	-	8	15	10 10	1	9	4	44	6	-	1	4
Mass.	38	-	19	-	Ż	49	64	-	11	7	226	233	-	2	1
R.I. Conn.	17	-	27 183	:	3	41 213	12 31	-	5 11	1	4 18	11 16	-	1	
	188	Q	1 1 2 1		154	939	272	٩	262	5	415	177		11	31
Upstate N.Y.	39	-	200	-	110	140	102	9	115	5	288	77	-	10	13
N.Y. City	67 59	-	322		21	102	41	-	-	-	-	5	-	-	15
Pa.	23	9	375		9	263	67	-	85	-	106	29 66	-	1	-
E.N. CENTRAL	52	-	3,237	-	143	4,260	244	4	439	5	505	361	<u>.</u> .	31	25
Ohio	7	-	549	•	3	1,133	75	-	89	-	154	45	-	1	3
ING. III.	22		323	:	10	2.502	25 64	2	20 152	:	90 100	19 122	-	18	20
Mich.	16	•	348		125	321	59	4	132	5	70	35	-	9	1
Wis.	4	-	743	-	4	226	21	-	46	-	91	140	-	3	1
W.N. CENTRAL Minn	16 4	-	853 392	:	13	652	58	2	131	8	154	184	-	22	6
lowa	2		25	-	1	9	1	-	18	1	18	50 14	-	4	1
Mo. N. Dak	9	-	96	-	-	368	23	1	54	3	79	109	-	-	4
S. Dak.	-		15		8	-	1	-		2	2	2	-	1	
Nebr.	-	-	97	-	1	113	5	1	5		ż	5	-	-	-
Kans.	1	-	228	-	-	140	15	-	40	4	16	3	-	•	1
S. ATLANTIC Del	171	:	886 8		354	579	327	26	1,691	8	239	239	-	18	9
Md.	48	-	195	-	18	88	38	20	949	4	59	37		2	2
D.C. Va	10		15		7	40	11	1	33	-	14	-	-	1	-
W. Va.	2		6	-	-	51	40	1	95 41	:	17	28 24	-	1	
N.C.	13	-	9	•	15	171	50	1	281	2	64	48	-	-	1
Ga.	15	:	4 82		239	3	23 56	1	50 82		5 24		-	-	-
Fla.	37	-	485	-	70	162	93	2	156	2	34	68	-	14	6
E.S. CENTRAL	18	4	181	-	3	229	111	-	87	6	136	175	-	5	3
Ky. Tenn	2	-	40	-	1	40	33	-	-	:	-	1	-	1	-
Ala.	7	-	23	-	2	50	46 30	-	49 14	5	64 65	105 58	-	4	1
Miss.	-	-	26	•	-	-	2	-	24	-	7	11	-		-
W.S. CENTRAL	48	172	4,180	3	91	3,144	128	5	611	32	144	264	-	66	36
La.	3		16 10	:	28	15 11	16 29	1	134	1	14	21	-	3	5
Okla.	9	-	175	-	-	106	16	-	107	3	45	46	-	1	1
Tex.	32	172	3,979	3†	63	3,012	67	4	268	28	59	182	-	62	30
MOUNTAIN Mont.	22	12	823	-	99	391	60	1	311	15	240	545	-	109	35
Idaho	4	-	16		10	2	10	-	1 142	3	32 37	33	-	14 49	32
Wyo. Colo	1	-	-	•	15	-	-	-	2		-	-	-	-	1
N. Mex.	4		81	:	46	82 31	17	- N	23 N	11	74	45	-	4	-
Ariz.	9	7	291	-	12	145	5	1	119		49	357	-	32	-
Nev.	1	3	127		- 3	114	6	-	9	1	27	19	-	2	- 1
PACIFIC	268		8 680		156	2 1 2 0	406	-	15		4		-	0	153
Wash.	20	-	202	-	69	2,129	496	2	490 44	17 5	521 142	374	-	526	155
Oreg. Calif	12	-	168	-	44	29	54	Ň	N	5	63	12	-	10	4
Alaska	230	:	0,223 78		3/	2,018	366		423	7	274	194	-	503	127
Hawaii	4	-	9	-	4	30	5	1	19		38	16	-	13	22
Guam	3	U		U	1	4	-	U	3	U	-	1	υ	-	-
r.n. V.I.	2	4	1,640 21	:	- 2	513	10	-	7	-	6	4	-	-	8
Amer. Samoa	35	U	190	Ű	-	-	-	Ū	9 19	Ū.		-	ū	:	-
L.IN.M.I.	•	U	•	U	-	-	-	υ	8	Ũ	4		Ű	-	-

TABLE II. (Cont'd.) Cases of specified notifiable diseases, United States, weeks ending September 22, 1990, and September 23, 1989 (38th Week)

*For measles only, imported cases includes both out-of-state and international importations. [§]Out-of-state N: Not notifiable

Dum. Cum. Cum. <th< th=""><th>Reporting Area</th><th>Syphilis (Primary &</th><th>(Civilian) Secondary)</th><th>Toxic- shock Syndrome</th><th>Tuber</th><th>culosis</th><th>Tula- remia</th><th>Typhoid Fever</th><th>Typhus Fever (Tick-borne) (RMSF)</th><th>Rabies, Animal</th></th<>	Reporting Area	Syphilis (Primary &	(Civilian) Secondary)	Toxic- shock Syndrome	Tuber	culosis	Tula- remia	Typhoid Fever	Typhus Fever (Tick-borne) (RMSF)	Rabies, Animal
UNITED STATES 34,471 31,039 233 16,828 15,282 102 344 487 3,126 NEW ENGLAND 1,260 1,233 18 406 422 3 22 15 5 Name 7 9 7 8 12 - - - 2 Winn 4 10 1 8 16 21 3 221 13 - - - 2 3 </th <th></th> <th>Cum. 1990</th> <th>Cum. 1989</th> <th>Cum. 1990</th> <th>Cum. 1990</th> <th>Cum. 1989</th> <th>Cum. 1990</th> <th>Cum. 1990</th> <th>Cum. 1990</th> <th>Cum. 1990</th>		Cum. 1990	Cum. 1989	Cum. 1990	Cum. 1990	Cum. 1989	Cum. 1990	Cum. 1990	Cum. 1990	Cum. 1990
NEW ENGLAND 1.200 1.233 18 405 422 3 22 15 5 N.H. 40 10 1 3 19 - - - 2 N.H. 40 10 1 3 19 - - - 2 Mass 495 376 8 219 219 3 21 13 - - - 2 2 Conn 701 884 6.395 22 4.062 2.986 1 73 22 688 N.Y.GUN 3.314 2.662 5 2.535 1.642 - 42 - - - - 2.22 688 1 13 3 385 1 1 1 3 385 1 1 1 3 385 1 1 1 3 385 1 1 1 3 385 1 1 1 3 385 1 1 1 3 385 1 1 1 1	UNITED STATES	34,471	31,039	233	16,828	15,282	102	344	487	3,126
Maine 7 9 7 6 12 - Conn 701 810 1 120 1 120 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 <td< td=""><td>NEW ENGLAND</td><td>1,260</td><td>1,233</td><td>18</td><td>405</td><td>422</td><td>3</td><td>22</td><td>15</td><td>5</td></td<>	NEW ENGLAND	1,260	1,233	18	405	422	3	22	15	5
N.H. 40 10 1 3 19 - - - 2 RL 40 15 25 1 22 47 -	Maine	7	9	7	6	12	-	-	-	-
Mass. 4.96 378 8 216 219 3 21 13 - Conn. 701 810 1 120 117 - 1 2 638 Upstate N.Y. 644 6396 8 299 237 - 14 12 91 N.J. 1.144 1.011 - 674 553 1.644 5 3 365 Choin 365 1654 1.561 2 25 1.11 1 3 365 Ohio 365 165 1.634 1.561 2 25 44 140 Ohio 365 165 20 25 45 50 3 365 Wich. 746 249 312 339 5 5 45 500 Minn. 72 78 72 78 72 7 6 50 Mich. 32 37 2	N.H.	40	10	1	3	19		-	-	2
R.I. 15 25 1 52 47 -	Mass.	496	378	8	216	219	3	21	13	-
Conn. 701 810 1 120 117 - 1 2 63 Upster NY. 648 6396 22 4062 2968 1 73 22 68 Upster NY. 644 639 8 299 237 - 14 12 N. N. Chy 3,314 2,862 5 2,535 1,644 533 1 14 7 3 236 N. CENTRAL 2,414 1,289 55 1,634 1,551 2 25 44 140 S. CENTRAL 2,414 1,289 55 1,634 1,551 2 25 44 140 S. M. CENTRAL 2,414 1,289 55 2,183 1,51 1 3 1 8 Mich. 745 446 24 312 339 - 5 7 46 Wis. 240 117 - 64 89 - 1 3 2 25 Mich. 745 446 24 312 339 - 5 7 46 Wis. 240 117 - 64 89 - 1 1 - 51 W.N. CENTRAL 3,240 117 - 64 89 - 1 1 - 51 W.N. CENTRAL 3,240 117 - 64 89 - 1 1 1 17 Mo. 202 126 8 228 178 22 1 1 1 Mo. 202 126 8 228 178 22 - 1 1 1 17 Mo. 202 126 8 228 178 22 1 1 17 Mo. 202 126 4 2 1 3 14 18 3 - 1 147 Mo. 202 128 8 218 178 22 3 2 1 12 38 S.ATLANTC 11,349 11,166 21 3,182 3,255 3 60 212 379 S.ATLANTC 11,349 11,166 21 3,182 3,255 3 60 212 379 Del, 135 140 1 2 38 23 1 - 1 2 M. S.ATLANTC 11,349 11,166 21 3,182 3,255 3 60 212 379 S.ATLANTC 1,349 11,166 21 3,182 3,255 3 60 212 379 Del, 135 140 1 2 38 23 1 - 1 2 M. S.ATLANTC 1,349 11,166 21 3,182 3,255 3 60 212 379 S.ATLANTC 1,349 11,166 21 3,182 3,255 3 60 212 379 Del, 135 140 1 2 38 27 - 30 15 324 DC. 788 608 576 1 223 277 - 30 15 324 DC. 788 608 17 10 17 17 10 17 409 1 1 2 123 7 S.C. 761 622 2 355 363 1 1 3 3 160 Fa. 3,921 5,258 31 1 - M. S. ATLANTC 1,268 771 10 17 12 1,205 7 7 2 64 377 S.C. 761 622 2 355 363 1 1 3 35 100 Fa. 3,921 5,258 3 1,161 . 2 13 160 Mo. 1 2 12 7 M.S. CHTRAL 3,304 2,766 1 2 315 346 6 . 47 277 M.S. CHTRAL 5,306 4,279 12 1,190 1,205 7 7 2 64 377 M.S. CHTRAL 5,306 4,279 12 1,193 1,818 34 11 70 377 Ten. 46 400 1 2 12 379 M.S. CHTRAL 5,306 4,279 12 1,193 1,818 3,329 4 12 4 4 3 M.S. CHTRAL 5,306 4,279 13 2 4 25 415 344 11 3 18 10 Moth. 6 1 2 11 77 137 147 158 8 2 45 11 1 4 4 40 Moth. 6 1 1 2 11 4 46 Moth. 177 83 7 147 155 8 2 46 1 1 77 M.S. CHTRAL 5,306 4,279 17 12 1,193 1,818 3,329 4 128 8 Moth. 104 40 30 - 13 44 32 17 3 18 40 11 19 M.Y. CHTRAL 5,306 4,279 17 13 344 12 3 18 10 Moth. 12 37 5 5 30 - 2 16 20 Moh. 177 8 2 44 10	R.I.	15	25	1	52	47	-	:	:	
MD. ATLANTIC 6,884 6,396 22 4,062 2,868 1 73 22 688 N.Y. CiVy 3,314 2,862 5 2,855 1,842 - 42 - - N.J. 1,44 1,289 5 2,855 1,864 490 - 3 3 365 Fa. 1,874 1,884 9 564 490 - 3 3 365 Ohio 384 107 2 286 273 1 5 3 3 365 Mich. 246 137 2 286 273 1 5 7 46 Min. 980 644 8 817 1 1 1 7 25 Wis. 240 17 - 64 289 - 1 1 1 1 1 Wis. 240 17 - 72 6 3 29 20 1 1 1 1 1 1 1 1 <	Conn.	701	810	1	120	117	-	1	2	3
	MID. ATLANTIC	6,984	6,396	22	4,062	2,968	1	73	22	688
N.J. 1148 1011 - 674 1693 1 14 7 222 Pa, 1,874 1,884 9 554 496 - 3 3365 E.N.CENTRAL 2,414 1,289 55 1,834 1,861 2 25 41 140 Ohio 64 47 1 138 138 1 1 1 9 Ind. 64 47 1 138 138 1 1 1 9 Mich. 745 6446 24 312 339 - 5 7 46 Win. 82 37 2 78 72 - - - 189 Gova 56 29 6 44 28 - 1 1 17 7 Nak. 1 1 - 16 12 4 - - 2 2 2 2 2 1 12 38 3 - 1 1 1 7	NY City	648 3 314	2 862	8	299	23/		14	12	91
Pa. 1,874 1,884 9 554 496 - 3 3 365 DNio 24 1,861 22 25 41 140 Ohio 64 47 1 388 1 1 1 9 Ind. 64 47 1 388 1 1 1 9 Mich. 745 446 24 312 339 -5 5 455 500 Mich. 745 446 24 390 35 5 5 55 500 Mon. 202 126 8 228 178 26 3 29 20 Nova. 1 3 - 16 12 - - 119 48 Nob. 9 21 3,182 3,255 3 60 212 89 S.ATLANTIC 1,339 11,166 21 3,182 3,255	N.J.	1,148	1.011	- 5	674	593	1	14	7	232
E.N. CENTRAL 2.414 1.299 55 1.634 1.661 2 25 41 140 Ohio 64 47 1 138 138 1 1 1 9 Ind. 640 54 8 834 716 1 3 2 25 Mich. 745 446 24 312 339 - 5 7 46 Win. 282 37 2 78 72 - - - 189 1	Pa.	1,874	1,884	9	554	496	-	3	3	365
Ohio 385 105 22 286 279 1 5 31 9 Ind. 64 47 1 138 1 1 1 9 III. 980 544 8 834 716 - 13 2 25 Wis. 240 117 - 64 89 - 1 - 51 WN. CENTRAL 393 245 25 439 380 35 5 45 500 Mon. 82 37 2 78 72 - - 11 1 17 Mos. 1 3 - 16 12 - - 72 134 Vast. 1 3 - 16 12 32 318 3 - 112 38 Vast. 1 3 - 1 10 24 4 - 12 135<	E.N. CENTRAL	2,414	1,299	55	1,634	1,561	2	25	41	140
	Ohio	385	105	22	286	279	1	5	31	9
Mich. 346 46 312 339 1 1 2 46 Wis. 240 117 - 64 89 - 1 - 51 Wis. 240 117 - 64 89 - 1 - 51 Minn. 82 37 2 78 72 - - - 189 Kowa 50 29 6 44 26 1 1 17 Nok. 1 1 - 10 24 - 2 120 Nok. 9 21 3 142 3.25 3 60 212 289 S.ATLANTIC 11,349 11,166 21 3,182 3,255 3 60 212 89 D.C. 98 66 1 223 255 4 1 1 33 N.C. 1,268 771 10	Ind.	64	47	1	138	138	1	1	1	9
Wis. 240 17 - 64 69 - 1 - 51 WN, CENTRAL 333 245 25 439 380 35 5 45 500 Minn. 82 37 2 78 78 78 7 - - 180 Noba 502 26 6 448 2 1 1 10 24 - - 2 100 N. Dak 1 1 - 100 24 - - 1 20 22 2 22 2 2 2 2 2 2 2 2 100 Nok 2 1 12 38 3 60 212 879 2 2 131 - 1 11 136 2 2 131 2 1 131 1 121 38 3 1 1 121 38 1 1 121 38 3 1 1 1 1 1 1 1	Mich.	745	446	24	312	339	-	5	27	25 46
W.N.CENTRAL 383 245 25 439 380 35 5 45 500 Ninn. 56 29 6 44 28 - 1 1 17 Mo. 202 126 8 228 178 26 3 29 20 N. Dak. 1 1 - 10 24 - 2 72 S. Dak. 1 1 - 10 24 - 2 70 Nebr. 9 21 3 14 18 3 - 1 44 Kars. 42 28 6 49 48 2 1 12 38 S.ATLANTIC 11,349 11,166 21 2,312 5,255 3 60 1212 38 146 Dc. 788 606 1 118 138 - 1 1 33 160 6 121	Wis.	240	117		64	89	-	ĭ	-	51
$\begin{array}{llllllllllllllllllllllllllllllllllll$	W.N. CENTRAL	393	245	25	439	380	35	5	45	500
Iowa 56 29 6 44 28 - 1<	Minn.	82	37	2	78	72		-	-	189
NDD. 202 126 8 228 178 20 3 29 20 S. Dak. 1 1 - 10 24 4 - 2 72 S. Dak. 1 1 - 10 24 4 - 2 72 S. Dak. 1 11 11 11 11 14 18 3 - 1 4 Kans. 42 28 6 49 48 2 1 12 38 S. ATLANTIC 11,349 11,166 21 3,182 3,255 3 60 212 879 Del. 135 140 1 26 31 - 0 15 214 D.C. 788 608 1 118 138 - 1 1 33 N.C. 1,268 771 10 417 409 1 2 13 160 Ga. 2,940 2,786 1 541 510 - 2 13 160 Fla. 3,921 5,258 3 1,165 1,208 - 1 9 37 Ay. 6	lowa	56	29	6	44	28	-	1	1	17
S. Dak. 1 1 - 10 24 4 - 2 16 Nabr. 9 21 3 14 18 3 - 1 4 Kans. 42 28 6 49 48 2 1 12 38 S. ATLANTIC 11,349 11,166 21 3,182 3,277 - 30 15 324 Del. 135 140 1 26 31 - - 1 21 Md. 868 576 1 233 277 - 30 15 324 Va. 611 392 2 275 265 1 5 19 148 Va. 611 392 2 235 363 1 1 33 100 12 125 77 30 16 13 31 100 16 12 12 13 160 11 14 13 16 14 12 13 160 12 13	Mo. N. Dak	202	126	8	228	1/8	26	3	29	20
Nebr. 9 21 3 14 18 3 - 1 4 Kans. 42 28 6 49 48 2 1 12 38 S. ATLANTIC 11,349 11,166 21 3,182 3,255 3 60 212 879 Del. 135 140 1 26 31 - - 1 21 DC. 788 606 1 118 138 - - 1 1 33 N.C. 1,268 771 10 417 409 1 2 13 160 Ga. 2,940 2,786 1 541 510 - 2 13 160 Fla. 3,172 2,017 12 1,190 1,205 7 2 64 137 Ky. 67 41 2 1,190 1,205 7 2 64 2	S. Dak.	i	1	-	10	24	4	-	2	160
Kans. 42 28 6 49 48 2 1 12 38 SATLANTIC 11,349 11,166 21 3,182 3,255 3 60 212 879 Dal. 135 140 1 28 3,255 3 60 212 879 Md. 868 576 1 233 277 -30 15 324 D.C. 708 6008 1 118 138 - - 1 - Va. 57 13 - 52 54 - 1 1 133 106 S.C. 761 622 2 355 363 1 1 35 106 Ga. 2,940 2,786 1 541 510 - 2 80 377 Ga. 2,940 2,786 3 1,165 1,205 7 2 64 137 Tyrn. 1,259 821 2 315 314 1 1 9 <td< td=""><td>Nebr.</td><td>9</td><td>21</td><td>3</td><td>14</td><td>18</td><td>3</td><td>-</td><td>1</td><td>4</td></td<>	Nebr.	9	21	3	14	18	3	-	1	4
S. ATLANTIC 11,349 11,166 21 3,182 3,265 3 60 212 879 Del. 135 140 1 26 31 - 1 21 Md. 868 576 1 233 277 - 30 15 324 D.C. 768 608 1 118 138 - 1 1 3 N.C. 1,268 771 10 417 409 1 2 125 7 S.C. 761 622 2 355 363 1 1 35 106 Ga. 2,940 2,786 1 541 510 - 2 13 160 Fla. 3,921 5,258 3 1,165 1,208 - 19 2 80 Fla. 3,921 5,258 3 1,165 1,208 - 19 2 80 E.S. CENTRAL 3,172 2,017 12 1,190 1,205 7 2 64 137 Ky. 67 41 2 225 301 1 1 9 37 Tenn. 1,259 821 8 315 354 6 - 47 27 Ala. 1,004 655 2 375 344 - 1 8 70 Miss. 842 500 - 215 206 3 W.S. CENTRAL 5,306 4,274 11 1,963 1,818 34 11 70 377 Ark. 443 264 - 262 189 26 - 17 442 Cha. 1,77 83 7 147 155 8 2 45 100 Mont. 177 83 7 147 155 8 2 45 100 MONTAIN 628 474 25 415 344 13 18 10 162 Ota. 37 584 - 1 8 70 Miss. 842 201 - 22 11 2 28 Cha. 1,77 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 13 18 10 162 Ota. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 13 18 10 162 Ota. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 1,225 - 9 6 2002 MOUNTAIN 628 474 25 415 344 13 18 10 162 Ota. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 1,225 - 9 6 2002 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont. 1 2 21 1 4 50 Mont. 1 2 21 21 4 50 Mont. 2 1 2 21 3 4 50 Mont. 1 2 21 21 4 50 Mont. 2 1 2 21 3 3 3 N.M. 8 13 4 32 27 3 - 3 3 Mos. 134 42 27 3 - 3 3 Mos. 134 42 27 3 - 3 3 Mos. 135 - 2 - 16 Colo. 37 58 7 27 39 3 - 1 41 1 Mot. 1 2 21 1 4 50 Mont. 2 1 2 11 3 18 10 162 Mont. 4 1 2 12 21 3 3 4 32 27 3 - 3 3 Mos. 91 153 - 5 - 3 - 3 3 Mos. 91 153 - 5 - 2 - 16 Colo. 37 58 - 66 217 4 1 1 1 Mos. 27 33 36 - 2 - 16 Colo. 37 58 - 66 217 4 1 1 - 2 Mos. 91 153 - 5 - 2 Mos. 91 153 - 5 - 2 Mos. 91 160 - 1 327 - 5 - 2 Mos. 91 173 - 1 40 132 - 5 - 5 3 Mos. 91 173 - 1 40 132 - 5 - 5 3 Mos. 91 173 - 1 40 132 - 5 - 5 3 Mos. 91 173 - 1 40 132 - 5 - 5 3 Mos. 91 173 - 1 40 12 5 - 5 4 Mos. 14 3 3 8 - 40 19 - 4 4 Mo	Kans.	42	28	6	49	48	2	1	12	38
Del. 135 140 1 26 31 1 21 Md. 868 576 1 233 277 - 30 15 324 D.C. 788 608 1 118 138 1 1 . Va. 611 392 2 275 265 1 5 19 148 W.Va. 57 13 - 52 54 - 1 1 33 N.C. 1266 771 10 417 409 1 2 125 7 S.C. 761 622 2 355 363 1 1 355 166 Ga. 2.940 2.786 1 541 510 - 2 13 160 Fla. 3.921 5.258 3 1,165 1.208 - 19 2 80 E.S.CENTRAL 3.172 2.017 12 1,190 1.205 7 2 64 137 Tenn. 1.259 821 8 315 354 6 - 47 27 Ala. 1.004 665 2 375 344 - 1 8 70 Miss. 842 500 - 215 206 3 MS.CENTRAL 5.306 4.274 11 1.963 1.818 34 11 70 377 Ark. 443 264 - 262 189 26 - 17 42 La. 1.171 1.032 1 170 249 - 2 285 105 Tex. 3.515 2.895 3 1.384 1.225 - 9 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Cha. 1 2 11 2 1 - 4 400 Mont 1 - 22 11 - 4 400 Mont 1 - 22 5 - 3 - 5 46 - 1 8 Ariz. 454 221 7 183 148 4 13 18 10 162 Mont 1 - 22 11 - 4 400 Mont 1 - 22 11 - 4 400 Mont 1 - 22 5 - 3 - 4 Ariz. 454 221 7 183 148 - 16 1 27 Va. 6 2 474 25 415 344 - 1 8 Ariz. 454 221 7 183 148 - 16 127 Cola. 37 58 7 27 39 3 - 1 111 PACIFIC 2.965 3.935 44 2.25 - 9 6 202 MOUNTAIN 628 474 25 415 344 3 - 1 8 Ariz. 454 221 7 183 148 - 16 127 Cola. 37 58 7 27 39 3 - 1 111 PACIFIC 2.965 3.935 44 3.518 3.329 4 128 8 Ariz. 454 221 7 183 148 - 16 1 27 Otha 8 13 4 32 27 3 - 3 - 46 Colo. 37 58 7 27 39 3 - 1 111 PACIFIC 2.965 3.935 44 3.518 3.329 4 128 8 Ariz. 454 221 7 183 148 - 16 1 27 Aris. 252 338 4 205 774 1 20 - 2 Aris. 252 338 4 205 774 1 20 - 2 Aris. 252 338 4 205 774 1 20 - 2 Aris. 252 338 4 205 774 1 20 - 2 Aris. 252 338 4 205 774 1 20 - 2 Aris. 252 338 4 205 774 1 20 - 2 Aris. 252 338 4 205 774 1 20 - 2 Aris. 350 - 2 Aris. 350 - 2 Aris. 350 - 2 Aris. 360 4 1 - 1 Aris. 7 Aris. 454 221 7 3 84 3.329 4 155 - 5 Aris. 35 - 2 Aris. 350 - 2 Aris. 360 4 Aris. 37 58 7 7 Aris. 454 221 7 7 3 Aris 46 3 - 2 Aris. 37 58 7 7 Aris 46 3 - 2 Aris. 38 4 205 774 1 20 - 7 Aris. 454 32 27 3 - 3 Aris. 454 32 - 3 Aris	S. ATLANTIC	11,349	11,166	21	3,182	3,255	3	60	212	879
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	Del. Md	135	140	1	26	31	-	-	1	21
Va. 611 392 2 275 2265 1 5 9 448 W.Va. 57 13 $ 52$ 54 $ 1$ 1 33 N.C. 1288 771 10 417 409 1 2 125 76 S.C. 761 622 2 355 363 1 1 35 106 Ga. $2,940$ $2,786$ 3 $1,165$ $1,208$ $ 19$ 2 80 ES. CENTRAL $3,172$ $2,017$ 12 $1,90$ $1,205$ 7 2 64 137 Tenn. $1,259$ 821 8 315 354 6 $ 47$ 27 Ark. 423 264 $ 262$ 11 70 377 Ark. 413 264 $ 262$ 17 49 6 202 28 Otia. $1,711$	D.C.	788	608	1	233	138	-	30	15	- 324
	Va.	611	392	2	275	265	1	5	19	148
N.C. 1,268 //1 10 41/ 409 1 2 125 // Ga. 2,940 2,786 1 541 510 - 2 13 160 Fla. 3,921 5,258 3 1,165 1,208 - 19 2 80 E.S. CENTRAL 3,172 2,017 12 1,190 1,205 7 2 64 137 Ky. 67 41 2 285 301 1 1 9 37 Tenn. 1,259 821 8 315 354 6 - 47 27 Ala. 1,004 655 2 375 344 - 1 8 70 Miss. 842 500 - 215 206 3 3 W.S. CENTRAL 5,306 4,274 11 1,963 1,818 34 11 70 377 Ark. 443 264 - 262 189 26 - 17 42 La. 1,171 1,032 1 170 249 - 2 2 28 Okla. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 1,225 - 9 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont 1 - 22 11 - 4 4 40 Idaho 6 1 2 11 21 - 4 4 40 Idaho 6 1 2 11 21 - 4 4 40 Idaho 6 1 2 11 21 - 4 5 5 Mont 1 - 22 11 - 4 4 40 Idaho 6 1 2 11 21 - 4 6 2 Mont 1 - 22 11 - 4 4 40 Idaho 6 1 2 11 21 - 4 7 5 Mont 1 - 22 11 - 4 4 40 Idaho 6 1 2 11 21 - 4 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont 1 - 22 11 - 4 4 40 Idaho 6 1 2 11 21 4 4 40 Idaho 6 1 2 11 21 4 6 40 Idaho 6 1 2 11 21 4 6 40 Idaho 6 1 2 11 21 4 6 40 Idaho 6 1 2 11 21 4 7 5 N. Mex. 32 21 3 84 63 4 - 1 8 Ariz. 454 221 7 183 148 - 16 1 27 N. Mex. 32 21 3 84 63 4 - 1 8 Ariz. 454 221 7 183 148 - 16 1 27 VIah 8 13 4 32 27 3 - 3 9 Nev. 91 153 - 51 35 - 2 - 16 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 PACIFIC 2,369 3,407 37 3,068 2,872 - 99 2 2,215 Alaska 14 3 PACIFIC 2,369 3,407 37 3,068 2,872 - 99 2 2,215 Alaska 14 3 PACIFIC 2,369 3,407	W. Va.	57	13	-	52	54	:	1	1	33
Ga Dob DOS I DOS I DOS IOC Fla. 3,921 5,258 3 1,165 1,208 - 19 2 80 Fla. 3,921 5,258 3 1,165 1,205 7 2 64 137 Ky. 67 41 2 285 301 1 1 9 37 Tenn. 1,259 821 8 315 354 6 - 47 27 Ala. 1,004 655 2 375 344 - 1 8 70 Miss. 842 500 - 215 206 - - 73 Ark. 443 264 - 202 189 26 - 17 42 La. 1,171 1,032 1 170 249 - - 2 280 Okla. 177 83 <td< td=""><td>S.C.</td><td>1,268</td><td>622</td><td>10</td><td>417</td><td>409</td><td>1</td><td>2</td><td>125</td><td>106</td></td<>	S.C.	1,268	622	10	417	409	1	2	125	106
Fla.3,9215,25831,1651,208-19280E.S. CENTRAL3,1722,017121,1901,2057264137Tenn.1,25982183153546-4727Ala.1,0046552375344-1870Miss.842500-2152063W.S. CENTRAL5,3064,274111,9631,818341170377Ark.443264-26218926-1742La.1,1711,0321170249228Okla.1,778371471558245106Tex.3,5152,89531,3841,225-96202MOUNTAIN62847425415344131810162Mont1221440Idaho6121146Colo.3758727393-111N. Mex.3221384634-118MOUNTAIN6283,935443,5383,2941288238Myo6<	Ga.	2,940	2,786	1	541	510		2	13	160
E.S. CENTRAL3,1722,017121,1901,2057264137Ky.6741228530111937Tenn.1,25982183153546-4727Ala.1,00466552375344-1870Miss.842500-2152063W.S. CENTRAL5,3064,274111,9631,818341170377Ark.443264-26218926-1742La.1,1711,0321170249228Okla.1778371471558245105Tex.3,5152,88531,3841,225-96202MOUNTAIN62847425415344131810162Mont1-2211440Idaho612112-3-111Mont12112-399111127Mont1213864-1127Mont1333329412	Fla.	3,921	5,258	3	1,165	1,208	-	19	2	80
Ky. 67 41 2 285 301 1 1 9 37 Ala. 1,004 655 2 375 344 - 1 8 70 Miss. 842 500 - 215 206 - - - 3 W.S. CENTRAL 5,306 4,274 11 1,963 1,818 34 11 70 377 Ark. 443 264 - 262 189 26 - 17 42 La. 1,171 1,032 1 170 249 - - 2 28 Okla. 177 83 7 147 155 8 2 45 106 Mont. - 1 - 12 - - 4 40 Mont. - 1 2 11 21 - - 4 40 Idaho 6 1 2 1 31 43 32 7 3 - 3	E.S. CENTRAL	3,172	2,017	12	1,190	1,205	7	2	64	137
Ienn. 1,259 821 8 315 354 6 - 47 27 Ala. 1,004 655 2 375 344 - 1 8 70 Miss. 842 500 - 215 206 - - - 3 W.S. CENTRAL 5,306 4,274 11 1,965 1818 34 11 70 377 Ark. 443 264 - 262 189 26 - 17 42 La. 1,171 1,032 1 170 249 - - 2 28 Okla. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 1,225 - 9 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont. - 1 2 11 21 - - 46	Ky.	67	41	2	285	301	1	1	9	37
India 1,004 003 2 373 344 - 1 0 70 Miss. 842 500 - 215 206 - - - 3 W.S. CENTRAL 5,306 4,274 11 1,963 1,818 34 11 70 377 Ark. 443 264 - 262 189 26 - 17 42 La. 1,171 1,032 1 170 249 - - 2 28 Okla. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 1,225 - 9 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont. - 1 2 11 21 - - 46 Colo. 37 58 7 27 39 - 1 11 <t< td=""><td>lenn. Δla</td><td>1,259</td><td>821</td><td>8</td><td>315</td><td>354</td><td>6</td><td>-</td><td>47</td><td>27</td></t<>	lenn. Δla	1,259	821	8	315	354	6	-	47	27
W.S. CENTRAL 5,306 4,274 11 1,963 1,818 34 11 70 377 Ark. 443 264 - 262 189 26 - 17 42 La. 1,171 1,032 1 170 249 - - 2 28 Okla. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 1,225 - 9 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont. - 1 - 2 11 21 - - 4 400 Idaho 6 1 2 11 21 - - 4 40 Mont. 32 21 7 33 4 32 3 - 1 11 N.Mex. 32 21 7 183 148 - 16	Miss.	842	500	-	215	206	-		-	3
Ark. 443 264 - 262 189 26 - 17 42 La. 1,171 1,032 1 170 249 - - 2 28 Okla. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 1,225 - 9 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont. - 1 - 22 11 - - 4 40 Idaho 6 1 2 11 21 - - 4 40 Idaho 6 1 2 11 21 - - 46 60 11 27 Woo. - 6 2 5 - 3 - 16 1 27 N Mex. 32 271 3 84 3 27 39 3<	W.S. CENTRAL	5 306	A 27A	11	1 963	1 818	34	11	70	377
La. 1,171 1,032 1 170 249 - - 2 28 Okla. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 1,225 - 9 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont. - 1 - 22 11 - - 4 40 Idaho 6 1 2 11 21 - - - 56 Wyo. - 6 2 5 - 3 - 1 11 N.Mex. 32 211 3 84 63 4 - 1 8 Ariz. 454 221 7 183 148 - 16 1 27 Nex. 91 153 - 51 35 - 2 - 16 PAC	Ark.	443	264		262	189	26		17	42
Ukia. 177 83 7 147 155 8 2 45 105 Tex. 3,515 2,895 3 1,384 1,225 - 9 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont. - 1 2 211 - - 4 40 Idaho 6 1 2 11 21 - - 4 40 Idaho 6 1 2 11 21 - - 4 40 Idaho 6 2 5 - 3 - - 1 11 N.Mex. 32 21 7 83 48 - 16 1 27 Nev. 91 153 - 51 35 - 2 - 16 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash.	La.	1,171	1,032	1	170	249	-	-	2	28
MOLINTAIN 6,5,15 2,695 3 1,364 1,225 - 3 6 202 MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont. - 1 - 22 11 - - 4 400 Idaho 6 1 2 11 21 - - 4 400 Idaho 6 1 2 11 21 - - 4 400 Value 6 2 5 - 3 - - 1 11 Nex. 32 21 3 84 63 4 - 1 18 Ariz. 454 221 7 183 148 - 16 1 27 Value 91 153 - 51 35 2 - 16 28 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 <tr< td=""><td>Ukla. Tev</td><td>177</td><td>83</td><td>7</td><td>147</td><td>155</td><td>8</td><td>2</td><td>45</td><td>105</td></tr<>	Ukla. Tev	177	83	7	147	155	8	2	45	105
MOUNTAIN 628 474 25 415 344 13 18 10 162 Mont. - 1 - 22 11 - - 4 400 Idaho 6 1 2 11 21 - - 4 400 Valo - 6 2 5 - 3 - - 56 Vyo. - 6 2 5 - 3 - 1 11 N. Mex. 32 21 3 84 63 4 - 1 88 Ariz. 454 221 7 183 148 - 16 1 27 Utah 8 13 4 32 27 3 - 3 9 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash 252		3,515	2,095	3	1,304	1,220		9	0	202
idaho 6 1 2 11 21 - - - 5 Wyo. - 6 2 5 - 3 - - 46 Colo. 37 58 7 27 39 3 - 1 11 N. Mex. 32 21 3 84 63 4 - 1 88 Ariz. 454 221 7 183 148 - 16 1 27 Utah 8 13 4 32 27 3 - 3 9 PACIFIC 2.965 3.935 44 3.538 3.329 4 128 8 238 Wash. 252 338 4 205 174 1 20 - - 0reg. 102 178 2 94 105 - 4 1 1 1 Calif. 2.599 3.407 37 3.068 2.872 - 99 2 215 A	MOUNTAIN	628	474	25	415	344	13	18	10	162
Wyo. - 6 2 5 - 3 - - 46 Colo. 37 58 7 27 39 3 - 1 111 N. Mex. 32 21 3 84 63 4 - 1 81 Ariz. 454 221 7 183 148 - 16 1 27 Utah 8 13 4 32 27 3 - 3 9 Nev. 91 153 - 51 35 - 2 - 16 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 - - - Caiff. 2,589 3,407 37 3,068 2,872 - 99 2 215 Alaska 14 3 - 31 46 3 - - 22	Idaho	6	1	2	11	21		-	-	40
Colo. 37 58 7 27 39 3 - 1 11 N. Mex. 32 21 3 84 63 4 - 1 81 Ariz. 454 221 7 183 148 - 16 1 27 Utah 8 13 4 32 27 3 - 3 9 Nev. 91 153 - 51 35 - 2 - 16 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 - - Calif. 2,589 3,407 37 3,068 2,872 99 2 215 Alaska 14 3 - 31 46 3 - - 22 Hawaii 8 9 1 140 132 - 5 5 - -	Wyo.	-	6	2	5	-	3	-	-	46
Ariz. 32 21 3 84 63 4 - 1 8 Ariz. 454 221 7 183 148 - 16 1 27 Utah 8 13 4 32 27 3 - 3 99 Nev. 91 153 - 51 35 - 2 - 16 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 - - - Oreg. 102 178 2 94 105 - 4 1 1 1 Calif. 2,589 3,407 37 3,068 2,872 - 99 2 212 Alaska 1 1 140 132 - 5 5 - - 22 Hawaii 8 9 1 140 132 - -	Colo.	37	58	7	27	39	3	-	1	11
Utah 8 12 1 100 100 1 100 Nev. 91 153 - 51 35 - 2 - 16 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 - - Oreg. 102 178 2 94 105 - 4 1 1 Calif. 2,589 3,407 37 3,068 2,872 - 99 2 215 Alaska 14 3 - 31 46 3 - - 22 Hawaii 8 9 1 140 132 - 5 5 - Guam 2 4 - 33 60 - - - - P.R. 223 385 - 66 217 - - - 33 VI. 10 8 - 4 4 - - - - CN.M.L. 3 8 - 40 19 - <td< td=""><td>Ariz.</td><td>32 454</td><td>21</td><td>3</td><td>84 183</td><td>03 148</td><td>4</td><td>16</td><td>1</td><td>27</td></td<>	Ariz.	32 454	21	3	84 183	03 148	4	16	1	27
Nev. 91 153 - 51 35 - 2 - 16 PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 - - Oreg. 102 178 2 94 105 - 4 1 1 1 Calif. 2,589 3,407 37 3,068 2,872 - 99 2 215 Alaska 14 3 - 31 46 3 - - 222 14 Hawaii 8 9 1 140 132 - 5 5 - - - 223 385 - 66 217 - - - 33 14 4 - - - - 33 14 4 - - - - 33 15 <td>Utah</td> <td>8</td> <td>13</td> <td>4</td> <td>32</td> <td>27</td> <td>3</td> <td>-</td> <td>3</td> <td>- 9</td>	Utah	8	13	4	32	27	3	-	3	- 9
PACIFIC 2,965 3,935 44 3,538 3,329 4 128 8 238 Wash. 252 338 4 205 174 1 20 - 122 4 - 31 46 3 - - - - - 22 4 - 33 60 - - - - - - - - - - - - - - - - - - -	Nev.	91	153	-	51	35	-	2	-	16
Wash. 252 338 4 205 174 1 20 - - - Oreg. 102 178 2 94 105 - 4 1 1 Calif. 2,589 3,407 37 3,068 2,872 - 99 2 215 Alaska 14 3 - 31 46 3 - - 22 Hawaii 8 9 1 140 132 - 5 5 - Guam 2 4 - 33 60 - - - 33 V.I. 10 8 - 44 4 - - - 33 Amer. Samoa - - 11 7 - 1 - - - C.N.M.I. 3 8 - 40 19 - 4 - - -	PACIFIC	2,965	3,935	44	3,538	3,329	4	128	8	238
Oreg. 102 178 2 94 105 - 4 1 1 Calif. 2,589 3,407 37 3,068 2,872 - 99 2 215 Alaska 14 3 - 31 46 3 - - 22 Hawaii 8 9 1 140 132 - 5 5 - Guam 2 4 - 33 60 -	Wash. Orog	252	338	4	205	174	1	20	:	-
Alaska 14 3 - 31 46 3 - 22 218 Hawaii 8 9 1 140 132 - 5 5 - 22 Guam 2 4 - 33 60 - - - - - P.R. 223 385 - 66 217 - - - 33 VI. 10 8 - 4 4 - - - - Amer. Samoa - - 11 7 - 1 - -	Calif.	102	178	2	94	105	-	4	1	1 21=
Hawaii 8 9 1 140 132 - 5 5 - 6 Guam 2 4 - 33 60 - 33 00 - - - - - - - - - - - - - 33 00 - - - - - - 33 33 - - 66 217 - <td>Alaska</td> <td>14</td> <td>3,407</td> <td>-</td> <td>3,000</td> <td>2,072</td> <td>3</td> <td>33</td> <td>2</td> <td>215</td>	Alaska	14	3,407	-	3,000	2,072	3	33	2	215
Guam 2 4 - 33 60 - - - - - - - - - - - - - 33 60 - - - - - - - - - 33 7 - 1 - - 33 33 7 - 1 - - 33 33 8 - 11 7 - 1 - - - 33 33 60 - - - 33 33 33 33 33 33 33 33 34 - - 33 34 - - 33 34 - 11 7 1 1 34 - 34 34 - 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 <th1< th=""> <th1< th=""> <th1< th=""> <t< td=""><td>Hawaii</td><td>8</td><td>9</td><td>1</td><td>140</td><td>132</td><td>-</td><td>5</td><td>5</td><td></td></t<></th1<></th1<></th1<>	Hawaii	8	9	1	140	132	-	5	5	
P.R. 223 385 - 66 217 - - 33 V.I. 10 8 - 4 4 - - - 33 Amer. Samoa - - 11 7 - 1 - - - 33 C.N.M.I. 3 8 - 40 19 - 4 - - - - - - 33	Guam	2	4	-	33	60	-	-	-	
v.i. 10 8 - 4 4	P.R.	223	385	-	66	217	-	-	-	33
C.N.M.L 3 8 - 40 19 - 4	v.i. Amer. Samoa	10	8	-	4	4	-	-	-	-
	C.N.M.I.	3	8	-	40	19		4	-	-

TABLE II. (Cont'd.) Cases of specified notifiable diseases, United States, weeks ending September 22, 1990, and September 23, 1989 (38th Week)

U: Unavailable

								-							
		All Cau	ises, B	y Age (Years)		P&I**			All Cau	ises, B	y Age (Years)		P&I**
Reporting Area	All Ages	≥65	45-64	25-44	1-24	<1	Total	Reporting Area	All Ages	≥65	45-64	25-44	1-24	<1	Total
NEW ENGLAND	622	417	116	58	14	17	36	S. ATLANTIC	1,071	627	238	130	36	40	50
Boston, Mass.	168	92	38	24	6	8	9	Atlanta, Ga.	152	89	34	20	5	4	5
Bridgeport, Conn.	47	33	9	3	1	1	3	Baltimore, Md.	175	101	46	18	7	3	11
Fall River Mass	23	17	ź	1	-		2	Charlotte, N.C.	96	53	17	17	4	5	5
Hartford, Conn.	67	47	13	Ġ	1	-	8	Miami Fla	103	54	20	16	3	6	8
Lowell, Mass.	26	18	4	1	1	2	-	Norfolk, Va.	53	30	12	6		5	2
Lynn, Mass.	18	15	2	1	-	-	2	Richmond, Va.	67	44	14	8	1	-	5
New Bedford, Mass.	21	14	7	2	-	-	-	Savannah, Ga.	43	30	8	5	-	-	3
Providence P I	4/	33	0	5		2	2	St. Petersburg, Fla.	68	47	12	2	2	5	6
Somerville, Mass.	10	- 23	1	1		-	-	Mashington DC	/5	44	18	10	4	1	4
Springfield, Mass.	44	25	11	8	-		1	Wilmington Del	20	14	25	19	•	5	
Waterbury, Conn.	28	21	3	1	3	-	1	E C CENTRAL	755	470	170	-			20
Worcester, Mass.	74	57	7	6	-	4	8	Birmingham Ala	/55	4/0	1/2	66 16	21	26	36
MID. ATLANTIC	2,613	1,680	485	274	78	96	119	Chattanooga Tenn	64	38	19	3	4	1	5
Albany, N.Y.	55	37	9	3	2	4	-	Knoxville, Tenn.	85	63	12	7	ž	i	5
Allentown, Pa.	27	21	5	1	-	-	2	Louisville, Ky.	115	72	31	6	1	5	5
Camdon N I	150	98	33	15	1	3	5	Memphis, Tenn.	157	97	32	16	4	8	11
Elizabeth, N.J.	19	12	5	1	-	1	3	Mobile, Ala.	76	43	20	6	5	2	-
Erie, Pa.†	30	24	3	3	-		ž	Nashville Tenn	123	91	27	11	- 2	2	6
Jersey City, N.J.	47	29	10	6	-	2	-		123		21			~ ~	
N.Y. City, N.Y.	1,288	805	250	156	44	33	44	W.S. CENTRAL	1,721	1,047	374	188	70	41	72
Newark, N.J.	65	29	11	18	3	4	8	Baton Bouge La	39	25	9	10	4	1	5
Philadelphia Pa	393	224	83	28	19	20	21	Corpus Christi, Tex.	31	23	6	-	2	-	-
Pittsburgh, Pa.†	73	53	13	-4	1	2	6	Dallas, Tex.	217	126	51	24	11	5	3
Reading, Pa.	35	32	2	1	-		5	El Paso, Tex.	56	33	14	6	2	-	5
Rochester, N.Y.	114	85	16	11	1	1	10	Fort Worth, Tex	107	64	22	14	-	7	6
Schenectady, N.Y.	22	20	1	1	-	•	-	Houston, Tex.s	/34	436	169	89	24	10	18
Svracuse NY	114	19	15	3 7	1	-	1	New Orleans, La.	113	66	27	10	9	1	-
Trenton, N.J.	22	15	2	3	-	2	2	San Antonio, Tex.	158	103	31	13	6	5	14
Utica, N.Y.	14	11	2	ĩ	-	-	2	Shreveport, La.	57	37	7	9	2	2	10
Yonkers, N.Y.	42	33	6	2	1	-	3	Tulsa, Okla.	69	49	9	6	4	1	2
E.N. CENTRAL	2,300	1,512	457	182	59	90	91	MOUNTAIN	640	411	131	49	25	23	27
Akron, Ohio	48	39	5	2	1	1	-	Albuquerque, N. Me	x. 65	34	18	9	3	1	-
Canton, Ohio	48	39	8	.1	-	-	2	Colo. Springs, Colo.	35	25	5	3	2	-	2
Cinciggo, III.s	504 140	362	125	45	10	22	16	Las Venas Nev	89	60	16	4	7	1	ģ
Cleveland, Ohio	179	109	28	15	12	15	9 5	Ogden, Utah	22	15	3	1	-	3	2
Columbus, Ohio	172	103	44	15	4	6	3	Phoenix, Ariz.	156	96	34	11	4	11	7
Dayton, Ohio	112	75	26	4	4	3	ž	Pueblo, Colo.	28	19	5	3	1	-	3
Detroit, Mich.	271	153	56	39	12	11	12	Salt Lake City, Utah	35	20	.7	1	5	2	-
Evansville, Ind.	39	22	13	2	-	2	-	rucson, Ariz.	90	63	17	0	2	2	2
Garv. Ind.	18	12	1	23	1	1	2	PACIFIC Barbalau Calif	1,771	1,113	335	208	66	45	125
Grand Rapids, Mich.	53	37	10	5	-	i	ż	Frespo, Calif.	1/0	12	21	+	2	8	16
Indianapolis, Ind.	153	93	38	7	4	11	7	Glendale, Calif.	10	9		1	-		-
Madison, Wis.§	35	26	5	4	-	-	2	Honolulu, Hawaii	87	64	14	8	-	1	11
Milwaukee, Wis.	124	95	19	6	1	3	4	Long Beach, Calif.	78	51	11	10	3	3	12
Bockford III	30	20	6	4	1	4	4	Los Angeles Calif.	391	216	83	61	18	10	17
South Bend, Ind.	42	36	3	2	2	1	2	Oakland, Calif.	60	36	13	4	5	2	2
Toledo, Ohio	100	69	15	10	2	4	5	Portland Oreg	121	83	21	4 9	5	3	6
Youngstown, Ohio	66	52	13	1	-	-	5	Sacramento, Calif.	146	93	30	14	ĕ	3	14
W.N. CENTRAL	677	474	117	49	25	12	41	San Diego, Calif.	122	74	27	16	5	-	10
Des Moines, Iowa	58	48	6	3	1	-	4	San Francisco, Calif.	165	84	40	32	6	2	5
Duluth, Minn.	34	27	6	-	1	-	4	San Jose, Calif.	159	99	36	18	4	2	11
Kansas City, Kans.	22	14	6	2	2	-		Spokane Wash	161	112	22	16	3	ช 1	כ 7
Lincoln Nebr	125	84 2F	23	12	5	1	13	Tacoma, Wash	43	49	⊿	2	2	-	6
Minneapolis. Minn	101	25	15	3 8	2	1	4		10 170 **		2 425	1 004	20.4	200	507
Omaha, Nebr.	57	41	.9	3	3	i	3		12,170	1,151	2,425	1,204	394	390	59/
St. Louis, Mo.	124	75	32	8	3	6	ž								
St. Paul, Minn.	60	50	1	4	4	1	-								
vvicnita, Kans.	58	36	12	6	3	1	1								

TABLE III. Deaths in 121 U.S. cities,* week ending September 22, 1990 (38th Week)

*Mortality data in this table are voluntarily reported from 121 cities in the United States, most of which have populations of 100,000 or more. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included.

**Pneumonia and influenza.

Because of changes in reporting methods in these 3 Pennsylvania cities, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks.

††Total includes unknown ages.

\$Data not available. Figures are estimates based on average of past available 4 weeks.

						Ciga	rette bra	nd (%)						
Characteristic	No.	Marlboro	Winston	Salem	Kool	Newport	Benson and Hedges	Camel	Merit	Pall Mall	Vantage	Virginia Slims	Kent	Total
Self-reported use*						· · · · · ·					Ÿ			
Overall	4700	24.1	9.6	8.2	5.6	4.5	4.3	4.0	3.8	2.9	2.8	2.7	2.2	74.7
Sex														
Male	2348	28.2	11.5	6.5	6.0	4.9	2.9	6.5	3.6	3.4	2.8	0.3	1.9	78.4
Female	2352	19.4	7.5	10.1	5.1	4.1	5.9	1.3	4.1	2.4	2.9	5.3	2.5	70.4
Race [†]														
White	4125	26.2	9.9	7.4	3.7	2.3	3.8	4.4	4.4	3.1	3.2	2.7	2.4	73.5
Black	438	6.0	5.7	15.3	18.8	21.0	8.1	0.8	0.1	2.0	0.4	3.0	0.5	81.5
Age (yrs)														
17–24	587	54.4	4.0	3.3	4.6	10.5	4.3	2.7	1.9	0.0	0.6	2.4	0.2	88.7
25-44	2434	25.9	9.5	10.0	7.0	5.5	4.1	4.2	5.0	1.0	2.7	3.3	1.4	79.7
45-64	1264	8.7	12.7	7.8	4.4	0.5	5.0	4.3	2.7	7.1	4.0	1.9	4.0	63.0
≥65	415	6.8	10.4	7.5	2.8	0.0	3.7	4.4	3.1	6.3	4.1	2.0	4.7	55.8
Education (yrs)														
<12	954	24.4	11.1	7.9	5.3	5.5	4.0	3.9	1.4	3.8	3.1	2.0	1.0	73.4
12	1961	24.6	9.6	8.3	6.3	4.8	3.0	4.6	3.8	3.0	2.8	3.0	2.4	76.0
13–15	1140	24.0	7.2	9.1	5.2	4.0	6.9	3.2	5.3	1.8	2.4	2.8	2.6	74.5
≥16	645	21.4	8.8	7.5	4.3	1.4	6.2	3.5	8.4	1.7	2.6	3.4	4.1	73.5
Market share														
1986		23.0	11.2	7.8	6.3	3.8	4.4	2.6	4.0	0.6	3.2	2.9	2.8	72.6
1989 [§]		26.5	9.2	6.2	6.0	4.7	3.7	2.7	3.8	0.6	2.6	3.1	2.0	71.1

TABLE 1. Self-reported cigarette brand use among current cigarette smokers – Adult Use of Tobacco Survey, 1986; and cigarette market shares – 1986 and 1989

*Data weighted to represent the civilian, noninstitutionalized, adult (≥17 years of age) U.S. population.

[†]Excludes the racial category "other" (n = 137).

[§]Preliminary data.

Brand Use – Continued

Brand Use - Continued

Brand use varied by smoker's sex, race, and age. Differences by race in part reflected increased use of mentholated cigarettes by blacks (4,5). Fifty-five percent of all black smokers reported using one of three brands that were available only in mentholated form (Newport, Kool, and Salem). Fifty-four percent of smokers 17–24 years of age used Marlboro, more than twice the proportion in older age groups or the entire population (Table 1). The use of Merit and Kent varied directly with increasing level of education; in comparison, the use of Newport and Pall Mall varied inversely with level of education (Table 1).

Reported by: A Anderson, Case Western Reserve Univ School of Medicine, Cleveland, Ohio. Epidemiology Br, Office on Smoking and Health, Center for Chronic Disease Prevention and Health Promotion, CDC.

Editorial Note: Unlike market share data, the AUTS data allow analysis of brand use by sociodemographic variables or other characteristics. Overall, self-reported brand use from the AUTS is consistent with market share data for 1986 (Table 1) (3). Discrepancies between the sales-based and self-reported data may reflect differences in the number of cigarettes smoked by users of different brands, differences in brand use between current smokers and former smokers who had quit in 1986 before the AUTS, and errors in measurement (e.g., use by a smoker of more than one brand). The similarity in market shares between 1986 and 1989 (Table 1) (3) suggests that the self-reported 1986 data on brand use may also represent more recent cigarette use.

Factors that may affect smokers' use of a brand of cigarettes include cost, the "taste" of the cigarette, the perceived harmfulness of the cigarette, and the image of those who smoke a particular brand as projected through its advertising. Assessing sociodemographic differences among smokers by brand use and determining reasons for those differences may help in developing and targeting effective interventions for reducing smoking among specific population subgroups. For example, local surveys have found that the proportion of teenaged smokers who use Marlboro is substantially higher than the brand's market share (6,7) – a finding consistent with the AUTS data for persons aged 17-24 years. As a result, a school curriculum designed in California is being used in several states to counter the advertised image of Marlboro smokers as strong, rugged, and independent (8). The key component of the curriculum, a British documentary film entitled Death in the West, features six real cowboys in the American West who were dying from lung cancer or emphysema. Although 26.2% of white smokers used Marlboro, only 6.0% of black smokers used that brand; therefore, a health education program based on the Marlboro image may have a greater impact among whites than among blacks.

Several brands have been marketed primarily or exclusively to women (9); for example, Virginia Slims (used by 5.3% of female smokers) advertising promotes the image of the independent or "liberated" female smoker. However, more than one quarter of female smokers use either Marlboro (19.4%) or Winston (7.5%), which have been depicted primarily as "male brands"; some women may smoke "male brands" because of the implication of gender equality (10).

AUTS data show that 76% of blacks but only 23% of whites smoked mentholated brands (5). Increased understanding of why blacks use mentholated brands may assist in designing smoking-prevention and smoking-cessation interventions targeted to blacks.

AUTS data (5) also indicate that more highly educated smokers were more likely to use brands with a low-tar yield (\leq 15 mg per cigarette). This finding suggests that

Vol. 39 / No. 38

MMWR

Brand Use - Continued

this group may be more receptive to the message that the benefits of quitting substantially exceed the benefits of switching from high- to low-tar brands (11,12).

By tracking trends in use of brands of cigarettes, the role of cigarette advertising in smoking initiation may be more clearly understood. For example, recent advertising campaigns for Camel cigarettes featuring the "Old Joe" dromedary cartoon character may "reposition" the brand into a younger population (13). An increase in the use of Camel cigarettes by young persons, particularly teenagers, would suggest that the Camel advertising campaign is stimulating the recruitment of new smokers. CDC's 1989 Teenage Attitudes and Practices Survey will provide national data on use of brands of cigarettes and smokeless tobacco among persons 12–18 years of age who use such products.

References

- 1. CDC. US Department of Health and Human Services report of the 1986 Adult Use of Tobacco Survey. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1989.
- 2. CDC. Cigarette smoking in the United States, 1986. MMWR 1987;36:581-5.
- Maxwell JC. The Maxwell consumer report: preliminary year end 1989 report. Richmond, Virginia: Wheat First Securities, 1989.
- Cummings KM, Giovino GA, Mendicino AJ. Cigarette advertising and black-white differences in brand preferences. Public Health Rep 1987;102:698–701.
- CDC. Reducing the health consequence of smoking: 25 years of progress a report of the Surgeon General, 1989. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1989; DHHS publication no. (CDC)89-8411.
- Hunter SM, Croft JB, Burke GL, Parker FC, Webber LS, Berenson GS. Longitudinal patterns of cigarette smoking and smokeless tobacco use in youth: the Bogalusa Heart Study. Am J Public Health 1986;76:193–5.
- 7. Goldstein AO, Fischer PM, Richards JW, Creten D. Relationship between high school student smoking and recognition of cigarette advertisements. J Pediatr 1987;110:488–91.
- 8. Glantz SA. Death in the West curriculum project [Letter]. N Y State J Med 1985;85:470-1.
- 9. Ernster VL. Mixed messages for women: a social history of cigarette smoking and advertising. N Y State J Med 1985;85:335–40.
- 10. Davis RM. Current trends in cigarette advertising and marketing. N Engl J Med 1987;316: 725–32.
- CDC. The health consequences of smoking: the changing cigarette a report of the Surgeon General, 1981. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1981; DHHS publication no (PHS)81-50156.
- 12. Davis RM, Healy P, Hawk SA. Information on tar and nicotine yields on cigarette packages. Am J Public Health 1990;80:551–3.
- Richards JW Jr, Fischer PM. Smokescreen: how tobacco companies market to children. World Smoking and Health 1990;15:12–4.

Progress in Chronic Disease Prevention

Evaluation of an Employee Smoking Policy – Pueblo, Colorado, 1989–90

In December 1988, the Colorado Department of Health and CDC were asked to help evaluate a planned worksite policy banning employee smoking for the Colorado State Hospital, a psychiatric hospital in Pueblo, Colorado. Purposes of the evaluation were to 1) determine whether implementation of the policy reduced the exposure of hospital employees to environmental tobacco smoke (ETS) in the workplace; 2) assess the acceptance of the policy among employees; and 3) assess the effect of the policy

Employee Smoking Policy - Continued

on the smoking behavior of employees who smoked. This report presents findings from employee surveys at three time periods: before, and at 3 and 12 months after policy implementation.

Before February 1, 1989 (the day the policy was implemented), employees were allowed to smoke in designated areas within the hospital. After February 1, smoking by employees was prohibited indoors; hospitalized patients were permitted to continue smoking in designated areas on patient-care wards.

Self-administered questionnaires were distributed to all 1400 hospital employees in January (before the policy change) and May 1989 and in February 1990. The questionnaire asked employees to provide information about their exposure to ETS at work. Other questions elicited attitudes and opinions about the new hospital smoking policy.

The questionnaires were analyzed as cross-sectional samples of the hospital work force. A cohort analysis was done of 73 smokers who voluntarily identified themselves on the questionnaire and responded to the two follow-up surveys; this analysis permitted assessment of individual behavioral changes. All analyses were stratified by smoking status.

"Ever smokers" were defined as persons who had smoked \geq 100 cigarettes in their lifetimes, including both current smokers (who continued to smoke at the time of the surveys) and former smokers who did not smoke. "Never smokers" were defined as persons who had smoked <100 cigarettes in their lifetimes. Smokers were asked how many cigarettes they smoked during work hours and in a 24-hour day.

In January 1989, 1032 (74%) employees responded to the questionnaire; in May 1989, 762 (54%) employees responded, and in February 1990, 745 (53%) employees responded to the follow-up survey. Age, sex, and ethnicity of respondents to each survey were similar to the demographic distribution of the entire hospital workforce (Colorado State Personnel Office, unpublished data).

In January 1989, before the employee smoking ban took effect, 41.5% of employees reported working in a smoke-free work area. In May, 3 months after the ban, 72.1% reported their work area was smoke-free (p<0.01, chi-square test); in February 1990, 80.5% reported their work area was smoke-free. The percentage of employees reporting smoke-free worksites did not vary by smoking status.

From January 1989 to February 1990, overall employee support for the smoking ban increased from 59% to 68%, respectively (p<0.01, controlled for smoking status, Mantel-Haenszel chi-square test); the greatest change occurred among former smokers. Support for the ban was greatest among never smokers and least among current smokers (Table 1).

The reported prevalence of current smoking varied little during the evaluation. In January 1989, 29% of respondents were current smokers, compared with 24% in May and 25% in February 1990. Among the cohort of 73 smokers, the average daily number of cigarettes smoked at work declined from 7.7 in January 1989 to 4.2 in February 1990; during the same period, however, the number of cigarettes smoked after work increased from 8.6 to 10.3. The net average change in cigarettes smoked in a 24-hour day declined by 1.8 cigarettes, from 16.3 to 14.5.

Reported by: GS Mayo, Colorado State Hospital, Pueblo; JA Pritzl, Colorado Dept of Administration; WF Young, RE Hoffman, MD, State Epidemiologist, Colorado Dept of Health. Program Svcs Activity, Office on Smoking and Health, Center for Chronic Disease Prevention and Health Promotion; Div of Field Svcs, Epidemiology Program Office, CDC.

Employee Smoking Policy – Continued

Smoking <u>Tot</u> status No.		Janu	ary 1989		May 1989				February 1990			
	otal	Support policy		Total		Support policy		Total		Support policy		
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Current*	300	(29)	60	(20)	184	(24)	39	(21)	187	(25)	45	(24)
Former [†]	334	(32)	234	(70)	260	(34)	211	(81)	237	(32)	194	(82)
Never⁵	398	(39)	317	(80)	318	(42)	272	(86)	321	(43)	269	(84)
Total	1032	(100)	611	(59)	762	(100)	522	(69)	745	(100)	508	(68)

TABLE 1. Opinions expressed by employees about worksite smoking ban before and 3 months and 12 months after implementation of the ban, by cigarette smoking status – Colorado State Hospital, 1989–90

*Smoked ≥100 cigarettes and continued to smoke at the time of the surveys.

[†]Smoked ≥100 cigarettes in their lifetimes but did not smoke at the time of the surveys.

[§]Smoked <100 cigarettes in their lifetimes.

Editorial Note: Smoke-free worksite policies decrease the exposure of nonsmokers to ETS (1). The American Medical Association, the American Hospital Association, and other groups have advocated smoke-free hospitals (2,3). However, psychiatric hospitals present special challenges to administrators attempting to prevent the exposure of employees and patients to ETS through the creation of smoke-free hospital environments. The prevalence of smoking among psychiatric patients appears to be substantially higher than among the general population (4), and the concept of the smoke-free psychiatric facility has not yet been widely accepted by hospital administrators and staff (5). For these reasons, policies that restrict smoking in psychiatric facilities have been difficult to enact. However, smoke-free policies for psychiatric hospitals should benefit patients served by these facilities in ways other than reducing risk for smoking-related disease. For example, patients who are smokers may require higher doses of therapeutic drugs than do patients who are nonsmokers (6), and some psychiatric patients may be at increased risk for fatal and nonfatal injuries from fire caused by cigarettes (7).

This evaluation indicates that employee acceptance of smoking restrictions can be sustained in a psychiatric facility, even after being in place 12 months. These findings are similar to those reported in other worksites (\mathcal{B}). Because inpatients were permitted to smoke indoors, approximately 20% of employees reported exposures to ETS at the worksite after policy implementation. Additional studies of smoke-free policies that benefit both patients and staff are under way at this facility. Through a combination of employee education and cooperation of all management levels, worksite policies can be implemented with minimal conflict and enforcement difficulty (\mathcal{G}).

In Colorado, only modest short-term changes in smoking behavior (e.g., fewer cigarettes smoked at work) occurred among current smokers, but these were partially offset by an increase in smoking after working hours. Long-term changes in the smoking practices of employees may produce health and economic benefits for smoking and nonsmoking employees, as well as for employers. *References*

- 1. CDC. Passive smoking: beliefs, attitudes, and exposures United States, 1986. MMWR 1988; 37:239–41.
- Ernster VL, Wilner SI. Nonsmoking policies in hospitals. J Public Health Policy 1985;6: 197–203.
- 3. Becker DM, Conner HF, Waranch R, et al. The impact of a total ban on smoking in the Johns Hopkins Children's Center. JAMA 1989;262:799–802.

Employee Smoking Policy – Continued

- 4. Hughes JR, Hatsukami DK, Mitchell JE, Dahlgren LA. Prevalence of smoking among psychiatric outpatients. Am J Psychiatry 1986;143:993–7.
- Resnick M, Bosworth EE. A survey of smoking policies in Oregon psychiatric facilities. Hosp Community Psychiatry 1988;39:313–5.
- Vinarova E, Vinar O, Kalvach Z. Smokers need higher doses of neuroleptic drugs. Biol Psychiatry 1984;19:1265–8.
- 7. Gaston EH. Solving the smoking problem on a chronic ward. J Psychiatr Treatment and Evaluation 1982;4:387-401.
- 8. Borland R, Owen N, Hill D, Chapman S. Changes in acceptance of workplace smoking bans following their implementation: a prospective study. Prev Med 1990;19:314–22.
- 9. Millar WJ. Smoke in the workplace and evaluation of smoking restrictions. Ottawa: Health and Welfare Canada, Minister of Supply and Services, 1988.

Cessation of Cigarette Smoking – United States, 1989

Smoking-initiation and smoking-cessation interventions are important in reducing the prevalence of cigarette smoking in the United States. However, progress in smoking cessation has varied appreciably by smokers' age, race, sex, educational attainment, and state of residence (1,2). To monitor progress in smoking cessation in relation to these factors, data from the 1989 Behavioral Risk Factor Surveillance System (BRFSS) were analyzed.

In 1989, health departments from 39 states and the District of Columbia participated in the BRFSS, a monthly random-digit-dialed telephone interview survey of adults aged \geq 18 years, to obtain information on selected health behaviors (3). Respondents were asked if they had ever smoked at least 100 cigarettes and if they currently smoked. The "quit ratio" was the percentage of ever smokers who were former smokers when interviewed. Ratios were weighted to represent the adult population of each participating state. To compare quit ratios between states, the weighted state-specific ratios were standardized for the age, race, sex, and educational attainment of the 1980 U.S. population. Quit ratios for subgroups (age, race, sex, and educational attainment) were standardized by adjusting for the other three variables.

The weighted quit ratio varied from 43% in Kentucky to 59% in Montana (median: 51%), and the standardized quit ratio from 41% in Oklahoma to 55% in Hawaii (Table 1). In general, standardized ratios were lowest in states in the Ohio River Valley and the south and highest in states in the Rocky Mountain and mid-central regions (Figure 1). The standardized quit ratio was also greater in persons >35 years of age, whites, men, and persons with high school education or more (Table 2).

Reported by: the following state BRFSS coordinators: L Eldridge, Alabama; J Contreras, Arizona; W Wright, California; M Adams, Connecticut; A Peruga, District of Columbia; S Hoecherl, Florida; J Smith, Georgia; A Villafuerte, Hawaii; J Mitten, Idaho; B Steiner, Illinois; S Joseph, Indiana; S Schoon, Iowa; K Bramblett, Kentucky; J Sheridan, Maine; A Weinstein, Maryland; L Koumjian, Massachusetts; J Thrush, Michigan; N Salem, Minnesota; J Jackson-Thompson, Missouri; M McFarland, Montana; S Spanake, Nebraska; K Zaso, L Powers, New Hampshire; L Pendley, New Mexico; J Marin, O Munshi, New York; C Washington, North Carolina; M Maetzold, North Dakota; E Capwell, Ohio; N Hann, Oklahoma; J Grant-Worley, Oregon; C Becker, Pennsylvania; R Cabral, Rhode Island; M More, South Carolina; S Moritz, South Dakota; D Ridings, Tennessee; J Fellows, Texas; L Post-Nilson, Utah; J Bowie, Virginia; K Tollestrup, Washington; D Porter, West Virginia; M Soref, Wisconsin. Office of Surveillance and Analysis, Div of Chronic Disease Control and Community Intervention, and Office on Smoking and Health, Center for Chronic Disease Prevention and Health Promotion, CDC.

Smoking Cessation - Continued

		Wei	ghted qu	it ratio	Standardized [§] quit ratio			
State	No.	Rank	%	95% CI	Rank	%	95% CI	
Alabama	695	35	45.0	±4.1	35	43.8	±4.1	
Arizona	743	14	51.7	±4.2	20	47.0	±4.4	
California	1017	6	55.1	±3.3	15	47.9	±3.6	
Connecticut	747	12	52.8	±4.1	10	48.5	±4.3	
District of Columbia	566	39	43.8	±5.0	9	48.5	±8.7	
Florida	887	10	53.8	±3.6	19	47.0	±3.8	
Georgia	720	28	47.9	±4.1	16	47.5	±3.9	
Hawaii	390	11	52.8	±3.8	1	55.2	±5.6	
ldaho	723	9	54.0	±4.1	5	52.3	±4.3	
Illinois	874	27	48.5	±3.6	29	45.6	±3.6	
Indiana	1068	34	45.0	±3.1	34	43.8	±3.1	
lowa	615	18	51.0	±4.5	12	48.2	±4.0	
Kentucky	909	40	43.0	±3.7	39	42.1	±3.2	
Maine	685	20	50.7	±3.9	18	47.3	±4.0	
Maryland	758	19	50.9	±4.1	17	47.4	±4.1	
Massachusetts	648	3	56.9	±4.2	4	52.3	±4.2	
Michigan	1178	30	46.5	±3.2	36	43.5	±3.1	
Minnesota	1674	2	57.4	±2.5	8	49.3	±2.7	
Missouri	710	25	49.0	±4.1	33	44.1	±4.2	
Montana	577	1	59.0	±4.3	2	54.3	±4.7	
Nebraska	634	21	50.5	±4.2	26	46.0	±4.2	
New Hampshire	756	5	55.3	±3.9	7	50.3	±4.2	
New Mexico	567	8	54.5	±4.4	11	48.2	±4.7	
New York	633	17	51.2	±4.7	25	46.0	±4.2	
North Carolina	832	31	45.9	±4.0	28	45.7	±3.7	
North Dakota	739	4	56.5	±3.9	3	52.7	±3.9	
Ohio	694	32	45.3	±4.3	32	45.0	±3.9	
Oklahoma	562	37	44.6	±4.6	40	41.2	±4.4	
Oregon	855	7	55.1	±3.6	13	48.2	±4.0	
Pennsylvania	917	29	47.0	±3.5	27	46.0	±3.6	
Rhode Island	922	22	49.9	±3.5	30	45.5	±3.1	
South Carolina	826	36	44.8	±3.8	31	45.1	±3.4	
South Dakota	730	24	49.4	±3.8	22	46.6	±3.8	
Tennessee	1129	38	44.0	±3.2	37	42.6	±3.1	
Texas	655	16	51.3	±4.3	14	48.1	±4.7	
Utah	588	15	51.5	±4.6	6	50.4	±4.6	
Virginia	686	26	48.9	±4.3	23	46.3	±4.3	
Washington	744	13	52.6	±3.8	21	46.9	±4.1	
West Virginia	879	33	45.1	±3.9	38	42.4	±3.7	
Wisconsin	656	23	49.7	±4.1	24	46.2	±4.0	

TABLE 1. Quit ratio* of ever smokers, by state[†] – Behavioral Risk Factor Surveillance System (BRFSS), 1989

*The percentage of ever smokers (those who had ever smoked ≥100 cigarettes) who were former smokers when interviewed.

[†]For the BRFSS, the District of Columbia is considered a state.

⁵Standardized for the distribution of the 1980 U.S. population by age, race, sex, and educational attainment.

[®]Confidence interval.

Smoking Cessation - Continued

FIGURE 1. Smoking quit ratios* in selected states[†], by tercile – Behavioral Risk Factor Surveillance System (BRFSS), 1989



*The percentage of ever smokers (those who had ever smoked ≥100 cigarettes) who were former smokers when interviewed.

[†]For the BRFSS, the District of Columbia is considered a state.

		Standardize	ed⁺ quit ratio	_
Characteristic	No.	%	95% Cl⁵	
Age (yrs)				
18–34 [¶]	9,440	32.3	±1.5	
35–54	11,843	43.5**	±1.5	
≥55	9,905	64.5**	±1.5	
Race				
Black [¶]	2,461	39.1	±2.8	
White	28,727	47.0**	±0.9	
Sex				
Female [¶]	16,073	43.3	±1.2	
Male	15,115	49.8**	±1.2	
Education (yrs)				
<12¶	5,688	36.0	±1.8	
12	11,424	43.2**	±1.3	
>12	14,076	55.9**	±1.3	

TABLE 2. Quit ratio* of ever smokers, by age, race, sex, and educational attainment – Behavioral Risk Factor Surveillance System, 1989

*The percentage of ever smokers (those who had ever smoked ≥100 cigarettes) who did not smoke at the time of the survey.

[†]Standardized by adjusting for other sociodemographic variables in the 1980 U.S. population (e.g., age was standardized for race, sex, and educational attainment).

[§]Confidence interval.

Referent group.

**Quit ratio is significantly higher than the referent group (p<0.05).

Smoking Cessation - Continued

Editorial Note: The differences between states in the weighted quit ratio can be explained only in part by state-specific differences in age, race, sex, and educational attainment of the populations, since these differences persisted after standardization for differences in sociodemographic composition. Other factors affecting smoking cessation that may explain the variations in smoking cessation by state include the percentage of heavy smokers (1), societal norms and attitudes about smoking cessation services (4). Restrictions on smoking also may play a role in the variations by state in smoking cessation (1). In general, states with the lowest quit ratios have the highest prevalence of current cigarette smoking (2).

Concerns about the health effects of smoking (5) and the occurrence of smokingrelated illnesses (6) may contribute to the higher quit ratios for persons aged >35years. Because continuing smokers are less likely than former smokers to survive to older ages, this differential mortality contributes to the higher quit ratios observed for older age groups (7). In addition, the higher quit ratios for older than for younger age groups may represent a longer opportunity to quit.

Findings in this and other reports (8) show that blacks were less likely than whites to be former smokers regardless of educational attainment. Limited use of established smoking cessation programs by blacks contributes to these racial differences (9). Nonetheless, trend data suggest that the rate of increase in the quit ratio since 1974 has been similar for whites and blacks (1,7).

Although men were more likely than women to be former smokers, the rate of increase in quit ratios over time has been similar for men and women (1,7). This finding is consistent with a diffusion phenomenon (i.e., quitting activity adopted initially by men that later diffused into the female population where it follows a pattern similar to that for men). Additionally, more men than women who quit cigarette smoking begin using cigars, pipes, or snuff or chewing tobacco (7). Thus, differences in smoking cessation by sex are smaller when use of other forms of tobacco are considered (7).

Greater difficulty in quitting among persons of low socioeconomic status may contribute to the lower quit ratios among persons with high school education or less (1). These and other findings suggest that smoking cessation interventions should target younger persons and persons of low socioeconomic status. In addition, such interventions should be aimed at blacks, who in general have a lower rate of smoking cessation than do whites (10).

Continued efforts are essential to motivate smokers to quit. Growth in tobacco-use prevention and control coalitions, which bring together a broad range of persons and organizations with the common goal of reducing the prevalence of tobacco use (11), will likely strengthen smoking cessation efforts by fostering a social climate that motivates smokers to quit. The American Stop Smoking Intervention Study, a planned 7-year project of the National Cancer Institute and the American Cancer Society, will substantially increase resources for tobacco control coalitions in the United States (12) and may accelerate progress in smoking cessation.

References

 CDC. Reducing the health consequences of smoking: 25 years of progress – a report of the Surgeon General. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1989; DHHS publication no. (CDC)89-8411.

Smoking Cessation - Continued

- 2. CDC. Regional variation in smoking prevalence and cessation: behavioral risk factor surveillance, 1986. MMWR 1987;36:751-4.
- 3. Remington PL, Smith MY, Williamson DF, Anda RF, Gentry EM, Hogelin GC. Design, characteristics, and usefulness of state-based behavioral risk factor surveillance: 1981–87. Public Health Rep 1988;103:366–75.
- CDC. Smoking and health: a national status report. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1988; DHHS publication no. (CDC)87-8395.
- 5. Pechacek TF, Danaher BG. How and why people quit smoking: a cognitive-behavioral analysis. In: Kendall PC, Hollon SD, eds. Cognitive-behavioral interventions: theory, research and procedures. Vol 21. New York: Academic Press, 1979.
- 6. Ockene JK, Hosmer D, Rippe J, et al. Factors affecting cigarette smoking status in patients with ischemic heart disease. J Chronic Dis 1985;38:985–94.
- CDC. The health benefits of smoking cessation: a report of the Surgeon General, 1990. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1990; DHHS publication no. (CDC)90-8416.
- 8. Novotny TE, Warner KE, Kendrick JS, Remington PL. Smoking by blacks and whites: socioeconomic and demographic differences. Am J Public Health 1988;78:1187–9.
- Orleans CT, Schoenbach VJ, Salmon MA, et al. A survey of smoking and quitting patterns among black Americans. Am J Public Health 1989;79:176–81.
- 10. Fiore MC, Novotny TE, Pierce JP, et al. Methods used to quit smoking in the United States: do cessation programs help? JAMA 1990;263:2760–5.
- 11. CDC. State coalitions for prevention and control of tobacco use. MMWR 1990;39:476-7, 483-5.
- McKenna J, Carbone E. Huge tobacco control project begun by NCI, ACS. J Natl Cancer Inst 1989;81:93–4.

Perspectives in Disease Prevention and Health Promotion

Smoking-Attributable Mortality – Kentucky, 1988

Smoking is the single most important preventable cause of death in the United States (1). Among states participating in the Behavioral Risk Factor Surveillance System (BRFSS), Kentucky has consistently ranked at or near the top in prevalence of smoking (2). In 1988, the BRFSS indicated that 34% of adults in Kentucky were current smokers, compared with a median prevalence of 24% for all states surveyed (3). To better characterize the public health burden of smoking in Kentucky, the Kentucky Department for Health Services recently estimated smoking-attributable mortality (SAM) and years of potential life lost (YPLL) in that state during 1988. This report summarizes results from that analysis.

SAM and YPLL were calculated using SAMMEC II (Smoking-Attributable Mortality, Morbidity, and Economic Costs) computer software (4). Calculations were made for 22 smoking-related diseases among adults aged \geq 35 years (Table 1). The analysis also included smoking-related burn fatalities for persons of all ages and four perinatal conditions related to maternal smoking (5). Age- and sex-specific mortality data for 1988 were obtained from the state's vital records system. Age- and sex-specific smoking prevalence rates for 1988 were obtained from the state's BRFSS. YPLL were calculated to life expectancy using 1985 data from CDC's National Center for Health Statistics (6).

The smoking-attributable fraction (SAF) was derived from age- and sex-specific relative risks of death (based on the American Cancer Society's Cancer Prevention Study II [1]) and prevalence data for current and former smokers from the 1988

Smoking Mortality – Continued

Cause of death (ICD-9-CM rubric)	Age group (yrs)	No. deaths	Crude SAF [†]	SAM
Neoplasms				
Lip, oral cavity, pharynx (140–149)	≥35	120	0.80	96
Esophagus (150)	≥35	102	0.79	81
Pancreas (157)	≥35	345	0.28	96
Larynx (161)	≥35	70	0.83	58
Trachea, bronchus, lung (162)	≥35	2,718	0.86	2,338
Cervix uteri (180)	≥35	101	0.31	31
Urinary bladder (188)	≥35	177	0.42	74
Kidney, other unspecified urinary organs (189)	≥35	148	0.35	52
Cardiovascular diseases				
Rheumatic heart disease (390–398)	≥35	54	0.17	9
Hypertensive disease (401–404)	≥35	396	0.19	77
lschemic heart disease (410–414)	≥35	8,393	0.24	2,034
Pulmonary circulation disease (415–417)	≥35	249	0.21	52
Other heart disease (420–429)	≥35	3,637	0.20	734
Cerebrovascular disease (430–438)	≥35	2,546	0.19	496
Atherosclerosis (440)	≥35	430	0.41	177
Aortic aneurysm (441)	≥35	235	0.50	118
Other arterial disease (442–448)	≥35	128	0.43	55
Respiratory diseases				
Respiratory tuberculosis (010–012)	≥35	28	0.29	8
Pneumonia, influenza (480–487)	≥35	1,324	0.28	367
Chronic bronchitis, emphysema (491–492)	≥35	300	0.82	246
Asthma (493)	≥35	54	0.28	15
Chronic airway obstruction (496)	≥35	1,132	0.82	924
Perinatal conditions				
Short gestation/low birth weight (765)	<1	70	0.21	15
Respiratory distress syndrome (769)	<1	36	0.19	7
Other respiratory condition of fetus and newborn (770)	<1	27	0.22	6
Sudden infant death syndrome (798.0)	<1	121	0.15	18
Other conditions				
Burn deaths (E890–E899)	>0	105	0.45	47
All other causes	>0	14,246	0.00	0
Total	>0	37,292	0.22	8,230

TABLE 1. Estimated smoking-attributable mortality (SAM),* by cause – Kentucky, 1988

*Total SAM was calculated by multiplying the number of deaths in each disease category by the specific smoking-attributable fraction (SAF). Because of rounding, SAM may not equal the product of SAF times the number of deaths.

[†]Derived from age- and sex-specific relative risks of death (based on the American Cancer Society's Cancer Prevention Study II [1]) and prevalence data for current and former smokers from the 1988 Behavioral Risk Factor Surveillance System.

Smoking Mortality - Continued

BRFSS. Total SAM was calculated by multiplying the number of deaths in each disease category by the specific SAF. Total smoking-attributable YPLL was calculated by multiplying the age-specific SAM by YPLL for each premature death.

In 1988, 8230 deaths in Kentucky were attributable to smoking, accounting for 22% of all deaths in the state during the year. Fifty-three percent of smoking-attributable deaths were from lung cancer and ischemic heart disease (Table 1). Sixty-eight percent of SAM occurred among men (Table 2). Sixty-seven percent of deaths occurred in persons \geq 65 years of age. However, when smoking-attributable deaths were calculated as a percentage of total deaths, persons aged 45–64 years had a higher percentage of deaths caused by smoking than did persons aged \geq 65 years (Figure 1). For men aged 55–64 years, 41% of all deaths were attributable to smoking. When considered as a separate cause of death, SAM was the most common cause of death in men, the third most common cause in women, and, for both sexes, the second most common cause in Kentucky (Table 2).

TABLE 2. Deaths from selected causes, including smoking, by sex – Kentucky, 1988

	М	ale	Fer	nale	Total		
Underlying cause of death	No.	(%)	No.	(%)	No.	(%)	
Diseases of the heart*	4,950	(25.1)	5,305	(30.2)	10,255	(27.5)	
Smoking-attributable mortality	5,589	(28.4)	2,642	(15.0)	8,230	(22.1)	
Malignant neoplasms*	2,665	(13.5)	2,950	(16.8)	5,615	(15.1)	
Cerebrovascular diseases*	746	(3.8)	1,404	(8.0)	2,150	(5.8)	
Unintentional injuries*	1,201	(6.1)	552	(3.1)	1,753	(4.7)	
Influenza and pneumonia*	431	(2.2)	581	(3.3)	1,012	(2.7)	
All other causes*	4.132	(21.0)	4.144	(23.6)	8,276	(22.2)	
Total	19,714	(100.0)	17,578	(100.0)	37,292	(100.0)	

*Excludes smoking-attributable deaths.





Smoking Mortality – Continued

In 1988, 115,458 YPLL before life expectancy in Kentucky were attributable to smoking. Fifty-five percent of smoking-attributable YPLL occurred in persons aged <65 years. The mean YPLL was 14 years per smoking-attributable death.

Reported by: R Finger, MD, State Epidemiologist, Dept for Health Svcs, Kentucky Cabinet for Human Resources. JM Shultz, PhD, Dept of Epidemiology and Public Health, Univ of Miami School of Medicine, Miami, Florida. Program Svcs Activity, Office on Smoking and Health, Center for Chronic Disease Prevention and Health Promotion; Div of Field Svcs, Epidemiology Program Office, CDC.

Editorial Note: This analysis quantifies the premature mortality caused by smoking in a state with a historically high prevalence of tobacco use. The high prevalence of smoking among middle-aged persons in Kentucky (38.4% among those aged 35–49 years and 34.9% among those aged 50–64 years) (2) is of special concern. The data indicate a need to intensify cessation efforts among these persons before the onset of chronic diseases associated with smoking. *The Health Benefits of Smoking Cessation: A Report of the Surgeon General, 1990,* describes the important reductions in risk that may be associated with smoking cessation at any age (7).

To reduce the burden of SAM in Kentucky, greater efforts are also necessary to prevent smoking among young persons. During the 1990 legislative session in Kentucky, the legislature enacted a law prohibiting the sale of tobacco products to all persons <16 years of age. This law also established fines for vendors who sell tobacco products to persons aged <16 years and requires that signs stating the age limit for purchase of tobacco be posted at the point of sale. Enforcement of laws such as this is critical to reducing tobacco use (8).

SAMMEC II software can be used to estimate the effects of smoking and has been distributed to all 50 states and the District of Columbia. Additional state-specific estimates may be made using this software to provide public health workers and policymakers with important updated information regarding the impact of smoking in their respective states (9).

References

- CDC. Reducing the health consequences of smoking: 25 years of progress a report of the Surgeon General. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1989; DHHS publication no. (CDC)89-8411.
- Anda RF, Waller MN, Wooten KG, et al. Behavioral risk factor surveillance, 1988. In: CDC surveillance summaries, June 1990. MMWR 1990;39(no. SS-2):1–21.
- 3. CDC. Cigarette smoking-Behavioral Risk Factor Surveillance System, 1988. MMWR 1989; 38:845-8.
- Shultz JM, Novotny TE, Rice DP. SAMMEC II: computer software and documentation. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, CDC, April 1990.
- 5. McIntosh ID. Smoking and pregnancy: attributable risks and public health implications. Can J Public Health 1984;75:141–8.
- 6. NCHS. Vital statistics of the United States, 1985: life tables. Hyattsville, Maryland: US Department of Health and Human Services, Public Health Service, CDC, 1988; DHHS publication no. (PHS)88-1108.
- CDC. The health benefits of smoking cessation: a report of the Surgeon General, 1990. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1990; DHHS publication no. (CDC)90-8416.
- 8. CDC. State laws restricting minors' access to tobacco. MMWR 1990;39:349-53.
- CDC. State-specific estimates of smoking-attributable mortality and years of potential life lost-United States, 1985. MMWR 1988;37:689-93.

The Morbidity and Mortality Weekly Report is prepared by the Centers for Disease Control, Atlanta, Georgia, and is available on a paid subscription basis from the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402, (202) 783-3238.

The data in this report are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the succeeding Friday. Accounts of interesting cases, outbreaks, environmental hazards, or other public health problems of current interest to health officials, as well as matters pertaining to editorial or other textual considerations should be addressed to: Editor, Morbidity and Mortality Weekly Report, Mailstop C-08, Centers for Disease Control, Atlanta, Georgia 30333; telephone (404) 332-4555.

Director, Centers for Disease Control William L. Roper, M.D., M.P.H. Director, Epidemiology Program Office Stephen B. Thacker, M.D., M.Sc.



Editor, *MMWR* Series Richard A. Goodman, M.D., M.P.H. Managing Editor Karen L. Foster, M.A.

☆U.S. Government Printing Office: 1990-731-103/22024 Region IV

Public Health Service HEALTH AND HUMAN SERVICES DEPARTMENT OF Atlanta, Georgia 30333 Penalty for Private Use \$300 Official Business Centers for Disease Control ŝ HBE Z × POSTAGE & FEES PAID FIRST-CLASS MAIL Permit No. G-284 PHS/CDC