



Published in final edited form as:

Microb Drug Resist. 2013 June ; 19(3): 175–184. doi:10.1089/mdr.2012.0177.

Antimicrobial resistance genes in multi-drug resistant *Salmonella enterica* serovars isolated most frequently from animals, retail meat, and humans in the U.S. and Canada

LaShanda M. Glenn¹, Rebecca L. Lindsey¹, Jason P. Folster², Jean M. Whichard², Gary Pecic², Patrick Boerlin³, Mathew W. Gilmour⁴, Patrick F. McDermott⁵, Heather Harbottle⁵, Paula J. Fedorka-Cray¹, and Jonathan G. Frye^{1,*}

¹Bacterial Epidemiology and Antimicrobial Resistance Research Unit, Agricultural Research Service, U.S. Department of Agriculture, Athens, Georgia ²Centers for Disease Control and Prevention, Atlanta, Georgia ³Ontario Veterinary College, Guelph, ON, Canada ⁴Public Health Agency of Canada, Winnipeg MB, Canada ⁵Center for Veterinary Medicine, U.S. Food and Drug Administration, Laurel, Maryland

Abstract

Salmonella enterica is a prevalent food-borne pathogen which can carry multi-drug resistance (MDR) and could pose a threat to human health. Identifying the genetic elements associated with MDR in *Salmonella* isolated from animals, food, and humans can help determine the sources of MDR in food animals and their impact on human health. Representatives of MDR *S. enterica* serovars most frequently isolated from healthy animals, retail meat, and human infections in the U.S. and Canada were subjected to detailed genetic analysis (n=56). These included U.S. slaughter (n=12), retail (n=9), and human (9) isolates, and Canadian slaughter (n=9), retail (n=9), and human (n=8) isolates. These isolates were assayed by microarray for antimicrobial resistance and MDR plasmid genes. Genes detected encoded resistance to aminoglycosides (alleles of *aac*, *aad*, *aph*, *strA/B*); beta-lactams (*bla*_{TEM}, *bla*_{CMY}, *bla*_{PSE-1}); chloramphenicol (*cat*, *flo*, *cmlA*); sulfamethoxazole (*sulI*); tetracycline (*tet*(A, B, C, D) and *tetR*); and trimethoprim (*dfrA*). Similar resistance genes were detected regardless of serovar, source, or location. Hybridization with IncA/C plasmid gene probes indicated that 27/56 isolates carried a member of this plasmid family; however these plasmids differed in several highly variable regions. Cluster analysis based on genes detected separated most of the isolates into two groups, one with IncA/C plasmids and one without IncA/C plasmids. Other plasmid replicons were detected in all but one isolate, and included I1 (25/56), N (23/56) and FIB (10/56). The presence of different mobile elements along with similar resistance genes suggest that these genetic elements may acquire similar resistance cassettes, and serve as multiple sources for MDR in *Salmonella* from food animals, retail meat, and human infections.

*Corresponding author. Bacterial Epidemiology and Antimicrobial Resistance Research Unit, USDA, Agricultural Research Service, Richard B. Russell Research Center, 950 College Station Rd., Athens, GA 30605. USA. Phone: (706) 546-3677. Fax: (706) 546-3172. jonathan.frye@ars.usda.gov.

Note: The mention of trade names or commercial products in this manuscript is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture, CDC, FDA, or PHAC.

Keywords

Salmonella; multi-drug resistance; antimicrobial resistance; plasmid

Introduction

Salmonella enterica is a leading cause of food borne illness worldwide.^{2,5,6,12,29} *Salmonella* can be transferred to humans through contaminated water and food products, leading to food-borne illness. In most cases, salmonellosis in an otherwise healthy individual results in a self-limiting infection that does not require antimicrobial treatment. Invasive *Salmonella* infections are more severe and require antimicrobial therapy.^{2,8} Antimicrobial resistant *Salmonella* are a concern, as infections caused by these microorganisms may be more difficult to treat compared to their susceptible counterparts.⁸ Both the U.S. and Canada developed initiatives to monitor the dissemination of antimicrobial resistance in *Salmonella* and other enteric pathogens isolated from animals, food, and humans.¹⁸ The National Antimicrobial Resistance Monitoring System (NARMS) was developed in 1996 to monitor U.S. antimicrobial resistance trends in human diagnostic isolates (Centers for Disease Control and Prevention, CDC), food animals (U.S. Department of Agriculture, Agricultural Research Service, USDA-ARS) and retail meats (U.S. Food and Drug Administration, Center for Veterinary Medicine, FDA-CVM) (<http://www.fda.gov/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/NationalAntimicrobialResistanceMonitoringSystem/default.htm>). In Canada, the Canadian Integrated Program for Antimicrobial Resistance Surveillance (CIPARS) was developed in 2002 to monitor similar antimicrobial resistance trends (<http://www.phac-aspc.gc.ca/cipars-picra/index-eng.php>). In these programs, isolates are collected and tested for resistance to antimicrobials used in both human and veterinary medicine.

Multi-drug resistant (MDR) *Salmonella* which are resistant to two or more classes of antimicrobials, are frequently encountered and may reduce the effectiveness of treatments.²¹ In 2007, 13% of Canadian *Salmonella* isolated from chickens and swine during slaughter (abattoir) and 3% isolated from retail chicken were resistant to five or more antimicrobials and consequently MDR. Almost 10% of *Salmonella* isolated in Canada from humans between 2004 and 2006 also exhibited antimicrobial resistance (<http://www.phac-aspc.gc.ca/cipars-picra/index-eng.php>). In 2007, 10% of *Salmonella* isolated in the U.S. from chicken, turkey, cattle and swine slaughter isolates were resistant to five or more antimicrobials while 5.6% of retail meat isolates exhibited similar resistances. In U.S. human isolates, 6.9% were resistant to five or more antimicrobials (<http://www.fda.gov/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/NationalAntimicrobialResistanceMonitoringSystem/ucm209340.htm>).

Plasmids are often associated with antimicrobial resistance in *Salmonella* and may also contain additional genes that provide heavy metal resistance, sanitizer resistance, or that aid in virulence and environmental adaptability.^{8,21,24,29,30} These plasmids are small, circular pieces of DNA that are often self-transmissible to other bacteria through conjugation. The transfer of plasmids from one bacterial cell to another results in the horizontal transfer of

genetic material, allowing acquisition of multiple genes by the recipient.^{10,27,30} Plasmids with the IncA/C replicon have been linked to multiple drug resistance in *Salmonella enterica* associated with food animals as well as the fish pathogen *Yersinia ruckeri* and *Y. pestis*, the causative agent of plague.⁴² In addition to IncA/C, several other plasmid replicons have been associated with MDR *Salmonella*, including B/O, HI1, HI2, I1, N, F, and P many of which are found to be co-resident in some MDR *Salmonella*.³⁷

This study was conducted to identify the antimicrobial resistance genes and genetic elements associated with MDR in *Salmonella* isolated from humans, animals, and retail meat by NARMS and CIPARS. Genetic analyses included detection of antimicrobial resistance and MDR plasmid genes by DNA microarrays and PCR detection of plasmid replicons.^{9,23} The detailed analysis of a collection of *Salmonella* representative of the most frequently isolated MDR serovars, identified antimicrobial resistance and MDR plasmid genes present in U.S. and Canadian isolates. Cluster and linkage analysis of this data was used to identify isolates with significantly similar profiles of antimicrobial resistance and plasmid genes from the different surveillance programs. Several genetic commonalities such as the presence of IncA/C plasmids were detected in MDR *Salmonella* isolated from different sources including humans, retail meat, and animals at slaughter. Comparable MDR genetics were also identified in isolates from similar animal, meat, and human sources sampled by the U.S. and Canadian surveillance programs.

Materials and Methods

Isolate selection, culture conditions, and antimicrobial susceptibility testing

Salmonella enterica chosen for this study were isolated from various sources by the U.S. Food Safety and Inspection Service (slaughter isolates, n=12), the U.S. Food and Drug Administration, Center for Veterinary Disease (retail isolates, n=9), the Centers for Disease Control and Prevention (human isolates, n=9), the Canadian Integrated Program for Antimicrobial Resistance Surveillance (slaughter isolates, n=9; retail isolates, n=9) and the Public Health Agency of Canada (human isolates, n=8). Isolates were obtained, cultivated, maintained, and stored as frozen stock cultures using standard methods. Culture media was obtained from Difco™ (Becton Dickinson and Company, Sparks, MD). All *Salmonella* isolates were subjected to testing via the Sensititre™ semi-automated antimicrobial susceptibility system following the manufacturers instructions (TREK Diagnostic Systems, Inc., Westlake, OH) for susceptibility to amikacin, gentamicin, kanamycin, streptomycin, ampicillin, amoxicillin-clavulanic acid, ceftiofur, ceftriaxone, cefoxitin, sulfamethoxazole/sulfisoxazole, trimethoprim-sulfamethoxazole, chloramphenicol, ciprofloxacin, nalidixic acid and tetracycline. Serovars with the highest frequency of MDR for each source agency were determined (MDR defined as resistance to two or more classes of antimicrobials). Isolates were then chosen from these serovars that were resistant to the maximum number of antimicrobial compounds (described below). Prevalence data, detailed isolation and testing methods available in U.S. NARMS reports: <http://www.fda.gov/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/NationalAntimicrobialResistanceMonitoringSystem/ucm209340.htm>, and Canadian CIPARS reports: <http://www.phac-aspc.gc.ca/cipars-picra/pubs-eng.php>

Microarray design and construction

The DNA microarrays and methods used for this analysis have been previously reported and validated.^{23,31,32} Briefly, the DNA microarrays consisted of 1267 gene probes. Of those probes, 775 were designed to detect antimicrobial resistance genes found in the National Center for Biotechnology Information (NCBI) database. The other 487 gene probes were designed to detect plasmid genes identified in six strains of bacteria.^{31,32} IncA/C gene probes on the microarray were designed from plasmid sequences identified in *Yersinia ruckeri* str. YR71 pYR1, *Y. pestis* biovar Orientalis str. IP275 pIP1202, *Photobacterium damsela* subsp. *piscicida* pP99-018, *Salmonella enterica* subsp. *enterica* serovar Newport str. SL254 pSN254, *Photobacterium damsela* subsp. *piscicida* pP91278 and *Escherichia coli* p1658/97. Gene sequences from the plasmid pHCM1 found in *S. enterica* subsp. *enterica* serovar Typhi str. CT18 were used to design probes for the genes in this IncHI1 plasmid. The microarrays were constructed by spotting probes onto Corning UltraGAPS amino-silane coated slides (Corning Inc., Life Sciences, Acton, MA) with an Qarray mini robot (Genetix, Hampshire, UK) and post processed following the manufacturers recommendations (Corning, Inc.) as previously described.^{31,32}

DNA isolation, labeling, microarray hybridization, and scoring

Total DNA from the isolates was prepared with the GenElute Bacterial Genomic DNA kit (Sigma, St Louis, MO) following instructions for Gram-negative bacteria from 5 mL of overnight cultures grown in LB broth at 37°C with shaking as previously described. Cy5 dye-labeled dCTP (Amersham, Piscataway, NJ), Klenow fragment (New England Biolabs, Beverly, MA) and random primers were used to label *Salmonella* genomic DNA overnight in 37°C water bath.²³ Labeled DNA was purified using Qiagen PCR clean-up kit (Qiagen, Valencia, CA), and hybridized to the microarray over night at 42°C in a Corning hybridization chamber as previously described. Microarrays were washed following the manufacturers protocol for hybridization with formamide buffer (Corning, Inc.). Slides were scanned using ScanArray Lite microarray analysis system and images were analyzed using the ScanArray Express software version 1.1 (Packard BioChip Technologies, Billerica, MA). Positive hybridizations were determined and scored as previously described.^{23,32}

Plasmid replicon typing

Three panels of multiplex PCR were used to determine the presence of 18 plasmid replicons commonly found in *Enterobacteriaceae*. PCR primers and parameters were previously described.^{9,30}

Cluster analysis of isolates

Relationships between isolate based on gene content was determined by hierarchical cluster analysis based on antimicrobial resistance and plasmid genes detected by microarray analysis. Open source CLUSTER 3.0 was used for this analysis with Euclidean distance for gene content.^{13,16} Dendrogram was viewed using Java TreeView version 1.1.4r3³⁸ (<http://jtreeview.sourceforge.net>).

Statistical analysis

Linkage disequilibrium (LD) was calculated as an extension of Fisher's exact probability test on contingency tables³⁹ as instituted by the program Arlequin¹⁷. Standard settings were used, 10,000 steps in the Markov chain and 1,000 dememorization steps; and calculations of D , D' , and r^2 coefficients were made with a significance level of 0.05.

Results

MDR in U.S. NARMS and Canadian CIPARS *Salmonella enterica* isolates selected for the study

The *Salmonella enterica* serovars chosen for the study were the most frequently isolated MDR serovars found in each source. For U.S. NARMS this was *Salmonella* serovars Typhimurium, Heidelberg, and Newport for both human (CDC1 – CDC9) and animal (USDA1 – USDA12) isolates. U.S. retail meat isolates (FDA1 – FDA9) were chosen from *Salmonella* serovars Typhimurium, Heidelberg, and Hadar. Isolates from Canadian CIPARS were serovars Typhimurium, Heidelberg, and Enteritidis from humans (CH1 – CH8), Typhimurium, Heidelberg, Hadar, and Saintpaul, from animals (CA1 – CA9), and serovars Typhimurium, Heidelberg and Hadar from retail meat (CR1 – CR9). Isolates resistant to the greatest number of antimicrobial compounds were chosen from those *Salmonella* serovars for each agency's collection from that source. The numbers of antimicrobial compounds the isolates were resistant to range in number from three compounds to 13, with an average of resistance to seven antimicrobials (Table 1 and Figure 1).

Detection of antimicrobial resistance and plasmid genes by microarray analysis

Resistance genes consistent with the observed phenotypes were detected in the isolates. A summary of the genes detected most frequently in the isolates is in Figure 1 (complete hybridization data available in supplemental table S1); an exception was isolate CH1, where only a *strA/B* gene was detected. The most prevalent aminoglycoside resistance genes detected were alleles of *aac(3)* (24/56), *aac(6')* (22/56), *aadA* (34/56), *aadB* (27/56), *aadE* (14/56), *aph* (21/56), and *strAB* (39/56). Detected β -lactam resistance associated genes included *bla_{TEM}* (32/56), *bla_{CMY-2}* (30/56), *bla_{PSE-1}* (36/56), and the β -lactamase resistance transcriptional regulatory gene *ampR* (5/56). Chloramphenicol resistance genes included *cat* (36/56), *floR* (27/56) and *cmlA* (7/56). Sulfamethoxazole resistance genes in the isolates include *sulI* (26/56) and *sulII* (23/56). Frequently detected tetracycline resistance associated genes were *tet(A)* (45/56), *tet(B)* (8/56), *tet(C)* (8/56), *tet(D)* (7/56) and the regulatory gene *tetR* (50/56). Also detected in several isolates were alleles of trimethoprim resistance genes, such as *dfr* (18/56).

Many of the isolates hybridized to multiple probes for IncA/C and HI1 plasmid genes. Twenty-seven isolates hybridized to a large number of IncA/C plasmid genes including genes in core regions used to define the IncA/C plasmid backbone (Table 1).^{31,42} Detection of these core regions, based upon sequence data from the pYR1 and pSN254 plasmids⁴², indicate that IncA/C plasmids are likely present in these 27 isolates. Eight of nine U.S. human isolates had IncA/C plasmid genes detected, three of which contained genes in all 12 core regions. The remaining isolates were missing genes in core regions 6–7, 6–9 or 6–10,

which have been identified as a region of insertions and deletions (indels) in the family of IncA/C plasmids.^{24,31,32,42} Three of nine U.S. retail meat isolates contained IncA/C plasmid genes but none of the isolates contained genes in all 12 core regions. One isolate, FDA9, is missing genes in region 10, another, FDA8, is missing regions 7–10 and another isolate, FDA2, is missing genes in regions 7–12. Four of the 10 U.S. animal isolates that contain IncA/C plasmids hybridized to genes in all 12 core regions. The remaining six isolates are missing genes in either regions 1–2, 7–9, or region 10–12. Five of 8 Canadian isolates from humans likely contain IncA/C plasmids. Three of the Canadian IncA/C positive human isolates contained genes in all 12 core regions and two isolates were missing genes in the indel at regions 6–9 or 7–9. Canadian Human Isolate CH1 had a different pattern of core IncA/C genes detected than all isolates in that group, with IncA/C regions 1, 4, 6–9, and 11 being missing. None of the Canadian retail meat or animal isolates had IncA/C genes detected in a large enough number to indicate the presence of IncA/C plasmids. However, one of the Canadian retail meat isolates, CR2, was the only isolate in the study with numerous IncHI1 plasmid genes detected (183/206), most likely indicating the presence of an HI1 plasmid.²⁴

Plasmid replicon typing

Replicons were detected in all isolates except Canadian animal isolate CA4. The IncA/C (27/56), I1 (25/56), N (23/56) and FIB (10/56) were the most prevalent replicons detected among all isolates (Figure 1 and Table 1). The IncA/C replicon was detected in all of the 27 isolates where large numbers IncA/C genes (>52) were detected by microarray analysis, confirming the presence of IncA/C plasmids in these isolates. These included U.S. human, retail and slaughter isolates, as well as Canadian human isolates. However, the IncA/C replicon was not detected in Canadian retail or animal slaughter isolates. Plasmid replicons I1 and N were widespread and detected in isolates from all sources in both countries. Replicons FII, FIA, P, F, and HI1 were present in fewer isolates. Most isolates (n=51) contained one or two replicon types, three isolates were positive for three replicon types and a single isolate was positive for four replicons. Overall, half of these 18 replicon types associated with *Enterobacteriaceae* were detected in at least one of the isolates.⁹

Cluster and LD analysis

Almost all of the *Salmonella* isolates fell into two major groups based upon the antimicrobial resistance and plasmid genes detected. These were group A (n=27) with numerous IncA/C plasmid genes detected and group B (n=27) with very few IncA/C genes detected (Figure 1 and Table 1). Two isolates, CR2 and CA2, were an exception to this and did not group together or with any of the other isolates. For CR2, this was due to the detection of an almost complete HI1 plasmid making its genes detected unique from all other isolates in the study. While isolate CA2 had several IncA/C plasmid genes detected, it did not have enough to cluster with group A, and also was not PCR positive for the IncA/C replicon. However, CA2 had too many IncA/C genes detected to cluster with group B and thus was a lone isolate. All 27 isolates of group A were IncA/C positive via microarray analysis as well as plasmid replicon typing. CH1, an isolate from a human in Canada, was different from other isolates in group A. This isolate did not hybridize with as many IncA/C gene probes as other isolates in group A, but five of the core regions were detected (2, 3, 5,

10, and 12) as was the IncA/C replicon by PCR analysis (Table 1). Group A consists mostly of U.S. slaughter, U.S. human and Canadian human isolates, although some U.S. retail meat isolates were present in this cluster. In addition to IncA/C, replicons I1, N and FIB were detected in group A. Cluster B contained 27 isolates in which IncA/C was not detected by microarray analysis or replicon typing. This cluster consists mostly of Canadian retail, Canadian animal slaughter and U.S. retail meat isolates. However, U.S. slaughter, U.S. human and Canadian human isolates were present in lower numbers. Replicons I1, FIB, N, P, FIA, and FII were also frequently detected in this cluster.

Linkage disequilibrium (LD) analysis detected significant linkage between the isolates' plasmid replicon, serovar, source agency, and animal source (cattle, chicken, swine, or turkey and their retail meat products, and also human isolates) (Figure 2). Significant linkage often corresponded with the separation of isolates into groups A and B by the cluster analysis of the microarray data. Most notably IncA/C had significant linkage with serovar, agency, and animal source, which are reflected in clusters A and B. All six *Salmonella* Newport isolates were in group A and contained IncA/C; while none of the six *Salmonella* Hadar isolates contained IncA/C and thus were in group B, leading to LD between IncA/C and these serovars. Significant positive LD was also detected for IncA/C to human isolates, with 14 out of 17 human isolates in group A; likewise, IncA/C was linked to cattle isolates, with all five in group A. The animal or animal meat product from which the isolate was collected also demonstrated a negative association with IncA/C. Only one of the 12 swine isolates and one of nine turkey isolates contained IncA/C plasmids and were in group A, while 10 of the remaining swine and all eight remaining turkey isolates were found in group B. These linkages suggest that the groups formed by clustering isolates based upon antimicrobial and plasmid genes are significantly different from each other and reflect gene content associated with specific replicons, serovars, agencies, and animal sources.

Animal and IncI1 were linked due to eight out of nine turkey isolates having the IncI1 replicon detected; conversely, only four out of 12 swine isolates had the IncI1 replicon. IncI1 also had a negative relationship with IncN attributable to only one of 23 IncN positive isolates also had the IncI1 replicon detected. Replicon FIB showed linkage with serovar, agency, and animal due to eight isolates being Typhimurium, only one isolate was from a U.S. agency (USDA), and six were from swine. Replicon FIB was also detected in eight of the isolates found in group B, and thus had a negative association with IncA/C and group A. However, some linkages are less clear due to the small number of isolates with the replicon detected. Linkages with IncF, FIA, FII, and P were detected, but are suspect due to only one or two of these replicons being detected in all of the isolates. For example, IncP is in disequilibrium with serovar and IncN, but only one P replicon was detected which is not a sampling large enough to determine if this association is representative of the bacterial population. Animal and agency also showed significant LD, which is likely an artifact due to no Canadian isolates were available from beef or cattle for this study.

Discussion

To investigate MDR in *Salmonella enterica* isolated from animals, retail meat, and infected humans, a small sampling of the most frequently isolated MDR serovars in the U.S. and

Canada were analyzed to determine genetic mechanisms associated with their antimicrobial resistance. Overall, many of the antimicrobial resistance genes detected in the study have been previously found in MDR isolates from the U.S. and Canada in other studies using similar methods or other complementary methods.^{24,30,31,42} The large number of genes detected in this study and their combination with plasmid genes and plasmid replicon typing allows a better understanding of MDR in *Salmonella* isolates from these sources. Cluster analysis based on the genes detected divided the isolates into two large clusters: A, with IncA/C plasmids and B, without IncA/C plasmids. This reveals the impact of IncA/C plasmids on MDR in U.S and Canadian isolates, while also implicating other genetic elements that are also associated with MDR in *Salmonella*.

The resistance genes detected within group A and group B were shared at a high level, indicating the presence of common mechanisms. Antimicrobial resistance genes usually found in IncA/C plasmids (e.g. *bla*_{CMY-2}, *floR*, *aac(3)*, *aadA*, *aphA1*, *strA/B*, *sull*, *sullII*, *dfrA*, *tet(A)*, *tet(B)*, *tet(C)*, *tet(D)*, and *tetR*) were detected in most isolates of group A.^{24,30,31,42} Isolates in group B had resistance genes indicative of integrons like SGI-1 or other MDR plasmids (e.g. *bla*_{PSE-1}, *bla*_{TEM}, *floR*, *aac(3)*, *aadA*, *sull*, *dfrA*, *tet(A/G)*, and *tetR*).^{24,30,35} In Addition, many of the resistance genes found in isolates from both group A and B were also similar, in spite of the different genetic vehicles for these MDR genes.^{24,30,35} This may reflect the cassette nature of many of these resistance genes, where the genes encoding MDR can be transferred to plasmids, to integrons, and/or to the chromosome resulting in different genetic elements carrying similar resistance genes. For example, many IncA/C plasmids and a variety of *Salmonella* Genomic Island-1 (SGI-1) sequences have been shown to share resistance genes cassettes.^{24,30} In addition, IncA/C plasmids have been shown to mobilize genetic elements such as SGI-1 and antimicrobial resistance gene cassettes *in-trans*.¹⁴ This type of movement could explain many of the genes found in common throughout these MDR isolates.

Certain resistance genes were also found to be widespread among these isolates and between the two clusters. For example, the *bla*_{CMY-2} gene that encodes resistance to extended spectrum cephalosporins and other beta-lactams was found in isolates from both cluster A and B.^{11,21,22} MDR IncA/C plasmids that carry the *bla*_{CMY-2} gene have been reported and characterized as a major contributor to MDR in *Salmonella* from the U.S. and Canada.^{3,11,21,22} In IncA/C positive group A, 23/27 isolates have the *bla*_{CMY-2} gene detected. However, 7/27 isolates in group B also have the *bla*_{CMY-2} gene detected. Five of these are *Salmonella* Heidelberg isolates which also have the IncII plasmid replicon detected. IncII plasmids carrying a *bla*_{CMY-2} gene have recently been found in *Salmonella* Heidelberg human isolates in the U.S.^{19,20} Therefore, it is likely that there are two or more genetic elements carrying the *bla*_{CMY-2} genes in MDR *Salmonella* from in these isolates. The prevalence of MDR *Salmonella* resistant to cephalosporins due to *bla*_{CMY-2} encoding IncA/C plasmids has declined recently in animal isolates; however, the spread of a *bla*_{CMY-2} gene by other genetic elements, such as IncII, could cause increased prevalence of *Salmonella* resistant to extended spectrum cephalosporins in the future.

A large amount of variability was found in the core backbone regions of the IncA/C plasmids detected in group A. Most of the regions missing are previously described indels

within regions 6–10, 1–2 and 7–12.^{24,31,32,42} The genes encoded in these indels have been observed to have effects on a great diversity of plasmid functions. Some of these include: transmissibility of the plasmids, mobilization of other genetic elements, phenotypes conferred by the plasmids, response to selective pressures including heavy metals, chemical sanitizers, and traditional antimicrobials, as well as phylogenetic implications.^{24,31,32,42} These aspects of the IncA/C plasmid family have been well characterized in other studies of *Salmonella* as well as other bacteria containing IncA/C plasmids.^{24,31–33,41,42} Interestingly, while IncA/C is associated with almost half of the MDR *Salmonella* isolates in this study, there is a large amount of diversity within the structure of the plasmids in these isolates, possibly indicating several different phylogenetic origins of the IncA/C plasmids in these *Salmonella* isolates and even within a single serovar of *Salmonella*. This suggests that there are multiple sources of MDR IncA/C *Salmonella* in these isolates rather than the spread of a single IncA/C plasmid among the different MDR *Salmonella* found in animals, retail meat, and human infections. Alternatively, the plasmids may have spread some time ago and then undergone mutations and deletions within certain serovars or in certain environments resulting in the diversity we have observed.

Other plasmids that could potentially carry MDR genes were detected in each of the large clusters, including IncI1 which has been previously shown to carry *bla*_{CMY-2} or MDR in bacteria isolated from humans and animals.^{19,20,25,26} IncN, IncFIB, and other plasmid replicons were also detected demonstrating that the genetic elements carrying MDR, while dominantly associated with IncA/C and I1, are complex and could be associated with other plasmids and genetic elements such as integrons like *Salmonella* Genomic Island 1 (SGI1).^{24,30,35} This observation is also supported by the two lone MDR isolates that are genetically distinct from the two large clusters, including one carrying an IncHI1 plasmid which is often associated with MDR in Typhoid and Paratyphoid serovars of *Salmonella*.^{36,40}

Within both large clusters, isolates which grouped together into sub-clusters also had other similarities. The isolates were often the same serovar when isolated from the same type of animal or animal meat, or when isolated by the same agency or in the same country. This was especially true for human isolates which clustered most closely to other human isolates and in some cases human isolates from the U.S. and Canada clustered together. However, human isolates did not cluster with animal or meat isolates. The clustering did show that 14 out of 17 human isolates were associated with IncA/C, a relationship between source and IncA/C that exhibits significant LD. This likely reflects an association between MDR human salmonellosis and *Salmonella* serovars, such as Newport, Heidelberg, Typhimurium, and others that carry a plasmid from the MDR IncA/C family.^{19,20,24} This association has been described in several reports on MDR human isolates collected during this time period (2005–2007), and emphasizes the impact of IncA/C plasmids on U.S. and Canadian human MDR *Salmonella* infections.^{4,15,19,20,24} U.S. animal MDR *Salmonella* isolates were also found to be mostly IncA/C positive (10/12), as were three of the U.S. retail meat isolates. In contrast all Canadian animal and retail meat isolates lacked the IncA/C plasmid.^{7,21,22,24,28,30} Canadian monitoring of cattle and beef had previously detected MDR

Salmonella with the IncA/C plasmid, however sampling of cattle and beef in Canada was discontinued before this study.

Human infections caused by MDR *Salmonella* are usually thought to be foodborne, with animals serving as reservoirs of resistance and retail meats acting as a vector for human disease.^{1,2,43} Agencies in the U.S. and Canada have developed programs to monitor the levels of antimicrobial resistance in *Salmonella* found in food animals, retail meats, and humans. This study was designed to investigate MDR *Salmonella* collected by these programs and to determine the genetic elements associated with antimicrobial resistance. Genetic elements, such as IncA/C and IncI1 plasmids were found in most of the isolates, but other plasmids and genetic elements were also associated with MDR in many of these *Salmonella*. It was also observed that IncA/C was found in most human MDR isolates from both the U.S. and Canada. These results suggest that in addition to phenotypic monitoring of antimicrobial resistance in *Salmonella*, investigations of the genetic elements causing resistance is necessary to understand the cause, prevalence, and spread of MDR and determine the impact of food borne *Salmonella* on human health.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank Jennifer Turpin, Georgina Hidalgo, Jovita Haro, Benny Barrett, and Takiyah Ball at USDA for technical assistance. From CIPARS we thank Lucie Dutil for isolate selection, and Laura Martin and Emily Weir for DNA isolation. We would also like to dedicate this manuscript in memory of our colleague, Lucie Dutil, whom we had the privilege to know and work with through the NARMS and CIPARS collaborations.

Reference List

1. Alcaine SD, Soyer Y, Warnick LD, Su WL, Sukhnanand S, Richards J, Fortes ED, McDonough P, Root TP, Dumas NB, Grohn Y, Wiedmann M. Multilocus sequence typing supports the hypothesis that cow- and human-associated *Salmonella* isolates represent distinct and overlapping populations. *Appl Environ Microbiol.* 2006; 72:7575–7585. [PubMed: 17028236]
2. Alcaine SD, Warnick LD, Wiedmann M. Antimicrobial resistance in nontyphoidal *Salmonella*. *J Food Prot.* 2007; 70:780–790. [PubMed: 17388077]
3. Allen KJ, Poppe C. Occurrence and characterization of resistance to extended-spectrum cephalosporins mediated by beta-lactamase CMY-2 in *Salmonella* isolated from food-producing animals in Canada. *Can J Vet Res.* 2002; 66:137–144. [PubMed: 12146884]
4. Andrysiak AK, Olson AB, Tracz DM, Dore K, Irwin R, Ng LK, Gilmour MW. Genetic characterization of clinical and agri-food isolates of multi drug resistant *Salmonella enterica* serovar Heidelberg from Canada. *BMC Microbiol.* 2008; 8:89. [PubMed: 18538029]
5. Antunes P, Machado J, Sousa JC, Peixe L. Dissemination of sulfonamide resistance genes (*sul1*, *sul2*, and *sul3*) in Portuguese *Salmonella enterica* strains and relation with integrons. *Antimicrob Agents Chemother.* 2005; 49:836–839. [PubMed: 15673783]
6. Butaye P, Michael GB, Schwarz S, Barrett TJ, Brisabois A, White DG. The clonal spread of multidrug-resistant non-typhi *Salmonella* serotypes. *Microbes Infect.* 2006; 8:1891–1897. [PubMed: 16714135]
7. Call DR, Kang MS, Daniels J, Besser TE. Assessing genetic diversity in plasmids from *Escherichia coli* and *Salmonella enterica* using a mixed-plasmid microarray. *J Appl Microbiol.* 2006; 100:15–28. [PubMed: 16405681]

8. Carattoli A. Plasmid-mediated antimicrobial resistance in *Salmonella enterica*. *Curr Issues Mol Biol*. 2003; 5:113–122. [PubMed: 12921226]
9. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods*. 2005; 63:219–228. [PubMed: 15935499]
10. Carattoli A, Miriagou V, Bertini A, Loli A, Colinon C, Villa L, Whichard JM, Rossolini GM. Replicon typing of plasmids encoding resistance to newer beta-lactams. *Emerg Infect Dis*. 2006; 12:1145–1148. [PubMed: 16836838]
11. Carattoli A, Tosini F, Giles WP, Rupp ME, Hinrichs SH, Angulo FJ, Barrett TJ, Fey PD. Characterization of plasmids carrying CMY-2 from expanded-spectrum cephalosporin-resistant *Salmonella* strains isolated in the United States between 1996 and 1998. *Antimicrob Agents Chemother*. 2002; 46:1269–1272. [PubMed: 11959555]
12. D'Aoust, JY. *Salmonella* Species. In: Doyle, MP.; Beauchat, LR.; Montville, TJ., editors. *Food Microbiology: Fundamentals and Frontiers*. ASM Press; Washington D.C: 1997. p. 129-158.
13. de Hoon MJ, Imoto S, Nolan J, Miyano S. Open source clustering software. *Bioinformatics*. 2004; 20:1453–1454. [PubMed: 14871861]
14. Douard G, Praud K, Cloeckeaert A, Doublet B. The *Salmonella* genomic island 1 is specifically mobilized in trans by the IncA/C multidrug resistance plasmid family. *PLoS ONE*. 2010; 5:e15302. [PubMed: 21187963]
15. Dutil L, Irwin R, Finley R, Ng LK, Avery B, Boerlin P, Bourgault AM, Cole L, Daignault D, Desruisseau A, Demczuk W, Hoang L, Horsman GB, Ismail J, Jamieson F, Maki A, Pacagnella A, Pillai DR. Ceftiofur resistance in *Salmonella enterica* serovar Heidelberg from chicken meat and humans, Canada. *Emerg Infect Dis*. 2010; 16:48–54. [PubMed: 20031042]
16. Eisen MB, Spellman PT, Brown PO, Botstein D. Cluster analysis and display of genome-wide expression patterns. *Proc Natl Acad Sci U S A*. 1998; 95:14863–8. [PubMed: 9843981]
17. Excoffier L, Laval G, Schneider S. Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evol Bioinform Online*. 2005; 1:47–50. [PubMed: 19325852]
18. Fedorka-Cray PJ, Dargatz D, Petersen K, Tollefson L. National antimicrobial resistance monitoring system. *Veterinary strains*. FDA CVM. USDA. 8-16-2005. USDA, ARS
19. Folster JP, Pecic G, Bolcen S, Theobald L, Hise K, Carattoli A, Zhao S, McDermott PF, Whichard JM. Characterization of extended-spectrum cephalosporin-resistant *Salmonella enterica* serovar Heidelberg isolated from humans in the United States. *Foodborne Pathog Dis*. 2010; 7:181–187. [PubMed: 19785533]
20. Folster JP, Pecic G, McCullough A, Rickert R, Whichard JM. Characterization of bla(CMY)-Encoding Plasmids Among *Salmonella* Isolated in the United States in 2007. *Foodborne Pathog Dis*. 2011; 8:1289–1294. [PubMed: 21883005]
21. Frye JG, Fedorka-Cray PJ. Prevalence, distribution and characterisation of ceftiofur resistance in *Salmonella enterica* isolated from animals in the USA from 1999 to 2003. *Int J Antimicrob Agents*. 2007; 30:134–142. [PubMed: 17531447]
22. Frye JG, Fedorka-Cray PJ, Jackson CR, Rose M. Analysis of *Salmonella enterica* with Reduced Susceptibility to the Third-Generation Cephalosporin Ceftriaxone Isolated from U.S. Cattle During 2000–2004. *Microb Drug Resist*. 2008; 14:251–258. [PubMed: 19025468]
23. Frye JG, Lindsey RL, Rondeau G, Porwollik S, Long F, McClelland M, Jackson CR, Englen MD, Meinersmann RJ, Berrang ME, Davis JA, Barrett JB, Turpin JB, Thitaram SN, Fedorka-Cray PJ. Development of a DNA Microarray to Detect Antimicrobial Resistance Genes Identified in the National Center for Biotechnology Information Database. *Microb Drug Resist*. 2009; 16:9–19. [PubMed: 19916789]
24. Glenn LM, Lindsey RL, Frank JF, Meinersmann RJ, Englen MD, Fedorka-Cray PJ, Frye JG. Analysis of Antimicrobial Resistance Genes Detected in Multidrug-Resistant *Salmonella enterica* Serovar Typhimurium Isolated from Food Animals. *Microb Drug Resist*. 2011
25. Johnson TJ, Jordan D, Kariyawasam S, Stell AL, Bell NP, Wannemuehler YM, Alarcon CF, Li G, Tivendale KA, Logue CM, Nolan LK. Sequence analysis and characterization of a transferable hybrid plasmid encoding multidrug resistance and enabling zoonotic potential for extraintestinal *Escherichia coli*. *Infect Immun*. 2010; 78:1931–1942. [PubMed: 20160015]

26. Johnson TJ, Shepard SM, Rivet B, Danzeisen JL, Carattoli A. Comparative genomics and phylogeny of the IncII plasmids: A common plasmid type among porcine enterotoxigenic *Escherichia coli*. *Plasmid*. 2011; 66:144–151. [PubMed: 21843549]
27. Johnson TJ, Wannemuehler YM, Johnson SJ, Logue CM, White DG, Doetkott C, Nolan LK. Plasmid replicon typing of commensal and pathogenic *Escherichia coli* isolates. *Appl Environ Microbiol*. 2007; 73:1976–1983. [PubMed: 17277222]
28. Kang MS, Besser TE, Hancock DD, Porwollik S, McClelland M, Call DR. Identification of specific gene sequences conserved in contemporary epidemic strains of *Salmonella enterica*. *Appl Environ Microbiol*. 2006; 72:6938–6947. [PubMed: 16963552]
29. Lavigne JP, Blanc-Potard AB. Molecular evolution of *Salmonella enterica* serovar Typhimurium and pathogenic *Escherichia coli*: from pathogenesis to therapeutics. *Infect Genet Evol*. 2008; 8:217–226. [PubMed: 18226587]
30. Lindsey RL, Fedorka-Cray PJ, Frye JG, Meinersmann RJ. Inc A/C plasmids are prevalent in multidrug-resistant *Salmonella enterica* isolates. *Appl Environ Microbiol*. 2009; 75:1908–1915. [PubMed: 19181840]
31. Lindsey RL, Frye JG, Fedorka-Cray PJ, Meinersmann RJ. Microarray-based analysis of IncA/C plasmid-associated genes from multidrug-resistant *Salmonella enterica*. *Appl Environ Microbiol*. 2011; 77:6991–6999. [PubMed: 21841024]
32. Lindsey RL, Frye JG, Fedorka-Cray PJ, Welch TJ, Meinersmann RJ. An oligonucleotide microarray to characterize multidrug resistant plasmids. *J Microbiol Methods*. 2010; 81:96–100. [PubMed: 20138094]
33. Lindsey RL, Frye JG, Thitaram SN, Meinersmann RJ, Fedorka-Cray PJ, Englen MD. Characterization of multidrug-resistant *Escherichia coli* by antimicrobial resistance profiles, plasmid replicon typing, and pulsed-field gel electrophoresis. *Microb Drug Resist*. 2011; 17:157–163. [PubMed: 21332363]
34. McClelland M, Sanderson KE, Clifton SW, Latreille P, Porwollik S, Sabo A, Meyer R, Bieri T, Ozersky P, McLellan M, Harkins CR, Wang C, Nguyen C, Berghoff A, Elliott G, Kohlberg S, Strong C, Du F, Carter J, Kremizki C, Layman D, Leonard S, Sun H, Fulton L, Nash W, Miner T, Minx P, Delehaunty K, Fronick C, Magrini V, Nhan M, Warren W, Florea L, Spieth J, Wilson RK. Comparison of genome degradation in Paratyphi A and Typhi, human-restricted serovars of *Salmonella enterica* that cause typhoid. *Nat Genet*. 2004; 36:1268–1274. [PubMed: 15531882]
35. Mulvey MR, Boyd DA, Olson AB, Doublet B, Cloeckeaert A. The genetics of *Salmonella* genomic island 1. *Microbes Infect*. 2006; 8:1915–1922. [PubMed: 16713724]
36. Parkhill J, Dougan G, James KD, Thomson NR, Pickard D, Wain J, Churcher C, Mungall KL, Bentley SD, Holden MT, Sebahia M, Baker S, Basham D, Brooks K, Chillingworth T, Connor P, Cronin A, Davis P, Davies RM, Dowd L, White N, Farrar J, Feltwell T, Hamlin N, Haque A, Hien TT, Holroyd S, Jagels K, Krogh A, Larsen TS, Leather S, Moule S, O’Gaora P, Parry C, Quail M, Rutherford K, Simmonds M, Skelton J, Stevens K, Whitehead S, Barrell BG. Complete genome sequence of a multiple drug resistant *Salmonella enterica* serovar Typhi CT18. *Nature*. 2001; 413:848–52. [PubMed: 11677608]
37. Poole TL, Edrington TS, Brichta-Harhay DM, Carattoli A, Anderson RC, Nisbet DJ. Conjugative transferability of the A/C plasmids from *Salmonella enterica* isolates that possess or lack bla(CMY) in the A/C plasmid backbone. *Foodborne Pathog Dis*. 2009; 6:1185–1194. [PubMed: 19743926]
38. Saldanha AJ. Java Treeview—extensible visualization of microarray data. *Bioinformatics*. 2004; 20:3246–3248. [PubMed: 15180930]
39. Slatkin M. Linkage disequilibrium in growing and stable populations. *Genetics*. 1994; 137:331–336. [PubMed: 8056320]
40. Wain J, Diem Nga LT, Kidgell C, James K, Fortune S, Song DT, Ali T, Gaora O, Parry C, Parkhill J, Farrar J, White NJ, Dougan G. Molecular analysis of incHI1 antimicrobial resistance plasmids from *Salmonella* serovar Typhi strains associated with typhoid fever. *Antimicrob Agents Chemother*. 2003; 47:2732–2739. [PubMed: 12936967]
41. Welch TJ, Evenhuis J, White DG, McDermott PF, Harbottle H, Miller RA, Griffin M, Wise D. IncA/C plasmid-mediated florfenicol resistance in the catfish pathogen *Edwardsiella ictaluri*. *Antimicrob Agents Chemother*. 2009; 53:845–846. [PubMed: 19029319]

42. Welch TJ, Fricke WF, McDermott PF, White DG, Rosso ML, Rasko DA, Mammel MK, Eppinger M, Rosovitz MJ, Wagner D, Rahalison L, LeClerc JE, Hinshaw JM, Lindler LE, Cebula TA, Carniel E, Ravel J. Multiple antimicrobial resistance in plague: an emerging public health risk. *PLoS ONE*. 2007; 2:e309. [PubMed: 17375195]
43. Zhao S, Qaiyumi S, Friedman S, Singh R, Foley SL, White DG, McDermott PF, Donkar T, Bolin C, Munro S, Baron EJ, Walker RD. Characterization of *Salmonella enterica* serotype newport isolated from humans and food animals. *J Clin Microbiol*. 2003; 41:5366–5371. [PubMed: 14662912]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1. Cluster analysis of *Salmonella enterica* isolates based on results of microarray detection of antimicrobial resistance and plasmid genes. The IncA/C positive cluster, group A, and IncA/C negative cluster, group B are labeled and indicated by brackets. Complete cluster analysis data is available in Supplemental Figure S1. Resistance profiles of each isolate is listed with the following abbreviations for the antimicrobials: AMC, amoxicillin-clavulanic acid; AMK, amikacin; AMP, ampicillin; CHL, chloramphenicol; CIP, ciprofloxacin; FOX, cefoxitin; GEN, gentamicin; KAN, kanamycin; NAL, nalidixic acid; SMX,

sulfamethoxazole; STR, streptomycin; SXT, trimethoprim-sulfamethoxazole; TCY, tetracycline; TIO, ceftiofur. Antimicrobial resistance genes detected are summarized for each isolate. Genes and gene families with multiple probes are summarized and presented only once with different genes and alleles of related genes separated by a comma; full hybridization data are available in Supplementary Table S1. Replicons detected in each isolate by PCR analysis are also listed with (–) indicating that none of the replicons assayed were detected.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table of significant Linkage Disequilibrium (significance level = 0.05)

	A/C	P	FII	FIA	FIB	II	F	HI1	N	serovar	animal	agency
A/C	*											
P	-	*										
FII	-	-	*									
FIA	-	-	+	*								
FIB	-	-	-	-	*							
II	-	-	-	-	+	*						
F	-	-	+	-	-	-	*					
HI1	-	-	-	-	-	-	-	*				
N	-	+	-	-	+	+	-	-	*			
serovar	+	+	-	-	+	-	-	-	-	*		
animal	+	-	-	-	+	+	-	-	-	+	*	
agency	+	-	-	-	+	-	-	-	-	-	+	*

Figure 2.

Pairwise linkage disequilibrium based on nine plasmid replicon groups, serovar, phage type and animal source of *Salmonella* Typhimurium strains. (+) indicates a *P* value of 0.05 or less and significant linkage disequilibrium; and a (-) indicates a *P* value of greater than >0.05 and no significant linkage disequilibrium. Replicon groups B/O, FIC, T, K, W, Y, FIA, X, L/M were not present in any isolate and were subsequently removed from the analysis.

Summary of pYR1 and pSN254 IncA/C plasmid genes detected by microarray analysis and replicon types detected by PCR analysis. (Full hybridization data available in supplemental Table S1)

Table 1

Year	ID	Serovar	pYR1	pSN254	IncA/C	Core	Regions	Detected ^a	Plasmid	Replicons							
2007	CDC1	Typhimurium	132	36	1	2	3	4	5	6	7	8	9	10	11	12	A/C, II
2007	CDC2	Typhimurium	90	29	1	2	3	4	5	-	-	-	-	10	11	12	A/C, II
2007	CDC3	Typhimurium	1	12	-	-	-	-	-	-	-	-	-	-	-	-	FIA, FII
2007	CDC4	Heidelberg	105	32	1	2	3	4	5	-	8	9	10	11	12	A/C	
2007	CDC5	Heidelberg	105	31	1	2	3	4	5	-	8	9	10	11	12	A/C, N	
2007	CDC6	Heidelberg	103	32	1	2	3	4	5	-	8	9	10	11	12	A/C	
2007	CDC7	Newport	90	37	1	2	3	4	5	-	-	-	-	-	11	12	A/C, N
2007	CDC8	Newport	131	26	1	2	3	4	5	6	7	8	9	10	11	12	A/C, II
2007	CDC9	Newport	136	28	1	2	3	4	5	6	7	8	9	10	11	12	A/C
2007	FDA1	Heidelberg	15	8	-	-	-	-	-	-	-	-	-	-	-	-	II
2007	FDA2	Typhimurium	106	15	1	2	3	4	5	6	-	-	-	-	-	-	A/C, II
2007	FDA3	Heidelberg	13	19	-	-	-	-	-	-	-	-	-	-	-	-	II
2007	FDA4	Heidelberg	13	8	-	-	-	-	-	-	-	-	-	-	-	-	II
2007	FDA5	Hadar	10	22	-	-	-	-	-	-	-	-	-	-	-	-	II
2007	FDA6	Hadar	7	23	-	-	-	-	-	-	-	-	-	-	-	-	II
2007	FDA7	Hadar	4	3	-	-	-	-	-	-	-	-	-	-	-	-	N
2007	FDA8	Typhimurium	111	22	1	2	3	4	5	6	-	-	-	-	11	12	A/C, II
2007	FDA9	Typhimurium var. 5-	146	19	1	2	3	4	5	6	7	8	9	-	11	12	A/C, N
2007	USDA1	Newport	89	30	1	2	3	4	5	6	-	-	-	10	11	12	A/C, II
2007	USDA2	Newport	94	32	1	2	3	4	5	6	-	-	-	10	11	12	A/C, N
2007	USDA3	Newport	82	23	1	2	3	4	5	6	-	-	-	10	11	12	A/C
2007	USDA4	Heidelberg	128	36	1	2	3	4	5	6	7	8	9	10	11	12	A/C, II
2007	USDA5	Heidelberg	119	33	-	-	3	4	5	6	7	8	9	10	11	12	A/C, N
2007	USDA6	Heidelberg	119	33	-	-	3	4	5	6	7	8	9	10	11	12	A/C, N
2007	USDA7	Typhimurium	135	36	1	2	3	4	5	6	7	8	9	10	11	12	A/C, II
2007	USDA8	Typhimurium	135	40	1	2	3	4	5	6	7	8	9	10	11	12	A/C, N
2007	USDA9	Typhimurium	127	33	1	2	3	4	5	6	7	8	9	10	11	-	A/C, FIB, II

Year	ID	Serovar	pYRI	pSN254	IncA/C Core Regions Detected ^a	Plasmid Replicons
2007	USDA10	Heidelberg	7	13	- - - - 5 - - - - -	- - - - - II
2007	USDA11	Heidelberg	14	5	- - - - -	- - - - - II
2007	USDA12	Heidelberg	144	36	1 2 3 4 5 6 7 8 9 10 11 12	A/C, II
2004	CHI	Heidelberg	44	8	- 2 3 - 5 - - - - -	- 10 - 12 A/C
2005	CH2	Typhimurium	133	29	1 2 3 4 5 6 7 8 9 10 11 12	A/C, II
2007	CH3	Typhimurium	96	13	1 2 3 4 5 6 - - - - -	- 10 11 12 A/C, N
2005	CH4	Heidelberg	80	19	1 2 3 4 5 - - - - -	- 10 11 12 A/C, N
2005	CH5	Heidelberg	130	29	1 2 3 4 5 6 7 8 9 10 11 12	A/C, N
2005	CH6	Enteritidis	134	31	1 2 3 4 5 6 7 8 9 10 11 12	A/C, FIB, N
2006	CH7	Typhimurium	1	4	- - - - -	- - - - - II
2006	CH8	Enteritidis	1	4	- - - - -	- - - - - FIB, N
2007	CR1	Typhimurium	17	22	- 2 - - - - -	- - - - - FIB, N
2007	CR2	Heidelberg	9	6	- - - - -	- - - - - HII
2007	CR3	Typhimurium	1	6	- - - - -	- - - - - FIB, N
2007	CR4	Hadar	9	22	- - - - -	- - - - - II, N
2007	CR5	Typhimurium	1	6	- - - - -	- - - - - FIB, N
2007	CR6	Hadar	4	4	- - - - -	- - - - - N
2007	CR7	Heidelberg	0	14	- - - - -	- - - - - II
2007	CR8	Heidelberg	0	19	- - - - -	- - - - - II
2007	CR9	Typhimurium	0	8	- - - - -	- - - - - N
2005	CA1	Saintpaul	9	27	- - 3 - - - - -	- - - - - P, N
2007	CA2	Heidelberg	28	24	- 2 - - - - - 9 - - - - -	- - - - - II
2007	CA3	Typhimurium	2	10	- - - - -	- - - - - FIB, FII, N, F
2007	CA4	Hadar	4	7	- - - - -	- - - - - <i>nd</i>
2007	CA5	Heidelberg	8	4	- - - - -	- - - - - FII, II, F
2007	CA6	Typhimurium var. 5-	2	7	- - - - -	- - - - - FIB
2007	CA7	Typhimurium	3	7	- - - - -	- - - - - II
2007	CA8	Typhimurium	1	10	- - - - -	- - - - - FIB, N
2007	CA9	Typhimurium	3	9	- - - - -	- - - - - FIB, N
Total on array			180	47		

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

IncA/C plasmid core regions are based upon the assay by Welch *et al.*, 2007. Core regions are numbered in order 1 to 12 around the plus strand of the *IncA/C* plasmid sequence. Regions are scored as present if all genes within them were detected by microarray analysis of the isolate. These regions are identified with the full hybridization data in supplemental Table S1.

nd: none detected