



Published in final edited form as:

Lancet. 2012 August 11; 380(9841): 560. doi:10.1016/S0140-6736(12)61322-1.

New global estimates of malaria deaths

Naman K Shah, Ashwani Kumar, and Neena Valecha

National Institute of Malaria Research, New Delhi 110077, India

Naman K Shah: namankshah@gmail.com

India figures prominently as the largest contributor of malaria deaths outside Africa in new global estimates by Christopher Murray and colleagues.¹ However, India is often the greatest source of uncertainty in such estimates because of its sheer scale. With 85% of the 1.2 billion population at risk of malaria from both major parasite species, across varied transmission settings, and via six primary vectors, representative empirical data are hard to produce.

If Murray and colleagues' estimates are accurate, the historic decrease in malaria deaths in India since 1980 was indeed driven by general development, improvements in the health system, and better case management rather than the reduction in malaria transmission, since cases of falciparum malaria increased between 1980 and 2000 in the context of stable reporting (table). However, to conclude that socioeconomic change and improved quality of care led to the apparent decline in fatal malaria would assume (1) that *Plasmodium vivax*, whose reported cases substantially decreased during the same period, contributes little mortality in India—a view under increasing attack in the face of recent evidence,² and (2) that resistance to both first-line and second-line antimalarial drugs, which increased between 1978 and 2007,³ is not a major driver of malaria mortality, contrary to the results of other studies and Murray and colleagues' own findings.

Finally, the country perspective is frequently missing in global estimates of disease. Although the results can be important in setting international priorities and in affecting donor agencies, they cannot improve national malaria control programmes unless they can provide disaggregated, frequently updated estimates.⁴ Thus, the measurement of malaria and all-cause mortality must be improved through routine systems.

References

1. Murray CJ, Rosenfeld LC, Lim SS, et al. Global malaria mortality between 1980 and 2010: a systematic analysis. Lancet. 2012; 379:413–31. [PubMed: 22305225]
2. Kochar DK, Tanwar GS, Khatri PC, et al. Clinical features of children hospitalized with malaria—a study from Bikaner, northwest India. Am J Trop Med Hyg. 2010; 83:981–89. [PubMed: 21036824]
3. Shah NK, Dhillon GP, Dash AP, Arora U, Meshnick SR, Valecha N. Antimalarial drug resistance of *Plasmodium falciparum* in India: changes over time and space. Lancet Infect Dis. 2011; 11:57–64. [PubMed: 21183147]

Correspondence to: Naman K Shah, namankshah@gmail.com.

We declare that we have no conflicts of interest.

4. Shah NK. Defining and counting malaria deaths. *Indian J Med Res.* 2012; 135:270–72. [PubMed: 22561610]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table

Trends in malaria from routine surveillance in India, 1980–2010

| Population * | BSE* ^a | ABER | Malaria * | Falciparum * ^b | malaria | API | AFI |
|--------------|-------------------|------|-----------|---------------------------|---------|-----|-----|
| 1980 687322 | 67170 | 9.8% | 2898 | 588 (20.3%) | | 4.2 | 0.9 |
| 1990 849515 | 74420 | 8.8% | 2019 | 752 (37.3%) | | 2.4 | 0.9 |
| 2000 1015900 | 86790 | 8.5% | 2032 | 1047 (51.5%) | | 2.0 | 1.0 |
| 2002 1048600 | 91618 | 8.7% | 1842 | 897 (48.7%) | | 1.8 | 0.9 |
| 2010 1170900 | 108679 | 9.3% | 1600 | 834 (52.1%) | | 1.4 | 0.7 |

Population data from World Bank. BSE=blood smears examined. ABER=annual blood examination rate in the population. API=annual parasite incidence per 1000. AFI=annual falciparum incidence per 1000.

* Thousands.