



HHS Public Access

Author manuscript

Pediatr Obes. Author manuscript; available in PMC 2017 June 01.

Published in final edited form as:

Pediatr Obes. 2016 June ; 11(3): 187–193. doi:10.1111/ijpo.12044.

Food insecurity and dietary intake among U.S. youth, 2007–2010

Lauren M. Rossen, PhD, MS and Emily K. Kobernik, MPH

Centers for Disease Control and Prevention, National Center for Health Statistics, Office of Analysis and Epidemiology, Infant, Child, and Women's Health Statistics Branch, Hyattsville, Maryland, 20782

Abstract

Background—There is limited research describing associations between food insecurity and dietary intake.

Objective—To examine differences in dietary intake by food security status among a nationally representative sample of children and adolescents in the U.S.

Methods—The sample included 5136 children, ages 2–15 years, from the National Nutrition and Health Examination Survey, 2007–2010. Propensity-score weighting was used to improve covariate balance between food secure and food insecure (marginal, low or very low food security) participants. Multivariate measurement error models were used to model usual intake of various dietary components and assess differences by food security status.

Results—Initial analyses using multivariate measurement error models determined there were no differences between food insecure and food secure children across several dietary components. In sensitivity analyses, children experiencing very low food security consumed fewer whole grains and more solid fats and added sugars compared to their food secure counterparts. Some of these differences were attenuated after propensity-score weighting, though intake of whole grains and added sugars remained significantly different.

Conclusions—Food insecurity was largely not associated with dietary intake among 2–15 year old U.S. children, though some differences were observed comparing food secure children to those experiencing very low food security.

Keywords

nutrition; survey; children; adolescents; propensity score methods

Corresponding Author: Lauren M. Rossen, National Center for Health Statistics, 3311 Toledo Rd, Room 6114, Hyattsville, MD 20782, [lrossen@cdc.gov], 301-458-4256.

Disclaimer: The findings and conclusions in this paper are those of the authors and do not necessarily represent the views of the National Center for Health Statistics, Centers for Disease Control and Prevention

Conflict of Interest: The Authors have no conflicts of interest to disclose.

Financial Disclosure: The Authors have no financial relationships relevant to this article to disclose.

INTRODUCTION

In 2012, approximately 8.3 million children (11.3% of children in the U.S.) lived in households in which at least one child experienced food insecurity, a condition characterized by a lack of consistent, dependable access to enough food for active and healthy living. (1) Many studies have evaluated associations between food insecurity and weight outcomes among children and adults in the U.S., often producing conflicting findings. (2–4) Different associations have been reported by sex, race or ethnicity, or by participation in food assistance programs. (2–5) A major limitation of these studies and a potential explanation for inconsistencies is the lack of evidence describing the pathways by which food insecurity may influence weight outcomes: namely, dietary intake.

Of the few studies that have been published to date, some have reported minimal differences between food secure and food insecure children's dietary patterns, (6) while others have reported that food insecurity is linked with lower fruit and vegetable intake, (7, 8) and higher fat and juice intake. (9) One previous study using NHANES reported that food insecurity was associated with reduced dairy and calcium intake and bone mineral content among adolescent boys, but not girls. (10) Existing studies of food insecurity and dietary intake have been limited by small and/or convenience samples, and narrow assessments of dietary intake.

While prior studies have examined poverty and/or participation in food assistance in relation to dietary outcomes, these concepts are somewhat distinct from food insecurity; (11) less than 60% of food insecure households participate in food assistance programs. (1) A recent study found that children who receive Supplemental Nutrition Assistance (SNAP) consumed a greater amount of sugar-sweetened beverages, high-fat dairy and processed meats than comparably low-income non-participants. (12) Although this study controlled for food insecurity, differences across food security categories were not explicitly examined.

Food insecurity is associated with income and participation in food assistance programs, but other factors, such as neighborhood context (socioeconomic factors [SES], urbanization) may influence food security, as they are associated with access to food and other resources. (1, 3, 4, 13, 14) These contextual factors have been found to be associated with dietary intake, particularly among low-income populations, (4, 13, 15) and have generally not been accounted for in studies of food security and dietary intake or related health outcomes. (14)

The objective of this study was to examine differences in dietary intake by food security status, controlling for various individual-level and contextual confounders such as neighborhood SES and urban/rural status.

METHODS

Study population

Data were from the 2007–2010 National Health and Nutrition Examination Survey (NHANES). The study population consisted of children and adolescents (2–15 years) who completed at least one 24-hour dietary recall in the mobile examination center. Pregnant

adolescents and breastfeeding children were excluded leaving a sample of 5183 children and adolescents, 5136 of whom had data on food security (99%). As this analysis used secondary data, it was not subject to review by the National Center for Health Statistics Ethics Review Board. The overall response rates for the NHANES MEC ranged from 82 to 87% for youth 2–15 years in 2007–2010.

Dietary Intake

Dietary intake was assessed via 24-hour recall completed by a trained interviewer with a second recall conducted for most participants via telephone 3–10 days later. Proxy respondents (e.g., primary caregiver) complete recalls for children younger than 6 years. Children 6–11 years complete the recalls with assistance from a proxy, and children 12 years and older complete the recalls for themselves. Due to these differences in reporting method, children were grouped by age into the following categories: 2–5 years, 6–11 years, and 12–15 years. Data on dietary intake was obtained from the individual foods files and the total nutrients files, which include nutrient information based on the USDA's Food and Nutrient Database for Dietary Studies (FNDDS). (Version 4.1. U.S. Department of Agriculture, Food Surveys Research Group. Beltsville, MD)

Unlike the FNDDS, the Food Patterns Equivalent Database (FPED) disaggregates food into its component parts to produce estimates of several dietary outcomes including: intact/whole fruit, fruit juice, total vegetables (which can be calculated to exclude white potatoes), whole grains, refined grains, solid fats (g), and added sugars (tsp). (Version 2007–2008 and 2009–2010. U.S. Department of Agriculture, Food Surveys Research Group. Beltsville, MD) For example, using the FNDDS, the apples in apple pie would be coded as a grain (e.g. pie with apples), whereas the FPED differentiates between the ingredients, coding apples as a fruit. Grams of solid fats and teaspoons of added sugars were converted to their caloric equivalents and summed to create a category referred to as Solid Fats and Added Sugars (SoFAS); this category typically also contains alcohol, but as this study was limited to children, that item was not examined. The total nutrient file provides estimates of total kilocalories (kcal), protein (g), carbohydrates (g), number of foods, and sodium (mg). Energy density (kcal/food item) was also calculated as a study of homeless youth suggested that food insecure individuals may rely on more calorically dense foods as a consequence of food scarcity. (16) Finally, the number of foods was included as a measure of dietary variety because food insecure households might ration food, and children might be subject to lower dietary variety. (11)

Food Security and Other Covariates

During the household interview, an adult respondent completed the U.S. Food Security Survey Scale, which consists of 18 items and is consistent with the USDA method of assessing food insecurity. (17) Children in the household are assigned a score based on the 8 items specific to households with children and characterized as “food secure” (score of 0), “marginal food security” (1), “low food security” (2–4) or “very low food security” (5–8). For the main analysis, the marginal, low, or very low food security categories (unweighted n's were 473, 633, and 92, respectively) were combined into one ‘food insecure’ group due to the relatively small numbers of children falling into each of the three categories.

Sensitivity analyses examined alternative categorizations of food insecurity, including looking at household food insecurity, which was categorized in the same way as the main analysis (grouping marginal, low and very low household food security). Because children over the age of 15 completed the questionnaire themselves, the analysis was restricted to children 2–15 to avoid reporting bias.

Respondents also provided information about participation in a variety of food assistance programs including SNAP, receipt of free or reduced price school lunch and/or breakfast, and participation in the Women, Infant and Children’s Nutrition (WIC) program (for children 0–5 years). Additional individual or household-level covariates included: whether the child had public or private health insurance, whether the child was foreign-born, caregiver marital status (i.e., single, married/cohabitating, divorced/separated/widowed), caregiver education level (i.e., less than High School, High School/GED, some college, college or higher) and age, household size, race/ethnicity, income-to-poverty ratio (PIR), and household smoking.

Restricted data files with geographic identifiers were used to link participating children to auxiliary data based on their census tract and county of residence. These data sources and covariates included: tract- and county-level socio-demographic and economic characteristics (e.g., racial and ethnic population distribution, population size, % poverty, median household income, % of residents with less than High School education); (18) residential racial segregation; (19) the number of arrests per 100,000 county residents; (20) the urban-rural designation of each county; (21) county-level food store density (e.g., the number of grocery stores, convenience stores, fast food restaurants), food price (e.g., the price of milk, soda, fruit), and food assistance participation rates. (22)

Statistical Analyses

Propensity Score Models—If food secure and food insecure children are not ‘exchangeable’ or comparable with respect to all characteristics except for food security status, then estimation of the effect of food insecurity will be biased. For example, if food insecure children are more likely to reside in disadvantaged neighborhoods than food secure children, and residing in these neighborhoods is related to dietary intake, then differences in dietary intake by food security status will be confounded by neighborhood SES. Propensity score methods can be used to ensure samples are balanced on potential confounders; (23) in this case, propensity score weighting was used to ensure food secure and food insecure children are otherwise comparable. (24) All relevant individual-level and contextual confounders were included in the propensity-score estimation model, (23) which was estimated in Stata 12.1 SE. (Release 12.1, StataCorp. College Station, TX) Methods are described in Appendix B. Consistent with recent guidelines on using propensity-score methods with complex surveys, inverse probability of treatment weights (IPTW) were created from the propensity scores, and combined with the day one dietary survey weights in order to provide unbiased estimates of the average treatment effect (ATE) that are generalizable to the survey’s target population. (24) In this case, the ATE represents the effect of food insecurity and the target population is U.S. children aged 2–15 years.

Outcome Models—Usual intake of various dietary components was modeled using methods developed by the National Cancer Institute. (25, 26) These methods use multivariate Markov Chain Monte Carlo models to estimate usual intake of several, possibly correlated, dietary components. (25, 26) Multipart nonlinear mixed models with correlated random effects account for skewed data, sequence and weekend effects; and can be used to estimate the usual intake of episodically (e.g., whole fruit) or ubiquitously (e.g., kcal) consumed foods. Post-stratified balanced repeated replicate (BRR) weights were used to account for the complex survey design.

Two sets of multivariate measurement error models were run. The first set incorporated just the survey weights, and adjusting only for age group, race/ethnicity and sex (i.e., the “regular” model). The second set of models incorporated both the propensity and survey weights, thereby controlling for potential confounders included in the propensity score estimation models (i.e., the “IPTW” model). For both the regular and IPTW models, mean intake of various dietary components produced by the models were compared by food security status and differences were assessed based on *t* tests. (25, 26) Outcome model analyses were conducted in SAS 9.3. (Version 9.3, SAS Institute Inc. Cary, NC)

Sensitivity analyses examined two different categorizations of child food insecurity. First, children with either low or very low food security (9.8% of children, unweighted *n* = 725) were compared to food secure children (84.3% of children, unweighted *n* = 4447). Second, children experiencing very low food security (1.1% of children, unweighted *n*=92) were compared to food-secure children. The first sensitivity analysis produced results similar to the main analyses, thus only the findings for the main analysis and second sensitivity analysis are shown. A final sensitivity analysis examined household food insecurity, which is also reported for comparison. Household food insecurity was categorized in a similar way to the main results, children in food secure households (unweighted *n*=3,033) were compared to those with either moderate, low, or very low household food security (unweighted *n*=2,103). These results are described as applying to “children in food insecure households” where all other results apply to child-level food security status.

RESULTS

Approximately 16% of children 2–15 years old experienced food insecurity (i.e., marginal, low, or very low food security) in the prior 12 months. There were substantial differences between food secure and food insecure children across several demographic and socioeconomic characteristics (see Appendix B). Food insecure children were more likely to be non-Hispanic black or Mexican-American, fall below the federal poverty threshold, and have caregivers with less than a High School education who were not married or cohabiting. Food insecure children also lived in larger households than children who were food secure. Household smoking was reported for a greater proportion of food insecure children, and nearly all food insecure children participated in at least one food assistance program (e.g., SNAP, WIC, free or reduced price school lunch or breakfast) in the prior year (94%) compared to only 67% of food secure children. A greater proportion of food insecure children were uninsured or on public insurance. There were several significant differences with respect to neighborhood or contextual covariates. Food insecure children were more

likely to live in census tracts with high deprivation, as well as experience both individual-level poverty and neighborhood deprivation. Food insecure children also resided in smaller, more urban counties with higher levels of crime compared to food secure children.

After propensity-score weighting, food secure and insecure children were balanced with respect to included covariates. Appendix A illustrates the standardized percent bias (i.e., the mean difference between groups expressed as a percentage of the average standard deviation) between food insecure and food secure children pre- and post-propensity score weighting.

Dietary Intake

No differences in dietary intake patterns were observed between food secure and food insecure children in either the regular or IPTW models. Mean intake of various dietary components can be seen in Table 1. Table 2 illustrates results of sensitivity analyses. In regular models, children experiencing very low food-security consumed fewer whole grains (0.20 fewer servings, SE 0.09, $P<0.05$), and a greater amount of solid fats and added sugars (SoFAS; 137.47 kcal more, SE 51.23, $P<0.05$) as compared to food secure children. In IPTW models, differences between children experiencing very low food security and their food secure counterparts were evident only for whole grain consumption (0.24 fewer servings, SE 0.09, $P<0.05$), and teaspoons of added sugars (6.90 more teaspoons, SE 3.36, $P<0.05$). Table 3 highlights differences by household food security status before and after propensity-score weighting. In regular models, children in households that were food secure consumed 0.10 fewer cups of juice (SE 0.04, $P<0.05$), 0.12 more servings of whole grains (SE 0.04, $P<0.05$), 1.82 fewer teaspoons of added sugar (SE 0.61, $P<0.05$), 42.19 fewer kcal from SoFAS (SE 18.89, $P<0.05$), and a greater variety of food items (0.85 more foods, SE 0.21, $P<0.05$) with lower average caloric density (by 12.34 kcal, SE 3.02, $P<0.05$) compared to children in food insecure households (consisting of households reporting moderate, low or very low food security). In IPTW models, all of these differences were completely attenuated. Supplemental Appendix C presents differences in dietary intake by child food security status stratified by age group; these results were consistent with the main findings presented here.

DISCUSSION

In 2007–2010, no differences in dietary intake were observed by child food security status among children aged 2 to 15-year-old. These findings are consistent with at least one study that reported few differences between food secure and food insecure children. (6) However, other studies have reported that food insecurity is linked with lower fruit and vegetable intake (7), and higher fat and juice intake. (9) Inconsistencies could be due to different categorizations of food insecurity (e.g., grouping marginal, low, and very-low or breaking out these categories), the use of convenience samples in prior studies, or residual confounding. Results of the sensitivity analyses suggest that residual confounding may play a large role. When children were grouped by household food security status, several differences in intake observed in the regular models (i.e., for fruit juice, whole grains, added

sugars, SoFAS, caloric density and dietary variety) were completely attenuated after adjusting for potential confounders in the IPTW models.

In support of the notion that categorization of food security status may affect findings, sensitivity analyses suggested that children experiencing very low food security consumed fewer whole grains and more calories from solid fats and added sugars than food secure children. In the IPTW models, which better accounted for various potential confounders, only reported consumption of whole grains and added sugars remained significantly different. Given the very small sample size of children in the very low food security group, conclusions are limited. However, more research is needed to further elucidate these potential relationships with larger samples of children reporting very low food security, particularly as whole grains and SoFAS each have been noted by the Dietary Guidelines for Americans as food items to encourage (whole grains) or reduce (SoFAS) due to their respective associations with chronic disease and cardiometabolic risk factors. (27)

While not the focus of this study, we observed no overall differences in weight status by child food security (data not shown) in regular or IPTW models. Kohn et al. (5) used propensity-score methods and found that food assistance program participation was associated with higher weight among food-secure children in NHANES, but not among food-insecure children. As that study did not include various contextual confounders in the estimation of the propensity-scores, residual confounding may have played a role.

There are a few limitations of this study worth noting. The measurement of food insecurity is challenging. Children may report the experience of food insecurity differently than adults in a household. (28) Food insecurity is not a static condition over time and may affect household members differentially by age, as children may be protected while adult or adolescent members experience the effects of food insecurity. (1, 11) Moreover, the relationship between food insecurity and participation in food assistance programs is complex and bidirectional. (1) Controlling for participation in food assistance programs may be controlling for a mediator, though some research has indicated that the effect of SNAP participation on food security and dietary intake may not be large. (29, 30) More research is needed to understand these complex relationships. Propensity-score methods only control for observed confounders, and residual confounding may still be present. Finally, we did not look at energy-adjusted macronutrients, but given the lack of differences in energy intake and macronutrient amounts, it is unlikely that differences would have been observed if energy-adjusted macronutrient intake had been included as an outcome. As with all other studies that rely on dietary recall data, findings are subject to biases and misreporting of dietary intake.

The strengths of this study are the use of a nationally representative sample of children, the utilization of sophisticated methods developed by the National Cancer Institute to assess usual dietary intake, and the employment of propensity score methods to more rigorously control for potential confounders vis-à-vis prior studies.

Conclusions

This analysis provides a description of dietary intake patterns among U.S. children by food security status. Food insecurity was not associated with intake of several dietary components. Propensity score methods could be readily applied to examine associations between participation in food assistance programs, dietary intake, and other health outcomes such as obesity. Further research could help clarify some of these associations given the conflicting findings with respect to food insecurity, food assistance program participation, and obesity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding Source: This work was performed under employment of the U.S. Federal government and the authors did not receive any outside funding.

Lauren M. Rossen: Dr. Rossen conceptualized and designed the study, carried out and interpreted the analyses, drafted the initial manuscript and approved the final manuscript as submitted. Emily K. Kobernik: Ms. Kobernik carried out and interpreted the analyses, reviewed and revised the manuscript and approved the final manuscript as submitted.

Abbreviations

NHANES National Health and Nutrition Examination Survey

References

1. Coleman-Jensen, A.; Nord, M.; Singh, A. Economic Research Report No. (ERR-155). Sep. 2013 Household food security in the United States in 2012; p. 41
2. Eisenmann JC, Gundersen C, Lohman BJ, Garasky S, Stewart SD. Is food insecurity related to overweight and obesity in children and adolescents? A summary of studies, 1995–2009. *Obes Rev.* 2011; 12(5):e73–83. [PubMed: 21382151]
3. Franklin B, Jones A, Love D, Puckett S, Macklin J, White-Means S. Exploring mediators of food insecurity and obesity: a review of recent literature. *J Community Health.* 2012; 37(1):253–264. [PubMed: 21644024]
4. Larson NI, Story MT. Food insecurity and weight status among U.S. children and families: a review of the literature. *Am J Prev Med.* 2011; 40(2):166–173. [PubMed: 21238865]
5. Kohn MJ, Bell JF, Grow HM, Chan G. Food insecurity, food assistance and weight status in US youth: new evidence from NHANES 2007–08. *Pediatr Obes.* 2013; 9(2):155–66. [PubMed: 23364918]
6. Grutzmacher S, Gross S. Household food security and fruit and vegetable intake among low-income fourth-graders. *J Nutr Educ Behav.* 2011; 43(6):455–463. [PubMed: 21855417]
7. Dave JM, Evans AE, Saunders RP, Watkins KW, Pfeiffer KA. Associations among food insecurity, acculturation, demographic factors, and fruit and vegetable intake at home in Hispanic children. *J Am Diet Assoc.* 2009; 109(4):697–701. [PubMed: 19328265]
8. Pilgrim A, Barker M, Jackson A, et al. Does living in a food insecure household impact on the diets and body composition of young children? Findings from the Southampton Women’s Survey. *J Epidemiol Community Health.* 2012; 66(6):e6. [PubMed: 21652519]

9. Mello JA, Gans KM, Risica PM, Kirtania U, Strolla LO, Fournier L. How is food insecurity associated with dietary behaviors? An analysis with low-income, ethnically diverse participants in a nutrition intervention study. *J Am Diet Assoc.* 2010; 110(12):1906–1911. [PubMed: 21111099]
10. Eicher-Miller HA, Mason AC, Weaver CM, McCabe GP, Boushey CJ. Food insecurity is associated with diet and bone mass disparities in early adolescent males but not females in the United States. *J Nutr.* 2011; 141(9):1738–1745. [PubMed: 21795427]
11. Cook JT, Frank DA. Food security, poverty, and human development in the United States. *Annals of the New York Academy of Sciences.* 2008; 1136:193–209. [PubMed: 17954670]
12. Leung CW, Blumenthal SJ, Hoffnagle EE, et al. Associations of food stamp participation with dietary quality and obesity in children. *Pediatrics.* 2013; 131(3):463–472. [PubMed: 23439902]
13. Larson NI, Story MT, Nelson MC. Neighborhood environments: disparities in access to healthy foods in the U.S. *Am J Prev Med.* 2009; 36(1):74–81. [PubMed: 18977112]
14. National Research Council and Institute of Medicine of the National Academies. *Research Opportunities Concerning the Causes and Consequences of Child Food Insecurity and Hunger: A Workshop Summary.* Washington, D.C: The National Academies Press; 2013.
15. Lovasi GS, Hutson MA, Guerra M, Neckerman KM. Built environments and obesity in disadvantaged populations. *Epidemiol Rev.* 2009; 31:7–20. [PubMed: 19589839]
16. Smith C, Richards R. Dietary intake, overweight status, and perceptions of food insecurity among homeless Minnesotan youth. *Hum Biol.* 2008; 20(5):550–563.
17. Bickel, G.; Nord, M.; Price, C.; Hamilton, W.; Cook, J. *Guide to Measuring Household Food Security, Revised 2000.* Alexandria, VA: U.S. Department of Agriculture, Food and Nutrition Service; 2000.
18. U.S. Department of Commerce Bureau of the Census. *Census of Population and Housing, 2000: Summary File 3.* Washington, DC: U.S. Department of Commerce, Bureau of the Census; 2002.
19. Escarce, J.; Lurie, N.; Jewell, A. *RAND Center for Population Health and Health Disparities (CPHHD) Data Core Series: Segregation Indices, 1990–2000.* Ann Arbor, MI: Inter-university Consortium for Political and Social Research; 2011.
20. Federal Bureau of Investigation. *Uniform Crime Reports.* Washington, D.C: Federal Bureau of Investigation; 2000.
21. Ingram DD, Franco SJ. NCHS urban-rural classification scheme for counties. *Vital Health Stat 2.* 2012; 154:1–65. [PubMed: 22783637]
22. U.S. Department of Agriculture Economic Research Service. *Food Environment Atlas.* 2011. Retrieved April 25, 2014 from: www.ers.usda.gov/data-products/food-environment-atlas.aspx
23. Stuart EA. Matching methods for causal inference: A review and a look forward. *Stat Sci.* 2010; 25(1):1–21. [PubMed: 20871802]
24. Dugoff EH, Schuler M, Stuart EA. Generalizing Observational Study Results: Applying Propensity Score Methods to Complex Surveys. *Health Serv Res.* 2014; 49:284–303. [PubMed: 23855598]
25. National Cancer Institute. *SAS macros for Fitting Multivariate Measurement Error Models and Estimating Multivariate Usual Intake Distributions.* 2013. Retrieved August 30, 2014 from: http://appliedresearch.cancer.gov/diet/usualintakes/macros_multiple.html
26. Guenther PM, Kirkpatrick SI, Reedy J, et al. The Healthy Eating Index-2010 is a valid and reliable measure of diet quality according to the 2010 dietary guidelines for Americans. *J Nutr.* 2014; 144(3):399–407. [PubMed: 24453128]
27. U.S. Department of Agriculture and U.S. Department of Health and Human Services. *Dietary Guidelines for Americans, 2010.* 7. Washington, DC:
28. Fram MS, Frongillo EA, Jones SJ, et al. Children are aware of food insecurity and take responsibility for managing food resources. *J Nutr.* 2011; 141(6):1114–1119. [PubMed: 21525257]
29. Leung CW, Cluggish S, Villamor E, Catalano PJ, Willett WC, Rimm EB. Few changes in food security and dietary intake from short-term participation in the Supplemental Nutrition Assistance Program among low-income Massachusetts adults. *J Nutr Educ Behav.* 2014; 46(1):68–74. [PubMed: 24238909]
30. Fox, MK.; Hamilton, W.; Lin, BH. *Effects of Food Assistance and Nutrition Programs on Nutrition and Health.* Vol. 4. Beltsville, MD: Economic Research Service; 2004.

What is already known on this subject

- Many studies have evaluated associations between food insecurity and weight among U.S. children, but findings are inconsistent.
- One limitation is the lack of evidence describing the pathways by which food insecurity may influence weight outcomes: namely, dietary intake.

What this study adds

- Food insecurity was largely not associated with dietary intake among 2–15 year old U.S. children.
- Some differences in intake of whole grains and added sugars were observed comparing food secure children to those experiencing very low food security.

Dietary intake by food security status among children ages 2–15 years (n=5,136) in NHANES, 2007–2010.

Table 1

	Regular Survey Weights		IPTW+Survey Weights	
	Food Secure Mean (S.E.)	Insecure Mean (S.E.)	Food Secure Mean (S.E.)	Insecure Mean (S.E.)
Fruit juice (cups)	0.51 (0.11)	0.42 (0.15)	0.48 (0.08)	0.51 (0.11)
Whole fruit [‡]	0.63 (0.17)	0.76 (0.22)	0.66 (0.08)	0.61 (0.10)
Vegetables excl white potatoes [‡]	0.58 (0.04)	0.56 (0.06)	0.56 (0.03)	0.60 (0.05)
Starchy vegetables [‡]	0.30 (0.07)	0.25 (0.07)	0.26 (0.02)	0.28 (0.07)
Whole grains	0.55 (0.07)	0.60 (0.08)	0.59 (0.05)	0.54 (0.15)
Refined grains	5.64 (0.10)	5.64 (0.31)	5.40 (0.09)	5.56 (0.34)
Solid fats (g)	34.66 (0.87)	34.38 (1.38)	34.23 (0.51)	34.70 (1.78)
Added sugars (tsp)	17.56 (0.92)	16.89 (1.67)	15.84 (0.23)	16.57 (1.90)
SoFAS (kcal)	592.91 (20.11)	579.68 (27.78)	561.46 (6.61)	577.40 (36.53)
Kcal	1826.36 (29.15)	1809.70 (51.95)	1760.91 (30.45)	1804.98 (66.73)
Protein (g)	65.05 (2.06)	66.52 (2.57)	65.32 (1.83)	65.76 (2.65)
Carbohydrate (g)	246.98 (6.71)	242.14 (7.80)	233.92 (3.73)	241.57 (9.73)
Fiber (g)	12.80 (0.24)	12.87 (0.56)	12.70 (0.35)	12.91 (0.63)
Caloric density (kcal/food item)	136.42 (5.47)	132.25 (5.56)	129.75 (3.82)	130.51 (6.93)
Sodium (mg)	2905.15 (45.95)	2882.04 (126.88)	2784.09 (83.74)	2886.72 (168.86)
Dietary variety (# foods)	14.30 (0.40)	14.61 (0.31)	14.38 (0.75)	14.70 (0.34)

[‡]cup equivalent

Note: Regular models used only the first day dietary recall survey weights, while IPTW models used propensity-score weights. There were no significant differences between the groups in either the regular or IPTW models.

Dietary intake among children ages 2–15 years (n=5,136) in NHANES 2007–2010, comparing food secure children (unweighted n=4449) to those with very low food-security (unweighted n=92). Significant differences shown in bold-type ($P<0.05$).

Table 2

	Regular Survey Weights		IPTW+Survey Weights	
	Food Secure Mean (S.E.)	Very low food-security Mean (S.E.)	Food Secure Mean (S.E.)	Very low food-security Mean (S.E.)
Fruit juice (cups)	0.51 (0.11)	0.37 (0.06)	0.48 (0.08)	0.37 (0.07)
Whole fruit [‡]	0.63 (0.17)	0.59 (0.11)	0.66 (0.08)	0.53 (0.26)
Vegetables excl white potatoes [‡]	0.58 (0.04)	0.67 (0.09)	0.56 (0.03)	0.60 (0.21)
Starchy vegetables [‡]	0.30 (0.07)	0.26 (0.05)	0.26 (0.02)	0.26 (0.05)
Whole grains	0.55 (0.07)^a	0.35 (0.06)^b	0.59 (0.05)^a	0.35 (0.09)^b
Refined grains	5.64 (0.10)	6.17 (0.35)	5.40 (0.09)	6.34 (1.01)
Solid fats (g)	34.66 (0.87)^a	41.72 (3.21)^b	34.23 (0.51)	42.52 (5.99)
Added sugars (tsp)	17.56 (0.92)^a	22.18 (2.08)^b	15.84 (0.23)^a	22.74 (3.42)^b
SoFAS (kcal)	592.91 (20.11)^a	730.39 (56.42)^b	561.46 (6.61)	746.43 (101.57)
Kcal	1826.36 (29.15)	1920.12 (90.50)	1760.91 (30.45)	1940.47 (175.72)
Protein (g)	65.05 (2.06)	68.49 (3.89)	65.32 (1.83)	68.16 (5.35)
Carbohydrate (g)	246.98 (6.71)	254.01 (12.11)	233.92 (3.73)	259.18 (21.39)
Fiber (g)	12.80 (0.24)	12.91 (0.98)	12.70 (0.35)	12.83 (1.02)
Caloric density (kcal/food item)	136.42 (5.47)	133.74 (7.59)	129.75 (3.82)	134.25 (11.6)
Sodium (mg)	2905.15 (45.95)	3024.12 (175.02)	2784.09 (83.74)	3065.27 (287.88)
Dietary variety (# foods)	14.30 (0.40)	15.25 (0.70)	14.38 (0.75)	15.26 (0.92)

[‡]cup equivalent

^{a, b} indicate significantly different, $P<0.05$

Note: Regular models used only the first day dietary recall survey weights, while IPTW models used propensity-score weights.

Dietary intake among children ages 2–15 years (n=5,136) in NHANES 2007–2010, comparing children in food secure households (unweighted n=3,033) to those in food insecure households (unweighted n=2,103). Significant differences shown in bold-type ($P<0.05$).

Table 3

	Regular Survey Weights		IPTW+Survey Weights	
	Food Secure Mean (S.E.)	Food-Insecure Mean (S.E.)	Food Secure Mean (S.E.)	Food-Insecure Mean (S.E.)
Fruit juice (cups)	0.41 (0.03) ^a	0.49 (0.03) ^b	0.47 (0.07)	0.52 (0.19)
Whole fruit [‡]	0.74 (0.05)	0.62 (0.04)	0.66 (0.09)	0.63 (0.15)
Vegetables excl white potatoes [‡]	0.58 (0.02)	0.57 (0.02)	0.56 (0.03)	0.56 (0.04)
Starchy vegetables [‡]	0.24 (0.02)	0.27 (0.02)	0.25 (0.02)	0.26 (0.03)
Whole grains	0.64 (0.03) ^a	0.52 (0.03) ^b	0.60 (0.06)	0.50 (0.08)
Refined grains	5.34 (0.11)	5.70 (0.14)	5.16 (0.22)	5.69 (0.66)
Solid fats (g)	33.82 (0.72)	35.28 (0.73)	33.65 (0.67)	34.70 (4.01)
Added sugars (tsp)	15.60 (0.26) ^a	17.42 (0.58) ^b	15.41 (0.29)	16.91 (3.21)
SoFAS (kcal)	553.97 (9.26) ^a	596.17 (15.00) ^b	549.32 (9.38)	582.90 (86.27)
Kcal	1766.69 (22.91)	1837.27 (28.21)	1727.70 (41.56)	1810.13 (199.55)
Protein (g)	66.17 (0.74)	66.68 (1.05)	64.14 (2.02)	65.46 (6.74)
Carbohydrate (g)	234.20 (3.37)	246.21 (4.40)	229.56 (5.52)	243.17 (29.38)
Fiber (g)	12.79 (0.32)	12.91 (0.21)	12.40 (0.59)	12.84 (1.35)
Caloric density (kcal/food item)	124.47 (1.15) ^a	136.81 (2.60) ^b	126.28 (1.90)	135.29 (13.55)
Sodium (mg)	2819.39 (50.00)	2916.69 (44.77)	2728.64 (97.28)	2876.76 (249.33)
Dietary variety (# foods)	15.15 (0.15) ^a	14.30 (0.17) ^b	14.55 (0.54)	14.16 (0.66)

[‡] cup equivalent

^{a,b} indicate significantly different, $P<0.05$

Note: Regular models used only the first day dietary recall survey weights, while IPTW models used propensity-score weights.