



Published in final edited form as:

Prev Med. 2015 September ; 78: 9–13. doi:10.1016/j.yjmed.2015.06.004.

Smoke-free home and vehicle rules by tobacco use status among US adults

Judy Kruger^{a,*}, Amal Jama^b, David M. Homa^a, Stephen D. Babb^a, and Brian A. King^a

^aOffice on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, GA 30341-3717, United States

^bD.B. Consulting Group, Atlanta, GA 30341-3717, United States

Abstract

Objective—To assess the prevalence and characteristics of smoke-free home and vehicle rules by tobacco use.

Methods—Data came from the 2012–2013 National Adult Tobacco Survey, a telephone survey of adults aged 18. Respondents who reported smoking is ‘never allowed’ inside their home or any family vehicle were considered to have smoke-free home and vehicle rules, respectively. Prevalence and characteristics of smoke-free rules were assessed overall and by current tobacco use (combustible only, noncombustible only, combustible and noncombustible, no current tobacco use). Assessed characteristics included: sex, age, race/ethnicity, education, marital status, income, region, and sexual orientation.

Results—Nationally, 83.7% of adults ($n = 48,871$) had smoke-free home rules and 78.1% ($n = 46,183$) had smoke-free vehicle rules. By tobacco use, prevalence was highest among nonusers of tobacco (homes: 90.8%; vehicles: 88.9%) and lowest among combustible-only users (homes: 53.7%; vehicles: 34.2%). Prevalence of smoke-free home and vehicle rules was higher among males, adults with a graduate degree, and adults living in the West.

Conclusions—Most adults have smoke-free home and vehicle rules, but differences exist by tobacco use. Opportunities exist to educate adults about the dangers of secondhand smoke and the benefits of smoke-free environments, particularly among combustible tobacco users.

Keywords

Smoke-free transactions; Nonsmoking homes; Vehicles; Public places

*Corresponding author at: Office on Smoking and Health, Centers for Disease Control and Prevention, 4770 Buford Highway, NE, F-79, Atlanta, GA 30341-3724, United States. ezk0@cdc.gov.

Conflict of interest statement

The authors declare that there are no conflicts of interests.

Author contributions

All authors contributed to the writing of this paper. JK drafted the manuscript. AJ analyzed the data. JK, DH, SB, BK designed the study and interpreted the data. All authors approved the final version to be submitted for consideration of publication.

Introduction

Secondhand smoke (SHS) exposure causes disease and death among nonsmoking adults and children (USDHHS, 2006, 2012, 2014). The Surgeon General has concluded there is no risk-free level of SHS exposure, and that completely eliminating smoking in indoor settings is the only way to fully protect nonsmokers (USDHHS, 2006). In the past decade, considerable progress has occurred in the implementation of comprehensive state and local smoke-free laws that prohibit smoking in worksites, bars, and restaurants; as of January 2015, 26 states and the District of Columbia had implemented such laws (CDC, 2015).

Historically, the voluntary adoption of smoke-free policies has increased, however, homes and vehicles remain significant sources of SHS exposure, particularly for children (USDHHS, 2006). One international study from New Zealand demonstrated that youth SHS exposure rates in vehicles and homes decreased between 2006 and 2012 (Healey et al., 2015). A US study from 2009–2010 found the prevalence of SHS exposure in vehicles to be 73.6% (King et al., 2013) and another national study reported the prevalence of smoke-free home rules increased from 43.0% in 1992–1993 to 83.0% in 2010–2011 (CDC, 2014a).

Although cigarette smoking has declined (USDHHS, 2014), more Americans are using other tobacco products, including combustible (e.g., cigars, cigarillos, and little cigars), noncombustible (e.g., chewing tobacco and snus), and emerging products (e.g., electronic cigarettes) (USDHHS, 2014; CDC, 2014b). As highlighted in the 2014 Surgeon General's report, this diversification of the tobacco product landscape has made it increasingly important to assess tobacco control interventions in the context of all forms of tobacco use (USDHHS, 2014).

In addition to protecting nonsmokers from SHS (USDHHS, 2006, 2014), implementing smoke-free rules in private settings can encourage smoking cessation (USDHHS, 2012), prevent relapse, and reduce the social acceptability of tobacco use (Hopkins et al., 2010). To date, studies of smoke-free home and vehicle rules have been assessed by cigarette smoking; however, no studies have assessed such rules by other tobacco use (King et al., 2013). Therefore, we assessed the prevalence and characteristics of smoke-free home and vehicle rules among US adults by tobacco use.

Methods

Data source

Data came from the 2012–2013 National Adult Tobacco Survey (NATS), a landline and cellular telephone survey of noninstitutionalized civilian US adults aged ≥18 (CDC, 2013). The sampling design was composed of independent samples drawn from households in the 50 US states and District of Columbia. During October 2012 to July 2013, 60,192 interviews were completed (landline: 57,999; cellular: 2193); the response rate was 44.9% (landline: 47.2%; cellular: 36.3%).

Measures

Smoke-free rules—Smoke-free home rules were determined by the question, “Not counting decks, porches, or garages, inside your home, is smoking always allowed, allowed only at some times or in some places, or *never allowed*?” Respondents who selected *never allowed* were classified as having a smoke-free home rule. Smoke-free vehicle rules were determined by the question, “Not counting motorcycles, in the vehicles that you or family members who live with you own or lease, is smoking always allowed in all vehicles, sometimes allowed in at least one vehicle, or *never allowed in any vehicle*?” Respondents who selected *never allowed in any vehicle* were classified as having a smoke-free vehicle rule. Missing responses, refusals, and responses of ‘*don’t know*’ or ‘*family doesn’t own or lease a vehicle*’ (vehicle only) were excluded (homes: $n = 2402$; vehicles: $n = 3220$).

Tobacco use—Tobacco use was categorized as: 1) any tobacco; 2) combustible only; 3) noncombustible only; 4) both combustible and noncombustible; and 5) no current tobacco use. Current combustible use was defined as smoking 100 cigarettes, cigars/cigarillos/filtered little cigars 50 times, regular pipes 1 time, or water pipes/hookahs 1 time during their lifetime, and now using these respective products ‘everyday’ or ‘some days’. Current noncombustible use was defined as using chewing tobacco, snuff, or dip 20 times during their lifetime, or snus or dissolvable tobacco products on 1 or more days, and now using these products ‘everyday’ or ‘some days.’ Any tobacco use was defined as current combustible use, and/or noncombustible use, and/or electronic cigarettes use (use 1 time during lifetime and now use ‘everyday’ or ‘some days’). Based on conclusions of the 2014 Surgeon General’s report, electronic cigarettes were categorized as a noncombustible product as they provide doses of nicotine and other additives to the user in an aerosol nicotine delivery systems (USDHHS, 2014). No current tobacco use was defined as not currently using combustible tobacco, noncombustible tobacco, or electronic cigarettes.

Sociodemographics—Sociodemographics included: sex, age, race/ethnicity, education, marital status, annual household income, US region, and sexual orientation.

Analysis—Descriptive statistics were calculated using SAS-callable SUDAAN v10 (SAS Institute Inc.) to assess smoke-free home and vehicle rule prevalence by sociodemographics and tobacco use. Chi-square tests were performed to determine overall significance for each tobacco use category (i.e., any tobacco; combustible only; noncombustible only; both combustible and noncombustible; and no current tobacco use) and sociodemographic variable ($p < 0.05$). Data were weighted to adjust for selection and nonresponse.

Results

Smoke-free home rules

Overall, 83.7% of adults had smoke-free home rules. By tobacco use, prevalence was highest among nonusers (90.8%), followed by noncombustible-only users (82.5%), both combustible and noncombustible users (54.9%), and combustible-only users (53.7%) (Table 1). Irrespective of tobacco use, prevalence was generally higher among males and

respondents aged 44 years and less, married or partnered, with a graduate degree, income of \$100,000, or living in the West (Table 1).

Chi-square findings revealed prevalence differences among adults who reported having 100% smoke-free home rules by tobacco use status and selected sociodemographic characteristics. Significant differences were noted in any tobacco users, combustible-only tobacco users, non-combustible-only tobacco users, and no current tobacco users by sex, age, marital status, annual household income, region and sexual orientation. Among any tobacco users, the prevalence of smoke-free rules in homes was greater for: men, married or partnered individuals, those with an annual household income of \$100,000, and residents of states in the West ($p < 0.05$). Among no current tobacco users, the prevalence of smoke-free rules in homes was greater for women, adults 25–44 years of age, married or partnered individuals, those with an annual household income of \$100,000, residents of states in the West, and heterosexual or straight individuals ($p < 0.05$).

Smoke-free vehicle rules

Overall, 78.1% of US adults had smoke-free vehicle rules (Table 2). By tobacco use, prevalence was highest among nonusers (88.9%), followed by noncombustible-only users, (69.1%), combustible-only users (34.2%), and both combustible and noncombustible users (24.2%). Irrespective of tobacco use, prevalence was generally higher among males and respondents aged 65, with a graduate degree, living in the West, or with unspecified sexual orientation.

Chi-square findings revealed prevalence differences among adults who reported having 100% smoke-free vehicle rules by tobacco use status and selected sociodemographic characteristics. Significant differences were noted in any tobacco users, combustible-only tobacco users, and no current tobacco users by sex, age, educational attainment, marital status, annual household income, region and sexual orientation. Among any tobacco users, the prevalence of smoke-free rules in vehicles was greater for men, residents of states in the West, and those with an unspecified sexual orientation ($p < 0.05$). Among no current tobacco users, the prevalence of smoke-free rules in vehicles was greater for women, adults aged 65 years of age, those with a graduate degree, married or partnered individuals, those with an annual household income of \$100,000, residents of states in the West, and those with an unspecified sexual orientation ($p < 0.05$).

Discussion

This study reveals that approximately 8 in 10 US adults report smoke-free rules at home (83.7%) and in their vehicle (78.1%). However, variations in such rules exist by tobacco use; about one-half of any tobacco users reported having smoke-free rules, while about one-third reported smoke-free vehicle rules. These findings indicate that opportunities exist to educate all tobacco users about the dangers of SHS and to promote voluntary adoption of smoke-free rules in private settings, particularly among combustible tobacco users. Additionally, jurisdictions with comprehensive policies prohibiting smoking in public places and worksites could extend SHS protections to include areas that are typically excluded from these policies. Continued efforts are critical to increase awareness of the hazards of

SHS exposure in US adults and the benefits of smoke-free home and vehicle rules to help inform state and local level capacity.

Among combustible-only users, 53.7% reported having smoke-free home rules (Table 1) and 34.2% reported having smoke-free vehicle rules (Table 2), both of which are higher than previously reported estimates among cigarette smokers (King et al., 2013; Ayo-Yusuf et al., 2014). Reasons for the high level of prevalence of smoke-free homes rules among respondents 44 years of age and less may be related to shifting social norms around the acceptability of smoke-free policies in indoor settings. The higher prevalence of smoke-free rules may be due to the proliferation of comprehensive smoke-free laws, declines in cigarette smoking, and changes in public attitudes regarding the acceptability of smoking near nonsmokers (IARC, 2009; Cheng et al., 2011, 2015; Hyland et al., 2009; Callinan et al., 2010).

Differences in the prevalence of smoke-free rules were apparent by tobacco use status. For example, smoke-free home and vehicle rule prevalence was higher among nonusers than any tobacco users. Since research suggests that smoke-free homes can enhance cessation among adult smokers and prevent relapse among former smokers (Hyland et al., 2009), efforts to encourage the adoption of voluntary smoke-free rules in private settings, including homes and vehicles, could reduce tobacco use. These findings suggest that although a larger proportion of any tobacco users have implemented voluntary smoke-free rules in homes and household vehicles, within the subsample, disparity in voluntary smoke-free rules in private settings exists. Sociodemographic differences were also observed, which could be due to varying smoking rates among these groups, factors related to social disapproval of smoking, or receptivity toward tobacco-related health messages and understanding of the dangers of SHS (CDC, 2011; Siahpush et al., 2006).

Strengths of this study include the use of a large, nationally representative sample and the assessment of multiple tobacco products. Limitations include the use of cross-sectional data, which are subject to recall bias. Additionally, variables of interest that may affect smoke-free rules in homes and cars, such as presence of children in the home or vehicle, were not included in the NATS questionnaire. Finally, limited sample size prevented the presentation of estimates for certain groups (e.g., noncombustible-only users).

Conclusion

Most US adults have implemented smoke-free home and vehicle rules; however, variations exist by tobacco use status. Completely eliminating smoking in indoor spaces is the only way to fully protect non-smokers from SHS; separating smokers from nonsmokers, cleaning the air, and ventilating buildings cannot completely eliminate exposure (2). Continued efforts to educate all tobacco users about the importance of voluntary smoke-free rules in homes and vehicles are warranted as part of a comprehensive approach to reducing tobacco use and SHS exposure, particularly among combustible tobacco users.

Acknowledgments

The findings and conclusions in this report are those of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention.

References

- Ayo-Yusuf OA, Olufajo O, Agaku IT. Exposure to secondhand smoke and voluntary adoption of smoke-free home and car rules among non-smoking South African adults. *BMC Public Health*. 2014; 14 <http://www.biomedcentral.com/1471-2458/14/580>.
- Callinan JE, Clarke A, Doherty K, Kelleher C. Legislative smoking bans for reducing secondhand smoke exposure, smoking prevalence and tobacco consumption. *Cochrane Libr*. 2010; (6) <http://dx.doi.org/10.1002/14651858.CD005992.pub2>.
- CDC. Vital signs: current cigarette smoking among adults aged 18 years—United States, 2005–2010. *MMWR Morb. Mortal. Wkly Rep*. 2011; 60:1207–1212. [PubMed: 21900875]
- CDC. Office on Smoking and Health and ICF. 2012–2013 National Adult Tobacco Survey Sample Design and Methodology Summary. 2013.
- CDC. Prevalence of smoke-free home rules—United States, 1992–1993 and 2010–2011. *MMWR Morb. Mortal. Wkly Rep*. 2014a; 63:765–769. [PubMed: 25188494]
- CDC. Tobacco product use among adults—United States, 2012–2013. *MMWR Morb. Mortal. Wkly Rep*. 2014b; 63:542–547. [PubMed: 24964880]
- CDC. State Tobacco Activities Tracking and Evaluation (STATE) System. 2015. <http://www.cdc.gov/tobacco/statesystem>
- Cheng K, Glantz SA, Lightwood JM. Association between smokefree laws and voluntary smokefree-home rules. *Am. J. Prev. Med*. 2011; 41:566–572. [PubMed: 22099232]
- Cheng K, Okechukwu ICA, McMillen R, Glantz SA. Association between clean indoor air laws and voluntary smokefree rules in homes and cars. *Tob. Control*. 2015; 24:168–174. <http://dx.doi.org/10.1136/tobaccocontrol-2013-051121>. [PubMed: 24114562]
- Healey B, Hoek J, Wilson N, Thomson G, Taylor S, Edwards R. Youth exposure to in-vehicle second-hand smoke and their smoking behaviors: trends and associations in repeated national surveys (2006–2012). *Tob. Control*. 2015; 24:146–152. <http://dx.doi.org/10.1136/tobaccocontrol-2013-051124>. [PubMed: 24046210]
- Hopkins DP, Razi S, Leeks KD, et al. Smoke-free policies to reduce tobacco use: a systematic review. *Am. J. Prev. Med*. 2010; 38:S275–S289. <http://dx.doi.org/10.1016/j.amepre.2009.10.029>. [PubMed: 20117612]
- Hyland A, Higbee C, Travers MJ, et al. Smoke-free homes and smoking cessation and relapse in a longitudinal population of adults. *Nicotine Tob. Res*. 2009; 11:614–618. [PubMed: 19346505]
- IARC. Evaluating the Effectiveness of Smoke-free Policies. Vol. 13. Lyon, France; 2009.
- King BA, Dube SR, Homa DM. Smoke-free rules and secondhand smoke exposure in homes and vehicles among US adults, 2009–2010. *Prev. Chronic Dis*. 2013;20218. <http://dx.doi.org/10.5888/pcd10.120218>.
- Siahpush M, McNeill A, Hammond D, Fong GT. Socioeconomic and country variations in knowledge of health risks of tobacco smoking and toxic constituents of smoke: results from the 2002 International Tobacco Control (ITC) Four Country Survey. *Tob. Control*. 2006; 15:S65–S70.
- USDHHS. The health consequences of involuntary exposure to tobacco smoke: a report of the Surgeon General. Department of Health and Human Services, Centers for Disease Control and Prevention; Atlanta, GA: 2006.
- USDHHS. Preventing tobacco use among young people: a report of the Surgeon General. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; Atlanta, GA: 2012.
- USDHHS. The health consequences of smoking—50 years of progress: a report of the Surgeon General. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; Atlanta, GA: 2014.

Percentage of adults who reported having 100% a smoke-free home rule, by tobacco use and selected sociodemographic characteristics—NATS, 2012–2013.

Table 1

Characteristics	Any tobacco ^d (n = 4791)	Combustible-only ^b (n = 3558)	Noncombustible-only ^c (n = 618)	Both combustible and noncombustible ^d (n = 155)	No current tobacco use ^e (n = 43,717)	Overall/ (n = 48,871)
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
<i>Overall</i>	56.4 (55.1–57.8)	53.7 (52.2–55.3)	82.5 (79.0–86.0)	54.9 (47.5–62.4)	90.8 (90.4–91.2)	83.7 (83.2–84.1)
<i>Sex</i>						
Male	60.2 (58.4–62.0) ^g	56.7 (54.6–58.9)	82.2 (78.6–85.7)	53.7 (45.9–61.5)	90.2 (89.5–90.8) ^g	82.3 (81.6–83.0)
Female	50.6 (48.4–52.7)	49.7 (47.4–52.1)	<i>h</i>	<i>h</i>	91.3 (90.8–91.8)	85.0 (84.4–85.6)
<i>Age group (years)</i>						
18–24	63.1 (58.8–67.3)	59.5 (54.3–64.7)	82.8 (72.8–92.8)	60.0 (46.6–73.4)	85.7 (83.9–87.5) ^g	80.4 (78.6–82.1)
25–44	65.4 (63.1–67.8)	63.6 (60.9–66.3)	89.0 (84.1–94.0)	59.4 (48.3–70.5)	92.6 (91.9–93.4)	85.9 (85.1–86.8)
45–64	46.8 (44.8–48.8)	44.4 (42.1–46.6)	76.7 (70.6–82.8)	36.3 (22.1–50.4)	91.6 (91.1–92.2)	81.9 (81.1–82.6)
65	43.4 (40.2–46.7)	39.8 (36.2–43.3)	78.8 (70.6–87.0)	<i>h</i>	89.6 (89.0–90.2)	85.3 (84.6–85.9)
<i>Race/Ethnicity</i>						
White, non-Hispanic	57.7 (56.0–59.3)	54.7 (52.8–56.6)	82.6 (78.8–86.4)	55.8 (47.4–64.1)	91.4 (91.0–91.9)	84.4 (83.9–84.9)
Black, non-Hispanic	41.0 (36.4–45.5)	39.7 (34.9–44.5)	<i>h</i>	<i>h</i>	86.6 (84.9–88.3)	76.4 (74.5–78.2)
Asian, non-Hispanic	80.6 (69.0–92.2)	76.8 (63.1–90.5)	<i>h</i>	<i>h</i>	92.6 (90.6–94.6)	91.5 (89.2–93.4)
Other non-Hispanic	49.1 (44.9–53.3)	45.5 (40.7–50.3)	82.0 (71.2–92.8)	46.2 (26.8–65.5)	88.7 (86.8–90.6)	75.6 (73.5–77.6)
Hispanic	69.5 (65.0–73.9)	68.1 (63.2–73.0)	<i>h</i>	<i>h</i>	91.0 (89.8–92.3)	87.7 (86.3–88.9)
<i>Educational attainment</i>						
0–12 years (no diploma)	46.5 (42.5–50.6)	45.4 (40.9–49.9)	77.3 (65.4–89.2)	<i>h</i>	86.6 (84.9–88.3)	75.1 (73.3–76.9)
General educational diploma	44.0 (37.1–50.8)	44.9 (37.4–52.4)	<i>h</i>	<i>h</i>	85.3 (81.7–88.9)	67.6 (63.5–71.4)
High school graduate	54.9 (52.4–57.5)	51.0 (48.1–54.0)	82.2 (76.2–88.1)	53.7 (41.5–65.9)	88.0 (87.0–89.0)	80.0 (79.0–81.0)
Some college (no degree)	59.6 (56.7–62.4)	56.9 (53.6–60.1)	84.6 (77.3–91.9)	<i>h</i>	90.8 (89.8–91.8)	83.4 (82.3–84.4)
Associate degree	60.5 (57.4–63.7)	57.7 (54.0–61.4)	84.7 (76.4–92.9)	<i>h</i>	92.0 (91.1–92.9)	85.2 (84.2–86.2)
Undergraduate degree	70.1 (67.0–73.2)	66.7 (63.0–70.4)	87.7 (81.5–93.9)	<i>h</i>	94.2 (93.6–94.7)	91.5 (90.9–92.1)
Graduate degree	71.3 (66.3–76.3)	70.6 (65.1–76.1)	<i>h</i>	<i>h</i>	94.6 (94.0–95.2)	93.1 (92.4–93.8)
<i>Marital status</i>						
Married/living with partner	61.7 (59.8–63.6) ^g	58.9 (56.7–61.2) ^g	84.0 (79.7–88.2)	61.6 (51.1–72.1)	93.0 (92.6–93.4) ^g	87.4 (86.9–87.9)
Single/separated/divorced/widowed	51.4 (49.4–53.4)	49.1 (46.8–51.4)	79.9 (74.0–85.9)	49.5 (39.2–59.8)	87.6 (86.9–88.4)	78.8 (78.0–79.6)

Characteristics	Any tobacco ^d	Combustible-only ^b	Noncombustible-only ^c	Both combustible and noncombustible ^d	No current tobacco use ^e	Overall ^f
	(n = 4791) % (95% CI)	(n = 3558) % (95% CI)	(n = 618) % (95% CI)	(n = 155) % (95% CI)	(n = 43,717) % (95% CI)	(n = 48,871) % (95% CI)
<i>Annual household income</i>						
<\$20,000	39.4 (36.0–42.8) ^g	37.0 (33.3–40.8) ^g	71.6 (57.3–86.0) ^g	<i>h</i>	86.3 (84.7–87.8) ^g	72.3 (70.5–73.4)
\$20,000–\$49,999	50.1 (47.7–52.5)	48.5 (45.8, 51.2)	75.6 (68.1–83.2)	50.7 (38.4–62.9)	88.0 (87.1–88.9)	78.3 (77.3–79.3)
\$50,000–\$99,999	65.2 (62.5–67.9)	63.0 (59.8, 66.1)	83.4 (77.4–89.4)	55.7 (40.6–70.7)	92.2 (91.6–92.9)	87.0 (86.2–87.8)
\$100,000	77.1 (73.8–80.5)	75.3 (71.2, 79.5)	92.2 (87.0–97.5)	71.1 (55.0–87.2)	95.3 (94.7–95.9)	93.0 (92.2, 93.6)
Unspecified	58.6 (54.6–62.5)	56.5 (52.1, 60.8)	85.6 (75.9–95.3)	<i>h</i>	90.0 (88.9–91.1)	84.7 (83.5–85.9)
<i>US census regionⁱ</i>						
Northeast	48.8 (45.2–52.5) ^g	45.5 (41.5–49.5) ^g	70.7 (57.4–84.0)	<i>h</i>	90.4 (89.5–91.4) ^g	82.8 (81.7–83.9)
Midwest	51.0 (48.2–53.9)	48.9 (45.6–52.2)	80.4 (73.0–87.8)	54.3 (38.9–69.7)	89.1 (88.2–90.0)	80.4 (79.3–81.4)
South	57.2 (55.0–59.4)	54.2 (51.6–56.7)	84.0 (79.3–88.8)	50.1 (39.7–60.5)	90.4 (89.7–91.1)	82.9 (82.1–83.6)
West	67.4 (64.6–70.3)	65.7 (62.4–68.9)	88.0 (81.0–94.9)	65.6 (47.0–84.2)	93.1 (92.3–93.8)	88.6 (87.8–89.4)
<i>Sexual orientation</i>						
Heterosexual/Straight	56.9 (55.5–58.4)	54.1 (52.4–55.8)	82.4 (78.9–86.0)	55.7 (48.1–63.4)	91.1 (90.7–91.5) ^g	84.0 (83.5–84.5)
Lesbian/Gay/Bisexual	51.0 (44.8–57.2)	49.1 (42.2–56.1)	<i>h</i>	<i>h</i>	85.6 (82.6–88.7)	74.9 (71.8–77.8)
Unspecified	54.4 (47.0–61.8)	53.0 (45.0–60.9)	<i>h</i>	<i>h</i>	89.6 (87.7–91.5)	85.0 (82.9–86.9)

Abbreviations: CI= confidence interval; NATS = National Adult Tobacco Survey.

^a Defined as ‘every day’ or ‘some day’ use of cigarettes, cigars/cigarillos/filtered little cigars, regular pipes, and water pipes/hookah, electronic cigarettes, or chewing tobacco/snuff/dip, snus, or dissolvable tobacco.

^b Defined as smoking at least 100 cigarettes during their lifetime and now smoking ‘every day’ or ‘some day’ and/or used 1 of the following tobacco product types and now use ‘everyday’ or ‘some day’: cigars/cigarillos/filtered little cigars, regular pipes, and water pipes/hookah 50 times in their lifetime and now smoked the product ‘every day’ or ‘some days’.

^c Defined as using chewing tobacco, snuff or dip 20 times in their lifetime, snus or dissolvable tobacco products on 1 day, and used these products ‘every day’ or ‘some days’.

^d Defined as ‘every day’ or ‘some day’ use of any combustible tobacco product and any noncombustible tobacco product.

^e Defined as not currently using electronic cigarettes, combustible tobacco, or noncombustible tobacco.

^f Includes persons whose tobacco use status is unknown.

^g Significant chi-square test ($p < 0.05$) indicated difference across groups within the specified characteristic.

^h Excluded because relative standard error = 30%.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

i Northeast: Connecticut, Maine, Massachusetts, New Jersey, New Hampshire, New York, Pennsylvania, Rhode Island, and Vermont; *Midwest*: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin; *South*: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia; *West*: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

Percentage of adults who reported having 100% smoke-free vehicle rules, by tobacco use and selected sociodemographic characteristics—NATS, 2012–2013.

Table 2

Characteristics	Any tobacco ^d (n = 3186)	Combustible-only ^b (n = 2290)	Noncombustible-only ^c (n = 537)	Both combustible and noncombustible ^d (n = 75)	No current tobacco use ^e (n = 42,649)	Overall/ tobacco use ^f (n = 46,183)
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
<i>Overall</i>	36.5 (35.1–37.8)	34.2 (32.7–35.8)	69.1 (64.7–73.6)	24.2 (18.0–30.4)	88.9 (88.4–89.3)	78.1 (77.6–78.6)
<i>Sex</i>						
Male	38.5 (36.7–40.3) ^g	35.5 (33.4–37.6)	68.7 (64.1–73.2)	22.9 (16.5–29.2)	88.3 (87.6–89.0) ^g	75.4 (74.6–76.1)
Female	33.2 (31.1–35.3)	32.5 (30.3–34.7)	80.2 (63.3–97.2)	<i>h</i>	89.4 (88.8–89.9)	80.7 (80.1–81.4)
<i>Age Group (years)</i>						
18–24	35.4 (31.2–39.6)	33.1 (28.1–38.0)	69.9 (57.3–82.6)	<i>h</i>	80.7 (78.8–82.7) ^g	70.3 (68.3–72.2)
25–44	37.7 (35.3–40.0)	36.1 (33.4–38.7)	69.2 (61.4–77.0)	26.2 (16.5–35.8)	89.6 (88.8–90.5)	76.9 (75.9–77.9)
45–64	34.0 (32.0–35.9)	31.2 (29.0–33.3)	67.0 (60.2–73.7)	29.3 (16.5–42.2)	89.7 (89.1–90.3)	77.8 (77.0–78.5)
65	43.2 (39.9–46.5)	40.6 (36.9–44.3)	75.4 (66.6–84.1)	<i>h</i>	91.2 (90.6–91.7)	86.7 (86.0–87.3)
<i>Race/Ethnicity</i>						
White, non-Hispanic	32.4 (30.9–34.0)	28.3 (26.6–30.1)	68.6 (63.7–73.5)	26.4 (19.0–33.7)	88.3 (87.8–88.8)	76.8 (76.2–77.4)
Black, non-Hispanic	42.1 (37.4–46.7)	40.6 (35.8–45.5)	86.8 (72.6–100.9)	<i>h</i>	87.2 (85.5–88.9)	77.3 (75.4–79.1)
Asian, non-Hispanic	51.9 (36.0–67.8)	53.2 (35.5–70.9)		<i>h</i>	94.5 (92.7–96.2)	90.7 (87.7–93.0)
Other non-Hispanic	31.4 (27.5–35.3)	29.6 (25.1–34.0)	65.3 (50.7–79.9)	<i>h</i>	86.4 (84.4–88.5)	68.5 (66.2–70.7)
Hispanic	58.4 (53.6–63.1)	59.7 (54.6–64.8)	64.7 (37.2–92.1)	<i>h</i>	91.9 (90.7–93.1)	86.7 (85.3–87.9)
<i>Educational Attainment</i>						
0–12 years (no diploma)	39.6 (35.6–43.7)	40.0 (35.5–44.5)	58.8 (45.1–72.6)	<i>h</i>	87.1 (85.4–88.8) ^g	73.6 (71.7–75.5)
General educational diploma	28.1 (21.9–34.3)	27.2 (20.5–34.0)	46.1 (8.5–83.7)	<i>h</i>	84.9 (81.1–88.8)	60.3 (55.9–64.5)
High school graduate	36.1 (33.5–38.6)	33.4 (30.6–36.3)	67.2 (59.3–75.1)	20.4 (10.3–30.4)	86.1 (85.0–87.1)	74.0 (72.9–75.1)
Some college (no degree)	32.7 (29.8–35.5)	30.6 (27.4–33.7)	67.1 (56.1–78.0)	<i>h</i>	87.5 (86.4–88.6)	74.6 (73.3–75.8)
Associate degree	34.5 (31.3–37.6)	30.6 (27.1–34.2)	72.0 (61.2–82.8)	<i>h</i>	88.6 (87.6–89.7)	77.0 (75.8–78.3)
Undergraduate degree	43.2 (39.7–46.7)	37.5 (33.5–41.4)	80.2 (71.9–88.5)	<i>h</i>	91.7 (91.0–92.3)	86.4 (85.6–87.2)
Graduate degree	48.7 (43.1–54.3)	46.1 (39.9–52.3)	92.1 (84.8–99.5)	<i>h</i>	94.1 (93.5–94.8)	91.2 (90.4–92.0)
<i>Marital Status</i>						
Married/living with partner	38.1 (36.2–40.0)	35.1 (32.9–37.3)	69.0 (63.3–74.6)	28.1 (19.0–37.2)	90.8 (90.3–91.3) ^g	81.4 (80.8–82.0)

Characteristics	Any tobacco ^d	Combustible-only ^b	Noncombustible-only ^c	Both combustible and noncombustible ^d	No current tobacco use ^e	Overall ^f
	(n = 3186)	(n = 2290)	(n = 537)	(n = 75)	(n = 42,649)	(n = 46,183)
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Single/separated/divorced/widowed	34.7 (32.8–36.7)	33.4 (31.2–35.6)	68.7 (61.4–76.1)	21.0 (12.5–29.6)	86.1 (85.3–86.9)	73.7 (72.8–74.6)
<i>Annual Household Income</i>						
<\$20,000	36.3 (32.8–39.8)	35.9 (32.1–39.7)	55.1 (38.7–71.5)	<i>h</i>	85.0 (83.4–86.7) ^g	70.7 (68.9–72.5)
\$20,000 - \$49,999	33.0 (30.7–35.3)	31.6 (29.0–34.2)	64.3 (55.5–73.2)	20.5 (10.7–30.3)	87.0 (86.1–87.9)	73.2 (72.2–74.2)
\$50,000 - \$99,999	35.3 (32.6–37.9)	31.8 (28.7–34.8)	63.5 (55.3–71.6)	21.0 (10.4–31.7)	88.5 (87.7–89.4)	78.3 (77.3–79.2)
\$100,000	45.0 (41.0–48.9)	39.8 (35.2–44.4)	84.6 (77.0–92.2)	41.0 (23.7–58.3)	92.9 (92.2–93.7)	86.8 (85.8–87.7)
Unspecified	41.1 (37.1–45.1)	40.1 (35.6–44.5)	77.4 (66.4–88.4)	<i>h</i>	89.6 (88.5–90.7)	81.6 (80.3–82.8)
<i>US Census Regionⁱ</i>						
Northeast	39.0 (35.4–42.7) ^g	37.2 (33.2–41.2)	78.7 (66.6–90.7)	<i>h</i>	89.1 (88.1–90.1) ^g	80.1 (78.9–81.2)
Midwest	29.6 (26.9–32.3)	27.5 (24.5–30.6)	58.6 (49.1–68.2)	<i>h</i>	86.3 (85.3–87.4)	73.4 (72.2–74.6)
South	35.8 (33.6–37.9)	32.3 (29.8–34.7)	70.2 (64.0–76.4)	26.9 (17.8–35.9)	87.8 (87.0–88.5)	76.0 (75.1–76.9)
West	43.7 (40.6–46.7)	43.2 (39.8–46.7)	76.4 (65.8–86.9)	<i>h</i>	92.5 (91.7–93.2)	84.2 (83.2–85.0)
<i>Sexual Orientation</i>						
Heterosexual/Straight	35.8 (34.4–37.3) ^g	33.4 (31.8–35.1) ^g	68.5 (63.9–73.1)	24.4 (17.9–30.9)	88.9 (88.5–89.4) ^g	78.1 (77.5–78.6)
Lesbian/Gay/Bisexual	31.7 (25.8–37.6)	30.5 (23.9–37.1)	80.3 (52.5–108.2)	<i>h</i>	81.8 (78.4–85.3)	66.5 (63.1–69.7)
Unspecified	58.3 (50.9–65.7)	56.6 (48.5–64.7)	78.3 (59.7–96.9)	<i>h</i>	92.2 (90.6–93.7)	87.6 (85.7–89.2)

Abbreviations: CI= confidence interval; NATS = National Adult Tobacco Survey.

^aDefined as ‘every day’ or ‘some day’ use of cigarettes, cigars/cigarillos/filtered little cigars, regular pipes, and water pipes/hookah, electronic cigarettes, or chewing tobacco/snuff/dip, snus, or dissolvable tobacco.

^bDefined as smoking at least 100 cigarettes during their lifetime and now smoking ‘every day’ or ‘some day’ and/or used 1 of the following tobacco product types and now use ‘everyday’ or ‘some day’: cigars/cigarillos/filtered little cigars, regular pipes, and water pipes/hookah 50 times in their lifetime and now smoked the product ‘every day’ or ‘some days’.

^cDefined as using chewing tobacco, snuff or dip 20 times in their lifetime, snus or dissolvable tobacco products on 1 day, and used these products ‘every day’ or ‘some days’.

^dDefined as ‘every day’ or ‘some day’ use of any combustible tobacco product and any noncombustible tobacco product.

^eDefined as not currently using electronic cigarettes, combustible tobacco, or noncombustible tobacco.

^fIncludes persons who have unknown tobacco use status.

^gSignificant chi-square test ($p < 0.05$) indicated difference across groups within the specified characteristic.

^hExcluded because relative standard error > 30%.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

i Northeast: Connecticut, Maine, Massachusetts, New Jersey, New Hampshire, New York, Pennsylvania, Rhode Island, and Vermont; *Midwest*: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin; *South*: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia; *West*: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.