



HHS Public Access

Author manuscript

Cancer Epidemiol Biomarkers Prev. Author manuscript; available in PMC 2015 December 17.

Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2013 December ; 22(12): 2148–2160. doi:
10.1158/1055-9965.EPI-13-0591.

Collaborative Cancer Epidemiology in the 21st Century: the Model of Cancer Consortia

Michael R. Burgio^{(1),(3)}, John P.A. Ioannidis⁽⁴⁾, Brett M. Kaminski⁽¹⁾, Eric DeRycke⁽¹⁾, Scott Rogers⁽¹⁾, Muin J. Khoury^{(1),(2)}, and Daniela Seminara⁽¹⁾

⁽¹⁾Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda, Maryland, USA

⁽²⁾Office of Public Health Genomics, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

⁽³⁾Scientific Consulting Group, Inc., Gaithersburg, Maryland, USA

⁽⁴⁾Stanford Prevention Research Center, Department of Medicine, and Department of Public Health and Policy, Stanford University School of Medicine, and Department of Statistics, Stanford University School of Humanities and Sciences, Stanford, California, USA

Abstract

During the last two decades, epidemiology has undergone a rapid evolution toward collaborative research. The proliferation of multi-institutional, interdisciplinary consortia has acquired particular prominence in cancer research. Herein, we describe the characteristics of a network of 49 established cancer epidemiology consortia (CEC) currently supported by the Epidemiology and Genomics Research Program (EGRP) at the National Cancer Institute (NCI). This collection represents the largest disease-based research network for collaborative cancer research established in population sciences. We describe the funding trends, geographic distribution and areas of research focus. The CEC have been partially supported by 201 grants and yielded 3876 publications between 1995 and 2011. We describe this output in terms of interdisciplinary collaboration and translational evolution. We discuss challenges and future opportunities in the establishment and conduct of large-scale team science within the framework of CEC, review future prospects for this approach to large scale, interdisciplinary cancer research and describe a model for the evolution of an integrated Network of Cancer Consortia optimally suited to address and support 21st century epidemiology.

Keywords

consortium; epidemiology; cancer; interdisciplinary research; translation

Corresponding Author: Michael R. Burgio, National Institutes of Health, National Cancer Institute, Epidemiology and Genomics Research Program, Division of Cancer Control and Population Sciences, 9609 Medical Center Drive, Rm. 4E320, MSC 9763, Rockville, MD 20850, telephone: (240) 276-6883, fax: (240) 276-7921, burgiom@mail.nih.gov.

Conflict of Interest: The authors declare no conflicts of interest. The findings and conclusions in this report are those of the authors and do not necessarily reflect the views of the Department of Health and Human Services.

Introduction

The transition towards large scale collaborations has been a hallmark in many fields of research in the last few decades (1). In epidemiology, and especially in the genomic epidemiology of complex diseases, this trend has been supported by a convergence of factors, including the rapid development of increasingly sophisticated genomic technologies (2), the progressive building of large population resources such as cohorts and biobanks (3, 4), and the requirement for larger sample sizes to address small effects. As a consequence, epidemiologic research on the genetic and environmental determinants of complex diseases has experienced a paradigm shift toward “Big and Bigger Science”, embodied by the rise of consortia as “hubs” of collaborative and interdisciplinary research within the framework of population sciences (5). Cancer epidemiology has been markedly affected by the advent of such collaborative infrastructures. Here, we examine the impact of research originating from interdisciplinary cancer consortia from 1995 to 2011, based on our experience with a network of 49 cancer-related consortia supported by the Epidemiology and Genomics Research Program (EGRP) of the National Cancer Institute (NCI). To our knowledge, this is the largest disease-based research network for collaborative, population-based research currently in existence. The reported analysis offers insights about the growth, impact, and future prospects of CECs as well as their role in supporting high-impact interdisciplinary research.

Definition

We define a consortium as “*A group of scientists from multiple institutions who have agreed to cooperative research efforts involving, but not limited to, pooling of information from more than one study for the purpose of combined analyses and collaborative projects. Such consortia are geared to address scientific questions that cannot otherwise be addressed through the effort of a team of investigators at a single institution due to scope, resources, population size, or the need for an interdisciplinary approach*” (6). The Cancer Epidemiology Consortia (CEC) network is a group of eligible cancer consortia which have received different forms of support by the National Cancer Institute (NCI) since 1995.

Consortia included in this network have either contacted the NCI/EGRP to be listed or have been specifically solicited through a targeted initiative. When applying to be listed on the EGRP website, each consortium is required to provide the following to be eligible: a description of cancer-related research questions that can be uniquely addressed by that consortium because of its characteristics (e.g. size and characteristics of population, enrolled biobank infrastructure, involvement of an interdisciplinary team of scientists from multiple institutions, etc.); an existing or proposed organizational structure and leadership plan; and a statement of commitment to data sharing within and outside the consortium. Since emerging consortia can experience a substantial lag time before beginning to publish in a substantive manner, we have limited the reported analyses to established cancer epidemiology consortia (CEC) that were launched before 2010. A complete list of the 49 established EGRP consortia included in the analyses, along with descriptive information on each appears in Table 1.

Characteristics of the Cancer Epidemiology Consortia

A common characteristic of the 49 established CEC is that each team has not assembled uniquely to execute one project, but their collaborations extend through time and across research projects that vary in design and complexity. Only 3 of these CEC had assembled in response to solicitations by the NCI or other NIH institutes (7–9), with the remainder coalescing spontaneously to address diverse research agendas. Funding for consortia-related research is provided by EGRP/NCI through a variety of investigator-initiated mechanisms or through support for communication and research networking activities such as meetings or teleconferences. Most consortia (n=41, 84%) focus on a single cancer type. The remaining 8 (16%) CEC study multiple cancers, specific translational topics (e.g. radiogenomics), or focus on diverse ethnic populations (Hispanics, African, Asian and Caribbean) in US and abroad (Table 1). Most CECs are international in nature, with the largest distribution of collaborating groups in high and mid-level income countries (Supplemental Figure 1). All 49 CECs have some type of associated biorepository.

Websites for CEC and Consortium Policies

Public websites are an essential tool for communication and global sharing of study results, dissemination of research tools, and provide a conduit to data sharing and research opportunities for large collaborative groups. Forty (82%) CEC have developed publicly available websites (reviewed from March 4–14, 2013): thirty-four (85%) included information on CEC leadership and twenty (50%) detailed the CEC organizational structure. Eighteen (45%) of the CEC websites included information on consortium membership requirements, nine (23%) included submission guidelines for new project proposals, and seven (18%) had eligibility requirements and contact information for participant enrollment. Twenty-one (53%) of the CEC websites included a restricted access area (portal) reserved for consortia members communication and internal data sharing.

CEC websites (accessed May 13–16, 2013), associated grant applications, and descriptive manuscripts were reviewed to determine if CEC had established data sharing policies, as was intended. In cases where no policy was found the CEC liaison or lead investigator was contacted and asked if the CEC had a data sharing policy in place. Overall, 29 (59%) had data sharing policies, 3 (6.1%) were in the process of developing them, 10 (20.4%) did not have policies in place, and for 8 (16.3%) CEC we were unable to confirm whether or not they had data sharing policies. Consortia supported entirely or in part through NCI-awarded grants and cooperative agreements are mandated to comply with the NIH data and resource sharing policies for what concerns the specific aims listed on the funded grants or cooperative agreements (10).

Grants Funding for CEC

In order to evaluate the investment in terms of funding support and the scientific productivity of CEC, we reviewed all the EGRP grants that were related to the 49 established CEC. Overall, 201 grants, funded by EGRP between fiscal year (FY) 1995 and 2011, were identified as consortium related by searching the NCI Portfolio Management Application (PMA) database (v14.0.3). Grant coding is conducted by EGRP program staff

and consortium codes were confirmed through a manual review of the EGRP grant portfolio. A grant was defined as consortium- related if it directly supported the main research activities and/or infrastructure of the consortium, or if it explicitly relied on the consortium's resources to conduct the proposed research project. An analysis of CEC-related grants shows a linear increase of investment in consortial research by EGRP from FY 1995–2011 (Supplemental Figures 2A and 2B). The average yearly increase in total number and total direct cost for EGRP CEC associated grants was 5.3 and \$4.2 million respectively; in contrast the total number of grants funded by EGRP has been flat since 1997 and the total direct cost of the whole portfolio has increased at a much slower rate (Supplemental Figures 3A and 3B). Starting in 2002, funding for projects that are collaborations between CEC increased over time. This reflects the increasing number of CEC as well as increases in the size and complexity of the collaborative network. While the total costs of CEC associated grants has been increasing, the average cost per grant has been relatively flat since 1998 and the fraction of small grants (<\$250K in direct costs per year) increased substantially from 2002 to 2007. This trend towards smaller grants is not seen in the overall EGRP portfolio (Supplemental Figures 4A and 4B).

The success rate of CEC-related grants (percentage of reviewed applications that receive funding) was compared to the success rate of EGRP-funded and NIH-funded grants (all mechanisms). Data were extracted from NIH RePORTER (11) (accessed 5/6/2013) and the NCI Portfolio Management Application (PMA) database (v14.0.3, accessed 4/30/2013). The success rate of CEC-related grants (48%) was consistently higher than the success rates for EGRP grants (28%) and NIH grants (25%) since FY 2000 (Table 2). This may reflect many factors, including the ready availability of resources and infrastructure in established consortia, increased communication across participating scientists and groups (12) as well as a more intense pre-submittal review of grants applications by the multiple participating investigators.

Productivity and Impact

To measure CEC scientific productivity, 3,876 CEC-related manuscripts were identified using three different methods. First, the CEC websites (if available) were searched for listings of manuscripts (websites initially accessed the week February 20, 2012 and checked for updates on 1/21/2013). Second, the CEC names and abbreviations were used as search terms for PubMed (13) queries (search performed on April, 30 2012); the titles and abstracts of these manuscripts were reviewed and results returned due to ambiguity in the search terms (such as a different organization having the same abbreviation as the CEC) were excluded. Finally, the NCI code and serial numbers of the 201 consortium-related EGRP grants (as identified above) were used as search terms in PubMed (search performed on May 13, 2012) to identify grant-related manuscripts. Results of these searches were combined and duplicate manuscripts and manuscripts published before the initiation year of the oldest, associated consortium and after 2011 were removed. The ascertainment and censoring, of EGRP CEC related publications, is summarized in Supplemental Figure 5.

The number of CEC associated papers published each year has increased linearly since 1998 (Figure 1A). Furthermore, a PubMed search for the terms “consortia OR consortium” NOT

“bacteria OR microbe OR microbial” (to exclude papers on microbial consortia; searched on 1/24/13) reveals an exponential increase in the number of papers containing these terms in their titles, abstracts, or authorship from 1985 to 2012 (Figure 1B). Further refining those search results by searching for the terms “cancer OR tumor” also reveal an exponential increase in cancer consortia papers since 1985. The rise of genome-wide association studies and the consequent need for extremely large numbers to reach adequate power has been cited as a major impetus for the formation of CEC (14, 15), however the trend towards team science in cancer research began well before publication of the first GWAS. Twenty-one (43%) of the CEC in the EGRP network were initiated before the first GWAS in the NHGRI GWAS Catalogue (16, 17).

CEC and Interdisciplinary Research

It has been proposed that team science, and therefore CEC-related science, should be ideally geared to support interdisciplinary research (18, 19). We define interdisciplinary research as “*a mode of research by teams or individuals that integrates information, data, techniques, tools, perspectives, concepts, and/or theories from two or more disciplines or bodies of specialized knowledge to advance fundamental understanding or to solve problems whose solutions are beyond the scope of a single discipline or area of research practice*” (20). To ascertain whether the publications produced by the 49 CEC examined are reflecting the CEC capability to support interdisciplinary research within an epidemiologic framework, the titles and abstracts for each of the 3,876 papers identified through the literature search described above were reviewed and each paper was assigned a primary scientific area. Papers were randomly assigned to 1 of 3 reviewers, with 10% of the papers assigned to all 3, and reviews were conducted over 14 rounds. Between each round, reviewers convened to discuss difficult to categorize papers, resolve discrepancies, and refine definitions. The scientific areas considered and their definitions are presented in Supplementary Table 1. A total of 3,729 papers were assigned a primary scientific area. (See supplemental methods for exclusion criteria.) 56% of the papers being scored fell into the environmental, lifestyle, and genomic epidemiology categories. 11.9% of the papers involved development of new methods and technologies, 9.4% focused on clinical and translational research, and 11% were classified as biology which encompasses basic laboratory research, including studies in cell lines and animal models (Supplemental Figure 6). Considering the CEC consortia were primarily designed to address questions within the framework of population sciences, the diversity of the associate literature is striking and demonstrates the flexibility of interdisciplinary nature of the CECs.

The types of research projects being undertaken by CECs have extended their interdisciplinary scope, and evolved with the recent “genomic revolution” as shown by the trends in the CEC publication’s scientific areas over time (Figure 2). Classic epidemiology studies evaluating environmental and lifestyle exposures represent a large proportion of this literature and increase over time (1995–2011). However the number of CEC papers in genomics areas, including candidate genes (CG), gene characterization (GC), genome-wide association (GWA), linkage (LK), loss of heterozygosity (LOH), or next generation sequencing (NGS) has grown significantly since 2000 and since 2010 genomics papers represented the largest category of EGRP CEC-associated publications. The growth in

genomics publications is largely driven by the exponential growth of GWAS, which (as defined here) include initial scans as well as replication studies (Supplemental Figure 7). This trend mirrors the decline of linkage publications as the search for genetic determinants of cancer risk shifted from searching for rare, highly penetrant variations in family studies to the search for common variants with small effect sizes, primarily through association studies (Supplemental Figure 8). Publications pertaining to methods and technologies, biology, molecular epidemiology and clinical and translation areas show a more modest but steady increase.

Team Science and Translation

To evaluate the contribution of EGRP CECs to translational research 3,363 papers from the literature search were reviewed and coded by translational research phase. (See supplemental methods for exclusion criteria.) The phases of the translational research continuum have been previously described (21–24) and are summarized with example papers from our literature search in Table 3 (25–34). Coding for translational research phase was conducted concurrently with coding for primary scientific area following the same procedures. Overall, 2645 (79%) papers were coded as T0, 582 (17%) as T1, 112 (3.3%) as T2, 18 (0.5%) as T3, and only 6 (0.2%) as T4.

Previous analyses have observed that T2 and above studies account for a small fraction of NIH-funded cancer genetics research (24). It has been estimated that only 0.64% of cancer genetics papers published in 2007 would be scored T2 or above (23). In our database of CEC publications 136 (3.5%) were scored T2+ and when we limited CEC publications to cancer genetics papers published in 2007 we found 3.4% (6 out of 171) were scored T2 or above. The increased proportion of T2 or above papers among CEC associated cancer genetics papers was statistically significant (P -value 2.4×10^{-6} ; See supplemental methods for details of this analysis). This enrichment of papers further down the translational continuum indicates that the collaborative and interdisciplinary infrastructure of CEC may specifically facilitate translational research.

Challenges and Future Prospects

The extended CEC network examined does not include all consortia focusing on cancer research, but is highly representative of that segment of cancer research focusing on human populations to understand the causes of cancer and related outcomes. The CEC network is international in scope, allowing for the study of populations with diverse genetic background and lifestyles, and encompasses studies with a variety of designs, from familial to case-control to prospective cohorts. The geographic distribution of the participating teams is still showing under-representation in low income regions, reflecting the need for infrastructure building in order to enrich the network to include populations with diverse genetic backgrounds, lifestyles and cultures. This may also reflect the fact that NIH funds primarily USA-based investigators and a more in-depth analysis of other sources of funding for international groups may reveal a more comprehensive panorama of international consortia. Tools of virtual communication (websites, portals) are widely used, but public dissemination of internal policies and processes and membership/participation criteria are still somewhat

limited in this Network. In this regard, complete transparency could greatly facilitate scientific exchange and rapid replication of initial results. Increasing the public posting of clear data sharing policies would facilitate not only collaborative projects among the investigators within the consortium, but also ready access to the consortia resources from new investigators interested in initiating collaborations thereby extending the consortia network. Consortia investigators are usually funded through a mosaic of mechanisms awarded by different funding agencies, and only rarely through special initiatives. For grants funded through NCI, adherence of the Principal Investigators to the NIH and NCI policies of data sharing is monitored through the lifetime of the grant and the application of current NIH/NCI policies on data sharing is mandatory and a condition for funding. For example, for some EGRP-funded consortia, applications for collaborative projects are tracked through a process of review and approval, and followed for productivity until completion (35, 36). It is to be noted that the data-sharing policies implemented by other national and international agencies vary considerable both in content and implementation (37). The NCI is in the process of establishing a database of consortia including the public posting of the internal and agency-mandated data sharing policies and to encourage maximum transparency. Increased transparency could greatly facilitate scientific exchange and rapid replication of initial results. A funding trend towards smaller CEC associated grants may indicate that while CEC require an initial substantial expenditure to establish the infrastructure necessary to conduct large-scale studies, their subsequently established resources and collaborative culture can be leveraged to support cost-efficient research effectively. Increased collaborations and synergism across consortia is also shown by the involvement of multiple CEC in individual grants. The increased success rate for consortia-related grant applications, as compared to the success rate for applications submitted to EGRP and NCI, may be symptomatic of the leverage provided by the extended consortia infrastructure and the extensive pre-application scrutiny and constructive pre-review usually provided by the consortia teams.

Our ascertainment of CEC-related publications has some limitations. Publications listed on a consortium's website are likely to be truly consortia-associated. However, 9 of the consortia did not have public websites, and of the remaining 40 websites, only 31 (62%) displayed a publication list. The completeness and update frequencies of the website publication lists were also variable. Publications captured by querying PubMed for the CEC name or abbreviation (followed by confirmation through independent review) are, almost by definition, CEC-related. However, CEC publication policies vary considerably and it is not uncommon for contributing CEC to only be mentioned in the methods, acknowledgments, or supplementary materials of a paper, i.e. sections that are not queried in PubMed searches, leading to under-ascertainment. Examination of the manuscripts associated with a sample of four established large-scale consortia funded by NIH (two of which were not included in the analyses presented in this manuscript as they did not respond to the selection criteria used) shows that acknowledgment of publications in consortia-related manuscripts, cited by their websites or listed in the associated grants, varied considerably across consortia, from 100% to 52%, (see Supplemental Table 2). This may be due to journal policies on acknowledgment format, to the lack of standard acknowledgement language distributed to the collaborators or to the absence of appropriate consortium acknowledgment requirements.

CEC-associated grants often encompass multiple specific aims, which may or may not require CEC resources. Therefore, when searching by acknowledged CEC-associated grant numbers we may be over-ascertaining CEC papers; this is in contrast to the website review and name/abbreviation searches which usually under-ascertain CEC related publications. Another possible source of measurement error is that CEC are typically supported through a mosaic of funding mechanisms, including grants from funding sources other than EGRP. Our grant number searches were limited to manuscripts citing EGRP-funded grants. While each individual search has limitations, we combined several strategies, with different strengths and weaknesses, to obtain an overall picture of the scientific productivity of this network of CECs. Results not only show a consistent increase in scientific output, after an initial lag period for the establishment of the needed infrastructures, but also the capability of consortia to support interdisciplinary science beyond the domains of classic epidemiology. The enrichment in the production of publications further down the translational continuum (T2 and higher) may also indicate the potential for unique contributions of CEC to translational research as a result of their interdisciplinary, team science approach. Interdisciplinary science is the first step in the path to translation and large consortia network may in the future provide an accelerated avenue to the development of preventive intervention and new therapeutic strategies.

We have described an extensive collection of cancer consortia which is showing the initial characteristics of an emerging interactive network as also shown by extensive co-authorship and co-membership across consortia (Abstract under review). Public posting of internal policies and processes, especially for what concerns data sharing and publications, and public availability of descriptive data on existing CECs resources (populations characteristics, protocols, questionnaires, publications, etc.), could considerably expedite collaborations across the consortia and with the scientific community at large. The combined expertise and infrastructure represented within this established network could also be of use in developing training approaches for young investigators across the spectrum of cancer epidemiology and related disciplines. This emerging Network of Cancer Consortia (NOCC) has demonstrated the capability to incorporate novel genomics technologies (genome-wide genotyping and next-generation sequencing technologies), and has the potential to be a fertile ground for the high-throughput application of different 'omics' approaches (38). Publication output shows that multi-level data sets are being assembled, integrated and analyzed to address hypotheses of increasing complexity.

It has been proposed that 21st century epidemiology will be driven by four overlapping drivers in the production of new knowledge and its translation: acceleration of trends toward multiple group interactive networks; rapid incorporation of emerging technologies into large-scale population studies; the building of infrastructure through which to assess factors and interventions at multiple levels and the capability of effectively integrating multi-level data sets for increasingly complex analyses (39). The extensive population resource, the reliance on interdisciplinary teams, and the facilitation of the translational pipeline by this emerging network of consortia may offer a supportive infrastructure to begin implementing these transformative goals.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Wuchty S, Jones BF, Uzzi B. The increasing dominance of teams in production of knowledge. *Science*. 2007; 316(5827):1036–9. Epub 2007/04/14. [PubMed: 17431139]
2. Caporaso NE, Bennett SN. The Rise of Consortia. *Nature Precedings* [Internet]. 2011
3. Boffetta P, Colditz GA, Potter JD, Kolonel L, Robson PJ, Malekzadeh R, et al. Cohorts and consortia conference: a summary report (Banff, Canada, June 17–19, 2009). *Cancer causes & control: CCC*. 2011; 22(3):463–8. Epub 2011/01/05. [PubMed: 21203821]
4. Vegvari A, Welinder C, Lindberg H, Fehniger TE, Marko-Varga G. Biobank resources for future patient care: developments, principles and concepts. *Journal of clinical bioinformatics*. 2011; 1(1): 24. Epub 2011/09/20. [PubMed: 21923917]
5. Hoover RN. The evolution of epidemiologic research: from cottage industry to “big” science. *Epidemiology*. 2007; 18(1):13–7. Epub 2006/12/21. [PubMed: 17179754]
6. EGRP-Supported Cancer Epidemiology Consortia. Bethesda, MD: National Institutes of Health(US); Available from: <http://epi.grants.cancer.gov/Consortia/>
7. Breast Cancer And The Environment Research Centers. Bethesda, MD: National Institutes of Health (US); 2002. Available from: <http://grants.nih.gov/grants/guide/rfa-files/RFA-ES-03-001.html>
8. Cooperative Family Registry for Epidemiologic Studies of Breast Cancer. Bethesda, MD: National Institutes of Health (US); 1994. Available from: <http://grants.nih.gov/grants/guide/rfa-files/RFA-CA-95-003.html>
9. Cooperative Family Registry for Epidemiologic Studies of Colon Cancer. Bethesda, MD: National Institutes of Health (US); 1996. Available from: <http://grants.nih.gov/grants/guide/notice-files/not96-192.html>
10. NIH Sharing Policies and Related Guidance on NIH-Funded Research Resources. Bethesda, MD: National Institutes of Health (US); Available from: <http://grants.nih.gov/grants/sharing.htm>
11. NIH RePORTER: Research Project Success Rates by Type and Activity. Bethesda, MD: National Institutes of Health(US); Available from: http://report.nih.gov/success_rates/Success_ByActivity.cfm
12. Hall KL, Feng AX, Moser RP, Stokols D, Taylor BK. Moving the science of team science forward: collaboration and creativity. *American journal of preventive medicine*. 2008; 35(2 Suppl):S243–9. Epub 2008/08/23. [PubMed: 18619406]
13. PubMed. Bethesda, MD: U.S. National Library of Medicine; Available from: <http://www.ncbi.nlm.nih.gov/pubmed>
14. Austin MA, Hair MS, Fullerton SM. Research Guidelines in the Era of Large-scale Collaborations: An Analysis of Genome-wide Association Study Consortia. *American journal of epidemiology*. 2012; 175(9):962–9. Epub 2012/04/12. [PubMed: 22491085]
15. Thun MJ, Hoover RN, Hunter DJ. Bigger, better, sooner--scaling up for success. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive . Oncology*. 2012; 21(4):571–5. Epub 2012/03/01.
16. Hindorff LA, Sethupathy P, Junkins HA, Ramos EM, Mehta JP, Collins FS, et al. Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proceedings of the National Academy of Sciences of the United States of America*. 2009; 106(23):9362–7. Epub 2009/05/29. [PubMed: 19474294]
17. A Catalog of Published Genome-Wide Association Studies. [database on the Internet]. Available from: www.genome.gov/gwastudies
18. Disis ML, Slattery JT. The road we must take: multidisciplinary team science. *Science translational medicine*. 2010; 2(22):22cm9. Epub 2010/04/09.

19. Rebbeck TR, Paskett E, Sellers TA. Fostering transdisciplinary science. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive . Oncology*. 2010; 19(5):1149–50. Epub 2010/05/08.
20. Facilitating Interdisciplinary Research. The National Academies Press; 2004.
21. Khoury MJ, Coates RJ, Evans JP. Evidence-based classification of recommendations on use of genomic tests in clinical practice: dealing with insufficient evidence. *Genetics in medicine: official journal of the American College of Medical Genetics*. 2010; 12(11):680–3. Epub 2010/10/27. [PubMed: 20975567]
22. Khoury MJ, Gwinn M, Yoon PW. The continuum of translation research in genomic medicine: how can we accelerate the appropriate integration of human genome discoveries into health care and disease prevention. *Genetics in medicine: official journal of the American College of Medical Genetics*. 2007; 9:665–74. [PubMed: 18073579]
23. Schully SD, Benedicto CB, Gillanders EM, Wang SS, Khoury MJ. Translational research in cancer genetics: the road less traveled. *Public health genomics*. 2011; 14:1–8. [PubMed: 20051673]
24. Schully SD, Benedicto CB, Khoury MJ. How can we stimulate translational research in cancer genomics beyond bench to bedside? *Genetics in medicine: official journal of the American College of Medical Genetics*. 2012; 14(1):169–70. Epub 2012/01/13. [PubMed: 22237448]
25. Freedman ND, Murray LJ, Kamangar F, Abnet CC, Cook MB, Nyren O, et al. Alcohol intake and risk of oesophageal adenocarcinoma: a pooled analysis from the BEACON Consortium. *Gut*. 2011; 60(8):1029–37. Epub 2011/03/17. [PubMed: 21406386]
26. Fabbri M, Bottoni A, Shimizu M, Spizzo R, Nicoloso MS, Rossi S, et al. Association of a microRNA/TP53 feedback circuitry with pathogenesis and outcome of B-cell chronic lymphocytic leukemia. *JAMA: the journal of the American Medical Association*. 2011; 305(1):59–67. Epub 2011/01/06.
27. Rebbeck TR, Wang Y, Kantoff PW, Krithivas K, Neuhausen SL, Godwin AK, et al. Modification of BRCA1- and BRCA2-associated breast cancer risk by AIB1 genotype and reproductive history. *Cancer research*. 2001; 61(14):5420–4. Epub 2001/07/17. [PubMed: 11454686]
28. Kastrinos F, Steyerberg EW, Mercado R, Balmana J, Holter S, Gallinger S, et al. The PREMM(1,2,6) model predicts risk of MLH1, MSH2, and MSH6 germline mutations based on cancer history. *Gastroenterology*. 2011; 140(1):73–81. Epub 2010/08/24. [PubMed: 20727894]
29. Domchek SM, Friebel TM, Singer CF, Evans DG, Lynch HT, Isaacs C, et al. Association of risk-reducing surgery in BRCA1 or BRCA2 mutation carriers with cancer risk and mortality. *JAMA: the journal of the American Medical Association*. 2010; 304(9):967–75. Epub 2010/09/03.
30. Kefford RF, Newton Bishop JA, Bergman W, Tucker MA. Counseling and DNA testing for individuals perceived to be genetically predisposed to melanoma: A consensus statement of the Melanoma Genetics Consortium. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology*. 1999; 17(10):3245–51. Epub 1999/10/03. [PubMed: 10506626]
31. Wirtzfeld DA, Mikula L, Gryfe R, Ravani P, Dicks EL, Parfrey P, et al. Concordance with clinical practice guidelines for adjuvant chemotherapy in patients with stage I–III colon cancer: experience in 2 Canadian provinces. *Canadian journal of surgery Journal canadien de chirurgie*. 2009; 52(2): 92–7. Epub 2009/04/29. [PubMed: 19399202]
32. Watkins KE, Way CY, Fiander JJ, Meadus RJ, Esplen MJ, Green JS, et al. Lynch syndrome: barriers to and facilitators of screening and disease management. *Hereditary cancer in clinical practice*. 2011; 9:8. Epub 2011/09/09. [PubMed: 21899746]
33. Ramsey SD, Clarke L, Etzioni R, Higashi M, Berry K, Urban N. Cost-effectiveness of microsatellite instability screening as a method for detecting hereditary nonpolyposis colorectal cancer. *Annals of internal medicine*. 2001; 135(8 Pt 1):577–88. Epub 2001/10/17. [PubMed: 11601929]
34. Newcomb PA, Storer BE, Morimoto LM, Templeton A, Potter JD. Long-term efficacy of sigmoidoscopy in the reduction of colorectal cancer incidence. *Journal of the National Cancer Institute*. 2003; 95(8):622–5. Epub 2003/04/17. [PubMed: 12697855]
35. John EM, Hopper JL, Beck JC, Knight JA, Neuhausen SL, Senie RT, et al. The Breast Cancer Family Registry: an infrastructure for cooperative multinational, interdisciplinary and translational

- studies of the genetic epidemiology of breast cancer. *Breast cancer research: BCR*. 2004; 6(4):R375–89. Epub 2004/06/26. [PubMed: 15217505]
36. Newcomb PA, Baron J, Cotterchio M, Gallinger S, Grove J, Haile R, et al. Colon Cancer Family Registry: an international resource for studies of the genetic epidemiology of colon cancer. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2007; 16(11):2331–43. Epub 2007/11/06.
 37. Knoppers BM, Harris JR, Tasse AM, Budin-Ljosne I, Kaye J, Deschenes M, et al. Towards a data sharing Code of Conduct for international genomic research. *Genome medicine*. 2011; 3(7):46. Epub 2011/07/27. [PubMed: 21787442]
 38. Verma M, Khoury MJ, Ioannidis JP. Opportunities and challenges for selected emerging technologies in cancer epidemiology: mitochondrial, epigenomic, metabolomic, and telomerase profiling. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2013; 22(2):189–200. Epub 2012/12/18.
 39. Lam TK, Spitz M, Schully SD, Khoury MJ. “Drivers” of Translational Cancer Epidemiology in the 21st Century: Needs and Opportunities. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2013 Epub 2013/01/17.
 40. Onar A, Ramamurthy U, Wallace D, Boyett JM. An operational perspective of challenging statistical dogma while establishing a modern, secure distributed data management and imaging transport system: the Pediatric Brain Tumor Consortium phase I experience. *Clinical and translational science*. 2009; 2(2):143–9. Epub 2010/05/07. [PubMed: 20443880]
 41. Poussaint TY, Phillips PC, Vajapeyam S, Fahey FH, Robertson RL, Osganian S, et al. The Neuroimaging Center of the Pediatric Brain Tumor Consortium-collaborative neuroimaging in pediatric brain tumor research: a work in progress. *AJNR American journal of neuroradiology*. 2007; 28(4):603–7. Epub 2007/04/10. [PubMed: 17416804]
 42. Cai Q, Long J, Lu W, Qu S, Wen W, Kang D, et al. Genome-wide association study identifies breast cancer risk variant at 10q21.2: results from the Asia Breast Cancer Consortium. *Human molecular genetics*. 2011; 20(24):4991–9. Epub 2011/09/13. [PubMed: 21908515]
 43. Chenevix-Trench G, Milne RL, Antoniou AC, Couch FJ, Easton DF, Goldgar DE. An international initiative to identify genetic modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: the Consortium of Investigators of Modifiers of BRCA1 and BRCA2 (CIMBA). *Breast cancer research: BCR*. 2007; 9(2):104. Epub 2007/05/01. [PubMed: 17466083]
 44. Neuhausen SL, Ozelik H, Southey MC, John EM, Godwin AK, Chung W, et al. BRCA1 and BRCA2 mutation carriers in the Breast Cancer Family Registry: an open resource for collaborative research. *Breast cancer research and treatment*. 2009; 116(2):379–86. Epub 2008/08/16. [PubMed: 18704680]
 45. Spurdle AB, Healey S, Devereau A, Hogervorst FB, Monteiro AN, Nathanson KL, et al. ENIGMA--evidence-based network for the interpretation of germline mutant alleles: an international initiative to evaluate risk and clinical significance associated with sequence variation in BRCA1 and BRCA2 genes. *Human mutation*. 2012; 33(1):2–7. Epub 2011/10/13. [PubMed: 21990146]
 46. Hunter DJ, Riboli E, Haiman CA, Albanes D, Altshuler D, Chanock SJ, et al. A candidate gene approach to searching for low-penetrance breast and prostate cancer genes. *Nature reviews Cancer*. 2005; 5(12):977–85. Epub 2005/12/13.
 47. Jia WH, Zhang B, Matsuo K, Shin A, Xiang YB, Jee SH, et al. Genome-wide association analyses in East Asians identify new susceptibility loci for colorectal cancer. *Nature genetics*. 2013; 45(2):191–6. Epub 2012/12/25. [PubMed: 23263487]
 48. Olson SH, Chen C, De Vivo I, Doherty JA, Hartmuller V, Horn-Ross PL, et al. Maximizing resources to study an uncommon cancer: E2C2--Epidemiology of Endometrial Cancer Consortium. *Cancer causes & control: CCC*. 2009; 20(4):491–6. Epub 2009/01/10. [PubMed: 19132539]
 49. Conway DI, Hashibe M, Boffetta P, Wunsch-Filho V, Muscat J, La Vecchia C, et al. Enhancing epidemiologic research on head and neck cancer: INHANCE - The international head and neck

- cancer epidemiology consortium. *Oral oncology*. 2009; 45(9):743–6. Epub 2009/05/16. [PubMed: 19442571]
50. Boffetta P, Armstrong B, Linet M, Kasten C, Cozen W, Hartge P. Consortia in cancer epidemiology: lessons from InterLymph. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive. Oncology*. 2007; 16(2):197–9. Epub 2007/02/16.
 51. Kricker A, Armstrong BK, Goumas C, Kanetsky P, Gallagher RP, Begg CB, et al. MC1R genotype may modify the effect of sun exposure on melanoma risk in the GEM study. *Cancer causes & control: CCC*. 2010; 21(12):2137–47. Epub 2010/08/20. [PubMed: 20721616]
 52. Ragin CC, Taioli E, McFarlane-Anderson N, Avery G, Bennett F, Bovell-Benjamin A, et al. African-Caribbean cancer consortium for the study of viral, genetic and environmental cancer risk factors. *Infectious agents and cancer*. 2007; 2:17. Epub 2007/09/26. [PubMed: 17892589]
 53. Rolland B, Smith BR, Potter JD. Coordinating centers in cancer epidemiology research: the Asia Cohort Consortium coordinating center. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive. Oncology*. 2011; 20(10):2115–9. Epub 2011/08/02.
 54. Song M, Rolland B, Potter JD, Kang D. Asia Cohort Consortium: challenges for collaborative research. *Journal of epidemiology/Japan Epidemiological Association*. 2012; 22(4):287–90. Epub 2012/06/08. [PubMed: 22672913]
 55. Anton-Culver H, Ziogas A, Bowen D, Finkelstein D, Griffin C, Hanson J, et al. The Cancer Genetics Network: recruitment results and pilot studies. *Community genetics*. 2003; 6(3):171–7. Epub 2004/07/09. [PubMed: 15237201]
 56. Brown RC, Dwyer T, Kasten C, Krotoski D, Li Z, Linet MS, et al. Cohort profile: the International Childhood Cancer Cohort Consortium (I4C). *International journal of epidemiology*. 2007; 36(4):724–30. Epub 2007/01/27. [PubMed: 17255350]
 57. West C, Rosenstein BS, Alsner J, Azria D, Barnett G, Begg A, et al. Establishment of a Radiogenomics Consortium. *International journal of radiation oncology, biology, physics*. 2010; 76(5):1295–6. Epub 2010/03/27.
 58. Olsen CM, Nagle CM, Whiteman DC, Ness R, Pearce CL, Pike MC, et al. Obesity and risk of ovarian cancer subtypes: evidence from the Ovarian Cancer Association Consortium. *Endocrine-related cancer*. 2013; 20(2):251–62. Epub 2013/02/14. [PubMed: 23404857]
 59. Petersen GM, de Andrade M, Goggins M, Hruban RH, Bondy M, Korczak JF, et al. Pancreatic cancer genetic epidemiology consortium. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive. Oncology*. 2006; 15(4):704–10. Epub 2006/04/15.
 60. Doherty R, Lubinski J, Manguoglu E, Luleci G, Christie M, Craven P, et al. AIDIT and IMPACT: building research collaborations in targeted prostate cancer screening. *Journal of BUON: official journal of the Balkan Union of Oncology*. 2006; 11(4):415–8. Epub 2007/02/20.
 61. Mitra AV, Bancroft EK, Barbachano Y, Page EC, Foster CS, Jameson C, et al. Targeted prostate cancer screening in men with mutations in BRCA1 and BRCA2 detects aggressive prostate cancer: preliminary analysis of the results of the IMPACT study. *BJU international*. 2011; 107(1):28–39. Epub 2010/09/16. [PubMed: 20840664]
 62. Mitra AV, Bancroft EK, Eeles RA. A review of targeted screening for prostate cancer: introducing the IMPACT study. *BJU international*. 2007; 99(6):1350–5. Epub 2007/04/11. [PubMed: 17419707]
 63. Schaid DJ, Chang BL. Description of the International Consortium For Prostate Cancer Genetics, and failure to replicate linkage of hereditary prostate cancer to 20q13. *The Prostate*. 2005; 63(3):276–90. Epub 2004/12/16. [PubMed: 15599943]
 64. Xu J, Lange EM, Lu L, Zheng SL, Wang Z, Thibodeau SN, et al. HOXB13 is a susceptibility gene for prostate cancer: results from the International Consortium for Prostate Cancer Genetics (ICPCG). *Human genetics*. 2013; 132(1):5–14. Epub 2012/10/16. [PubMed: 23064873]
 65. Rebbeck TR, Devesa SS, Chang BL, Bunker CH, Cheng I, Cooney K, et al. Global patterns of prostate cancer incidence, aggressiveness, and mortality in men of african descent. *Prostate cancer*. 2013; 2013:560857. Epub 2013/03/12. [PubMed: 23476788]

66. Kote-Jarai Z, Olama AA, Giles GG, Severi G, Schleutker J, Weischer M, et al. Seven prostate cancer susceptibility loci identified by a multi-stage genome-wide association study. *Nature genetics*. 2011; 43(8):785–91. Epub 2011/07/12. [PubMed: 21743467]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1.

(A) The number of EGRP CEC associated papers by year, from 1996 – 2011. The linear trend line excludes data points from 1996 and 1997. (B) The number of papers retrieved, by year, from PubMed queries using the terms “consortia” or “consortium” on cancer and on all topics from 1985 to 2012.




Figure 2.

The number of papers in scientific areas by year; definitions of each scientific area are in Supplementary Table 1. Genomics=Candidate Gene, Gene Characterization, Genome-wide Association, Gene-Environment, Linkage, Loss of Heterozygosity, and Next-generation Sequencing papers combined; EL=Environment, Lifestyle & Descriptive Epidemiology; MT=Methods and Technologies; BIO=Biology; CT=Clinical & Translational; ME=Molecular Epidemiology; RS=Resources; BR=Behavioral.

Table 1

Descriptive information of the 49 network CEC.

CEC Name (Abbreviation)	Initiation Year	Cancer Site(s)	Study Participants	Component Groups*	Countries†	Website URL
International Consortium of Bladder Cancer (ICBC)	2005	Bladder	13,090 cases, 16,537 controls	23	BEL, CHN, DEU, DNK, ESP, FRA, GBR, ITA, JPN, NLD, ROK, SWE, TUR, USA	http://icbc.cancer.gov/
Brain Tumor Epidemiology Consortium (BTEC)	2003	Brain	~6,500 cases	2	AUT, DNK, GBR, SWE, USA	http://epi.grants.cancer.gov/btec/index.html
GLIOGENE	2007	Brain	4,269 cases; 2,834 controls	15	DNK, GBR, ISR, SWE, USA	http://www.gliogene.org/
Pediatric Brain Tumor Consortium (PBTC) (40, 41)	1999	Brain	1,330 controls	12	USA	http://www.pbtc.org/
Asia Breast Cancer Consortium(ABCC) (42)	2008	Breast	17,153 cases, 16,943 controls	10	CHN, JPN, ROK, TWN, USA	None
Breast Cancer and the Environment Research Program (BCERP)	2003	Breast	Cohort of 1,200 young girls	4	USA	http://www.beerc.org/index.htm
Breast Cancer Association Consortium (BCAC)	2005	Breast	77,741 cases, 83,362 controls	69	AUS, BEL, BLR, CAN, CHN, CYP, DEU, DNK, ESP, FIN, FRA, GBR, HKG, IRL, ITA, JPN, MEX, MYS, NLD, NOR, POL, PRT, ROK, RUS, SGP, SWE, THA, TWN, USA	http://www.stlcam.ac.uk/consortia/beac/
Breast Cancer Consortium for Outcomes and Survival (BC2OS)	2007	Breast	48,000 cases	32	AUS, CHN, GRC, MYS, NGA, NLD, NOR, ROK, SWE, TUR, USA	None
Consortium of Investigators of Modifiers of BRCA 1/2 (CIMBA) (43)	2005	Breast	More than 10,000 BRCA1 carriers and 5,000 BRCA2 carriers	59	AUS, AUT, BEL, CAN, COL, CZE, DEU, DNK, ESP, FIN, FRA, GBR, GRC, HKG, HUN, ISL, ISR, ITA, LTU, LVA, MYS, NLD, PAK, POL, ROK, PRT, RUS, SGP, SWE, USA, ZAF	http://www.stlcam.ac.uk/consortia/cimba/
African-American Breast Cancer Consortium	2008	Breast	6,671 cases, 15,314 controls	4	USA	None
Prevention and Observation of Surgical Endpoints and Modifiers of Cancer Risk in BRCA 1/2 Mutation Carriers (PROSE-Modifier Studies)	2000	Breast	5,187 BRCA1/2 carriers	23	AUT, GBR, NLD, USA	http://www.cceb.upenn.edu/pages/prose/index.html
Women, Cancer and Radiation Exposure (WECARE)	2002	Breast	700 asynchronous bilateral BC cases; 1400 unilateral BC cases	20	DNK, NOR, SWE, USA	http://sktweb.mskcc.org/WECARE/front.html

CEC Name (Abbreviation)	Initiation Year	Cancer Site(s)	Study Participants	Component Groups*	Countries [†]	Website URL
Breast Cancer Family Registry (BCFR) (35, 44)	1995	Breast, Ovary	14,605 cases, 51,801 controls	6	AUS, CAN, USA	http://epi.grants.cancer.gov/CFR/about_breast.html
Evidence-based Network for the Interpretation of Germline Mutant Alleles Consortium (ENIGMA) (45)	2009	Breast, Ovary, others	3,504 <i>BRCA1</i> UV/VUS carrier families; 5,717 <i>BRCA2</i> UV/VUS carrier families	43	AUS, AUT, BRA, CAN, CZE, DNK, ESP, FIN, FRA, GBR, GRC, HKG, ITA, MYS, NLD, SWE, USA	http://enigmaconsortium.org/
Breast and Prostate Cancer Cohort Consortium (BPC3) (46)	2003	Breast, Prostate	At least 10,500 PC cases, 8,500 BC cases; 22,700 controls	10	DNK, DEU, ESP, FRA, GBR, GRC, ITA, NLD, NOR, SWE, USA	http://epi.grants.cancer.gov/BPC3/
Asia Colorectal Cancer Consortium (ACCC) (47)	2009	Colon	At least 7,456 cases, 11,671 controls	6	CHN, JPN, ROK, USA	None
Genetics and Epidemiology of Colorectal Cancer Consortium (GECCO)	2008	Colon	~20,000 cases, ~20,000 controls	18	CAN, DEU, FRA, USA	http://www.fhcr.org/content/public/en/labs/phs/projects/cancer-prevention/projects/gecco.html
Molecular Epidemiology of Colorectal Cancer (MECC)	1998	Colon	3,330 cases, 3,330 controls	2	ISR, USA	http://itemaker.umich.edu/gruber.lab/gruber.research/mecc
Colon Cancer Family Registry (CCFR) (36)	1997	Colon	16,438 cases, 48,300 controls	6	AUS, CAN, ESP, NZL, USA	http://epi.grants.cancer.gov/CFR/about_colon.html
Epidemiology of Endometrial Cancer Consortium (E2C2) (48)	2006	Endometrium	4,523 cases, 25,217 controls from case-control studies; 5,818 cases from cohort studies	36	AUS, BEL, CAN, CHE, CHN, GBR, GRC, ITA, NLD, NOR, POL, SWE, USA	http://epi.grants.cancer.gov/eccc
Asian Barrett's Consortium	2008	Esophagus	Not A available	2	BRN, CHN, IDN, IND, JPN, MYS, ROK, SGP, THA, TWN, VNM	None
International Barrett's and Esophageal Adenocarcinoma Consortium (BEACON)	2005	Esophagus	1,585 EA cases, 2,736 BE cases; 4,274 controls	18	AUS, CAN, FRA, GBR, IRL, NLD, SWE, USA	http://beacon.tlvnet.net/
International Head and Neck Cancer Epidemiology Consortium (INHANCE) (49)	2004	Head, Neck	30,059 cases, 40,949 controls	43	BRA, CAN, CHE, CHN, DEU, FRA, IND, ITA, JPN, NLD, PRI, USA	http://inhance.iarc.fr/index.php
Childhood Leukemia International Consortium (CLIC)	2006	Hematopoietic	13,000 cases, 21,000 controls	22	AUS, BRA, CAN, CRI, DEU, EGY, FRA, GBR, GRC, ITA, NZL, USA	https://ccls.berkeley.edu/clic/
Chronic Lymphocytic Leukemia Research Consortium (CRC)	2006	Hematopoietic	5,300 cases	8	GBR, USA	http://cll.ucsd.edu/
International Lymphoma Epidemiology Consortium (InterLymph) (50)	2001	Hematopoietic	13,000 cases, 16,000 controls	18	AUS, CAN, CZE, DEU, DNK, EGY, ESP, FIN, FRA, GBR, IRL, ISR, ITA, JAM, JOR, SWE, USA	http://epi.grants.cancer.gov/InterLymph/
International Multiple Myeloma Consortium (IMMC)	2007	Hematopoietic	2,870 cases	10	AUS, CAN, GBR, IRL, ITA, USA	None

CEC Name (Abbreviation)	Initiation Year	Cancer Site(s)	Study Participants	Component Groups*	Countries [†]	Website URL
Genetic Epidemiology of Lung Cancer (GELCC)	1999	Lung	2,000 cases, 2,000 controls	5	USA	http://www.eh.ac.edu/gebcc/
International Lung Cancer Consortium (ILCCO)	2004	Lung	59,530 cases, 68,974 controls; 795,000 cohort participants	56	CAN, CHN, DEU, DNK, ESP, FRA, GBR, GRC, ISL, ISR, ITA, JPN, NLD, NOR, ROK, SGP, SWE, TWN, USA	http://ilcco.iarc.fr/
Genes, Environment and Melanoma (GEM) (51)	1999	Melanoma	~3,700 single or multiple melanoma cases	8	AUS, CAN, ITA, USA	http://gemstudy.org/main/index.html
Melanoma Genetics Consortium (GenoMEL)	1997	Melanoma	5,327 cases	22	ARG, AUS, BRA, CHL, COL, ESP, FRA, GBR, ISR, ITA, LVA, MEX, NLD, POL, SVN, SWE, URY, USA	http://www.genomel.org/
African-Caribbean Cancer Consortium (AC3) (52)	2006	Multi-site	1,677 cases, 2,638 controls	54	AIA, BHS, BRB, CYM, GHA, JAM, KEN, KNA, MTQ, NGA, NZL, TTO, USA	http://www.ac-ca-consortium.org/
Asia Cohort Consortium (ACC) (53, 54)	2004	Multi-site	1,185,000 cohort participants (40)	21	BGD, CHN, IND, JPN, MYS, ROK, SGP, THA, TWN	http://www.asiacohort.org/Pages/Default.aspx
Cancer Genetics Network (CGN) (55)	1998	Multi-site	26,271 enrollees with personal or family history of cancer	14	USA	http://www.cancergen.org/
Cohort Consortium (CoCo)	2000	Multi-site	>4,000,000 cohort participants; 481,371 cases	50	AUS, CAN, CHN, DEU, DNK, ESP, FIN, FRA, GBR, GRC, IRN, ITA, JPN, NLD, NOR, SGP, SWE, USA	http://epi.grants.cancer.gov/Consortia/cohort.html
International Childhood Cancer Cohort Consortium (I4C) (56)	2005	Multi-site	Cohort of ~1,000,000 births; 801 incident cases	11	AUS, CHN, CHN, DNK, ESP, FRA, GBR, ISR, ITA, JPN, NOR, ROK, USA	http://www.nationalchildrensstudy.gov/research/internationalinvolvement/pages/default.aspx
Latin American Cancer Epidemiology Consortium (LACE)	2008	Multi-site	17,842 cases; 22,709 controls	8	ARG, BRA, COL, ESP, GBR, MEX, PRI, PRT, USA	None
Radiogenomics Consortium (RGC) (57)	2009	Multi-site	14,790 cases	87	AUS, BEL, CAN, CHE, DEU, DNK, ESP, FRA, GBR, ISR, ITA, JPN, NLD, POL, ROK, TWN, USA	None
Ovarian Cancer Association Consortium (OCAC) (58)	2005	Ovary	13,548 cases; 19,913 controls	47	AUS, BEL, CAN, CHN, DEU, DNK, FIN, JPN, MYS, NLD, NOR, POL, USA	http://www.stlcam.ac.uk/consortia/ocac/index.html
Ovarian Cancer Cohort Consortium (OC3)	2009	Ovary	4,152 cases	22	AUS, BEL, CAN, CHE, DEU, DNK, FRA, GBR,	None

CEC Name (Abbreviation)	Initiation Year	Cancer Site(s)	Study Participants	Component Groups*	Countries [†]	Website URL
Pacific Ovarian Cancer Research Consortium (POCRC)	1999	Ovary	Screening cohort of >1,000 women	6	JPN, NLD, NOR, POL, SGP, USA JPN, NLD, NOR, POL, SGP, USA	http://www.pocrc.org/
Pancreatic Cancer Case Control Consortium (PANC4)	2006	Pancreas	10,991 cases, 16,358 controls	16	CAN, AUS, CHN, GBR, ITA, USA	http://panc4.org/
Pancreatic Cancer Cohort Consortium (PanScan)	2006	Pancreas	>1,900,000 cohort participants; 4,620 incident cases; 780 retrospective cases	20	AUS, CHN, DEU, DNK, ESP, FRA, GBR, GRC, ITA, NLD, NOR, SWE, USA	http://epi.grants.cancer.gov/PanScan/
Pancreatic Cancer Genetic Epidemiology Consortium (PACGENE) (59)	2002	Pancreas	11,587 cases; 7,417 familial controls	6	CAN, USA	http://mayoresearch.mayo.edu/petersen_lab/epidemiology.cfm
Identification of Men with a genetic predisposition to Prostate Cancer: Targeted screening in BRCA1 and BRCA2 mutation carriers and controls (IMPACT) (60–62)	2005	Prostate	205 BRCA1/2 carriers, 95 non-carriers; 11 cases	39	AUS, CAN, DNK, ESP, GBR, ISR, ITA, MYS, NOR, POL, SVK, SVN, SWE, USA	http://www.impact-study.co.uk/
International Consortium for Prostate Cancer Genetics (ICPCG) (63, 64)	1996	Prostate	6,400 cases and 6,000 controls from 2550 PCA families	15	AUS, CAN, DEU, FIN, FRA, GBR, NOR, SWE, USA	https://www.icpcg.org/
Men of African Descent and Prostate Cancer (MADCaP) (65)	2007	Prostate	4,703 cases, 5,327 controls	22	BHS, BRB, GBR, GHA, GUY, NGA, SEN, TTO, UGA, USA, ZAF	http://epi.grants.cancer.gov/madcap/index.html
Prostate Cancer Association Group to Investigate Cancer Associated Alterations in the Genome (PRACTICAL) (66)	2007	Prostate	At least 26,055 cases and 25,256 controls	66	AUS, BEL, BGR, CAN, CHE, CHN, DEU, DNK, ESP, FIN, FRA, GBR, IND, IRL, JPN, MYS, NOR, POL, PRT, ROU, SWE, THA, USA	http://www.stl.cam.ac.uk/consortia/practical/index.html
Prostate Cancer Transatlantic Consortium (CaPTC)	2005	Prostate	385 cases	8	BHS, NGA, USA	http://epi.grants.cancer.gov/captc/

* Component groups can include individual studies, institutions, sites, and regional centers.

[†] Three letter country codes: AIA=Anguilla, ARG=Argentina, AUS=Australia, AUT=Austria, BEL=Belgium, BGD=Bangladesh, BGR=Bulgaria, BHR=Belarus, BLR=Belarus, BRA=Brazil, BRB=Barbados, BRN=Brunei, CAN=Canada, CHE=Switzerland, CHL=Chile, CHN=China, COL=Colombia, CRI=Costa Rica, CYM=Cayman Islands, CYP=Cyprus, DEU=Germany, DNK=Denmark, EGY=Egypt, ESP=Spain, FIN=Finland, FRA=France, GBR=United Kingdom, GHA=Ghana, GRC=Greece, GUY=Guyana, HKG=Hong Kong, HUN=Hungary, IDN=Indonesia, IND=India, IRL=Ireland, IRN=Iran, ISL=Iceland, ISR=Israel, ITA=Italy, JAM=Jamaica, JOR=Jordan, JPN=Japan, KEN=Kenya, KNA=St Kitt and Nevis, LTU=Lithuania, LVA=Latvia, MEX=Mexico, MTQ=Martinique, MYS=Malaysia, NGA=Nigeria, NLD=Netherlands, NOR=Norway, NZL=New Zealand, PAK=Pakistan, POL=Poland, PRI=Puerto Rico, PRT=Portugal, ROK=South Korea, ROU=Romania, RUS=Russia, SEN=Senegal, SGP=Singapore, SVK=Slovakia, SVN=Slovenia, SWE=Sweden, THA=Thailand, TTO=Trinidad and Tobago, TUR=Turkey, TWN=Taiwan, UGA=Uganda, URY=Uruguay, USA=United States, VNM=Viet Nam, ZAF=South Africa.

Table 2

EGRP CEC grant success rate versus EGRP and NIH grant success rates.

	CONSORTIA			EGRP			NIH [†]		
	Awarded*	Submitted	Success Rate	Submitted	Awarded	Success Rate	Submitted	Awarded	Success Rate
Total	255	529	48%	5550	1571	28%	554981	136554	24.6%
By Year									
2011	21	51	41%	472	70	15%	49592	8765	17.7%
2010	25	66	38%	470	97	21%	45983	9455	20.6%
2009	27	64	42%	555	139	25%	43142	8881	20.6%
2008	29	52	56%	398	94	24%	43467	9460	21.8%
2007	22	44	50%	441	126	29%	47455	10100	21.3%
2006	25	47	53%	426	90	21%	45688	9128	20%
2005	18	36	50%	332	76	23%	43069	9599	22.3%
2004	14	34	41%	366	114	31%	40861	10052	24.6%
2003	16	25	64%	305	102	33%	34710	10393	29.9%
2002	6	15	40%	304	112	37%	30068	9396	31.2%
2001	11	15	73%	296	117	40%	28368	9098	32.1%
2000	4	9	44%	253	87	34%	27798	8765	31.5%
1999	9	11	82%	376	134	36%	26408	8556	32.4%
1998	16	41	39%	272	93	34%	24151	7518	31.1%
1997	3	7	43%	284	120	42%	24221	7388	30.5%
1996	0	2	0%						
1995	9	10	90%						
By Type									
New	191	421	45%	4953	1290	26%	459618	94598	20%
Renewal	64	108	59%	597	281	47%	92849	41083	44%

* There are 255 new and competitive renewals included in the consortia grant portfolio. Between 1995–2011, there were 161 grant ID's with one new/renewals (N=161), 27 grant ID's with two new/renewals (N=54), 12 grant ID's with 3 new/renewals (N=36), and 1 grant ID with 4 new/renewals (N=4), for a total of 201 unique grant ID's with 255 new or renewal grants awarded.

[†] Data from NIH RePORTER: (accessed 5/6/2013). Data not available before 1997; Success rate by grant size not listed.

Table 3

The number and percentage of CEC papers on the continuum of translational research with types of research and examples from the literature analysis.

Phase	# (%)	Notation	Types of Research	Examples from Literature Review
T0	2645 (79%)	Discovery research	Genome-wide association studies; candidate gene studies; environmental and lifestyle risk factor studies; biological studies of disease etiology	Alcohol intake and risk of oesophageal adenocarcinoma: a pooled analysis from the BEACON Consortium (25) Association of a microRNA/TP53 feedback circuitry with pathogenesis and outcome of B-cell chronic lymphocytic leukemia (26)
T1	582 (17%)	Discovery to candidate health application	Phases I and II clinical trials; observational studies to characterize genes and gene-environment interaction; pharmacogenomics	The PREMM(1,2,6) model predicts risk of MLH1, MSH2, and MSH6 germline mutations based on cancer history (28) Modification of BRCA1- and BRCA2-associated breast cancer risk by AIB1 genotype and reproductive history (27)
T2	112 (3.3%)	Health application to evidence-based practice guidelines	Phase III clinical trials; observational studies assessing the efficacy and utility of interventions; evidence synthesis and guidelines development	Association of risk-reducing surgery in BRCA1 or BRCA2 mutation carriers with cancer risk and mortality (29) Counseling and DNA testing for individuals perceived to be genetically predisposed to melanoma: A consensus statement of the Melanoma Genetics Consortium (30)
T3	18 (0.5%)	Practice guidelines to health practice	Phase IV clinical trials; dissemination research; implementation research; diffusion research	Lynch syndrome: barriers to and facilitators of screening and disease management (32) Concordance with clinical practice guidelines for adjuvant chemotherapy in patients with stage I-III colon cancer: experience in 2 Canadian provinces (31)
T4	6 (0.2%)	Practice to population health impact	Outcomes research; population monitoring of morbidity, mortality, benefits and risks; cost-effectiveness	Cost-effectiveness of microsatellite instability screening as a method for detecting hereditary nonpolyposis colorectal cancer (33) Long-term efficacy of sigmoidoscopy in the reduction of colorectal cancer incidence (34)