



HHS Public Access

Author manuscript

Hum Vaccin Immunother. Author manuscript; available in PMC 2016 January 19.

Published in final edited form as:

Hum Vaccin Immunother. 2014 ; 10(6): 1436–1448. doi:10.4161/hv.28857.

Rotavirus vaccines:

Current status and future considerations

Catherine Yen^{1,2,*}, Jacqueline E Tate¹, Terri B Hyde², Margaret M Cortese¹, Benjamin A Lopman¹, Baoming Jiang¹, Roger I Glass³, and Umesh D Parashar¹

¹Division of Viral Diseases; Centers for Disease Control and Prevention; Atlanta, GA USA

²Global Immunization Division; Centers for Disease Control and Prevention; Atlanta, GA USA

³Fogarty International Center; National Institutes of Health; Bethesda, MD USA

Abstract

Rotavirus is the leading cause of severe diarrhea among children <5 years worldwide. Currently licensed rotavirus vaccines have been efficacious and effective, with many countries reporting substantial declines in diarrheal and rotavirus-specific morbidity and mortality. However, the full public health impact of these vaccines has not been realized. Most countries, including those with the highest disease burden, have not yet introduced rotavirus vaccines into their national immunization programs. Research activities that may help inform vaccine introduction decisions include (1) establishing effectiveness, impact, and safety for rotavirus vaccines in low-income settings; (2) identifying potential strategies to improve performance of oral rotavirus vaccines in developing countries, such as zinc supplementation; and (3) pursuing alternate approaches to oral vaccines, such as parenteral immunization. Policy- and program-level barriers, such as financial implications of new vaccine introductions, should be addressed to ensure that countries are able to make informed decisions regarding rotavirus vaccine introduction.

Keywords

rotavirus; diarrhea; gastroenteritis; rotavirus vaccines; vaccine impact

Background

Diarrhea is a major cause of death among children <5 y of age globally.¹ Rotavirus is the leading cause of severe diarrhea, resulting in an estimated 453 000 deaths in 2008, most of which occurred in developing countries of sub-Saharan Africa and South-East Asia (Fig. 1).² Rotavirus also causes considerable morbidity, with global estimates of 2.3 million hospitalizations and 24 million outpatient visits annually among children aged <5 y.^{2,3} Data from the Global Rotavirus Surveillance Network of the World Health Organization (WHO),

*Correspondence to: Catherine Yen; cyen@cdc.gov.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Financial Support

The authors have indicated that they have no financial relationships relevant to this article to disclose.

a network of sentinel surveillance sites in over 50 countries, indicate that rotavirus is responsible for ~40% of acute gastroenteritis hospitalizations among children <5 y of age in regions without widespread rotavirus vaccine use.⁴ Since improvements in water and sanitation do not prevent the majority of rotavirus disease, rotavirus vaccines are an essential part of an integrated approach to the control of diarrhea that also includes interventions, such as access to safe drinking water, sanitation, and handwashing facilities, breast feeding, vitamin A and zinc supplementation, and appropriate case management.⁵ Since 2009, WHO, with support from its Strategic Advisory Group of Experts (SAGE) on Immunization, has recommended that rotavirus vaccines be included in all national immunization programs and considered a priority, particularly in countries with high diarrhea-related mortality.⁶ By April 2014, 56 countries had introduced rotavirus vaccines into their national immunization programs, and ~50 more likely will follow within the next several years.^{7,8}

This review will discuss the 2 globally licensed rotavirus vaccines and other vaccines locally available or in development, the current status of vaccine introduction, information on the impact of vaccine introduction on immunization systems and disease burden, vaccine safety issues, and future considerations for vaccine introduction.

Rotavirus Vaccines

Since 2006, 2 rotavirus vaccines have been licensed and used globally—Rotarix (GlaxoSmithKline) and RotaTeq (Merck and Co., Inc.). Rotarix is a live, attenuated vaccine containing a single G1P[8] human rotavirus strain. RotaTeq is a live, attenuated vaccine containing 5 human-bovine reassortant rotavirus strains—G1P7[5], G2P7[5], G3P7[5], G4P7[5], and G6P1A[8] (Table 1). Both vaccines are administered orally to infants starting at a minimum age of 6 wk, with a minimum 4 wk interval between doses (2 doses per Rotarix course, 3 doses per RotaTeq course).⁶ Previous WHO administration recommendations for upper age limits of 15 wk for the first dose of vaccine and 32 wk for the last dose of vaccine were removed in 2013.⁶ For both vaccines, clinical trials conducted in high and upper-middle-income countries in the Americas, Asia, and Europe demonstrated vaccine efficacy of 72–100% in preventing severe rotavirus disease during 1- to 3-y follow-up periods, while trials conducted in lower income countries in Africa and Asia demonstrated vaccine efficacy of 49–72% (Table 1).^{9–17} Although definite reasons for this lower efficacy are unknown, reasons proposed include factors that can result in interference of uptake of a live, oral vaccine, such as breast milk, stomach acid, maternal antibodies, and co-administration of oral poliovirus vaccine (OPV), and factors that may cause an impaired immune response to vaccine, such as malnutrition, and other infections (e.g., human immunodeficiency virus, malaria, and tuberculosis).^{18,19}

China and Vietnam have locally manufactured oral vaccines that are licensed for use only within these countries (Table 2). Available in China, the Lanzhou lamb rotavirus vaccine (LLR; Lanzhou Institute of Biological Products) is a live, attenuated vaccine containing a single G10P[12] lamb rotavirus strain. The recommended schedule of administration is one dose annually for children 2 mo to 3 y of age and one dose at age 3–5 y.²⁰ Since 2000, over 30 million doses have been sold, but little is known about vaccine efficacy, safety, and vaccination coverage since LLR is available only through the private market.²⁰ Available in

Vietnam, Rotavin-M1 (POLYVAC) is a live, attenuated vaccine containing a single G1P[8] human rotavirus strain similar to Rotarix. The recommended schedule of administration is 2 doses, starting at a minimum age of 6 wk, given at least 30 d apart.²¹ Phase I and II studies demonstrated immunogenicity and safety profiles similar to Rotarix;²¹ efficacy data currently are unavailable. Recently, India licensed its own locally manufactured oral vaccine, ROTAVAC (Bharat Biotech International, Ltd) (Table 2). ROTAVAC is a live, attenuated vaccine containing a single neonatal rotavirus G9P[11] strain, 116E. The recommended schedule of administration is 3 doses at 6, 10, and 14 wk. The phase III trial demonstrated vaccine efficacy of ~56% in preventing severe rotavirus diarrhea.²² Bharat Biotech is pursuing WHO pre-qualification of the vaccine so that it may be available globally.²³

Additional live oral and parenteral rotavirus vaccines are in development or clinical trial stages, but not yet licensed (Table 2). Five live, single to multi-strain vaccines, including 2 lamb-human reassortant vaccines from China (Lanzhou and NF-R7), the previously licensed rhesus-human reassortant vaccine RotaShield, a bovine-human reassortant vaccine (UK), and a neonatal human strain vaccine from Australia (RV3), currently are undergoing clinical trials. One subunit rotavirus vaccine currently is undergoing a phase I trial, while several other candidates, including an inactivated rotavirus vaccine (IRV), are in pre-clinical stages of development. Alternative schedules, such as administration of a neonatal dose (RotaShield, RV3) are also being explored.

Current Status of Rotavirus Vaccine Introduction

As of April 2014, 56 countries have introduced rotavirus vaccines into their national immunization programs; an additional 4 have introduced vaccines regionally, and 3 have widespread coverage through the private market (Fig. 2).⁷ Although Australia and countries in the Americas and Europe were the earliest to introduce after vaccine licensure, countries in other regions have followed, many with support from the GAVI Alliance, a public-private global health partnership with a mission to save children's lives and protect people's health by increasing access to immunization in poor countries. As of March 2014, 16 countries have GAVI Alliance support to introduce rotavirus vaccines nationally, 4 countries have GAVI applications under review, and 11 countries are planning to submit GAVI applications.^{7,8} In addition, 16 non-GAVI eligible countries are planning introductions, making a total of 47 more countries planning to introduce rotavirus vaccines within the next several years. However, despite this progress and a universal recommendation for inclusion of rotavirus vaccines in national immunization programs, 84 countries have no reported plans to introduce rotavirus vaccines. The majority (70%) of these countries are located in Europe, South-East Asia, and the Western Pacific.

Vaccine Impact

While data from clinical trials provide important information on vaccine efficacy and safety in controlled settings, program and field conditions may differ with regards to vaccine management and administration and thus impact both vaccine effectiveness and the immunization system. Evaluation of the impact of vaccine introduction on immunization

systems and disease burden is crucial for improving and sustaining immunization programs and stimulating introduction of additional vaccines.

To help evaluate the programmatic impact of new vaccine introduction on immunization systems, WHO recommends post introduction evaluations (PIEs) 6–12 mo after any new vaccine introduction.²⁴ Findings from these evaluations may provide countries with information regarding the impact of rotavirus vaccine introduction on various program topics (e.g., vaccination coverage recording and reporting, cold chain and vaccine management, monitoring and supervision, training and knowledge of health-care workers, waste management, adverse events following immunization, and advocacy and communication planning), which may be used to improve local immunization systems and inform subsequent vaccine introductions. From 2009–2013, rotavirus vaccine PIEs were conducted in 13 countries in Africa, Latin America, the Eastern Mediterranean region, and Europe.²⁵

Monitoring of disease trends and vaccine effectiveness studies can provide timely information necessary to evaluate the impact of vaccine introduction on disease burden. Findings generated from Rotarix and RotaTeq impact assessments, mostly in Australia and countries in Europe and the Americas, have demonstrated substantial declines of 22–50% in diarrhea-related mortality,²⁶⁻²⁹ 17–55% in diarrhea-related hospitalizations,^{27,30-45} and 49–91% in rotavirus-specific hospitalizations among children <5 y of age (Table 3).^{30-34,36,37,42-59} Many of these studies also have reported potential indirect benefits for unvaccinated older children and young adults, with reductions of 6–51% in diarrhea-related hospitalizations^{30-32,35,37,45,60-62} and 20–92% in rotavirus-specific hospitalizations (Table 4).^{31,32,34,37,42-45,49,50,54,56,5,61-63} Additional studies have demonstrated vaccine effectiveness in preventing rotavirus hospitalizations similar to vaccine efficacy observed in clinical trials. High and upper-middle-income countries including Australia (certain regions), Taiwan, Austria, Belgium, France, Germany, Northern Israel, Spain, Mexico, Brazil (certain regions), and the US have reported vaccine effectiveness estimates of 79–100%,^{37,38,53,60,64-83} while lower income countries including Bolivia, El Salvador, and Nicaragua have reported vaccine effectiveness estimates of 43–92% (Table 5).⁸⁴⁻⁸⁹

To date, both Rotarix and RotaTeq have provided protection against a range of rotavirus strains, as demonstrated by the Rotarix clinical trials conducted in Africa, for which vaccine efficacy was ~60–64% for both G1 (contained in Rotarix) and non-G1 (not contained in Rotarix) rotavirus types,⁹⁰ and by various vaccine effectiveness studies conducted in Australia, Europe, and the Americas, for which vaccine effectiveness estimates were 71–95% against rotavirus strains not contained in RotaTeq and/or Rotarix.^{38,66,70,78,80,83,89,91} Monitoring of rotavirus strains continues in order to detect any global changes in strain prevalence and any emergence of unusual strains, and allow for strain-specific measures of vaccine effectiveness in the event that there is concern about vaccine effectiveness against an emergent or novel strain.

Vaccine Safety

In 1999, the first licensed rotavirus vaccine, RotaShield (Wyeth), was withdrawn from US market within a year after licensure due to an association with intussusception, an obstruction of the small intestine that can require radiological or surgical intervention. It was estimated that ~1 excess case of intussusception occurred per 10 000 infants vaccinated with RotaShield.⁹² Pre-licensure clinical trials for Rotarix and RotaTeq that included 60 000–70 000 infants each did not demonstrate an increased risk for intussusception,^{14,16} However, post-licensure monitoring studies were recommended to detect a possible low risk that may not have been identified in the clinical trials. In 2011, findings from studies conducted in Mexico (Rotarix only) and Australia (Rotarix and RotaTeq) reported a low level risk of intussusception with both Rotarix and RotaTeq on the order of ~1–2 excess cases per 100 000 vaccinated infants, mostly within the first week after the first dose of vaccine.^{93–9} These risks were lower than the risk associated with RotaShield, and a subsequent review of available data and a risk-benefit analysis of rotavirus vaccination conducted by the WHO Global Advisory Committee on Vaccine Safety (GACVS) determined that the benefits of rotavirus vaccination of all infants greatly exceeded the risks of intussusception associated with vaccination (Table 6).⁹⁶ Additional analyses to examine the risk-benefit of rotavirus vaccination without upper age restrictions estimated that universal rotavirus vaccination in low- and low-middle-income countries could prevent an additional 47 200 (range: 18 000–63 700) rotavirus deaths while potentially causing an additional 294 (range: 161–471) intussusception deaths among a cohort of children <5 y of age.⁹⁸ These analyses contributed to the 2013 WHO recommendation that upper age restrictions for rotavirus vaccination be removed to allow for greater vaccination coverage and potentially greater reductions in the number of rotavirus deaths as the benefits of vaccination continue to outweigh the risks of intussusception.⁶

Recently, new data from the US also have demonstrated a low risk of intussusception with both vaccines. In 2 studies conducted among separate managed care populations, an approximate risk of 1–5 excess cases of intussusception per 100 000 infants vaccinated with rotavirus vaccine was reported.^{98,99} While the risk-benefit of rotavirus vaccination with these 2 vaccines remains approximately the same as that seen when using the earlier risk estimates from Mexico and Australia, questions remain as to whether this risk may be higher in any particular subgroup of infants and whether the level of risk seen in high- and middle-income countries will occur in low-income countries where vaccine efficacy is lower.¹⁰⁰ Additional special studies and continued post-introduction intussusception monitoring, especially in low-income countries, may help answer these questions.

Future Considerations

Countries worldwide have established estimates of the considerable diarrheal disease burden of rotavirus.^{4,6} Existing licensed rotavirus vaccines have proven to be efficacious in clinical trials and effective in post-introduction evaluations, with many countries demonstrating adequate capacity to introduce these vaccines into routine immunization programs and substantial declines in diarrheal and rotavirus-specific morbidity and mortality (Tables 3–5). Despite this, the full public health impact of these vaccines on rotavirus disease and child

mortality has not been realized as most countries, including some with the highest disease burden, have not yet introduced rotavirus vaccines into their national immunization programs. Why is this the case? What are the remaining questions to be answered? What are the potential barriers to introduction?

Several key research activities may help to address remaining questions about rotavirus vaccine use under field conditions and inform vaccine introduction decisions, especially in low-income countries. These include: (1) establishing effectiveness, impact, and post-licensure safety of the current WHO-prequalified rotavirus vaccines in low-income settings; (2) identifying potential strategies to improve performance of oral rotavirus vaccines in developing countries, such as zinc supplementation to potentially strengthen the immune response to vaccination, withholding breast feeding just before and after vaccine administration to prevent interference of vaccine uptake by maternal antibodies, or a neonatal dosing schedule to help increase the amount of rotavirus disease prevented in settings where children may acquire disease at an earlier age; and (3) pursuing alternate approaches to oral vaccines, such as parenteral vaccines to bypass possible interference of vaccine uptake by gastric acid and breast milk to improve vaccine efficacy in developing countries.¹⁸

To identify potential barriers to introduction at a policy level, one may consider key issues in the decision-making process to introduce a new vaccine.¹⁰¹ Once barriers are identified, solutions may be implemented (Table 7). While each country should have a mechanism for an evidence-based, informed-decision making process, which may include expert groups that provide technical advice to national immunization programs, such as National Immunization Technical Advisory Groups (NITAGs) (http://www.healthinternetwork.com/immunization/sage/national_advisory_committees/en/) or other advisory committees on immunization,¹⁰¹ barriers and concerns may differ by country or even region. For example, one country may be concerned with vaccine introduction costs while another may be concerned with the programmatic impact of vaccine introduction or vaccine efficacy and safety. Given the potential diversity of opinion, it will be crucial to understand issues at the local policy making level to best inform decision makers.

During the past several years, significant progress has been made in the prevention and control of rotavirus diarrhea. The introduction of rotavirus vaccines into the national immunization programs of over 50 countries has resulted in substantial declines in rotavirus-related morbidity and mortality. However, questions remain that will need to be answered by additional research, and policy- and program-level barriers should be removed to ensure that countries are able to make informed decisions regarding rotavirus vaccine introduction and to help realize the full potential impact of these vaccines.

Acknowledgments

Disclaimer

The findings and conclusions of this report are those of the authors and do not necessarily represent the official position of the US Centers for Disease Control and Prevention.

Abbreviations

GACVS	Global Advisory Committee on Vaccine Safety
LLR	Lanzhou lamb rotavirus vaccine
SAGE	Strategic Advisory Group of Experts on Immunization
WHO	World Health Organization

References

1. Liu L, Johnson HL, Cousens S, Perin J, Scott S, Lawn JE, Rudan I, Campbell H, Cibulskis R, Li M, et al. Child Health Epidemiology Reference Group of WHO and UNICEF. Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. *Lancet*. 2012; 379:2151–61. [http://dx.doi.org/10.1016/S0140-6736\(12\)60560-1](http://dx.doi.org/10.1016/S0140-6736(12)60560-1). [PubMed: 22579125]
2. Tate JE, Burton AH, Boschi-Pinto C, Steele AD, Duque J, Parashar UD, WHO-coordinated Global Rotavirus Surveillance Network. 2008 estimate of worldwide rotavirus-associated mortality in children younger than 5 years before the introduction of universal rotavirus vaccination programmes: a systematic review and meta-analysis. *Lancet Infect Dis*. 2012; 12:136–41. [http://dx.doi.org/10.1016/S1473-3099\(11\)70253-5](http://dx.doi.org/10.1016/S1473-3099(11)70253-5). [PubMed: 22030330]
3. Parashar UD, Hummelman EG, Bresee JS, Miller MA, Glass RI. Global illness and deaths caused by rotavirus disease in children. *Emerg Infect Dis*. 2003; 9:565–72. <http://dx.doi.org/10.3201/eid0905.020562>. [PubMed: 12737740]
4. World Health Organization. Global rotavirus information and surveillance bulletin. 2013; 7:1–11.
5. World Health Organization. United Nations Children’s Fund. Ending preventable child deaths from pneumonia and diarrhoea by 2025: the integrated Global Action Plan for Pneumonia and Diarrhoea (GAPPD). WHO Press; France: 2013.
6. World Health Organization. Rotavirus vaccines. WHO position paper – January 2013. *Wkly Epidemiol Rec*. 2013; 88:49–64. [PubMed: 23424730]
7. International Vaccine Access Center. VIMS report: global vaccine introduction. Johns Hopkins Bloomberg School of Public Health; Mar. 2014 Available from: <http://www.jhsph.edu/research/centers-and-institutes/ivac/vims/IVAC-VIMS-Report-2014-Mar.pdf>
8. PATH. Rotavirus vaccine introductions: worldwide. PATH; Apr. 2014 Available from: <http://sites.path.org/rotavirusvaccine/rotavirus-advocacy-and-communications-toolkit/country-introduction-maps-and-list/>
9. Armah GE, Sow SO, Breiman RF, Dallas MJ, Tapia MD, Feikin DR, Binka FN, Steele AD, Laserson KF, Anshah NA, et al. Efficacy of pentavalent rotavirus vaccine against severe rotavirus gastroenteritis in infants in developing countries in sub-Saharan Africa: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2010; 376:606–14. [http://dx.doi.org/10.1016/S0140-6736\(10\)60889-6](http://dx.doi.org/10.1016/S0140-6736(10)60889-6). [PubMed: 20692030]
10. Kawamura N, Tokoeda Y, Oshima M, Okahata H, Tsutsumi H, Van Doorn LJ, Muto H, Smolenov I, Suryakiran PV, Han HH. Efficacy, safety and immunogenicity of RIX4414 in Japanese infants during the first two years of life. *Vaccine*. 2011; 29:6335–41. <http://dx.doi.org/10.1016/j.vaccine.2011.05.017>. [PubMed: 21640780]
11. Linhares AC, Velázquez FR, Pérez-Schael I, Sáez-Llorens X, Abate H, Espinoza F, López P, Macías-Parra M, Ortega-Barría E, Rivera-Medina DM, et al. Human Rotavirus Vaccine Study Group. Efficacy and safety of an oral live attenuated human rotavirus vaccine against rotavirus gastroenteritis during the first 2 years of life in Latin American infants: a randomised, double-blind, placebo-controlled phase III study. *Lancet*. 2008; 371:1181–9. [http://dx.doi.org/10.1016/S0140-6736\(08\)60524-3](http://dx.doi.org/10.1016/S0140-6736(08)60524-3). [PubMed: 18395579]
12. Madhi SA, Cunliffe NA, Steele D, Witte D, Kirsten M, Louw C, Ngwira B, Victor JC, Gillard PH, Chevart BB, et al. Effect of human rotavirus vaccine on severe diarrhea in African infants. *N Engl J Med*. 2010; 362:289–98. <http://dx.doi.org/10.1056/NEJMoa0904797>. [PubMed: 20107214]

13. Phua KB, Lim FS, Lau YL, Nelson EA, Huang LM, Quak SH, Lee BW, Teoh YL, Tang H, Boudville I, et al. Safety and efficacy of human rotavirus vaccine during the first 2 years of life in Asian infants: randomised, double-blind, controlled study. *Vaccine*. 2009; 27:5936–41. <http://dx.doi.org/10.1016/j.vaccine.2009.07.098>. [PubMed: 19679216]
14. Ruiz-Palacios GM, Pérez-Schael I, Velázquez FR, Abate H, Breuer T, Clemens SC, Chevart B, Espinoza F, Gillard P, Innis BL, et al. Human Rotavirus Vaccine Study Group. Safety and efficacy of an attenuated vaccine against severe rotavirus gastroenteritis. *N Engl J Med*. 2006; 354:11–22. <http://dx.doi.org/10.1056/NEJMoa052434>. [PubMed: 16394298]
15. Vesikari T, Karvonen A, Prymula R, Schuster V, Tejedor JC, Cohen R, Meurice F, Han HH, Damaso S, Bouckennooghe A. Efficacy of human rotavirus vaccine against rotavirus gastroenteritis during the first 2 years of life in European infants: randomised, double-blind controlled study. *Lancet*. 2007; 370:1757–63. [http://dx.doi.org/10.1016/S0140-6736\(07\)61744-9](http://dx.doi.org/10.1016/S0140-6736(07)61744-9). [PubMed: 18037080]
16. Vesikari T, Matson DO, Dennehy P, Van Damme P, Santosham M, Rodriguez Z, Dallas MJ, Heyse JF, Goveia MG, Black SB, et al. Rotavirus Efficacy and Safety Trial (REST) Study Team. Safety and efficacy of a pentavalent human-bovine (WC3) reassortant rotavirus vaccine. *N Engl J Med*. 2006; 354:23–33. <http://dx.doi.org/10.1056/NEJMoa052664>. [PubMed: 16394299]
17. Zaman K, Dang DA, Victor JC, Shin S, Yunus M, Dallas MJ, Podder G, Vu DT, Le TP, Luby SP, et al. Efficacy of pentavalent rotavirus vaccine against severe rotavirus gastroenteritis in infants in developing countries in Asia: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2010; 376:615–23. [http://dx.doi.org/10.1016/S0140-6736\(10\)60755-6](http://dx.doi.org/10.1016/S0140-6736(10)60755-6). [PubMed: 20692031]
18. Patel M, Shane AL, Parashar UD, Jiang B, Gentsch JR, Glass RI. Oral rotavirus vaccines: how well will they work where they are needed most? *J Infect Dis*. 2009; 200(Suppl 1):S39–48. <http://dx.doi.org/10.1086/605035>. [PubMed: 19817613]
19. Patel M, Steele AD, Parashar UD. Influence of oral polio vaccines on performance of the monovalent and pentavalent rotavirus vaccines. *Vaccine*. 2012; 30(Suppl 1):A30–5. <http://dx.doi.org/10.1016/j.vaccine.2011.11.093>. [PubMed: 22520134]
20. Fu C, He Q, Xu J, Xie H, Ding P, Hu W, Dong Z, Liu X, Wang M. Effectiveness of the Lanzhou lamb rotavirus vaccine against gastroenteritis among children. *Vaccine*. 2012; 31:154–8. <http://dx.doi.org/10.1016/j.vaccine.2012.10.078>. [PubMed: 23127516]
21. Dang DA, Nguyen VT, Vu DT, Nguyen TH, Nguyen DM, Yuhuan W, Baoming J, Nguyen DH, Le TL, Rotavin-M1 Vaccine Trial Group. A dose-escalation safety and immunogenicity study of a new live attenuated human rotavirus vaccine (Rotavin-M1) in Vietnamese children. *Vaccine*. 2012; 30(Suppl 1):A114–21. <http://dx.doi.org/10.1016/j.vaccine.2011.07.118>. [PubMed: 22520120]
22. Department of Biotechnology GoI. [February 3, 2014] The need for rotavirus vaccines in India: understanding efficacy and impact. 2013. http://www.defeatdd.org/sites/default/files/node-images/ROTAVAC%20efficacy%20fact%20sheet_FINAL.pdf
23. Department of Biotechnology GoI. [February 3, 2014] International social innovation partnership leads to lifesaving vaccine. 2013. http://www.defeatdd.org/sites/default/files/node-images/ROTAVAC%20history_FINAL_0.pdf
24. World Health Organization. New vaccine post-introduction evaluation (PIE) tool. WHO Document Services; Geneva, Switzerland: 2010.
25. Personal communication with World Health Organization. Expanded Programme on Immunization;
26. Bayard V, DeAntonio R, Contreras R, Tinajero O, Castrejon MM, Ortega-Barría E, Colindres RE. Impact of rotavirus vaccination on childhood gastroenteritis-related mortality and hospital discharges in Panama. *Int J Infect Dis*. 2012; 16:e94–8. <http://dx.doi.org/10.1016/j.ijid.2011.09.003>. [PubMed: 22154592]
27. do Carmo GM, Yen C, Cortes J, Siqueira AA, de Oliveira WK, Cortez-Escalante JJ, Lopman B, Flannery B, de Oliveira LH, Carmo EH, et al. Decline in diarrhea mortality and admissions after routine childhood rotavirus immunization in Brazil: a time-series analysis. *PLoS Med*. 2011; 8:e1001024. <http://dx.doi.org/10.1371/journal.pmed.1001024>. [PubMed: 21526228]

28. Lanzieri TM, Linhares AC, Costa I, Kolhe DA, Cunha MH, Ortega-Barria E, Colindres RE. Impact of rotavirus vaccination on childhood deaths from diarrhea in Brazil. *Int J Infect Dis*. 2011; 15:e206–10. <http://dx.doi.org/10.1016/j.ijid.2010.11.007>. [PubMed: 21193339]
29. Richardson V, Hernandez-Pichardo J, Quintanar-Solares M, Esparza-Aguilar M, Johnson B, Gomez-Altamirano CM, Parashar U, Patel M. Effect of rotavirus vaccination on death from childhood diarrhea in Mexico. *N Engl J Med*. 2010; 362:299–305. <http://dx.doi.org/10.1056/NEJMoa0905211>. [PubMed: 20107215]
30. Bégué RE, Perrin K. Reduction in gastroenteritis with the use of pentavalent rotavirus vaccine in a primary practice. *Pediatrics*. 2010; 126:e40–5. <http://dx.doi.org/10.1542/peds.2009-2069>. [PubMed: 20587671]
31. Chang HG, Smith PF, Tserenpuntsag B, Markey K, Parashar U, Morse DL. Reduction in hospitalizations for diarrhea and rotavirus infections in New York state following introduction of rotavirus vaccine. *Vaccine*. 2010; 28:754–8. <http://dx.doi.org/10.1016/j.vaccine.2009.10.075>. [PubMed: 19896451]
32. Clarke MF, Davidson GP, Gold MS, Marshall HS. Direct and indirect impact on rotavirus positive and all-cause gastroenteritis hospitalisations in South Australian children following the introduction of rotavirus vaccination. *Vaccine*. 2011; 29:4663–7. <http://dx.doi.org/10.1016/j.vaccine.2011.04.109>. [PubMed: 21575665]
33. Cortes JE, Curns AT, Tate JE, Cortese MM, Patel MM, Zhou F, Parashar UD. Rotavirus vaccine and health care utilization for diarrhea in U.S. children. *N Engl J Med*. 2011; 365:1108–17. <http://dx.doi.org/10.1056/NEJMoa1000446>. [PubMed: 21992123]
34. Cortese MM, Tate JE, Simonsen L, Edelman L, Parashar UD. Reduction in gastroenteritis in United States children and correlation with early rotavirus vaccine uptake from national medical claims databases. *Pediatr Infect Dis J*. 2010; 29:489–94. [PubMed: 20354464]
35. Curns AT, Steiner CA, Barrett M, Hunter K, Wilson E, Parashar UD. Reduction in acute gastroenteritis hospitalizations among US children after introduction of rotavirus vaccine: analysis of hospital discharge data from 18 US states. *J Infect Dis*. 2010; 201:1617–24. <http://dx.doi.org/10.1086/652403>. [PubMed: 20402596]
36. Dey A, Wang H, Menzies R, Macartney K. Changes in hospitalisations for acute gastroenteritis in Australia after the national rotavirus vaccination program. *Med J Aust*. 2012; 197:453–7. <http://dx.doi.org/10.5694/mja12.10062>. [PubMed: 23072242]
37. Field EJ, Vally H, Grimwood K, Lambert SB. Pentavalent rotavirus vaccine and prevention of gastroenteritis hospitalizations in Australia. *Pediatrics*. 2010; 126:e506–12. <http://dx.doi.org/10.1542/peds.2010-0443>. [PubMed: 20732946]
38. Gurgel RG, Bohland AK, Vieira SC, Oliveira DM, Fontes PB, Barros VF, Ramos MF, Dove W, Nakagomi T, Nakagomi O, et al. Incidence of rotavirus and all-cause diarrhea in northeast Brazil following the introduction of a national vaccination program. *Gastroenterology*. 2009; 137:1970–5. <http://dx.doi.org/10.1053/j.gastro.2009.07.046>. [PubMed: 19632228]
39. Lanzieri TM, Costa I, Shafi FA, Cunha MH, Ortega-Barria E, Linhares AC, Colindres RE. Trends in hospitalizations from all-cause gastroenteritis in children younger than 5 years of age in Brazil before and after human rotavirus vaccine introduction, 1998-2007. *Pediatr Infect Dis J*. 2010; 29:673–5. <http://dx.doi.org/10.1097/INF.0b013e3181da8f23>. [PubMed: 20300045]
40. Molto Y, Cortes JE, De Oliveira LH, Mike A, Solis I, Suman O, Coronado L, Patel MM, Parashar UD, Cortese MM. Reduction of diarrhea-associated hospitalizations among children aged < 5 Years in Panama following the introduction of rotavirus vaccine. *Pediatr Infect Dis J*. 2011; 30(Suppl):S16–20. <http://dx.doi.org/10.1097/INF.0b013e3181fefc68>. [PubMed: 21183835]
41. Quintanar-Solares M, Yen C, Richardson V, Esparza-Aguilar M, Parashar UD, Patel MM. Impact of rotavirus vaccination on diarrhea-related hospitalizations among children < 5 years of age in Mexico. *Pediatr Infect Dis J*. 2011; 30(Suppl):S11–5. <http://dx.doi.org/10.1097/INF.0b013e3181fefb32>. [PubMed: 21183834]
42. Raes M, Strens D, Vergison A, Verghote M, Standaert B. Reduction in pediatric rotavirus-related hospitalizations after universal rotavirus vaccination in Belgium. *Pediatr Infect Dis J*. 2011; 30:e120–5. <http://dx.doi.org/10.1097/INF.0b013e318214b811>. [PubMed: 21436757]

43. Sáfadi MA, Berezin EN, Munford V, Almeida FJ, de Moraes JC, Pinheiro CF, Racz ML. Hospital-based surveillance to evaluate the impact of rotavirus vaccination in São Paulo, Brazil. *Pediatr Infect Dis J.* 2010; 29:1019–22. [PubMed: 20543761]
44. Yen C, Armero Guardado JA, Alberto P, Rodriguez Araujo DS, Mena C, Cuellar E, Nolasco JB, De Oliveira LH, Pastor D, Tate JE, et al. Decline in rotavirus hospitalizations and health care visits for childhood diarrhea following rotavirus vaccination in El Salvador. *Pediatr Infect Dis J.* 2011; 30(Suppl):S6–10. <http://dx.doi.org/10.1097/INF.0b013e3181fe05>. [PubMed: 21048524]
45. Yen C, Tate JE, Wenk JD, Harris JM 2nd, Parashar UD. Diarrhea-associated hospitalizations among US children over 2 rotavirus seasons after vaccine introduction. *Pediatrics.* 2011; 127:e9–15. <http://dx.doi.org/10.1542/peds.2010-1393>. [PubMed: 21172995]
46. Anderson EJ, Rupp A, Shulman ST, Wang D, Zheng X, Noskin GA. Impact of rotavirus vaccination on hospital-acquired rotavirus gastroenteritis in children. *Pediatrics.* 2011; 127:e264–70. <http://dx.doi.org/10.1542/peds.2010-1830>. [PubMed: 21262887]
47. BATTERY JP, Lambert SB, Grimwood K, Nissen MD, Field EJ, Macartney KK, Akikusa JD, Kelly JJ, Kirkwood CD. Reduction in rotavirus-associated acute gastroenteritis following introduction of rotavirus vaccine into Australia's National Childhood vaccine schedule. *Pediatr Infect Dis J.* 2011; 30(Suppl):S25–9. <http://dx.doi.org/10.1097/INF.0b013e3181fefdee>. [PubMed: 21183837]
48. Custodio H, Masnita-Iusan C, Wludyka P, Rathore MH. Change in rotavirus epidemiology in northeast Florida after the introduction of rotavirus vaccine. *Pediatr Infect Dis J.* 2010; 29:766–7. <http://dx.doi.org/10.1097/INF.0b013e3181dbf256>. [PubMed: 20661105]
49. Eberly MD, Gorman GH, Eide MB, Olsen CH, Rajnik M. The effect of rotavirus immunization on rotavirus gastroenteritis hospitalization rates in military dependents. *Vaccine.* 2011; 29:650–9. <http://dx.doi.org/10.1016/j.vaccine.2010.11.041>. [PubMed: 21129394]
50. Hemming M, Räsänen S, Huhti L, Paloniemi M, Salminen M, Vesikari T. Major reduction of rotavirus, but not norovirus, gastroenteritis in children seen in hospital after the introduction of RotaTeq vaccine into the National Immunization Programme in Finland. *Eur J Pediatr.* 2013; 172:739–46. <http://dx.doi.org/10.1007/s00431-013-1945-3>. [PubMed: 23361964]
51. Lambert SB, Faux CE, Hall L, Birrell FA, Peterson KV, Selvey CE, Sloots TP, Nissen MD, Grimwood K. Early evidence for direct and indirect effects of the infant rotavirus vaccine program in Queensland. *Med J Aust.* 2009; 191:157–60. [PubMed: 19645646]
52. Msimang VM, Page N, Groome MJ, Moyes J, Cortese M, Seheri M, Kahn K, Chagan M, Madhi SA, Cohen C. Impact of Rotavirus Vaccine on Childhood Diarrheal Hospitalization Following Introduction into the South African Public Immunization Program. *Pediatr Infect Dis J.* 2013; 32:1359–64. <http://dx.doi.org/10.1097/INF.0b013e3182a72fc0>. [PubMed: 24569308]
53. Paulke-Korinek M, Kollaritsch H, Aberle SW, Zwazl I, Schmidle-Loss B, Vécsei A, Kundi M. Sustained low hospitalization rates after four years of rotavirus mass vaccination in Austria. *Vaccine.* 2013; 31:2686–91. <http://dx.doi.org/10.1016/j.vaccine.2013.04.001>. [PubMed: 23597718]
54. Paulke-Korinek M, Kundi M, Rendi-Wagner P, de Martin A, Eder G, Schmidle-Loss B, Vecsei A, Kollaritsch H. Herd immunity after two years of the universal mass vaccination program against rotavirus gastroenteritis in Austria. *Vaccine.* 2011; 29:2791–6. <http://dx.doi.org/10.1016/j.vaccine.2011.01.104>. [PubMed: 21320539]
55. Paulke-Korinek M, Rendi-Wagner P, Kundi M, Kronik R, Kollaritsch H. Universal mass vaccination against rotavirus gastroenteritis: impact on hospitalization rates in austrian children. *Pediatr Infect Dis J.* 2010; 29:319–23. [PubMed: 19935446]
56. Payne DC, Staat MA, Edwards KM, Szilagyi PG, Weinberg GA, Hall CB, Chappell J, Curns AT, Wikswo M, Tate JE, et al. New Vaccine Surveillance Network (NVSN). Direct and indirect effects of rotavirus vaccination upon childhood hospitalizations in 3 US Counties, 2006-2009. *Clin Infect Dis.* 2011; 53:245–53. <http://dx.doi.org/10.1093/cid/cir307>. [PubMed: 21705316]
57. Tate JE, Panozzo CA, Payne DC, Patel MM, Cortese MM, Fowlkes AL, Parashar UD. Decline and change in seasonality of US rotavirus activity after the introduction of rotavirus vaccine. *Pediatrics.* 2009; 124:465–71. <http://dx.doi.org/10.1542/peds.2008-3528>. [PubMed: 19581260]
58. Zeller M, Rahman M, Heylen E, De Coster S, De Vos S, Arijis I, Novo L, Verstappen N, Van Ranst M, Matthijnsens J. Rotavirus incidence and genotype distribution before and after national

- rotavirus vaccine introduction in Belgium. *Vaccine*. 2010; 28:7507–13. <http://dx.doi.org/10.1016/j.vaccine.2010.09.004>. [PubMed: 20851085]
59. Zlomy M, Kofler S, Orth D, Würzner R, Heinz-Erian P, Streng A, Prelog M. The impact of Rotavirus mass vaccination on hospitalization rates, nosocomial Rotavirus gastroenteritis and secondary blood stream infections. *BMC Infect Dis*. 2013; 13:112. <http://dx.doi.org/10.1186/1471-2334-13-112>. [PubMed: 23452879]
 60. Cortese MM, Leblanc J, White KE, Jerris RC, Stinchfield P, Preston KL, Meek J, Odofin L, Khizer S, Miller CA, et al. Leveraging state immunization information systems to measure the effectiveness of rotavirus vaccine. *Pediatrics*. 2011; 128:e1474–81. <http://dx.doi.org/10.1542/peds.2011-1006>. [PubMed: 22084328]
 61. Gastañaduy PA, Curns AT, Parashar UD, Lopman BA. Gastroenteritis hospitalizations in older children and adults in the United States before and after implementation of infant rotavirus vaccination. *JAMA*. 2013; 310:851–3. <http://dx.doi.org/10.1001/jama.2013.170800>. [PubMed: 23982372]
 62. Lopman BA, Curns AT, Yen C, Parashar UD. Infant rotavirus vaccination may provide indirect protection to older children and adults in the United States. *J Infect Dis*. 2011; 204:980–6. <http://dx.doi.org/10.1093/infdis/jir492>. [PubMed: 21878425]
 63. Anderson EJ, Shippee DB, Weinrobe MH, Davila MD, Katz BZ, Reddy S, Cuyugan MG, Lee SY, Simons YM, Yogev R, et al. Indirect protection of adults from rotavirus by pediatric rotavirus vaccination. *Clin Infect Dis*. 2013; 56:755–60. <http://dx.doi.org/10.1093/cid/cis1010>. [PubMed: 23349228]
 64. Adlhoch C, Hoehne M, Littmann M, Marques AM, Lerche A, Dehnert M, Eckmanns T, Wichmann O, Koch J. Rotavirus vaccine effectiveness and case-control study on risk factors for breakthrough infections in Germany, 2010–2011. *Pediatr Infect Dis J*. 2013; 32:e82–9. <http://dx.doi.org/10.1097/INF.0b013e3182720b71>. [PubMed: 23334342]
 65. Boom JA, Tate JE, Sahni LC, Rench MA, Hull JJ, Gentsch JR, Patel MM, Baker CJ, Parashar UD. Effectiveness of pentavalent rotavirus vaccine in a large urban population in the United States. *Pediatrics*. 2010; 125:e199–207. <http://dx.doi.org/10.1542/peds.2009-1021>. [PubMed: 20083525]
 66. Boom JA, Tate JE, Sahni LC, Rench MA, Quaye O, Mijatovic-Rustempasic S, Patel MM, Baker CJ, Parashar UD. Sustained protection from pentavalent rotavirus vaccination during the second year of life at a large, urban United States pediatric hospital. *Pediatr Infect Dis J*. 2010; 29:1133–5. <http://dx.doi.org/10.1097/INF.0b013e3181ed18ab>. [PubMed: 20634776]
 67. Braeckman T, Van Herck K, Meyer N, Pirçon JY, Soriano-Gabarró M, Heylen E, Zeller M, Azou M, Capiou H, De Koster J, et al. RotaBel Study Group. Effectiveness of rotavirus vaccination in prevention of hospital admissions for rotavirus gastroenteritis among young children in Belgium: case-control study. *BMJ*. 2012; 345:e4752. <http://dx.doi.org/10.1136/bmj.e4752>. [PubMed: 22875947]
 68. Castilla J, Beristain X, Martínez-Artola V, Navascués A, García Cenoz M, Alvarez N, Polo I, Mazón A, Gil-Setas A, Barricarte A. Effectiveness of rotavirus vaccines in preventing cases and hospitalizations due to rotavirus gastroenteritis in Navarre, Spain. *Vaccine*. 2012; 30:539–43. <http://dx.doi.org/10.1016/j.vaccine.2011.11.071>. [PubMed: 22122860]
 69. Chang WC, Yen C, Wu FT, Huang YC, Lin JS, Huang FC, Yu HT, Chi CL, Lin HY, Tate JE, et al. Effectiveness of 2 rotavirus vaccines against rotavirus disease in Taiwanese infants. *Pediatr Infect Dis J*. 2014; 33:e81–6. <http://dx.doi.org/10.1097/INF.000000000000105>. [PubMed: 24569388]
 70. Correia JB, Patel MM, Nakagomi O, Montenegro FM, Germano EM, Correia NB, Cuevas LE, Parashar UD, Cunliffe NA, Nakagomi T. Effectiveness of monovalent rotavirus vaccine (Rotarix) against severe diarrhea caused by serotypically unrelated G2P[4] strains in Brazil. *J Infect Dis*. 2010; 201:363–9. <http://dx.doi.org/10.1086/649843>. [PubMed: 20047501]
 71. Cortese MM, Immergluck LC, Held M, Jain S, Chan T, Grizas AP, Khizer S, Barrett C, Quaye O, Mijatovic-Rustempasic S, et al. Effectiveness of monovalent and pentavalent rotavirus vaccine. *Pediatrics*. 2013; 132:e25–33. <http://dx.doi.org/10.1542/peds.2012-3804>. [PubMed: 23776114]
 72. Desai SN, Esposito DB, Shapiro ED, Dennehy PH, Vázquez M. Effectiveness of rotavirus vaccine in preventing hospitalization due to rotavirus gastroenteritis in young children in Connecticut, USA. *Vaccine*. 2010; 28:7501–6. <http://dx.doi.org/10.1016/j.vaccine.2010.09.013>. [PubMed: 20851087]

73. Donauer S, Payne DC, Edwards KM, Szilagyi PG, Hornung RW, Weinberg GA, Chappell J, Hall CB, Parashar UD, Staat MA. Determining the effectiveness of the pentavalent rotavirus vaccine against rotavirus hospitalizations and emergency department visits using two study designs. *Vaccine*. 2013; 31:2692–7. <http://dx.doi.org/10.1016/j.vaccine.2013.03.072>. [PubMed: 23583814]
74. Gagneur A, Nowak E, Lemaitre T, Segura JF, Delaperrière N, Abalea L, Poulhazan E, Jossens A, Auzanneau L, Tran A, et al. IVANHOE investigators. Impact of rotavirus vaccination on hospitalizations for rotavirus diarrhea: the IVANHOE study. *Vaccine*. 2011; 29:3753–9. <http://dx.doi.org/10.1016/j.vaccine.2011.03.035>. [PubMed: 21443962]
75. Justino MC, Linhares AC, Lanzieri TM, Miranda Y, Mascarenhas JD, Abreu E, Guerra SF, Oliveira AS, da Silva VB, Sanchez N, et al. Effectiveness of the monovalent G1P[8] human rotavirus vaccine against hospitalization for severe G2P[4] rotavirus gastroenteritis in Belém, Brazil. *Pediatr Infect Dis J*. 2011; 30:396–401. <http://dx.doi.org/10.1097/INF.0b013e3182055cc2>. [PubMed: 21150692]
76. Martínón-Torres F, Bouzón Alejandro M, Redondo Collazo L, Sánchez Lastres JM, Pértega Díaz S, Seoane Pillado MT, Martínón Sánchez JM, ROTACOST research team. Effectiveness of rotavirus vaccination in Spain. *Hum Vaccin*. 2011; 7:757–61. <http://dx.doi.org/10.4161/hv.7.7.15576>. [PubMed: 21521947]
77. Muhsen K, Shulman L, Kasem E, Rubinstein U, Shachter J, Kremer A, Goren S, Zilberstein I, Chodick G, Ephros M, et al. Effectiveness of rotavirus vaccines for prevention of rotavirus gastroenteritis-associated hospitalizations in Israel: a case-control study. *Hum Vaccin*. 2010; 6:450–4. <http://dx.doi.org/10.4161/hv.6.6.11759>. [PubMed: 20448471]
78. Payne DC, Boom JA, Staat MA, Edwards KM, Szilagyi PG, Klein EJ, Selvarangan R, Azimi PH, Harrison C, Moffatt M, et al. Effectiveness of pentavalent and monovalent rotavirus vaccines in concurrent use among US children <5 years of age, 2009–2011. *Clin Infect Dis*. 2013; 57:13–20. <http://dx.doi.org/10.1093/cid/cit164>. [PubMed: 23487388]
79. Snelling TL, Andrews RM, Kirkwood CD, Culvenor S, Carapetis JR. Case-control evaluation of the effectiveness of the G1P[8] human rotavirus vaccine during an outbreak of rotavirus G2P[4] infection in central Australia. *Clin Infect Dis*. 2011; 52:191–9. <http://dx.doi.org/10.1093/cid/ciq101>. [PubMed: 21288843]
80. Snelling TL, Schultz R, Graham J, Roseby R, Barnes GL, Andrews RM, Carapetis JR. Rotavirus and the indigenous children of the Australian outback: monovalent vaccine effective in a high-burden setting. *Clin Infect Dis*. 2009; 49:428–31. <http://dx.doi.org/10.1086/600395>. [PubMed: 19566443]
81. Staat MA, Payne DC, Donauer S, Weinberg GA, Edwards KM, Szilagyi PG, Griffin MR, Hall CB, Curns AT, Gentsch JR, et al. New Vaccine Surveillance Network (NVSN). Effectiveness of pentavalent rotavirus vaccine against severe disease. *Pediatrics*. 2011; 128:e267–75. <http://dx.doi.org/10.1542/peds.2010-3722>. [PubMed: 21768317]
82. Wang FT, Mast TC, Glass RJ, Loughlin J, Seeger JD. Effectiveness of the pentavalent rotavirus vaccine in preventing gastroenteritis in the United States. *Pediatrics*. 2010; 125:e208–13. <http://dx.doi.org/10.1542/peds.2009-1246>. [PubMed: 20100757]
83. Yen C, Figueroa JR, Uribe ES, Carmen-Hernández LD, Tate JE, Parashar UD, Patel MM, Richardson López-Collado V. Monovalent rotavirus vaccine provides protection against an emerging fully heterotypic G9P[4] rotavirus strain in Mexico. *J Infect Dis*. 2011; 204:783–6. <http://dx.doi.org/10.1093/infdis/jir390>. [PubMed: 21810915]
84. Cardellino A, Khawaja S, Sánchez Cruz E, Mast TC. Effectiveness of vaccination with the pentavalent rotavirus vaccine in Nicaragua as determined using the screening method. *Hum Vaccin Immunother*. 2013; 9:1449–53. <http://dx.doi.org/10.4161/hv.24338>. [PubMed: 23571175]
85. de Palma O, Cruz L, Ramos H, de Baires A, Villatoro N, Pastor D, de Oliveira LH, Kerin T, Bowen M, Gentsch J, et al. Effectiveness of rotavirus vaccination against childhood diarrhoea in El Salvador: case-control study. *BMJ*. 2010; 340:c2825. <http://dx.doi.org/10.1136/bmj.c2825>. [PubMed: 20551120]
86. Mast TC, Khawaja S, Espinoza F, Paniagua M, Del Carmen LP, Cardellino A, Sánchez E. Case-control study of the effectiveness of vaccination with pentavalent rotavirus vaccine in Nicaragua. *Pediatr Infect Dis J*. 2011; 30:e209–15. <http://dx.doi.org/10.1097/INF.0b013e31822a8527>. [PubMed: 21768920]

87. Patel M, Pedreira C, De Oliveira LH, Tate J, Orozco M, Mercado J, Gonzalez A, Malespin O, Amador JJ, Umaña J, et al. Association between pentavalent rotavirus vaccine and severe rotavirus diarrhea among children in Nicaragua. *JAMA*. 2009; 301:2243–51. <http://dx.doi.org/10.1001/jama.2009.756>. [PubMed: 19491186]
88. Patel M, Pedreira C, De Oliveira LH, Umaña J, Tate J, Lopman B, Sanchez E, Reyes M, Mercado J, Gonzalez A, et al. Duration of protection of pentavalent rotavirus vaccination in Nicaragua. *Pediatrics*. 2012; 130:e365–72. <http://dx.doi.org/10.1542/peds.2011-3478>. [PubMed: 22753550]
89. Patel MM, Patzi M, Pastor D, Nina A, Roca Y, Alvarez L, Iniguez V, Rivera R, Tam KI, Quaye O, et al. Effectiveness of monovalent rotavirus vaccine in Bolivia: case-control study. *BMJ*. 2013; 346:f3726. <http://dx.doi.org/10.1136/bmj.f3726>. [PubMed: 23783434]
90. Steele AD, Neuzil KM, Cunliffe NA, Madhi SA, Bos P, Ngwira B, Witte D, Todd S, Louw C, Kirsten M, et al. Human rotavirus vaccine Rotarix™ provides protection against diverse circulating rotavirus strains in African infants: a randomized controlled trial. *BMC Infect Dis*. 2012; 12:213. <http://dx.doi.org/10.1186/1471-2334-12-213>. [PubMed: 22974466]
91. Patel MM, Glass R, Desai R, Tate JE, Parashar UD. Fulfilling the promise of rotavirus vaccines: how far have we come since licensure? *Lancet Infect Dis*. 2012; 12:561–70. [http://dx.doi.org/10.1016/S1473-3099\(12\)70029-4](http://dx.doi.org/10.1016/S1473-3099(12)70029-4). [PubMed: 22742639]
92. Murphy TV, Gargiullo PM, Massoudi MS, Nelson DB, Jumaan AO, Okoro CA, Zanardi LR, Setia S, Fair E, LeBaron CW, et al. Rotavirus Intussusception Investigation Team. Intussusception among infants given an oral rotavirus vaccine. *N Engl J Med*. 2001; 344:564–72. <http://dx.doi.org/10.1056/NEJM20010223440804>. [PubMed: 11207352]
93. Buttery JP, Danchin MH, Lee KJ, Carlin JB, McIntyre PB, Elliott EJ, Booy R, Bines JE, PAEDS/ APSU Study Group. Intussusception following rotavirus vaccine administration: post-marketing surveillance in the National Immunization Program in Australia. *Vaccine*. 2011; 29:3061–6. <http://dx.doi.org/10.1016/j.vaccine.2011.01.088>. [PubMed: 21316503]
94. Patel MM, López-Collada VR, Bulhões MM, De Oliveira LH, Bautista Márquez A, Flannery B, Esparza-Aguilar M, Montenegro Renoier EI, Luna-Cruz ME, Sato HK, et al. Intussusception risk and health benefits of rotavirus vaccination in Mexico and Brazil. *N Engl J Med*. 2011; 364:2283–92. <http://dx.doi.org/10.1056/NEJMoa1012952>. [PubMed: 21675888]
95. Velázquez FR, Colindres RE, Grajales C, Hernández MT, Mercadillo MG, Torres FJ, Cervantes-Apolinar M, DeAntonio-Suarez R, Ortega-Barria E, Blum M, et al. Postmarketing surveillance of intussusception following mass introduction of the attenuated human rotavirus vaccine in Mexico. *Pediatr Infect Dis J*. 2012; 31:736–44. <http://dx.doi.org/10.1097/INF.0b013e318253add3>. [PubMed: 22695189]
96. World Health Organization. Rotavirus vaccine and intussusception: report from an expert consultation. *Wkly Epidemiol Rec*. 2011; 86:317–21. [PubMed: 21800466]
97. Patel MM, Clark AD, Sanderson CF, Tate J, Parashar UD. Removing the age restrictions for rotavirus vaccination: a benefit-risk modeling analysis. *PLoS Med*. 2012; 9:e1001330. <http://dx.doi.org/10.1371/journal.pmed.1001330>. [PubMed: 23109915]
98. Weintraub ES, Baggs J, Duffy J, Vellozzi C, Belongia EA, Irving S, Klein NP, Glanz JM, Jacobsen SJ, Naleway A, et al. Risk of intussusception after monovalent rotavirus vaccination. *N Engl J Med*. 2014; 370:513–9. <http://dx.doi.org/10.1056/NEJMoa1311738>. [PubMed: 24422678]
99. Yih WK, Lieu TA, Kulldorff M, Martin D, McMahon-Walraven CN, Platt R, Selvam N, Selvan M, Lee GM, Nguyen M. Intussusception risk after rotavirus vaccination in U.S. infants. *N Engl J Med*. 2014; 370:503–12. <http://dx.doi.org/10.1056/NEJMoa1303164>. [PubMed: 24422676]
100. Glass RI, Parashar UD. Rotavirus vaccines--balancing intussusception risks and health benefits. *N Engl J Med*. 2014; 370:568–70. <http://dx.doi.org/10.1056/NEJMe1315836>. [PubMed: 24422677]
101. World Health Organization. Vaccine introduction guidelines - adding a vaccine to a national immunization programme: decision and implementation. WHO Document Services; Geneva, Switzerland: 2003.
102. Australian Government Therapeutic Goods Administration. [August 24, 2011] Rotavirus vaccination and risk of intussusception. 2011. <http://www.tga.gov.au/safety/alerts-medicine-rotavirus-110225.htm>

103. Cortese, MM. Summary of intussusception risk and benefits of rotavirus vaccination in the United States; Adivsory Committee on Immunization Practices Meeting; Atlanta, GA. June 20, 2013; 2013.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1.

10 countries with the greatest number of rotavirus deaths in 2008. Adapted from: Tate et al. 2008 estimate of worldwide rotavirus-associated mortality in children younger than 5 y before the introduction of universal rotavirus vaccination programmes: a systematic review and meta-analysis. *Lancet Infect Dis* 2012; 12(2):136–141.




Figure 2.
 National rotavirus vaccine introductions, by geographic region—56 countries*

Table 1

Currently licensed and globally recommended rotavirus vaccines

Characteristic	Rotavirus vaccine	
Trade name	Rotarix®	RotaTeq®
Manufacturer	GlaxoSmithKline	Merck and Co., Inc.
Country of manufacture	Belgium	USA
Composition	Live-attenuated G1P1A[8] human rotavirus strain	Live human-bovine reassortant rotavirus strains: G1P7[5], G2P7[5], G3P7[5], G4P7[5], G6P1A[8]
Pharmaceutical form	Liquid, lyophilized + diluent	Liquid
Presentation	1 dose plastic tube or 1 dose applicator or 1 dose vial	1 dose tube
Route of administration	Oral	Oral
Recommended schedule of administration	Minimum age of 1st dose: 6 wk 2 doses at least 4 wk apart <i>Previous WHO recommendations for upper age limits of 15 wk for the first dose and 32 wk for second dose have been removed</i>	Minimum age of 1st dose: 6 wk 3 doses at least 4 wk apart <i>Previous recommendations for upper age limits of 15 wk for first dose and 32 wk for third dose have been removed</i>
Contraindications per manufacturer	Hypersensitivity to the active substance or to any of the excipients; hypersensitivity after previous administration of rotavirus vaccines; history of intussusception; history of uncorrected congenital malformation of the gastrointestinal tract that would predispose for intussusception; history of severe combined immunodeficiency disease (SCID)	A demonstrated history of hypersensitivity to the vaccine or any component of the vaccine; history of severe combined immunodeficiency disease (SCID); history of intussusception
Volume per dose	1.5 mL (tube and applicator) 1 mL (lyophilized + diluent)	2 mL
Cost per dose	GAVI-eligible: \$2.43 (2014 UNICEF pricing) Non-GAVI: variable	GAVI-eligible: \$3.50–5.00 (2014 UNICEF pricing) Non-GAVI: variable
Date of WHO prequalification	Mar 2009 (tube and applicator) Jan 2007 (vial)	Oct 2008
Licensure status	Licensed in >100 countries	Licensed in >100 countries
Full series efficacy against severe rotavirus gastroenteritis (95% confidence interval)	<u>3-y follow-up:</u> 97% (88–100) (Hong Kong/Singapore/Taiwan) <u>2-y follow-up:</u> 81% (71–87) (Latin America) 90% (85–94) (Europe) 72% (54–84) (China) 92% (62–99) (Japan) <u>1-y follow-up:</u> 49% (11–72) (Malawi) 72% (40–88) (South Africa)	<u>3-y follow-up:</u> 94% (89–97) (Finland) <u>1-y follow-up:</u> 98% (88–100) (US/Finland) 51% (13–73) (Bangladesh/Vietnam) 64% (40–79) (Kenya/Ghana/Mali) <u>Up to 1-y follow-up:</u> 100% (55–100) (Japan)

Table 2

Rotavirus vaccines that are regionally used, recently licensed, or in development

Name	Composition	Route of administration	Organization/Company	Stage of development
Lanzhou Lamb Rotavirus (LLR)	Live attenuated lamb rotavirus strain, G10P[12]	Oral	Lanzhou Institute of Biological Products, China	Licensed for use in China
Rotavin-M1	Live attenuated human rotavirus strain, G1P[8]	Oral	POLYVAC, Vietnam	Licensed for use in Vietnam
ROTAVAC	Live attenuated neonatal rotavirus strain, G9P[11] (aka 116E)	Oral	Bharat Biotech, India	Recently licensed for use in India; pursuing WHO pre-qualification
LLR reassortants	Live attenuated lamb-human reassortant rotavirus strains, G2, G3, G4	Oral	Lanzhou Institute of Biological Products, China	Phase III
RotaShield	Live attenuated rhesus-human reassortant rotavirus strains, tetravalent	Oral	International Medica Foundation, USA	Phase II complete, Phase III pending
RV3	Live attenuated neonatal rotavirus strain, G3P[6]	Oral	Biofarma, Indonesia	Phase II
UK reassortants	Live attenuated bovine-human reassortant strains, tetravalent to hexavalent	Oral	National Institutes of Health, USA; Instituto Butantan, Brazil; Serum Institute of India, India; Shantha Biotech, India; / Minghai Biotechnology Co., China; Wuhan Institute of Biological Products, China	Phase I and II, Research
NF-R7	Live attenuated lamb-human reassortant strain, G4	Oral	Shenzhen Kangtai Biological Products Company, China	Phase I
Subunit	Truncated VP8 of P4, P6, P8	Intramuscular	National Institutes of Health and PATH, USA	Phase I
IRV	Inactivated G1P[8], G2P[4]	Intramuscular or intradermal	US CDC; multiple (China, Europe, India)	Pre-clinical
Subunit	Virus-like particles: VP 2/6/7 and VP 2/4/6/7	To be determined	Baylor College of Medicine, USA	Research
Subunit	Truncated VP8 in norovirus P particles	To be determined	Cincinnati Children's Hospital Medical Center, USA	Research
Subunit	VP6 combined with norovirus G1 and GII VLPs	To be determined	University of Tampere School of Medicine, Finland	Research

Table 3Summary of rotavirus vaccine impact studies among children <5 y – disease trends^a

Location	Vaccine	% Reduction	References
Reduction in diarrheal death rates			
Brazil			
• Nationwide	Rotarix	22%	27
• Nationwide	Rotarix	33–39%	28
Mexico, Nationwide	Rotarix	46%	29
Panama, Nationwide	Rotarix	32–50%	26
Reduction in diarrhea hospitalizations			
Australia			
• National	Rotarix, RotaTeq	38%	36
• Queensland	RotaTeq	20–40%	37
• South Australia	RotaTeq	48% (0 to <6 y)	32
Belgium, Nationwide	Rotarix, RotaTeq	33% (0 to 2 y)	42
Brazil			
• Nationwide	Rotarix	26–48% (0 to 1 y)	39
• Nationwide	Rotarix	17%	27
• Sao Paolo	Rotarix	29%	43
• Sergipe	Rotarix	44–55%	38
El Salvador, Nationwide	Rotarix	28–37% (includes outpatient)	44
Mexico, Nationwide	Rotarix	40% (during rotavirus season)	41
Panama, Bocas del Toro, Chiriquí, Los Santos, San Miguelito, Metropolitan Region	Rotarix	37%	40
USA			
• Nationwide	RotaTeq	45% (during rotavirus season)	35
• Nationwide	RotaTeq	29–50%	45
• 3 regions (Northeast, Midwest, South)	RotaTeq	30–45%	34
• Privately insured children, 13 states ^b	RotaTeq	25–33%	33
• New Orleans, LA	RotaTeq	50%	30
• New York, NY	RotaTeq	40% (0 to <2 y)	31
South Africa, Gauteng and Mpumalanga provinces	Rotarix	32–33%	51
Reduction in rotavirus hospitalizations			
Australia			
• National	Rotarix, RotaTeq	71%	36
• New South Wales, Queensland, Victoria	Rotarix, RotaTeq	87%	47
• Queensland	RotaTeq	53–57%	51
• Queensland	RotaTeq	50%	37
• Queensland	RotaTeq	83% (0–71 mo)	32

Location	Vaccine	% Reduction	References
<ul style="list-style-type: none"> South Australia 			
Austria <ul style="list-style-type: none"> Nationwide Nationwide Nationwide Tyrol 	Rotarix, RotaTeq Rotarix, RotaTeq Rotarix, RotaTeq Rotarix, RotaTeq	70% (0 to 1 y) 76–79% (0 to 2 y) 73–74% 62–88%	55 54 53 59
Belgium <ul style="list-style-type: none"> Nationwide Leuven 	Rotarix, RotaTeq Rotarix, RotaTeq	58–77% 49–66%	42 58
Finland, Tampere	Rotarix, RotaTeq	91% (<2 y; incl. ER visits)	50
Brazil, Sao Paolo	Rotarix	59%	43
El Salvador, San Salvador, Santa Ana, San Miguel, La Libertad, La Paz	Rotarix	69–81%	44
USA <ul style="list-style-type: none"> Nationwide Nationwide 3 regions (Northeast, Midwest, South) Privately insured children, 13 states^b Chicago Cincinnati, OH, Nashville, TN, Rochester, NY Jacksonville, FL New Orleans, LA New York, NY US military dependents 	RotaTeq RotaTeq RotaTeq RotaTeq RotaTeq RotaTeq RotaTeq RotaTeq RotaTeq RotaTeq RotaTeq	67–69% 66–83% 69–81% 60–75% 62% (during viral season) 55–89% (0 to 35 mo) 72% 67% (includes ED visits) 85% (0 to <2 y) 62%	57 45 34 33 46 56 48 30 31 49
South Africa, Gauteng and Mpumalanga provinces	Rotarix	40–44%	52

^aEstimated rotavirus vaccination coverage ranges of 37–94% mostly among children 0–1 y or 0–2 y, if reported.

^bAlaska, Idaho, Massachusetts, Maine, North Dakota, New Hampshire, New Mexico, Oregon, Rhode Island, Vermont, Washington, Wisconsin, Wyoming.

Table 4

Summary of rotavirus vaccine impact studies – indirect benefits for unvaccinated individuals^a

Location	Vaccine	Age range	% Reduction	References
Reduction in diarrhea hospitalizations				
Australia				
• Queensland	RotaTeq	2 to 4 y	30–40%	37
• South Australia	RotaTeq	2 to <6 y	42–49%	32
USA				
• Nationwide	RotaTeq	2 to <5 y	35–41%	34
• Nationwide	RotaTeq	2 to <5 y	43–45%	35
• Nationwide	RotaTeq	5 to 24 y	17–48% (during rotavirus season)	45
• Nationwide	RotaTeq	5 to 44 y	8–29%	62
• Nationwide	Rotarix, RotaTeq	5 to 44 y	6–30%	61
• Nationwide	RotaTeq	2 to <5 y	51%	30
• New Orleans, LA	RotaTeq	2 to <3 y	31–36%	31
• New York, NY	RotaTeq	3 to <5 y	33–37%	31
• New York, NY	RotaTeq	5 to 18 y	9–12%	31
• New York, NY	RotaTeq			
Reduction in rotavirus hospitalizations				
Australia				
• Queensland	RotaTeq	2 to 19 y	30–70%	37
• South Australia	RotaTeq	2 to <6 y	50–83%	32
Austria				
• Nationwide	Rotarix, RotaTeq	2 to 5 y	35%	54
• Tyrol	Rotarix, RotaTeq	2 to 18 y	62–89%	59
Belgium, Nationwide	Rotarix, RotaTeq	2 to 5 y	20–64%	42
Finland, Tampere	RotaTeq	2 to <16 y	72%	50
Brazil, Sao Paolo	Rotarix	2 to <5 y	24%	43
El Salvador, Nationwide	Rotarix	2 to <5 y	41–81%	44
USA				
• Nationwide	RotaTeq	2 to <5 y	69–78%	34
• Nationwide	RotaTeq	2 to <5 y	41–80%	45
• Nationwide	RotaTeq	5 to 24 y	65–71%	62
• Nationwide	Rotarix, RotaTeq	5 to 44 y	43–70%	61
• Chicago, IL	RotaTeq	³ 18 y	48%	63
• Chicago, IL	RotaTeq	2 to <3 y	92%	56
• Cincinnati, OH, Nashville, TN, Rochester, NY	RotaTeq	2 to <3 y	70–76%	31
• Cincinnati, OH, Nashville, TN, Rochester, NY	RotaTeq	3 to <5 y	79–88%	31
• New York, NY	RotaTeq	5 to 18 y	70%	31
• New York, NY	RotaTeq	0 to 4 y	57% (unvaccinated individuals)	49
• New York, NY	RotaTeq			

Location	Vaccine	Age range	% Reduction	References
<ul style="list-style-type: none"> <li data-bbox="266 254 505 279">• US military dependents 				

^aEstimated rotavirus vaccination coverage ranges of 37–94% mostly among children 0–1 y or 0–2 y, if reported.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 5

Vaccine effectiveness (VE) against rotavirus hospitalizations

Location	Vaccine type	Vaccine effectiveness	Predominant strain	References
Australia <ul style="list-style-type: none"> Queensland Central Australia Central Australia 	RotaTeq Rotarix Rotarix	94% 84% 51%	G9P[8] G2P[4]	37 80 79
Taiwan, Linkou, Changhua, Kaohsiung	Rotarix, RotaTeq	92%, 97%	G1P[8]	69
Austria, Nationwide	Rotarix, RotaTeq	79–96%	G1P[8], G2P[4]	53
Belgium, Nationwide	Rotarix	90%	G2P[4]	67
France, Northwest Brittany	RotaTeq	98%		74
Germany, Mecklenberg-Western Pomerania	Rotarix, RotaTeq	80%	G1P[8], G9P[8]	64
Northern Israel	Rotarix, RotaTeq	89%		77
Spain <ul style="list-style-type: none"> Nationwide Navarre 	Rotarix, RotaTeq Rotarix, RotaTeq	97%, 95% 83%	G9	76 68
Bolivia	Rotarix	69–77%	G9P[8], G2P[4], G3P[8], G9P[6]	89
Brazil <ul style="list-style-type: none"> Northeast Brazil Recife Belem 	Rotarix Rotarix Rotarix	89–95% 80–81% 40–76%	G2P[4] G2P[4] G2P[4]	38 70 75
El Salvador, Nationwide	Rotarix	76%	G1P[8]	85
Mexico, Chiapas	Rotarix	94%	G9P[4]	83
Nicaragua, Nationwide <ul style="list-style-type: none"> Nationwide Managua, Jinotepe, Masaya, Matagalpa Western Region Western Region 	RotaTeq RotaTeq RotaTeq RotaTeq	43–49% 45–70% 64–87% (includes ED visits) 72–92% (includes ED visits)	G2P[4] G1P[8], G2P[4]	87 88 86 84
USA <ul style="list-style-type: none"> Nationwide Houston, TX Houston, TX Cincinnati, OH, Nashville, TN, Rochester, NY Cincinnati, OH, Nashville, TN, Rochester, NY CA,MO,NY,OH,TN,TX,WA New Haven, CT CT, GA 	RotaTeq RotaTeq RotaTeq RotaTeq RotaTeq RotaTeq Rotarix, RotaTeq RotaTeq	100% 100% 83–86% (includes ED visits) 95% 92% (includes ED visits) 86% 94–97% 94–98%, 97% 90–92%	G3P[8] G3P[8] G1-G3, G9, G12 G1P[8] G1P[8], G2P[4], G3P[8], G12P[8] G1-G4, G9 G1P[8], G2P[4]	82 65 66 81 73 78 72 71 60

Location	Vaccine type	Vaccine effectiveness	Predominant strain	References
• CT, GA, MN				

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 6Risk of intussusception and benefits of rotavirus vaccination in Mexico, Brazil, Australia, and the US^a

Country	Diarrhea hospitalizations (deaths) prevented by vaccination	Intussusception cases (deaths) potentially caused by vaccination	Reference
Mexico	11 600 (663)	41 (2)	94
Brazil	69 600 (640)	55 (3)	94
Australia	7000 (0)	6 (0)	102
US	53 444 (14)	45–213 (0.1–0.5)	103

^aData are for one fully vaccinated birth cohort followed to age 5 y.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 7

Key issues and potential barriers and related questions in the decision to introduce rotavirus vaccine

Key issues	Potential barriers/concerns	Related questions	Potential solutions
Policy-related			
Rotavirus/diarrheal disease as a public health priority	Competing priorities (e.g., other new vaccine introductions, other health interventions on the public health agenda)	Do policy makers have sufficient information to appropriately prioritize rotavirus vaccine?	Ensure that existing disease burden and vaccine information and impact data available to policy makers
Evidence of disease burden	<ul style="list-style-type: none"> Lack of local disease burden data Lack of regional disease burden data 	<ul style="list-style-type: none"> Is the local rotavirus disease burden similar to other countries? What is the rotavirus disease burden in other countries in the region? 	<ul style="list-style-type: none"> Establish sentinel rotavirus surveillance Communicate with regional partners to share available disease burden data
Vaccine efficacy, quality and safety	<ul style="list-style-type: none"> Lack of local efficacy/effectiveness data Concerns about intussusception 	<ul style="list-style-type: none"> Will the current vaccines be able to provide protection against the local rotavirus strains? Why is vaccine efficacy worse in lower income countries? Will vaccine use be associated with an increase in intussusception cases in my country? 	<ul style="list-style-type: none"> Provide existing rotavirus strain-specific effectiveness data Investigate strategies to improve vaccine uptake Develop vaccines with improved efficacy (e.g., parenteral vaccines) Establish baseline intussusception rate and/or post-introduction intussusception surveillance if possible
Existing interventions to prevent rotavirus	Lack of awareness regarding the role of rotavirus vaccine as the main means of prevention	Won't improve access to clean water and sanitation prevent rotavirus disease?	Improve communication about the role of rotavirus vaccines as the current mainstay for rotavirus prevention
Economic and financial issues (e.g., cost-effectiveness, fiscal impact, financial sustainability)	<ul style="list-style-type: none"> High cost of existing rotavirus vaccines Lack of cost-related analyses 	<ul style="list-style-type: none"> Will rotavirus vaccines ever be sold at a lower price? What is the potential cost-effectiveness of a rotavirus vaccination program? What is the potential fiscal impact of rotavirus vaccine introduction on the immunization program? 	<ul style="list-style-type: none"> Ensure affordable vaccine prices Conduct cost-effectiveness and fiscal impact evaluations in countries that have not introduced vaccine
Program-related			
Vaccine presentation	Vaccine packaging will take up too much cold chain space	Is there sufficient cold chain capacity for rotavirus vaccines? If not, how can we increase capacity?	Support cold chain assessments (i.e., Effective Vaccine Management) and expansion as needed
Vaccine supply availability	Available vaccine supply	Will there be enough vaccine	Work with vaccine manufacturers

Key issues	Potential barriers/concerns	Related questions	Potential solutions
	insufficient to accommodate national introduction	supply for the country's target population?	to ensure sufficient global vaccine supply
Strength of the existing immunization program	Current routine immunization program not ready to introduce a new vaccine (e.g., insufficient capacity to accommodate new vaccine, insufficient resources - funding and/or staff - to conduct training sessions and social mobilization)	How will we be able to acquire the resources and technical support required to introduce rotavirus vaccine?	Provide technical support for vaccine introduction planning and implementation, including potential means for acquiring necessary funding

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript