



# Pregnancy and Birth Defects Zika Action Plan Post-Summit Teleconference

Sonja A Rasmussen, MD, MS  
Center for Surveillance, Epidemiology, and Laboratory Services  
Centers for Disease Control and Prevention

Janet Cragan, MD, MPH  
National Center on Birth Defects and Developmental Disabilities  
Centers for Disease Control and Prevention

May 11, 2016

## After this teleconference learners will be able to

- Review what we know and don't know about Zika virus infection in pregnancy, and Zika-related pregnancy and infant outcomes.
- Describe CDC's response to the threats posed by the Zika virus, including monitoring pregnancy and infant outcomes.
- Review specific considerations regarding monitoring and response to Zika-associated infant outcomes, and what CDC is doing to support rapid case-finding and referral to services for affected families.

# Zika and Pregnancy

Sonja A Rasmussen, MD, MS

# History Making




"Never before in history has there been a situation where a bite from a mosquito could result in a devastating malformation ..."

– T. Frieden, *Fortune*—April 13, 2016

# Zika Virus Clinical Disease Course and Outcomes: Adults and Children

- Clinical illness usually mild
- Many infections asymptomatic
- When symptoms do occur, they last several days to a week
- Severe disease requiring hospitalization uncommon
- Fatalities are rare
- Guillain-Barré syndrome reported in patients following suspected Zika virus infection




# Zika Virus Infection in Pregnant Women

- Pregnant women can be infected
  - Through a mosquito bite
  - Through sex with an infected male
- If infected around conception
  - Zika might present risk to fetus
- If infected during pregnancy
  - Zika can be passed to the fetus during pregnancy or around the time of birth




# CDC Lab Confirms Zika In Fetal Tissues

- Evidence of Zika virus has been identified in:
  - Amniotic fluid
  - Placenta
  - Brain
  - Products of conception



# Infants with Microcephaly



Images courtesy of  
C. Moore and Nova  
Diagnóstico



# Zika is a cause of microcephaly


The NEW ENGLAND JOURNAL of MEDICINE

SPECIAL REPORT

**Zika Virus and Birth Defects — Reviewing the Evidence  
for Causality**

**April 13, 2016**

# Many Questions Remain

- What is the level of risk from a Zika virus infection during pregnancy?
  - When during pregnancy Zika virus infection poses the highest risk to the fetus?
  - What is the full range of potential health problems that Zika virus infection may cause?
  - What are other factors (e.g., co-occurring infection) that might affect the risk for birth defects?
- 

# Collecting data for action

US Zika Pregnancy  
Registry



Zika Active Pregnancy  
Surveillance System  
(Puerto Rico)



Proyecto Vigilancia de  
Embarazadas con Zika  
(Colombia)



# US Zika Pregnancy Registry



# US Zika Pregnancy Registry : Purpose

- **Purpose of registry:**

To monitor pregnancy and infant outcomes following Zika virus infection during pregnancy and to inform clinical guidance and public health response

- **How it works:**

The registry is a supplemental surveillance effort coordinated by CDC and dependent on the voluntary collaboration of the state, tribal, local, and territorial health departments



# US Zika Pregnancy Registry

- **Who is included:**

Pregnant women with laboratory evidence of Zika virus infection and exposed infants born to these women; infants with laboratory evidence of congenital Zika virus infection and their mothers

- **How can you support the registry?**

Spread the word about the US Zika Pregnancy Registry and assist with health department follow-up for pregnant women and infants who are part of the registry



# What CDC is Doing to Respond

- Monitoring spread of Zika virus through public health surveillance
- Increasing laboratory capacity for testing to identify Zika virus infection
- Assisting with the development of tests that can improve detection of previous infection with Zika virus
- Working with partners to improve mosquito control efforts
- Providing recommendations for prevention
- Promoting effective health communication strategies
- Focusing on supporting state, local, tribal, and territorial response efforts
- Building state capacity to identify babies with birth defects

# More information: Zika and Pregnancy

- More information is available on the US Zika Pregnancy Registry on [CDC's US Zika Pregnancy Registry webpage](#)
- To contact CDC Registry staff, call the CDC Emergency Operations Center watch desk at 770-488-7100 and ask for the Zika Pregnancy Hotline or email [ZIKApregnancy@cdc.gov](mailto:ZIKApregnancy@cdc.gov)
- More information on caring for pregnant women, infants, or children with Zika virus infection is available at [CDC's Zika website](#).

The screenshot shows the CDC website interface for Zika Virus. At the top, the CDC logo and name are visible, along with a search bar and a 'CDC A-Z INDEX' dropdown menu. The main heading is 'Zika Virus'. A left-hand navigation menu lists various topics such as 'What CDC is doing', 'About Zika Virus Disease', 'Prevention', 'Transmission', 'Symptoms, Diagnosis, & Treatment', 'Areas with Zika', 'Information for Specific Groups', 'For Health Care Providers', 'Clinical Evaluation & Disease', 'Diagnostic Testing', and 'Pregnancy Registry'. The 'Pregnancy Registry' item is highlighted. The main content area features the heading 'US Zika Pregnancy Registry' with social media icons for Facebook, Twitter, and YouTube. Below this, there is a section titled 'Need for a US Zika Pregnancy Registry' which contains text explaining the registry's purpose: 'Zika virus infection during pregnancy has been linked to adverse outcomes including pregnancy loss and microcephaly, absent or poorly developed brain structures, defects of the eye and impaired growth in fetuses and infants. Despite these observations, very little is known about the risks of Zika virus infection during pregnancy. Information about the timing, absolute risk, and spectrum of outcomes associated with Zika virus infection during pregnancy is needed to direct public health action related to Zika virus and guide testing, evaluation, and management.' At the bottom of the page, contact information is provided: 'For questions about the registry please email [ZikaPregnancy@cdc.gov](mailto:ZikaPregnancy@cdc.gov) or call 770-488-7100.



# Zika and Birth Defects

Janet Cragan, MD, MPH



# What is Microcephaly?


- Clinical finding of a small head when compared to infants of same sex and age
- Measured by head circumference (HC) or occipitofrontal circumference (OFC)
- Reliable assessment of intracranial brain volume
- Often leads to cognitive and/or neurologic issues
- Difficult birth defect to monitor because of inconsistent definition and use of terminology
  - Clinicians use different cut-points such as less than 3<sup>rd</sup>, 5<sup>th</sup>, or 10<sup>th</sup> percentile for age and sex



# Range of Microcephaly Severity



Baby with Typical Head Size



Baby with Microcephaly



Baby with Severe Microcephaly

[CDC's microcephaly webpage](#)



# Types of Congenital Microcephaly

- Disproportionate - Head is small out of proportion to the weight and length, which may be normal for age and sex
- Proportionate - Head size, weight and length all are small for age and sex but proportional to each other
- “Relative” microcephaly - Head size measures within the normal range for age and sex, but is small out of proportion to the weight and length

# Zika-Related Definition of Congenital Microcephaly

## Definite

- Live Births
  - Head circumference (HC) at birth <3rd percentile for gestational age and sex, OR
  - If HC at birth is not available, HC <3rd percentile for age and sex within the first 6 weeks of life, adjusted for gestational age if preterm
- Stillbirths and Elective Terminations
  - HC at delivery <3rd percentile for gestational age and sex

## Possible

- Live Births
  - If an earlier HC is not available, HC <3rd centile for age and sex beyond 6 weeks of life
- All Pregnancy Outcomes
  - Microcephaly diagnosed or suspected on prenatal ultrasound in the absence of available postnatal HC measurements

# Suggested Reference Charts for Head Circumference At Birth by Gestational Age

Gestational Age at Birth	Reference Chart	Web Link
33 to 43 Weeks	INTERGROWTH-21st Newborn Size at Birth Chart	<a href="https://intergrowth21.tghn.org/articles/intergrowth-21st-newborn-size-birth-chart/">https://intergrowth21.tghn.org/articles/intergrowth-21st-newborn-size-birth-chart/</a> A tool for calculating percentiles for head circumference for infants 33-42 weeks is available at: <a href="https://intergrowth21.tghn.org/global-perinatal-package/intergrowth-21st-comparison-application/">https://intergrowth21.tghn.org/global-perinatal-package/intergrowth-21st-comparison-application/</a>
34 to 32 Weeks	INTERGROWTH-21st Very Preterm Size at Birth References	<a href="https://intergrowth21.tghn.org/articles/intergrowth-21st-very-preterm-size-birth-references-and-z-scores-standard-deviations/">https://intergrowth21.tghn.org/articles/intergrowth-21st-very-preterm-size-birth-references-and-z-scores-standard-deviations/</a> A tool for calculating percentiles for head circumference for infants 24-32 weeks is also available from this site.
< 24 Weeks	INTERGROWTH-21st Fetal Growth Standards	<a href="https://intergrowth21.tghn.org/articles/intergrowth-21st-fetal-growth-standards/">https://intergrowth21.tghn.org/articles/intergrowth-21st-fetal-growth-standards/</a> A tool for calculating z-scores for fetal growth standards is also available from this site.

Intergrowth-21st Fetal Growth Standards are based on measurements in utero only. International standards for birth measurements in infants less than 24 weeks gestation are not available. For most elective pregnancy terminations and many stillbirths, accurate postnatal head circumference measurements are not possible.

For a study comparing head circumference measurements in utero to those obtained after birth, see: Melamed N, Yogev Y, Danon D, et al. Sonographic estimation of fetal head circumference: how accurate are we? Ultrasound Obstet Gynecol 2011

# Assessing Prevalence

- Subdivide the cases of microcephaly into groups
  - Severity of microcephaly
    - HC <3<sup>rd</sup> percentile for age and sex
    - HC between 3<sup>rd</sup> and 5<sup>th</sup> percentiles for age and sex
    - HC >5<sup>th</sup> percentile for age and sex
    - HC values missing
  - Known (documented) causes
    - Chromosomal or genetic abnormalities
    - Syndromes (diagnosed or suspected)
    - In utero infections and types (positive culture or antibody titers)
    - Exposure to a known teratogen (e.g., alcohol, hydantoin)
  - No documented cause
- Monitor the relative proportion of each group over time



# Brain Defects Linked with Zika in the Fetus

- Decreased total brain tissue with resulting microcephaly
- Calcium deposits in the brain indicating brain damage
- Excess fluid in the brain cavities and surrounding the brain
- Absent or poorly formed brain structures
- Abnormal eye development



# Fetal Brain Disruption Sequence

- First described in 1984 but noted in earlier literature
- Brain destruction resulting in collapse of the fetal skull, microcephaly, scalp rugae and neurologic impairment
- Photos and X-ray from 1990 series;\* phenotype appears to be present in affected babies in Brazil



\*Moore, et al. J Pediatr 1990;116:383-386.

# Adverse Outcomes and Zika Virus

- A range of other problems have been reported:
  - Eye abnormalities
  - Hearing impairment
  - Seizures
  - Swallowing impairment
  - Hypertonicity and posturing
  - Contractures, including club foot and curving of the joints
  - Severe irritability
  - Developmental delay
  - Growth abnormalities, including intrauterine growth restriction and disproportionate growth (head size alone affected)

# Draft Funding Opportunity Announcement

## #CDC-RFA-DD16-1605

- Surveillance, prevention, and intervention activities for infants with microcephaly and other adverse outcomes linked with the Zika virus
  - Establish, enhance, and maintain rapid population-based surveillance of microcephaly and other adverse outcomes (especially central nervous system defects) possibly linked to Zika virus infection during pregnancy using an active case-finding methodology
  - Participate in centralized pooled clinical and surveillance data projects
  - Ensure affected infants and families are referred to services
  - Assess health and developmental outcomes of these children
- Additional information is available on the HHS Grants Forecast website at [https://forecast.grantsolutions.gov/index.cfm?switch=grant.view&gff\\_grants\\_forecastInfoID=100001785](https://forecast.grantsolutions.gov/index.cfm?switch=grant.view&gff_grants_forecastInfoID=100001785)

# Sources for Data Collection

- Where deliveries occur
  - Birth hospitals, birthing centers/midwifery practices, home births
  - Where elective terminations are performed after prenatal diagnosis of defects
- Where children with microcephaly and central nervous system abnormalities are seen and evaluated
  - Pediatric and family practice clinics
  - Subspecialty clinics (neurology, genetics)
  - Developmental clinics, early intervention programs
- Reporting by healthcare providers and programs
  - May need to revise reporting forms to include information specific to abnormalities of interest

# Engage the Healthcare Community

- Need to educate the healthcare community about Zika-related outcomes and why reporting is important
- Provide a letter from the state health commissioner or someone in authority outlining the need for reporting and the circumstances that allow physicians to report patient information
- Collaborate with state professional societies (AAP, AAFP, ACOG, Hospital Association) to inform providers caring for infants and young children
- Provide feedback and ongoing updates to maintain reporting and ascertainment going forward

## More information: Zika and Birth Defects

- For clinical questions, please contact [ZikaMCH@cdc.gov](mailto:ZikaMCH@cdc.gov)
- Information about microcephaly, including webinar on conducting surveillance: <http://www.cdc.gov/ncbddd/birthdefects/microcephaly.html>
- National Birth Defects Prevention Network: <http://nbdpn.org/>

# Thanks to our many collaborators and partners!

**Sonja Rasmussen, MD, MS**

[skr9@cdc.gov](mailto:skr9@cdc.gov)

**Jan Cragan, MD, MPH**

[jdc9@cdc.gov](mailto:jdc9@cdc.gov)

For more information, contact CDC  
1-800-CDC-INFO (232-4636)  
TTY: 1-888-232-6348 [www.cdc.gov](http://www.cdc.gov)

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

