

CDC PUBLIC HEALTH GRAND ROUNDS

Addressing the Unique Challenges of Opioid Use Disorder in Women



January 17, 2017



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

The Opioid Overdose Crisis and the Federal Response



Karin A. Mack, PhD

Associate Director for Science

Division of Analysis, Research, and Practice Integration

National Center for Injury Prevention and Control



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

“We have to stop treating addiction as a moral failing, and start seeing it for what it is: a chronic disease that must be treated with urgency and compassion.”

– Dr. Vivek H. Murthy, United States Surgeon General

FACING ADDICTION IN AMERICA

The seal of the U.S. Department of Health & Human Services is faintly visible in the background. It features a central figure holding a staff with a snake, surrounded by the text "U.S. DEPARTMENT OF HEALTH & HUMAN SERVICES" and the year "1798".


*The Surgeon General's Report on
Alcohol, Drugs, and Health*

U.S. Department of Health & Human Services


Call 1-800-662-HELP or visit [samhsa.gov/find-help](https://www.samhsa.gov/find-help)

Changing Landscape of Prescription Drug Overdoses – Rise of an Epidemic

2002 Rapid Increase in Drug Overdose Death Rates by County





2007 Rapid Increase in Drug Overdose Death Rates by County



2014

Rapid Increase in Drug Overdose Death Rates by County





The amount of opioid prescriptions dispensed has
QUADRUPLED
since 1999



But the pain that
Americans report remains
UNCHANGED



A close-up photograph of a hand holding a black pen, poised to write on a prescription form. The form has a square box containing the letters 'RX' and a line labeled 'Prescriber:'. In the background, a blue stethoscope is visible on a white surface.

Quarter billion

opioid prescriptions in 2013

A Growing Epidemic Among Women



Every 3 minutes, a woman goes to the emergency department for prescription painkiller misuse or abuse.



Vital^{CDC}**signs**TM

www.cdc.gov/vitalsigns


Since 1999, there
have been more than



71,800
deaths among women from
overdoses related to prescription
opioids


Every year since 2007 more
women have died from
drug overdoses than from
motor vehicle crashes

Rise in Female Overdose Deaths and Recent Increase in Heroin and Fentanyl Deaths




National Vital Statistics System Mortality File at wonder.cdc.gov/mcd.html

Sharp Increase in Opioid Prescriptions Associated with Increase in Deaths



A Major Risk Factor for Heroin Use



7 out of 10 women
who used heroin in
the past year also
misused opioids in
the past year


Overview of Federal Response

Three Pillars of CDC's Opioid Prevention Work

- **Improve data quality and track trends**
- **Strengthen state efforts by scaling up effective interventions**
- **Support healthcare providers with resources**



CDC Guideline for Prescribing Opioids for Chronic Pain



Altmetric Score
altmetric.com/

Three Key Principles of the Guideline

- **Non-opioid therapy preferred for chronic pain**
(Outside of end-of-life care)
- **Lowest possible effective dosage should be prescribed**
- **Providers should closely monitor all patients prescribed opioids**
 - Use of prescription drug monitoring programs (PDMPs)

CDC Guideline Implementation



1

Translation and Communication



2

Clinical Training



3


Health System Implementation



4

Insurer Implementation

Educational Resources



➤ Patient materials

- Graphics and messages
- Fact sheets
- Posters
- Podcasts
- Infographics

Tools and Materials

Checklist for prescribing opioids for chronic pain

For primary care providers treating adults (18+) with chronic pain ≥ 3 months, excluding cancer, palliative, and end-of-life care.

CHECKLIST

When **CONSIDERING** long-term opioid therapy

- Set realistic goals for pain and function based on diagnosis (eg, walk around the block).
- Check that non-opioid therapies tried and optimized.
- Discuss benefits and risks (eg, addiction, overdose) with patient.
- Evaluate risk of harm or misuse.
 - Discuss risk factors with patient.
 - Check prescription drug monitoring program (PDMP) data.
 - Check urine drug screen.
- Set criteria for stopping or continuing opioids.
- Assess baseline pain and function (eg, PFG scale).
- Schedule initial reassessment within 1–4 weeks.
- Prescribe short-acting opioids using lowest dosage on product labeling; match duration to scheduled reassessment.

If **RENEWING** without patient visit

- Check that return visit is scheduled ≤ 3 months from last visit.

REFERENCE

EVIDENCE ABOUT OPIOID THERAPY

- *Benefits of long-term opioid therapy for chronic pain not well supported by evidence.*
- *Short-term benefits small to moderate for pain; inconsistent for function.*
- *Inconsistent evidence for long-term benefits in low back pain, headache, and fibromyalgia.*

NON-OPIOID THERAPIES

Use alone or combined with opioids, as indicated:

- Non-opioid medications (eg, NSAIDs, TCAs, SNRIs, anti-convulsants).
- Physical treatments (eg, exercise therapy, weight loss).
- Behavioral treatment (eg, CBT).
- Procedures (eg, intra-articular corticosteroids).

EVALUATING RISK OF HARM OR MISUSE

Known risk factors include:

- Illegal drug use; prescription drug use for nonmedical reasons.
- History of substance use disorder or overdose.
- Mental health conditions (eg, depression, anxiety).
- Sleep-disordered breathing.



**SAFER
PRESCRIBING
AT YOUR
FINGERTIPS.**

DOWNLOAD THE
OPIOID GUIDE
APP TODAY

www.cdc.gov

Training Resources: Webinars

Topics include:

Non-opioid treatments for chronic pain

Dosing and titration of opioids

Strategies to reduce opioid overdoses

Effectively communicating with patients




CDC Overdose Prevention for States Initiative: Components

- Prescription drug monitoring programs (PDMPs)
- Community, insurer, or health system interventions
- State policy evaluation
- Rapid response projects




CDC Overdose Prevention in States Initiatives



HHS Secretary's Opioid Initiative: Focus on Three Priority Areas to Save Lives



Provide resources to assist health professionals in making informed prescribing decisions



Increase use of naloxone (Narcan[®] and others)



Expand use of Medication-Assisted Treatment (MAT) (e.g., buprenorphine, methadone)

HHS Activities

White Paper: Opioid Use, Misuse, and Overdose in Women

This paper was prepared for the U.S. Department of Health and Human Services Office on Women's Health with contract support from NORC at the University of Chicago.
December 2016



Department of Health and Human Services
Substance Abuse and Mental Health Services
Administration

State Targeted Response to the Opioid Crisis Grants

Short Title: Opioid STR

(Initial Announcement)

Funding Opportunity Announcement (FOA) No. TI-17-014

Catalogue of Federal Domestic Assistance (CFDA) No.: 93.788

PART 1: Programmatic Guidance

Note to Applicants: This document MUST be used in conjunction with SAMHSA's "Funding Opportunity Announcement (FOA) PART II: General Policies and Procedures".

Key Dates:

Application Deadline	Applications are due by February 17, 2017.
----------------------	--



National Institutes of Health
Turning Discovery Into Health

HHS: U.S. Department of Health and Human Services
[womenshealth.gov/files/documents/white-paper-opioid-508.pdf](https://www.womenshealth.gov/files/documents/white-paper-opioid-508.pdf)
www.samhsa.gov/grants/grant-announcements/ti-17-014
sciencedirect.com/science/article/pii/S030646031630329X

Join The Movement: TurnTheTideRx.org



Surgeon General VADM
Vivek Murthy, MD

**T U R N
T H E
T I D E**



Calls to Action:

Pledge to combat opioid misuse

Recognize addiction as a chronic illness for which effective treatment exists

Comprehensive Approaches to Care of Women with Substance Use Disorders



Linda Frazier, RN, MCHES, CADC

*Chair, Alcohol, Tobacco, and Other Drugs Section,
American Public Health Association*

Director, Addictions Initiatives, Advocates for Human Potential



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

Women Need Gender-Responsive Approaches

- 1. Women benefit from approaches that account for both biological (sex) and psychosocial (gender) differences associated with their substance use**

Women Need Treatment, Behavioral Interventions and Supportive Environments

2. Women with opioid use disorders require access to

- Medication-assisted treatment

AND

- Effective, trauma-informed behavioral interventions
- Safe settings that allow them to continue to care for their children



Recovery Support Needs to Address Women's Multiple Roles in Society

3. Treatment and recovery support may include

- Care coordination
- Safe housing
- Domestic violence services
- Child care
- Transportation
- Parenting support
- Opportunities to connect with other women in recovery



Biologic Differences Create Increased Risks for Women

- **Substance use in women progresses more quickly to dependency and to the onset of medical problems and disorders**
 - Smaller body mass, and higher fat-to-water ratio
 - Differences in metabolism, absorption, and elimination
- **Women can have more severe withdrawal and have higher risk of opioid overdose**

Gateways to Initiation of Substance Use Are Different

- **Women more likely than men to define selves in terms of their relationships and obligations**
- **Influence of intimate partners in starting substance use**
- **Influence of relationships**
 - Family, friends, and peers who use
- **Women more likely to move in and out of periods of problematic use**
- **Trauma history often beginning in childhood**

Price A, Simmel C. *Partners' influence on women's addiction and recovery: The connection between substance abuse, trauma, and intimate relationships*. 2002, Berkeley, CA: University of California at Berkeley.

Downs W, Miller B. In Wekerle C, Wall, AM. *The violence and addiction equation: Theoretical and clinical issues in substance abuse and relationship violence*. 2002, New York, NY: Brunner-Routledge.

Girls and Drug Use



- **Drug use is increasing among adolescent girls and women at higher rates than for men and boys**
- **This increase is also reflected in increasing rates of criminal justice involvement and incarceration of girls and women**

Women's Pathways to Use

➤ **Co-occurring disorders**

- More likely for women than men
- Women have higher rates of depression, anxiety
- May use substances to relax, reduce stress, focus attention, increase confidence
- May use substances in relation to eating disorders or body image concerns
 - ▣ Seeking effects such as weight loss, increased energy

➤ **Heavy use of prescription medications**

- Especially for older women

Women and Prescription Drugs

- **Healthcare professionals tend to miss signs of addiction in females**
 - Especially in older women and younger girls
- **Females are more likely to**
 - Be prescribed a drug by a physician
 - Receive long-term prescriptions for sedatives and analgesics for depression, anxiety and other disorders



Screening Brief Intervention and Referral to Treatment (SBIRT)

- **Screening** assesses severity of use and identifies appropriate level of treatment
- **Brief intervention** raises awareness regarding use and motivates behavior change
- **Referral to treatment** provides access to treatment, recovery supports, and specialty care as needed

Gender Differences In Seeking Care

- **Unlike men, women commonly report stigma as one of the top reasons they do not seek treatment for substance use disorders**
- **Women seeking treatment report high rates of childhood victimization, histories of sexual abuse, and current danger (47%) from violent partners**
- **Women tend to enter treatment at a much later stage of addiction, with more serious health complications due to accelerated physiological damage**

womenshealth.gov/; https://archives.drugabuse.gov/NIDA_Notes/NN0013.html

store.samhsa.gov/product/TIP-51-Substance-Abuse-Treatment-Addressing-the-Specific-Needs-of-Women/SMA15-4426

Downs W, Miller B. In Wekerle C, Wall, AM. *The violence and addiction equation: Theoretical and clinical issues in substance abuse and relationship violence*. New York, NY: Brunner-Routledge.

Culture and Substance Use Disorders

“The complex interplay of culture and health—as well as the influence of differing attitudes toward, definitions of, and beliefs about health and substance use among cultural groups—affects the psychosocial development of women and their alcohol, drug, and tobacco use and abuse.”

**–*Addressing the Specific Needs of Women* (TIP No. 51),
SAMHSA’s Center for Substance Abuse Treatment**

Core Principles of Gender-Responsive Care


Gender-responsive care:

- 1. Addresses women's unique experience**
- 2. Is trauma-informed**
- 3. Uses a relational approach**
- 4. Is comprehensive**
- 5. Provides a safe healing environment**

Six Principles of Trauma-Informed Care

Trauma-informed care:

1. Safety
2. Trustworthiness and transparency
3. Peer support and mutual self-help
4. Collaboration and mutuality
5. Empowerment, voice, and choice
6. Cultural, historical, and gender issues



Selected Evidence-Based Practices

- **Pharmacotherapies (i.e., medication-assisted treatment)**
- **Screening Brief Intervention and Referral to Treatment (SBIRT)**
- **Motivational approaches**
- **Cognitive Behavioral Therapies**
- **Family therapies**
- **Contingency management**
- **Telehealth and technological applications**
- **Peer support**

Major Dimensions that Support Recovery

- **Health**—learning to manage one’s diseases or symptoms, abstain from use of alcohol, illicit drugs, and non-prescribed medications, and make healthy choices that support physical and emotional well-being
- **Home**—have a stable and safe place to live
- **Purpose**—meaningful activities, such as a job, family caretaking, or creative endeavors, and the independence, income, and resources to participate in society
- **Community**—having relationships and social networks that provide support, friendship, love, and hope

Recovery Support – Hope is Foundational

- **Knowing that someone else overcame similar challenges gives women the slightest bit of optimism that things could get better**
- **Developing self-efficacy and patient activation**
- **Success begets success**

Words and Actions Matter

*"Words are important.
If you want to care for something, you call it a flower,
if you want to kill something, you call it a weed."*

– Don Coyhis, Recovery Advocate, Educator, President and Founder of White Bison

Change How We Label Something and We Change How We Think of It

Say this ...

Substance use disorder

Resourceful

Woman with a violent partner

Neonatal exposure or abstinence
syndrome NAS

Not this ...

Alcohol or drug problem

Manipulative









Battered woman

Neonatal dependency

Putting Principles into Practice

- **Medications combined with behavioral therapies and ongoing recovery supports**
- **Further integration of primary care, mental health and SUD's services and supports**
- **Bringing comprehensive, gender-responsive, trauma-informed and family-centered services to scale**

SAMHSA Resources Sampling



Prevention of Substance Use Through Integration into Healthcare



Mishka Terplan, MD, MPH, FACOG, FASAM
Professor, Obstetrics and Gynecology, and Psychiatry
Virginia Commonwealth University



Sex and Gender Differences


**CAUTION
MEN AT WORK**



Women work all the time-
Men have to put up signs when they work.

Focus on Women's Needs Over Their Life Course

The typical woman spends 5 years pregnant, postpartum, or trying to get pregnant, and 30 years trying to avoid getting pregnant



Adapted from: Gold RB, Sonfield A, Frost JJ, et al. *Guttmacher Institute*, Feb 2009. guttmacher.org/sites/default/files/report_pdf/nextsteps.pdf


Gender and Behavioral Health Burden

Diagnosis	Percent Reporting	
	Female	Male
Serious Psychological Distress (past month)	6.0%	4.1%
Any Mental Illness (past year)	26.2%	17.3%
Serious Mental Illness (past year)	5.0%	3.0%
Major Depressive Episode (past year)	8.5%	4.7%


Opioid Epidemic Is Increasingly Young, White, and Female

- **Women of reproductive age (15–44 years) receive more prescription medications than men**
- **More women than men are initiating opioid misuse**
- **In 2015:**
 - 0.9 million males
 - 1.2 million females
- **3,300 women per day**

Reproductive Health along a Substance Use Continuum



Women Seeking Healthcare Have Concerns Across Multiple Health Domains




Contemporary Healthcare System: United States




- **Providing care in silos**
 - Does not meet the needs of individuals
 - Inefficient
 - Costly

Unmet Mental Health Needs Contribute to Increased Overall Healthcare Costs

Monthly Healthcare Expenditures for Chronic Conditions, with and without Comorbid Depression, 2005



Solution is Integration of Mental Health, Substance Use, and Primary Care Services



U.S. Healthcare Reform: The Affordable Care Act (ACA)

➤ Integration

- Behavioral health: mental health and addiction
- Somatic and behavioral health
- Reproductive health integration

➤ Parity

- Reimbursement for primary care and mental health-related services, including substance use treatment
- Medicaid expansion
 - Contraceptive coverage

Treating a Biobehavioral Disorder Must Go Beyond Just Fixing the Chemistry


We Need to Treat the Whole Person!

**Pharmacological
Treatments
(Medications)**

Behavioral Therapies

Medical Services

Social Services



In Social Context

Shift Focus to Entire Life Course, Not Just Pregnancy

Prevalence of Reproductive Health Hits in Search Engines

	Reproductive Health	Sexual Health	Contraception	HIV	Pregnancy
NIDA	21	22	17	125,000	19,800
SAMHSA	55	29	43	3910	1350
ASAM	6	3	7	179	121


NIDA: National Institute on Drug Abuse

SAMHSA: Substance Abuse and Mental Health Services Administration

ASAM: American Society of Addiction Medicine

Unpublished data

NIDA Principles of Drug Addiction Treatment



4. EFFECTIVE TREATMENT ATTENDS TO MULTIPLE NEEDS OF THE INDIVIDUAL, NOT JUST HIS OR HER DRUG ABUSE. To be effective, treatment must address the individual's drug abuse and any associated medical, psychological, social, vocational, and legal problems. It is also important that treatment be appropriate to the individual's age, gender, ethnicity, and culture.

13. TREATMENT PROGRAMS SHOULD TEST PATIENTS FOR THE PRESENCE OF HIV/AIDS, HEPATITIS B AND C, TUBERCULOSIS, AND OTHER INFECTIOUS DISEASES, AS WELL AS PROVIDE TARGETED RISK-REDUCTION COUNSELING, LINKING PATIENTS TO TREATMENT IF NECESSARY. Typically, drug abuse

Gap Between Principles and Practice

➤ **Recognition of**


- Need for holistic practice, but certain domains more included
- Importance of integration, but reproductive health and contraception lag

➤ **Inequities within reproductive health**


- Negatively affect women with substance use disorders
- Play out across the life course




**Reproductive
Health**



**Substance Use
Misuse
Addiction**



Opportunities for Public Health at the Intersection of Reproductive Health and Substance Use



Public Health at the Intersection of Reproductive Health and Substance Use

- **Bringing the silo of reproductive health into public health programming for people who use drugs**
- **Baltimore City Health Department**
 - Syringe Exchange Program



Public Health at the Intersection of Reproductive Health and Substance Use

- **Baltimore City Health Department Reproductive Health Project on the Van**
- **Integrating family planning with syringe exchange**



Injectable Contraceptive Continuation among Female Exotic Dancers Seeking Mobile Reproductive Health Services

Caitlin E. Martin, MD, MPH
Jennifer J. Han, ScM
Chris Serio-Chapman, BS
Patrick Chaulk, MD
Mishka Terplan, MD, MPH

Abstract: Objectives. We describe depot medroxyprogesterone acetate (DMPA) continuation patterns among female exotic dancers receiving reproductive health services at a mobile syringe exchange. **Methods.** Clients initiating DMPA between November 2009 and August 2012 were identified retrospectively *via* chart review. Life table analysis measured continuation. Client characteristics were compared using chi-square tests. **Results.** Sixty nine clients were identified; 72% were African American and 63% were younger than 25. At three months, 36% of the study sample continued DMPA; those continuing were more likely to be White ($p=0.01$) and receive other services ($p=.01$). The 12-month cumulative continuation probability was 0.09. Considering those who had received an injection, continuation proportions were higher (46% at 6; 71% at 12 months). **Conclusions.** A subset of female exotic dancers may favor DMPA as a long term contraceptive. Integrating mobile reproductive health services into public health programs can help fulfill the unique health needs of this high-risk population.

Contraception and Clean Needles: Feasibility of Combining Mobile Reproductive Health and Needle Exchange Services for Female Exotic Dancers

| Eva Moore, MD, MSPH, Jennifer Han, ScM, Christine Serio-Chapman, BS, Cynthia Mobley, MD, MPH, Catherine Watson, MSW, and Mishka Terplan, MD, MPH

Young women engaged in exotic dancing have a higher need for reproductive health services than women not in this profession, and many also use drugs or exchange sex for money or drugs. Few report receiving reproductive health services. We describe a public health, academic, and community partnership that provided reproductive health services on needle exchange mobile vans in the “red light district” in downtown Baltimore, Maryland. Women made 220 visits to the vans in the first 21 months of the program’s operation, and 65% of these visits involved provision of contraception. Programmatic costs were feasible. Joint provision of needle exchange and reproductive health services targeting exotic dancers has the potential to reduce unintended pregnancies and link pregnant, substance-abusing women to reproductive care, and such programs should be implemented more widely. (*Am J Public Health*. 2012;102:1833–1836. doi:10.2105/AJPH.2012.300842)

dancers and their need to engage in higher risk sex work to obtain money for drugs. Rates of HIV and other sexually transmitted infections (STIs) are high among sex trade workers, as are rates of unwanted pregnancies.⁵ Exotic dancers differ from other sex trade workers because of the club environment, which can be

Integrating Contraception with Syringe Exchange

- **Innovative prevention services in nontraditional setting**
- **\$85 per client, including clinician costs and supplies**
- **Potential to reduce unintended pregnancies**
- **Link pregnant women who use drugs to prenatal care**

Public Health at the Intersection of Reproductive Health and Substance Use

- **Addiction treatment is an opportunity to improve reproductive health**
 - Decrease infectious disease transmission (especially HIV and HCV)
 - Decrease unintended pregnancies
 - Prevent substance exposed pregnancies

- **Bringing the silo of reproductive health into addiction treatment**
 - Integrating family planning into treatment
 - Baltimore Reproductive Health Initiative

Baltimore Reproductive Health Initiative

- **Screening**
- **Education**
- **Service Delivery**

- **Funded by Abell Foundation**



Screening Tool for Family Planning Needs

**One Key Question:
Would you like to get
pregnant in the next year?**

**Question is
client-focused
nonjudgmental
closed-ended**

**Baltimore City – PSEP Reproductive Health Initiative
One Key Question® (OKQ) Client Screening Questionnaire**

Site:	<input type="checkbox"/> Recovery Network <input type="checkbox"/> Gaudenzia	Today's Date:
Birth Year:		
Sex (at birth):	<input type="checkbox"/> Female <input type="checkbox"/> Male <input type="checkbox"/> Intersex	Ethnicity: <input type="checkbox"/> Hispanic/Latin@ <input type="checkbox"/> Non-Hispanic/Latin@
Gender:	Race:	

1. **Would you like to get pregnant (or impregnate a partner) in the next year?**

No (If checked, go to question 2)

Yes (If checked, skip to question 3)

I'm OK either way or Unsure (If checked, skip to question 4)

Not applicable (menopause, sterilization, hysterectomy, currently pregnant/expecting, other: _____ (If checked, questionnaire ends here))

Declines to answer (If checked, questionnaire ends here)

If client answered "no" to question 1:

2. (A) Are you using a birth control method right now?

Yes. What type?: _____

No

(B) If yes, are you happy with your birth control method?

Yes

No and not interested in change

No and looking to change

If client answered "yes" to question 1:

3. Are you taking folic acid or a Prenatal Multivitamin?

Yes

No

4. Are you receiving pregnancy planning (preconception health) counseling with a clinician?

Yes

No

5. If no, would you like to be linked to a pregnancy planning (preconception health) counselor?

Yes

No

Service Delivery

Unique Clients	N=134	Percent
Contraceptive need	82	61%
Received contraception	68	83%
Overall LARC among those with contraceptive need	45	66%
<u>Method choice</u>		
Nexplanon - Implant	28	41%
Mirena IUD	15	22%
Combined oral contraceptive	13	19%
Depo Provera	8	12%
Paraguard/copper IUD	2	3%
Hormone patch	2	3%

IUD: Intrauterine device

LARC: Long-acting reversible contraceptive

Unpublished data

Conclusions

- **Opportunities for prevention at intersection of reproductive health and substance use, misuse, addiction**
- **Through integration – move toward greater equality and addressing injustices**



CDC PUBLIC HEALTH GRAND ROUNDS

Addressing the Unique Challenges of Opioid Use Disorder in Women



January 17, 2017



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention