Morbidity and Mortality

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

PUBLIC HEALTH SERVICE

Prepared by the

COMMUNICABLE DISEASE CENTER 634-5131

AUG I

For release August 2, 1963

ATLANTA, GEORGIA 30333

Vol. 12, No. 30

Weekly

Repor

PROVISIONAL INFORMATION ON SELECTED NOTIFIABLE DISEASES IN THE UNITED STATES AND ON DEATHS IN SELECTED CITIES FOR WEEK ENDED JULY 27, 1963

POLIOMYELITIS – The 11 cases of poliomyelitis reported for the week ending July 27, of which 10 were paralytic, continues well below the level of reported cases during 1961.

Cumulative poliomyelitis totals for the United States are compared with the previous four years in the table (right column).

California and Georgia each reported two cases this week, without geographic concentration in either State. New York, Pennsylvania, and Nebraska each reported one case. Four cases were reported from Alabama.

Three of the four Alabama cases were from Walker County, where a total of seven cases have occurred between late May and mid-July, 1%3. Additional suspect cases are under investigation, and Type I polio virus was recently isolated from two of the cases. A mass immunization program will be conducted on Saturday, August 3, in Walker County (approx. 65,000 population), with Type I oral polio vaccine from the CDC epidemic reserve.

Poliomyelitis (Cumulated Weekly) 1st Through 30th Week


	1963	1962	1961	1960	1959
Paralytic	109	230	213	609	1,342
Total	132	301	327	837	2,040
Poliomy	elitis (Six V	Veek Totalı	s) 25th Thro	ugh 30th We	ek
Paralytic	51	95	75	325	823
Total	65	126	124	450	1,304

Table 1. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES (Cumulative totals include revised and delayed reports through previous week)

Michaeld 1851 Provid La Meser 1		30th We	ek	Cumulative				
Disease	Ended	Ended	Median	First 30 weeks				
file the entry, but and the	$\begin{tabular}{ c c c c c c c c c c c c c c c c c c c$	Median 1958 - 1962						
Aseptic meningitis	50	87	and September .	806	000			
rucellosis	4		14					
rucellosis	6	2	6			448		
acephalitis, infectious	2/	33	4.2			360		
"Datitis, infectious and serum						921		
easies						21,573		
eningococcal infections						385,955		
pliomyelitis, total						1,480		
Paralutio						837		
Paralytic	10	17				609		
Nonparalytic	Server and a series	3		14	50	168		
Unspecified		-	6	9	21	60		
	3,403	3,326		229,382	210.357			
Stanue	4	9						
ularemia	9	14		160				
ularemia Yphoid fever Yphus fever, tick-borne,	10	19	19	236	308	395		
(Rocky Mountain spotted)	8	15		83	126			
abies in Animals	80	81	69	2,314	2,476	2,310		

Table 2. NOTIFIABLE DISEASES OF LOW FREQUENCY

Anthrax: Botulism: Malaria: La 1, N. C 1 Plague:	Cum. 2 Psittacosis: Mass 1, Mich 1, Ill 1 5 Rabies in Man: 53 Smallpox: - Typhus, murine:	Cum. 38 - - 12
---	---	----------------------------



EPIDEMIOLOGICAL REPORT

Salmonellosis – California

On April 19, 1963, a California County Health Department received a report that two teenagers, a 16-year-old boy and his girl friend, had become ill on April 15, one day after eating Easter Sunday dinner with another teenage couple at a local cafeteria. Three of the individuals, including the two with illness, had eaten turkey and dressing, while the other ate fish. The one person who ate turkey and dressing but did not become ill stated that the turkey and dressing served to her was from a new tray.

Investigations at the cafeteria disclosed that the management had received a report from another customer stating that three members of his party became ill on Monday after eating the turkey and dressing on Easter Sunday. No name or phone number had been obtained. Subsequently, three other individuals, in two separate groups, were reported as having developed illness on April 15, all with a history of having eaten the turkey and dressing at the cafeteria the day previously.

The five ill individuals who could be contacted had incubation periods ranging between 19 and 24 hours following consumption of the turkey and dressing dinner at the cafeteria on April 14, 1963. All patients were hospitalized and had stool cultures positive for *Salmonella schwarzengrund*. Clinical symptoms in these cases included nausea, vomiting, stomach cramps, diarrhea, chills and fever to 104° F.

Stool specimens were obtained from the 16 food handlers who had assisted in the preparation and serving of the turkey and dressing, and from the remaining 118 employees. Six of the 16 food handlers had stool cultures positive for Salmonella schwarzengrund. Among those positive was the head chef, who did most of the preparation and who stated that he had been ill with "intestinal flu" about the second week of March 1963. Positive cultures were also obtained from another cook who helped in preparing the turkey, and from two counter girls and two supply boys who assisted in serving. Ten additional positive cultures were obtained from the remaining 118 employees. Only one of the employees had complained of being ill, having had symptoms of stomach cramps, vomiting, and fever on Monday, April 15, after having eaten the turkey dinner on Easter Sunday. In addition to the 16 employees from whom Salmonella schwarzengrund was obtained, one employee was positive for Salmonella typhi.

Investigation of food handling facilities disclosed that the procedures used in preparing the dinners for Easter Sunday differed from the normal operating procedure in that preparation of the 20 frozen turkeys used for the meal had begun on April 10, four days before serving, in contrast to a 48-hour period ordinarily required. The turkeys had been cooked for adequate periods on April 11. Subsequent procedures including boning, preparation of dressing and slicing, at intervals over the subsequent 2½ days prior to serving, indicate the likelihood that contamination took place after cooking and that

246

나 없다. 그 또 남편한 태양에서 다 그렇게 만나 가 있어? 신하겠다. 이 이상하는 것 이 가 가지? 요구가 들는 것이 하네?

11717-012 - 227-01

adequate incubation times were present for growth of the organism.

Following identification of positive carriers in the kitchen staff, all employees with positive stool specimens were removed from work until their infections had cleared.

(Reported by F. A. Listick, Assistant Supervising Sanitarian, Los Angeles City Health Department, and Dr. Philip K. Condit, Chief, Bureau of Communicable Diseases, California State Department of Health.)

SURVEILLANCE SUMMARIES

Malaria Surveillance - United States - 1962

Reports of 119 cases of malaria, with onsets of illness during 1962, were received by the Malaria Surveillance Unit from 30 States, the District of Columbia and the military services. The table on page 252 shows the cases according to geographic distribution and classification.

One hundred of the 119 cases, in whom a positive blood smear had been obtained, were reported as "confirmed". Although confirmation by blood film was not obtained in the remaining 19 cases, these were classified as presumptive on the basis of clinical and epidemiological evidence suggesting malaria. *Plasmodium vivax* was confirmed in 72 cases; *Plasmodium /alciparum* in 20 cases; and *Plasmodium malariae* in 4 cases. The type of parasite was undetermined in 4 cases.

One hundred and ten of the 119 cases reported during 1962 were of foreign origin. Seventy-five of these occurred in military personnel, all of whom had served in areas of the world where malaria incidence is high. Among the 44 civilian cases, 24 occurred in Americans infected while traveling abroad, 11 among foreign visitors to the United States, and 9 were reported as indigenous to the United States.

The 9 cases of malaria classified as indigenous were reported from six States. Of these, a California case and one from Illinois occurred in persons with previous histories of malaria, who had no known recent exposure. One case (reported from Tennessee) was voluntarily infected while a prisoner. Multiple units of whole blood, the donors of which have not yet been completely studied, had been received by the case reported from New York, which is believed to have been transfusion-acquired. No definite source of infection has been found for the remaining five cases, reported from Arizona, Illinois, and New Hampshire.

The reported malaria morbidity and mortality in the United States for the period 1932-1962 is shown in the figure on pape 246. The moderate rise in total cases during 1962, as compared to the previous four years, is due to an increase in the number of cases among military personnel, especially those who served in Southeast Asia. Of the 119 cases reported during 1962, 75 were accounted for by military personnel, as compared to 45 for the year 1961.

(Case reports included in the Malaria Surveillance Summary for 1962 were submitted by State Health Departments and the military services. To facilitate reporting National Malaria Surveillance Report forms are available from the Surveillance Section, Communicable Disease Center. In addition a National Malaria Depository

(Continued on page 252)

SUMMARY OF PNEUMONIA AND INFLUENZA DEATHS

The weekly average number of pneumonia-influenza deaths for the four-week period ending July 27 was 410 as compared with an expected weekly average of 386.

PNEUMONIA-INFLUENZA	DEATHS IN	108 CITIES

		WEEK	ENDING	for several	4 Week	Weekly	
	7/6	7/13	7/20	7/27	Total	Average	
Observed	426	459	369	388	1,642	410	
Expected	393	388	384	380	1,545	386	
Excess	33	71	-15	8	97	24	




Table 3. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES FOR WEEKS ENDED

JULY 27, 1963 AND JULY 28, 1962

many that we half	Polio	myelitis,	total c	ases	Pol	iomyeliti	s, paraly	ytic	Poliomy nonpara		and the second sec	ptic ngitis
Area	20+1		and the second sec	ative		121.12		lative	1.34	Barry # 51	-011 (M	and the second
shing mounted at hour	30th			30 weeks	30th	1	First 3	30 weeks		week	30th	week
	1963	1962	1963	1962	1963	1962	1963	1962	1963	1962	1963	1962
UNITED STATES	11	20	132	301	10	17	109	230	1	3	50	87
NEW ENGLAND	-	-	1	4		1000	1	4	CTAB.	11100.010	00111	1
Maine New Hampshire	-	-	sata 🖓 Ma	10 10 10		/	-		-		-	-
Vermont.	1		-	the second second	-		-	-				-
Massachusetts	1.2	1.1	1	3				-	-		-	-
Rhode Island	-	-	-	2	1.2.1		1	3	-		-	1
Connecticut	-	-	-	1	5-E 1		-	1	. : : : : :	44023	DHOT	ELV PAR
MIDDLE ATLANTIC	2	-	35	39	1		26	26	1		3	3
New York.	1		6	35	-	N. 199	5	20	1	S [11]	3	2
New Jersey	-	1	1	3	-		1	3			-	-
Pennsylvania	1	111 34	28	1	1	1000	20	1	100	1.0		1
EAST NORTH CENTRAL	-	4	20	18		3	13	12	SIT-	1	8	15
Ohio Indiana.	-	1	7	6	-	1	3	6		-	-	5
Illinois	-	1	27	4	0.5.0	1	1	3	-		2	3
Michigan	-	2	2	0		1	6	3		-	3	3
Wisconsin	-	1	2	2	1.1	state-of	2	11.101	- VII -	0 <u>1</u> 1	3	4
WEST NORTH CENTRAL	1	-	5	13	1	1.1	5	7	binde	meed 1	3	3
Minnesota	-	-	3	3	1.2.3	10.00	3	3	111111111	HDO HUH	3	3
Iowa. Missouri	-	- 1		5	i - 1	-	100.000	2	- 93-au		-	-
North Dakota		lest-Tri	1	5	-	-	1	2	-	-	-	-
South Dakota	1.2	-	-				-	Proc. 201	-			-
Nebraska	1		ī				N	10 BT-14	P 1204	1.57 .00	-	-
Kansas	-	-		2. 22	1	-	1	280 - 21	1.00	100	1000	2.5
SOUTH ATLANTIC	2	2	17	18	2	2	14	15	1.	217.12		1
Delaware	-	_		-	-	-	14	15	1.1211	1.2.11	6	-
Maryland	0.00	parte 17	sait-in	- 12 m		1.1	2	1.1			1	125
District of Columbia Virginia	-	-	-	1	-	-	-					-
West Virginia.	-	1	2	3	-	1	1	3	-	1.1		-
North Carolina.	1		1 3	3			1	3		015.00		1.1.1.1.1.1.1
South Carolina	-	1	3	2		- 1	3	2		1 2110	100	
Georgia	2		3	2	2	-	2	1 2			- CT-	
Florida	-	-	5	6	-	1	5	4	100		6	1
EAST SOUTH CENTRAL	4	3	21	13	4	3	19	9	the second s	the state of the s	1	5
Kentucky.	-	1		6	-	1		4	-		1	-
Tennessee	-	-	4	3	-		4	1	_	1000	-	1
Alabama. Mississippi.	4	1	15	3	4	1	13	3		1.14.31	-12-10	
	-	1	2	1	- 1	1	2	1		1.00	1.0.2	4
WEST SOUTH CENTRAL	-	10	17	154		8	17	121		2	3	33
Arkansas	-		1	1		1.0-1.0	1	1	10.504	161.5214	201 10	0.004.12
Oklahoma		1	14	9		-	14	7	and there	1		-
Texas.	1	9	2	2 142	- 21 3	8	2	113		1	1 2	2 31
MOUNTAIN			N. Comment				day in		riet Inc.		in the	
Montana	-	-	-	9 3		-	-	7	-	-	2	-
Idaho		-		د		-		2			-	
Wyoming.	-	-	10 million (1997)	1	- 2 -			1			-	
Colorado.	-	-	and the second	1				1			1 -	-
New Mexico Arizona	-	-			C	-	-	-	1			-
Utah.	- 2	-	-	3			-	3		7,746	2	-
Nevada	-	-	- 1	1	1.1	1 .	100-107	1	The second	1412.23		-
PACIFIC	2	1	16			-					- opline	
Washington	-	-	16 -	33	2	1	14	29	12171	30173	24 2	26
Oregon		-	2	5	1.00		1	5	-		2	-
California. Alaska	2	1	14	28	2	1	13	24	-	-	22	26
Hawaii.						-	-			1.45.1		-
	-	-	-	-	-	-	/-	-	1-1-5	1000	-	-
Puerto Rico		-	4	9	-		4	9	-	1.503		-

Morbidity and Mortality Weekly Report

Table 3. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES FOR WEEKS ENDED

JULY 27, 1963 AND JULY 28, 1962 (Continued)

a history and are last	Brucel	losis	Diphth	eria	Encepha infect		infe	Hepati ctious a		n	Measles	
Area		Cumu- lative	17.3	Cumu- lative		0.12.09			week			
the state of the state	30th week	30 weeks	30th week		30th	week	Under 20 yr.	20 & over	To	tal	30th	week
2011 Cont 1 7 10 1	1963	1963	1963	1963	1963	1962	1963	1963	1963	1962	1963	1962
UNITED STATES	4	207	6	150	24	33	340	315	689	808	2,532	2,60
NEW ENGLAND		_		7		2	22	10	70	- 0		
Maine	- 1			1 <u>1</u>	1	-	33	40 12	78 28	58 15	95	2
New Hampshire	-					1.2	9	8	17	15	12	
Vermont				1 1 1 ×	-		2	1	3	2	16	
Massachusetts		- 1		5		2	5	16	21	34	40	
Rhode Island		-	-	2	1	-	-	S C	4	1	3	1.00
Connecticut	-	-				1.1	1	3	5	5	19	1000
MIDDLE ATLANTIC	- I II.	5	-	21	6	13	84	66	150	174	577	4
New York	- T - 1	3		13	5	10	48	40	88	96	362	3
New Jersey		-	-	3			16	9	25	29	143	1
Pennsylvania	1	2		5	1	3	20	17	37	49	72	
EAST NORTH CENTRAL	1 - L	18	5	22	1	4	64	60	127	151	638	6
Ohio		-		1		K.C 02	8	14	22	51	126	1.01
Indiana Illinois	1.20	5	1	5		-	5	3	8	12	43	125
Michigan		10	4	11 3 -	-	22	26	18	46	34	105	
Wisconsin	1 - L	1		2	1 -	2 -	22	23 2	45	46 8	200	2
EST NORTH CENTRAL	1	127		37						111.000		- 201
Minnesota	-	8		15	3		12	11	24	48 7	61	
Iowa	-	92	1.12 - 2.1	1		1	1	0.1	2	11	29	1.11
Missouri		7	- E - C	1	1		1	2	3	11	3	
North Dakota	-			1	1		1	4	5	10	24	1123
South Dakota		8		11	-	10.0	5	1	6	2	5	1100
Nebraska Kansas	1	6		8		10.70	3	i - 1	3	3	1.1 - T.A.	
Same a Street of Acres				1.1	1		1	4	5	4	NN	
SOUTH ATLANTIC Delaware	1	9	1	30	5	2	42	33	76	80	244	2
Maryland	ST 11.		1	-	-		1		1	2	11	
District of Columbia.	3 -				1	1	6	4	10	13	43	
Virginia		3		1.1.1.1.1			8	4	13	3 16	1 52	110
West Virginia				1	_		7	5	12	11	86	111
North Carolina	1	2		1	1		8	15	23	15	13	1
South Carolina		-		9			2	1	3	4	6	2010
Georgia Florida		3 1	- 1	10	- 3	1	3	1	4	4	1	
State of the second	21	1	1	9	3	1	4	2	6	12	31	
AST SOUTH CENTRAL		10		10	1	1	22	20	42	78	113	1
Kentucky		3	-		1		1	7	8	26	35	
Tennessee	- 1. TO	5		2		-	7	6	13	24	63	
Mississippi		2		8	1.15	1	6	3	9 12	19 9	12	
EST SOUTH CENTRAL				1.7						1	1-12-1	
Arkansas	2	24 4		17	1	2	39	24	64	54	145	2
Louisiana	2	5		2			8	1	13	8	2	1844
Oklahoma	-	5	_	6	1.1.1.1	1	1	-	13	11	3 15	
Texas	S 1	10		8	1	2	27	18	46	32	125	1
OUNTAIN	- 11	7	_	4	-	100	6	4	31	26	254	
Montana	- I - I	-		-	9		4	1	5	20	38	2
Idaho		-	-	-	1.48	2		-	6	4	24	1.000
Wyoming	I	1				-	-	-		3	1	1504
Colorado	-	T	-	3	100		-		4	5	67	- 19
New Mexico Arizona		-		1			2	-	2	2	NN	
Utah		3			-			-	11	2	94	
Nevada	-	5			-	-		3	3	8	30	April 1
ACIFIC		7		2		_		_				1
Washington	10.1	7		2	6	8	38	57	97	139	405	4
Oregon		2		-	1	1	4	3	7	37	34	
Californía	E 10	4		2	4	7	2 29	7 46	10 75	11 86	35 271	1 2
Alaska		· · · · ·		1.2	2	-	3	40	4	4	57	-
Hawaii	-	1	-			-	-	1	1	i	8	
uerto Rico	-						1				f	-

Table 3. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES

FOR WEEKS ENDED

Meningococcal Streptococcal Tickborne Typhus Infections Sore Throat & Tetanus Tularemia Typhoid Fever Rabies in Animals Scarlet Fever (Rcky Mt. Cumu-Spotted) Cumu-Cumu-Area lative lative lative 30th wk. 30+h week 30th wk. 30th wk. 30 weeks 30th wk. 30th wk. 30 weeks 30th week 30 weeks 1963 1963 1963 1962 1963 1963 1963 1963 1963 1963 1962 1963 UNITED STATES 34 1,581 3,403 3,326 4 8 9 10 80 81 236 2,314 NEW ENGLAND 4 99 254 171 6 24 1 17 Maine..... 9 -4 ----1 New Hampshire 4 _ -12 Vermont..... 3 4 2 1 -_ _ 7 --10 Massachusetts..... 2 46 53 49 -.... _ 4 -1 Rhode Island 9 . 4 11 4 _ _ -. -Connecticut..... 1 20 184 106 -.... 1 -MIDDLE ATLANTIC ... 7 220 109 76 -4 55 3 7 71 New York 4 100 -83 55 _ -1 19 2 5 53 New Jersey..... 1 32 14 9 -. 2 -Pennsylvania..... 2 88 12 _ 12 _ 3 34 1 2 18 EAST NORTH CENTRAL .. 6 219 201 247 -_ 1 16 16 34 388 Ohio..... 1 72 15 18 _ _ _ 6 4 24 231 Indiana..... 31 35 -23 -1 2 2 38 Illinois..... 1 42 50 30 -_ 1 6 1 55 Michigan..... 4 75 78 -2 81 -1 6 38 Wisconsin 27 41 . 49 7 . _ _ 2 4 26 92 77 WEST NORTH CENTRAL. -96 4 2 15 З 31 23 582 Minnesota..... 19 -..... -1 3 3 10 143 6 _ Iowa..... -36 19 5 1 2 12 224 8 Missouri..... -31 --1 1 8 102 5 3 North Dakota 4 40 43 _ . 22 3 1 South Dakota -5 -2 --1 ... 4 1 63 1 Nebraska..... 20 1 -17 Kansas..... 8 29 . 1 _ 1 -11 7 SOUTH ATLANTIC 298 263 454 6 1 7 9 -38 335 Delaware..... 2 2 3 -1 1 Maryland..... 3 47 38 11 2 1 -5 ----Dist. of Columbia... 4 I ---_ 1 67 72 86 -2 Virginia..... -5 1 2 121 West Virginia..... 16 70 134 -_ -6 2 2 98 North Carolina 1 _ _ 52 11 5 --5 1 -6 South Carolina 14 1 129 _ 1 30 --2 6 Georgia..... . -22 1 1 3 -_ 2 41 . Florida..... _ 1 69 26 102 -12 2 4 62 EAST SOUTH CENTRAL .. _ 119 737 751 1 1 21 7 3 188 Kentucky..... -25 8 34 1 -_ 3 88 -1 1 Tennessee..... -53 619 648 -4 14 4 1 84 -Alabama..... 21 21 8 1 L 4 1 16 _ Mississippi..... 20 89 61 2 WEST SOUTH CENTRAL .. 1 154 548 560 2 ÷ 4 1 39 8 445 -Arkansas..... 10 -1 _ -18 32 Louisiana..... 63 1 1 1 38 8 -Oklahoma. 29 12 1 -4 35 3 -Texas..... 1 52 535 559 1 10 7 340 -MOUNTAIN 1 1 57 53 757 -7 3 565 1 1 Montana..... -3 26 21 ------Idaho..... 4 26 63 -----Wyoming..... 4 ---13 8 _ -Colorado..... 12 319 194 1 _ 6 -3 -New Mexico..... 4 190 21 161 ----1 Arizona..... 1 108 9 . 70 --1 3 3 30 Utah..... 14 1 75 48 _ -1 --Nevada..... . 3 --..... ---299 PACIFIC 8 439 452 1 2 39 5 4 224 1 24 Washington..... 3 38 34 --1 20 18 Oregon..... 14 --4 2 1 1 5 1 California..... 236 347 390 -2 33 5 4 214 Alaska..... 11 29 12 4 9 ---1 -Hawaii..... 8 7 2 _

JULY 27, 1963 AND JULY 28, 1962 - (Continued)

Puerto Rico.

-

2

...

_

11

1

-

3

10

14

1

5

Morbidity and Mortality Weekly Report

Table 4 (B). REPORTED PNEUMONIA-INFLUENZA DEATHS IN REPORTING CITIES

(Tables 4(A), 4(B), 4(C), and 4(D) will be published in sequence covering a four-week period.)^o

Area	21	For week	s ending	1.11	Area	For weeks ending				
	7/6	7/13	7/20	7/27		7/6	7/13	7/20	7/2	
EW ENGLAND:		1.	198.10		SOUTH ATLANTIC:	1. 1.	1925	5 . Sel	2	
Boston, Mass	5	4	8	3		CIAME TO		BIROSI	1.1	
Bridgeport, Conn	13	1	4	2	Atlanta, Ga Baltimore, Md	1	1	1	5	
Cambridge, Mass	0	0	4	õ	Charlotte, N.C	5	6	7	6	
Fall River, Mass	ō	1	Ö	1	Jacksonville, Fla	0	1	1	3	
Hartford, Conn	0	1	ő	ô	Miami, Fla		1	1	0	
Lowell, Mass	0	3	1	2	Norfolk, Va	0	1	0	1	
Lynn, Mass	0	1	2	õ	Richmond, Va	6	4	0	2	
New Bedford, Mass	1	2	ĩ	ő	Savannah, Ga	1	3	0	4	
New Haven, Conn	1	3	õ	ő	St. Petersburg, Fla	0	4	1	2	
Providence, R.I	4	3	1	4	Tampa, Fla	4	6	1	4	
Somerville, Mass	2	5	1	1	Washington, D.C	3	5	1	5	
Springfield, Mass	6	8	7.	5	Wilmington, Del	8	11	7	7	
Waterbury, Conn	0	õ	0	õ	international period	2	4	0	0	
Worcester, Mass	7	2	ŏ	2	EAST SOUTH CENTRAL:		TANTH	and set	100.00	
second prior in a second second like	A. 191.	-	, in the second s	~	Birmingham, Ala					
LDDLE ATLANTIC:					Chattanooga, Tenn	1	3	3	1	
Albany, N.Y	3	1 1	3	0	Knoxville, Tenn	3	4	4	2	
Allentown, Pa	2	0	1	2	Louisville, Ky	0	0	1	1	
Buffalo, N.Y	9	3	3	6*	Memphis, Tenn	7	15	1	7	
Camden, N.J	5	1	2	1	Mobile, Ala	2	3	4	6	
Elizabeth, N.J	3	3	0	1	Montgomery, Ala	1	0	0	0	
Erie, Pa	5	ō	4	3		0	2	0	3	
Jersey City, N.J.	8	13	5	4	Nashville, Tenn	2	4	2	6	
Newark, N.J.	4	4	2		WEST SOUTH CENTRAL:		9.9			
New York City, N.Y	132	83	54	1				1.000		
Paterson, N.J.	5	4		59	Austin, Tex	3	3	9	6	
Philadelphia, Pa	3	13	3 28	3	Baton Rouge, La	1	3	2	1	
Pittsburgh, Pa	1	244 Bits		22	Corpus Christi, Tex	4	1	2	0	
Reading, Pa	5	4	5	6	Dallas, Tex	0	0	2	4	
			4	0	El Paso, Tex	2	1	0	4	
Rochester, N.Y	8	11	8	13	Fort Worth, Tex	1	2	4	1	
Schenectady, N.Y	0	0	0	0	Houston, Tex	0	5	6	3	
Scranton, Pa	2	2	0	2	Little Rock, Ark	1	5	4	5	
Syracuse, N.Y.	1	2	2	4	New Orleans, La	4	3	11	11	
Trenton, N.J.	2	8	4	2	Oklahoma City, Okla	0	1	1	1	
Utica, N.Y	3	3	3	1	San Antonio, Tex	4	7	4	2	
Yonkers, N.Y	2	3	2	1	Shreveport, La	7	4	4	4	
ST NORTH CENTRAL:		1.	-	1.00	Tulsa, Okla	0	3	0	2	
Akron, Ohio	0				MOUNTAIN:			dite		
Canton, Ohio	2	1	0	1	Albuquerque, N. Mex					
Chicago, Ill		4	2	0	Colorado Springs, Colo	4	1	2	0	
Cincinnati, Ohio	28	22	26	28	Denver, Colo	2	3	2	2	
Cleveland, Ohio	4	5	3	2	Ogden, Utah	4	8	6	9	
Columbus, Ohio	4	4	2	2		0	4	0	1	
Dayton, Ohio	2	2	1	3	Phoenix, Ariz	2	6	6	5	
Detroit, Mich	1	0	1	0	Pueblo, Colo	1	1	0	0	
Evansville, Ind	9	5	7	15	Salt Lake City, Utah	0	2	1	0	
	1	0	3	2	Tucson, Ariz	0	1	2	3	
Flint, Mich	3	5	2	2	PACIFIC		1.1		1.00	
Fort Wayne, Ind	3	1	3	1	PACIFIC: Berkeley Calif	0				
Gary, Ind	2	5	1	1	Berkeley, Calif	0	1	0	2	
Grand Rapids, Mich	1	0	7	1	Fresno, Calif	7	1	1	0	
Indianapolis, Ind	2	5	3	4	Glendale, Calif	1	1	0	0	
Madison, Wis	0	0	0	0	Honolulu, Hawaii	0	2	1	1	
Milwaukee, Wis	3	3	3	1	Long Beach, Calif	2	2	0	0	
Peoria, Ill.	0	0	0	0	Los Angeles, Calif	7	13	15	13	
Rockford, Ill	7	5	1	0	Oakland, Calif	0	5	0	2	
South Bend, Ind	2	4	1	3	Pasadena, Calif	0	0	0	1	
Toledo, Ohio	1	5	6	1	Portland, Oreg	2	0	2	0	
Youngstown, Ohio	1	0	0	1	Sacramento, Calif	0	1	1	1	
ST NODTH CENTRAL		1			San Diego, Calif	2	0	2	3	
ST NORTH CENTRAL:			S		San Francisco, Calif	2	6	3	3	
Des Moines, Iowa	0	2	1	1	San Jose, Calif	0	1	9	0	
Duluth, Minn	0	0	0	0	Seattle, Wash	4	5	3	0	
Kansas City, Kans	3	3	1	8	Spokane, Wash	0	0	0	2	
Kansas City, Mo	1	5	1	1	Tacoma, Wash	1	2	1	0	
Lincoln, Nebr	0	3	1	3				-		
Minneapolis, Minn	3	4	2	0	San Juan, P.R	1	2	1	2	
Omaha, Nebr	1	1	0	1						
St. Louis, Mo	6	10	4	8						
		1			Comment That March 17 has for	- 100 C	lasted (11 4 2		
St. Paul, Minn	4	7	1	2	^O Current Week Mortality fo	r 108 5	erected (ities		

*Estimate - based on average percent of divisional total. Totals for previous weeks include reported corrections.

388 732

6,143

is maintained at CDC under the direction of the Chief, Microbiology Section, CDC. Blood smears may be submitted either with the Surveillance Form or may be mailed directly to the National Malaria Depository, Communicable Disease Center, Atlanta, Georgia, 30333).

REPORTED CASES OF MALARIA - UNITED STATES - 1962

	Total	No.	No.	Fo	Indig		
State*		Confirmed		Military	Civil.	Visit.	
Alabama	2	2	: H 10	1	1	1	-
Arizona	2	1	1		-		2
California	8	6	2	2	3	2	1
Colorado	1	1	2		1.4	1	- 1
Connecticut	4	3	1	the last of	3	-	-
Dist. of Cal.	1	1	- 14 III		1	- 1	-
Florida	8	7	1	3	3	2	-
Georgia	2	2	- 16	2		121	-
Hawaii	14	6	8	14	-		_
Illinois	5	4	1	114-11	2	50 - T	3
Indiana	1		1	1	1.21	-	-
Kansas	1	1	- i= 10		-	1	
Kentucky	3	3		3	-	-	-
Maryland	2	1	1	1	1	-	
Massachusetts	: 3	3	- 14 DP	3	-	-	
Michigan	1	1	-		1	- 100	-
Minnesota	3	3		1	1	1	-
Mississippi	1	1		1	11-11		-
Missouri	4	4		3	1	1.00	_
Nebraska	1	1				1	- N.
New Hampshir	e 2	2		- 1	1.0	-	1
New Jersey	1	1	-		1	-	-
New Mexico	2	2	- L 10	2	2 - 1	-	
New York	6	6		100	3	2	1
North Caroling	32	31	1	32	1 L I	1.1	
North Dakota	1	1	1	_	1	-	-
Oregon	1	1			_	1	
Pennsylvania	1	1	_	1	1.1	_	
Rhode Island	1			1	1	3.2	
Tennessee	2		2	_	1	-	1
Virginia	3	3	- E	3	1.0	1.1	1.
TOTALS	119	100	19	75	24	11	9

INTERNATIONAL NOTES - QUARANTINE MEASURES

Smallpox - Poland

A total of 17 cases of smallpox, including 2 deaths and one imported case, have been reported to date in Wroclaw, Poland.





Official Business

POSTAGE AND FEES PAID U. S. DEPARTMENT OF H. E. W