



HHS Public Access

Author manuscript

J Acad Nutr Diet. Author manuscript; available in PMC 2018 February 01.

Published in final edited form as:

J Acad Nutr Diet. 2017 February ; 117(2): 214–227. doi:10.1016/j.jand.2016.07.021.

A comprehensive assessment of food parenting practices: development and psychometric testing of HomeSTEAD's family food practices survey

Amber E Vaughn, MPH, RD [Associate Director],

Children's Healthy Weight Research Group, Center for Health Promotion and Disease Prevention, University of North Carolina at Chapel Hill, 1700 Martin L. King Jr. Blvd., CB 7426, Chapel Hill, NC 27599-7426, telephone: (919) 843-0900, fax: (919) 966-6462

Tracy Dearth-Wesley, PhD [Researcher],

University of North Carolina at Chapel Hill, 1700 Martin L. King Jr. Blvd., CB 7426, Chapel Hill, NC 27599-7426, telephone/fax: (919) 338-0791

Rachel G. Tabak, PhD, RD [Research Assistant Professor],

Prevention Research Center in St. Louis, Brown School, Washington University in St. Louis, One Brookings Drive, CB 1196, St. Louis, MO 63130, telephone: (314) 935-0153, fax: (314) 935-0150

Maria Bryant, PhD [Senior Research Fellow and Diet & National Institute for Health Research (NIHR) Career Development Fellow, NIHR Research Design Service YH Advisor, Lead for Diet & Obesity], and

Institute of Clinical Trials Research, University of Leeds, Leeds, LS2 9JT, telephone: +44 (0) 113-343-7632, fax: +44 (0) 113-343-1471

Dianne S. Ward, EdD [Professor]

Department of Nutrition, Gillings School of Global Public Health, and Fellow, Center for Health Promotion and Disease Prevention, University of North Carolina at Chapel Hill, 2202 McGavran-Greenberg Hall, CB 7461, Chapel Hill, NC 27599-7461, telephone: (919) 843-0901, fax: (919) 966-6462

Abstract

Background—Parents' food parenting practices have a significant influence on children's dietary intake and risk for obesity and chronic disease. Understanding the impact and interactions between parents' practices and children's behavior is limited by a lack of development and psychometric testing and/or limited scope of current measures. HomeSTEAD (Home Self-administered Tool for Environmental assessment of Activity and Diet) was created to address this gap.

Correspondence to: Amber E Vaughn.

Conflict of Interest: No authors have any conflicts of interest to report.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Objective—This paper describes development and psychometric testing of HomeSTEAD's family food practices survey.

Participants/Design—Between August 2010 and May 2011, a convenience sample of 129 parents of children ages 3-12 years were recruited from central North Carolina and completed the self-administered HomeSTEAD survey on three occasions during a 12 to 18-day window. Demographics and child diet were assessed at Time 1. Child height and weight were measured during the in-home observations (following Time 1 survey).

Statistical analysis—Exploratory factor analysis (EFA) with Time 1 data was used to identify potential scales. Scales with more than three items were examined for scale reduction. Following, mean scores were calculated at each time points. Construct validity was assessed by examining Spearman rank correlations between mean scores (Time 1) and children's diet (fruits and vegetables, sugar-sweetened beverages, snacks, sweets) and BMI z-scores. Repeated measures analysis of variance was used to examine differences in mean scores between time points; and single-measure intraclass correlations (ICC) were calculated to examine test-retest reliability between time points.

Results—EFA identified 24 factors and retained 124 items; however, scale reduction narrowed items to 86. The final instrument captures five Coercive Control practices (16 items), seven Autonomy Support practices (24 items), and 12 Structure practices (46 items). All scales demonstrated good internal reliability ($\alpha > 0.62$), 18 factors demonstrated construct validity (significant association with child diet, $p < 0.05$), and 22 demonstrated good reliability (ICC > 0.61).

Conclusions—HomeSTEAD's family food practices survey provides a brief, yet comprehensive and psychometrically sound assessment of food parenting practices.

Keywords

children; parents; feeding practices; measurement; psychometric properties

Introduction

Dietary guidelines offer advice for a healthful diet — one that provides adequate nutrition, promotes a healthy weight, and prevents chronic disease.¹ Unfortunately, data from many countries around the world (Australia,² Europe,³ and North America^{4,5}) have demonstrated that children's eating patterns fail to meet these recommendations. For example, few children in the US consume recommended intakes of whole grains (<1%), vegetables (7%), fruit (29%), and milk (37%), and most exceed recommended limits for solid fats (97%) and added sugars (90%).⁴

Parents play an important role in children's socialization, including the norms and habits they adopt with regards to food and eating.⁶ Parents' behaviors shape the physical and social environment in which their children grow up, influencing their home environment as well as their children's interactions with the outside world. “Food parenting practices” refers to the behaviors or actions (intentional or unintentional) performed by parents for child rearing purposes that influence their children's attitudes, behaviors, or beliefs around food and eating.⁷ Literature in this area suggests that parent practices such as making healthy foods

more available, modeling healthy eating, and providing encouragement to eat healthy foods help promote children's consumption of those healthy foods.^{7,8} In comparison, practices such as restriction, pressure, and food bribes may inadvertently promote increased intake of unhealthy foods.⁹⁻¹⁴

One of the barriers to fully understanding how the home environment, and specifically food parenting practices, influences children's dietary intake is availability of appropriate measures.¹⁵ While there are many measures available, they are often limited in scope and have not undergone a comprehensive process of development.¹⁵ Development often lacks clear conceptualization of what the instrument is designed to measure, fails to use systematic or informed approaches to selecting and refining items, and includes incomplete reliability and validity testing.¹⁵ In order to advance our understanding of how the home environment influences children's dietary intake and eating behaviors, the field needs a comprehensive measure of food parenting practices with items that have undergone a rigorous development process including reliability and validity testing.

HomeSTEAD, the Home Self-administered Tool for Environmental assessment of Activity and Diet, is a newly developed instrument designed to address this gap in measurement by providing a comprehensive evaluation of home environmental factors thought to influence children's diet and physical activity.¹⁶ HomeSTEAD builds upon our previous work to develop the Healthy Home Survey,¹⁷ an assessment of the home environment related to children's eating and activity. While the Healthy Home Survey provided a useful pilot measure, a more expanded instrument was needed.

For HomeSTEAD, two frameworks were adopted to help guide identification of relevant constructs and ensure comprehensive coverage. The Analysis Grid for Environments Linked to Obesity (ANGELO) framework¹⁸ and the Model of the Home Food Environment Pertaining to Child Obesity¹⁹ both recognize multiple spheres of influence — physical, political, socio-cultural, economic — that influence child weight and weight-related behaviors. When considering their application to the home environment, it resulted in the development of a four-part instrument: a home food inventory (physical food environment), a family food practices survey (social food environment), a home physical activity and media equipment inventory (physical environment around physical activity), and a family physical activity and screen time practices survey (social environment around physical activity).¹⁶ The development of this new instrument, specifically the component for the social environment around food, afforded the opportunity to develop a comprehensive assessment of food parenting practices.

The purpose of this paper is to describe the development and psychometric testing of the scales related to the home's social food environment. This has been conceptualized primarily as food parenting practices.

Methods

Methods used to develop the HomeSTEAD tool are described in detail elsewhere, but the aspects most relevant to the development of its family food practices survey are provided

below.¹⁶ All protocols were approved by the Institutional Review Board at the University of North Carolina at Chapel Hill (09-1177), and all participants provided written informed consent.

HomeSTEAD Instrument Development

HomeSTEAD's family food practices survey was developed using a mixed methods approach, which began with identifying a theoretical framework and conducting a systematic review of the literature. Following the application of the ANGELO framework,^{18,19} a systematic review was conducted to identify current measures of food parenting practices.¹⁵ The review led to the refinement of constructs resulting in a content map of food parenting practices, which has recently been published.⁷ Concurrently, items and scales from existing measures identified in this review were cataloged into a database and categorized according to the content map. When existing items were available, the research team reviewed sets of similar items and selected those that were deemed to be the most relevant for that construct. When existing items were not available, the research team developed new items. Where possible, response options were standardized across sections of the HomeSTEAD survey. For example, food parenting-related items generally used 5-point likert-type response scales (e.g., 1 = never, 2 = rarely, 3 = sometimes, 4 = often, 5 = always; or 1 = strongly disagree, 2 = disagree, 3 = neutral, 4 = agree, 5 = strongly agree).

Content validity of this initial collection of items was assessed with the help of two expert reviewers. Experts were asked to provide feedback and suggestions related to content coverage, item relevance and intention, and question format and clarity. The instrument was refined based on this feedback.

Then, one-on-one guided cognitive interviews were conducted with parents of 3–12 years old children to examine clarity and comprehension of items. Participants for the cognitive interviews were recruited through newspaper advertisements, listserv notifications, and community postings. Each cognitive interview focused on just one of the four sections in HomeSTEAD, which reduced participant burden and limited interviews to 45 minutes or less. For the family food practices section, the first round of cognitive interviews included 11 parents. Content analysis of these interviews allowed problematic items to be identified, discussed by the team, and revised. Revised items underwent a second round of cognitive interviews with five additional parents, at which time items were deemed acceptable (no remaining issues regarding clarity or interpretation of items). Participants received \$15 as an incentive for participation.

At the end of this stage, HomeSTEAD included 214 items dedicated to assessment of food parenting practices.

Survey Administration

A convenience sample of 129 families with at least one child between the ages of 3 and 12 years old were recruited for instrument testing (August 2010–April 2011). Recruitment strategies employed were similar to those described above for cognitive interviews. The sample size was based on power calculations that would ensure adequate reliability evidence (assuming a kappa of 0.60, 80% trait prevalence, alpha of 0.05, and beta of 0.80).²⁰ While a

larger sample size would be desired for the exploratory factor analysis part of instrument testing,²¹ pragmatic constraints (e.g., funding, resources) prohibited it. When participants had more than one child in the target age group, one child was selected to serve as the “reference” child. Initially, selection was random. However, to ensure that there was an even distribution of children across the 3-12 year old age range, older children were often selected as the reference child for participants recruited later. In addition to having at least one child in the target age group, eligibility criteria also specified that the reference child could not have physical limitations that would impact their diet or physical activity behavior (e.g., extreme food allergies, physical disability), the family must reside in central North Carolina with no plans to move during the study period, and that the parent be able to speak English. Participants were offered up to \$100 for completing the full protocol.

Participants completed all four sections of the self-administered HomeSTEAD survey at three separate time points over a period of 12 to 18 days and allowed research staff to conduct an in-home observational assessment (see Figure 1 for timeline). The Time 1 HomeSTEAD survey was mailed to participants along with a demographic survey, a child diet screener,²² and a consent form. The survey instructions reminded parents to keep the reference child in mind when responding to questions and to select responses that best reflected what they actually do, think, and feel. Two or three days later, staff conducted a scheduled home visit. During these home visits, staff collected signed consent and all Time 1 surveys, completed an observational assessment of the home's physical environment (e.g., foods and beverages present in the home), and measured children's height and weight. Height was measured to the nearest 1/8 inch using a Shorr or Seca stadiometer (Shorr Productions, Olney, MD; Seca Corporation, Columbia, MD); weight was measured to the nearest 0.1 pounds using a Seca portable electronic scale (model 770 or 874, Seca Corporation, Columbia, MD). All measures were taken while participants were in light clothing and no shoes. Participants received the Time 2 HomeSTEAD survey at the conclusion of the home visit with instructions to return the survey via mail within 24 hours. Approximately 10 days later, participants were mailed the Time 3 HomeSTEAD survey with instructions to complete and return the survey within 4 days. If the third survey was not completed and returned within an additional 10 days (even after repeated reminder calls), that participant's data were not included in the analysis. All data were collected between August 2010 and May 2011.

Statistical Analysis

Using Time 1 data, responses to individual items were reviewed to evaluate missingness and variability in responses. Items were flagged if 80% or more of responses fell within two response options, or 75% or more of responses fell within one response option, indicating low variability and questioning the item's usefulness in distinguishing unique participant characteristics. Additionally, items were flagged if correlations between items were 0.75 or higher given that high correlations can suggest duplication of the concepts being assessed.

Following this preliminary review of the data, exploratory factor analyses (EFA) were used to identify potential food parenting practice scales. All EFAs were performed in MPlus Version 7.3²³ using Time 1 data. Given the limited sample size (n= 129) and the large

number of items (n=214), preliminary sorting of items was necessary. An emerging content map of food parenting practices (noted earlier)⁷ guided the sorting of items into three broad categories: Coercive Control, Autonomy Support, and Structure. Separate EFAs were performed on items from each broad category; however, some items where categorization was less clear were examined in more than one category. A weighted least squares minimum variance estimator and geomin rotated solution were used. Factor solutions were evaluated based on eigenvalues, scree test, and interpretability criteria (e.g., comparative fit index, root mean square of approximation).^{21,24} Items with low factor loadings (<0.40) were eliminated individually (item with lowest factor loading eliminated first); the EFA was then repeated. If an item cross-loaded (>0.40 on multiple factors), the item was included in the factor with the higher loading. Items that has been flagged as having high correlations (>0.75) were considered throughout this process of refining which items would be eliminated or retained. Once factors were identified, a composite score for each scale was calculated by averaging scores from its individual items, including reverse coding when necessary.

Scales with more than three items were examined for possible scale reduction. For each scale, multiple versions of reduced scales were examined, and several criteria were considered when deciding which items to include in the final reduced scale. Criteria included factor loadings of individual items (giving preference to items with higher loadings), comparability of the reduced scale's internal consistency to that of the original scale (giving preference to those maintaining a Cronbach's alpha >0.7),²⁵ and observed correlations between the full versus reduced scales and child dietary outcomes (with higher correlations suggesting greater construct validity). The latter was assessed using Spearman rank correlations between the reduced scales' composite scores and children's dietary outcomes (weekly intake of fruits and vegetables, sugar-sweetened beverages, snacks, and sweets intake from parent-completed screener of child diet) and BMI z-scores.

Following the scale reduction process, mean scores (SD) for the final reduced scales were determined from data taken at each time point (Time 1, 2 and 3). Mean differences were tested using repeated measures analysis of variance (ANOVA); single-measure intraclass correlations (ICC) were calculated to examine test-retest reliability. The single-measure ICC, ICC(1,1) from Shrout and Fleiss,²⁶ estimates reliability given a single random administration. ICCs of 0.61-0.80 indicate moderate agreement, and ICCs of 0.81-1.00 indicate substantial agreement.²⁷ Correlations, ANOVA, and ICC analyses were conducted using SAS[®] software, Version 9.4 of the SAS System for Windows (SAS Institute, Cary, NC, released July 2013).

Results

Parents in the study sample (n= 129) were predominately mothers (91%). The sample included a mix of racial and income groups. The majority was white (71%) or African American (25%), but very few were Hispanic or Latino (3%). The majority had a household income above the area's median (68% with annual household income \$50,000) and were well educated (79% college degree or higher). Children in the sample included similar numbers of boys and girls (51% vs 49%, respectively), were on average 7.1 ± 2.9 years old, and had a BMI percentile of 59.6 ± 27.1 . An additional 19 parents (separate from the 129

already described) completed the screening process but did not end up participating due to ineligibility (child too young (3), moving/distance from project office (2)), scheduling conflicts (11), or participation in other studies that would be disrupted by participation in this study (3). Compliance with study protocols was high with 125 parents (97%) completing the all three self-administrations and the home observation. Participants also completed the multiple self-administrations in a timely manner. On average, there were 3.9 ± 3.7 days between Time 1 and Time 2 self-administrations and 12.4 ± 5.6 days between Time 2 and Time 3 self-administrations.

The initial EFA identified 24 factors and retained 124 of the 214 items, including five factors (28 items) within Coercive Control, seven factors (34 items) within Autonomy Support, and 12 factors (62 items) within Structure. Nineteen of the 24 factors had greater than three items per factor, and therefore underwent examination for scale reduction. Following the scale reduction process, Coercive Control scales were reduced from 28 to 16 items; Autonomy Support scales were narrowed from 34 to 24 items; and Structure scales were trimmed down from 62 to 46 items; resulting in a final instrument with 86 items.

Coercive Control Scales

The Coercive Control scales included restriction, soothing with food, threats and bribes, “clean plate” policy, and pressure to eat. All scales had 3-4 items and acceptable internal consistency (Cronbach's α all 0.62). Table 1 provides the items, factor loadings, and Cronbach's alphas for each of these five scales. Composite scores for each of these scales are calculated by averaging the individual items (i.e., Likert responses) within each scale. For all Coercive Control scales, higher scores reflect greater use of those practices.

Autonomy Support Scales

Autonomy Support scales included encouragement, reasoning, praise, nutrition education, and guided choices around *when*, *what*, and the *amount* of food eaten. Each of the scales was reduced to 3-4 items. The final reduced scales had acceptable internal consistency (Cronbach's α all 0.66). Table 2 provides the items, factor loadings, and Cronbach's alphas for each of these seven scales.

Similar to the Coercive Control scales, composite scores are calculated by averaging the individual items, taking into consideration reverse scoring where noted. For most scales, higher scores reflect greater use of that specific practice. The practice of guided choices presents a challenge for this standard approach to measurement and interpretation. Guided choices represents a balance between parent and child control over when, what, and how much the child eats. For example, the “parent allows the child a choice in what he/she eats, but options from which the child must choose are determined by the parent.”⁷ Assessing this construct often requires measuring the two extreme behaviors – parents making decisions unilaterally vs. children having unrestricted choice in their eating. Currently, higher scores reflect greater child choice and control and lower scores reflect greater parent control. Scores closer to the middle may be a better reflection of this balance.

Structure Scales

Lastly, the Structure scales captured a broad array of parenting practices that build eating competence in children. These factors included monitoring of unhealthy foods, rules and limits around unhealthy foods, components of the child's eating environment (i.e., meal setting, family eating, atmosphere of meals, eating area, and distractions), planning and preparation of healthy meals, attractive presentation of healthy foods, availability and accessibility of healthy foods, modeling, and weight talk. Some of the final scales in this category were slightly longer (up to 6 items). Again, all had acceptable internal consistency (Cronbach's α all 0.62). Table 3 provides the items, factor loadings, and Cronbach's alphas for each of these 12 scales. One deviation in the standard factor analysis was the movement of an item (i.e., Do you have fruits and vegetables that your child likes available at home?) from planning and preparation of healthy meals (where its factor loading was 0.60) into availability and access (where its factor loading was 0.32). Movement of this item allowed both scales to be more consistent with how these practices have been conceptualized in the feeding literature.

As in other sections, composite scores for each of these scales are calculated by averaging the individual items, taking into consideration reverse scoring where noted. Higher scores generally reflect greater use of practices that provide positive structure. Two clarifications to note are the scoring for distractions and weight talk. For distractions, a higher score reflects greater presence of distractions during meals, which in turn reflects a lack of structure. Also, the factor analysis and scale reduction of the distractions scale also suggested that differentiating between weekday and weekend day practices may be unnecessary. While original items asked about weekday and weekend day practices separately, it is likely sufficient to ask about the practices across a typical week. In the current analysis, there was a high correlation between weekday and weekend versions of the item so only the weekday version was retained. Similar findings were also observed in the family eating scale. With regard to the weight talk scale, higher scores reflect greater discussion of weight and dieting by family members. Such practices have been shown to be associated with disordered eating.^{28,29} Therefore, for the distractions and weight talk scales, higher scores do not reflect positive structure.

Table 4 provides average composite scores for factors from each of the three sections described.

Reliability

Means (SD) for factor composite scores at Times 1, 2, and 3 and reliability testing (ICC) are shown in Table 4. ANOVA testing did not find any significant differences between the means from the three time points for any of the factors. Of the 24 factors, 22 (92%) had ICC scores greater than 0.61, indicating moderate or better agreement. Four of these 22 factors had ICC scores greater than 0.81, indicating substantial agreement: (1) meal setting, (2) family eating, (3) eating area/physical space, and (4) planning and preparation of healthy meals. The other two scales had ICC scores just slightly below 0.61 (soothing with food ICC = 0.60, guided choices: what food is eaten ICC 0.57).

Construct Validity

Correlations between factor composite scores and parent-reported children's weekly consumption of fruits and vegetables, sugar-sweetened beverages, snacks, and sweets are reported in Table 5. Observed intakes were as follows: 3.3 servings/day (± 1.7) of fruits and vegetables (servings of fruits, but not juices, and vegetables, but not fried potatoes), 0.6 servings/day (± 0.9) of sugar-sweetened beverages (fruit and sport drinks, flavored waters, regular soda), 0.8 servings/day (± 0.8) of snacks (potato chips, tortilla chips, cheese nibs, chex mix, etc.), and 1.1 servings/day (± 0.8) of sweets (candy, breakfast pastries, cookies, brownies, pies, cakes, and ice cream). Correlations between all factor composite scores and child BMI z-scores were also examined but were not significant (results not shown).

Three of the five Coercive Control scales were significantly associated with aspects of child dietary intake. These associations were in the predicted direction with more coercive controlling practices being associated with lower intakes of healthy foods and higher intakes of unhealthy foods. Specifically, more frequent parental restriction was associated with decreased consumption of fruits and vegetables ($r = -0.27$, $p < 0.01$) and increased consumption of sugar-sweetened beverages ($r = 0.49$, $p < 0.01$). Additionally, coercive control practices related to "clean plate" policies and pressure to eat were significantly associated with increased consumption of snacks ($r = 0.23$ and 0.22 , respectively, $p < 0.05$).

Most scales within the Autonomy Support category were significantly associated with child intake of fruits and vegetables, sweetened beverages, sweets, or snacks. In addition, observed associations were generally in the predicted directions, with autonomy supporting practices being associated with greater intakes of healthy foods and lower intakes of unhealthy ones. Specifically, more frequent encouragement and nutrition education were associated with increased consumption of fruits and vegetables ($r = 0.38$ and 0.36 , respectively, $p < 0.01$). Greater child vs. parent choice over the *amount* of food eaten was significantly associated with decreased consumption of sugar-sweetened beverages ($r = -0.28$, $p < 0.01$) and snacks ($r = -0.37$, $p < 0.01$). More frequent nutrition education was significantly associated with decreased consumption of sweets ($r = -0.20$, $p < 0.05$). However, a few unexpected correlations were observed between praise and guided choices over *what* the child eats and consumption of snacks.

Ten of the 12 Structure scales were significantly associated with various aspects of child intake. Once again, associations were in the predicted direction with greater structure being associated with higher intakes of healthy foods and lower intakes of unhealthy ones. Specifically, most of the Structure practices were significantly associated with increased consumption of fruits and vegetables ($r = 0.23$ to 0.50 , $p < .05$) and decreased consumption of sugar-sweetened beverages ($r = -0.18$ to -0.42 , $p < .05$). In addition, greater rules and limits around unhealthy foods, planning and preparation of healthy meals, and modeling were associated with decreased consumption of sweets and snacks ($r = -0.21$ to -0.31 , $p < .05$). Increased consumption of sweets and snacks was associated with more frequent use of TV during meals (i.e., distractions) ($r = 0.21$ and 0.25 , respectively, $p < .05$). In contrast, distractions or more frequent use of TV during meals was significantly associated with decreased consumption of fruits and vegetables and increased consumption of sugar-sweetened beverages ($r = -0.21$ and 0.39 , respectively, $p < .05$).

Discussion

The development of this instrument represents a significant advancement in the measurement of food parenting practices. Unlike many other measures to date, the instrument provides a comprehensive set of scales to measure food parenting practices created through a rigorous process of development. The Healthy Home Survey represented an initial attempt by this team to produce a comprehensive measure; however, it was guided more by the literature and factors that had already been identified as predictors of child diet (and physical activity) behaviors.¹⁷ For HomeSTEAD, development was guided more by theory, specifically the ANGELO framework. Its development was also informed by a systematic literature review¹⁵ and an emerging content map developed by experts in the field;⁷ keeping this instrument aligned with the most current thinking in the field. The development process also included a review of potential items by an expert panel, cognitive testing with a sample of parents, factor analyses of scales, assessment of construct validity, and evaluation of test-retest reliability. The resulting instrument offers a comprehensive assessment of the variety of food parenting practices that have been identified in the literature using scales with solid psychometric properties. HomeSTEAD's rigorous development process included five of the six key elements that were recommended as part of a recent review of measures of food parenting practices (conceptualization of instrument purpose, development and refinement of item pool, and assessment of reliability and validity).¹⁵

The scope and breadth of HomeSTEAD's family food practices survey will facilitate a greater understanding of how parents' practices influence child diet and weight. Prior to the development of the HomeSTEAD instrument, the most comprehensive measures captured only 10-12 practices.³⁰⁻³³ HomeSTEAD, in comparison, captures 24 practices. Also, there is a nice overlap between the scales assessed and the most recent food parenting practices content map (see Figure 2 for illustrated comparison). Having a comprehensive assessment of food parenting practices will allow future researchers to assess the relative importance of these practices and how the use of practices might interact to influence child eating habits and weight.

Assessment of construct validity (i.e., associations between food parenting practice scales and markers of child diet) provides some initial assessment of the relative importance of these practices. For example, restriction, nutrition education, guided choices, rules and limits, family eating, distractions, planning and preparation of healthy meals, and modeling appear to be more important, while atmosphere, eating area, and weight talk may be less important. Future studies are needed to confirm these associations using more rigorous diet assessment protocols (e.g., 24-hour diet recall, food diary) in larger, more diverse samples with longitudinal data.

The construct validity testing did produce some unexpected results, specifically for praise and guided choices, which may help the field refine how each of these practices are conceptualized. In the literature, verbal praise and encouragement have been shown to be associated with increased intake of healthy foods (e.g., fruits and vegetables) in children.^{8,34-36} However, similar associations were not observed in the current study. In

contrast, a significant positive association was observed between parent's use of praise and children's intake of snacks. These unexpected findings may suggest that the construct of praise, and the scale used to assess it, may be interpreted differently by different parents and would benefit from refinement to focus on praise offered for eating *healthy* foods (e.g., fruits and vegetables, whole grains, and lean proteins) and not just foods offered. Guided choices is a relatively new construct in the feeding literature; however, experts hypothesize that the practice of guided choices should promote healthier intakes in children.⁷ The current study showed mixed results around guided choices. When parents allow children greater choice in *what* foods are eaten, children appear to eat more snacks and sweets. However, when children are allowed greater choice in the *amount* of food eaten, children appear to consume fewer sugar sweetened beverages and snack foods. These findings appear to support Satter's Division of Responsibility in Feeding,³⁷⁻³⁹ which suggests that parents decide what foods are offered and children decide how much of those foods to eat. Therefore, the construct for guided choices may need refinement to emphasize how parents and children share responsibility in food choices – balancing greater parent choice in *what* is eaten with greater child choice over the *amount* eaten. Hence, measurement of guided choices may be improved by merging these two scales in a way that emphasizes this more specific division of responsibility in food choices. These unexpected findings around praise and guided choices offer an opportunity and guidance for advancing the conceptualization of these practices.

A few limitations to this study are important to acknowledge. First, a brief dietary screener that included only a few markers of child dietary intake was used to assess construct validity. Several additional markers of diet quality were not assessed, including intakes of dark green vegetables, whole grains, and lean proteins. Despite the limitations of the screener, the associations that were observed between food parenting practices and the markers of child diet were generally in the hypothesized direction. Our sample size, selected for psychometric evaluation, limited our ability to explore differences across age groups. The sample included only about 40 children in each age group (i.e., 3-4 years, 5-8 years, 9-12 years). Child age likely influences parent use of various food parenting practices and the impact those practices have on child intake (e.g., younger children need greater assistance and support, but older children can build off of basic skills and be more independent). While it was beyond the scope of the current study to explore these age-related differences, the final HomeSTEAD instrument is appropriate for the full age range and will be useful in future studies trying to understand the evolution of parents feeding practices. Our limited sample size also necessitated the presorting of items prior to the EFA. While an emerging content map of food parenting practices was used to guide this process, it did make certain assumptions about grouping of items and influenced the identification of factors. Confirmatory analysis in a larger sample is needed. Another limitation of the sample was the lack of socio-economic diversity. Future research will need to evaluate instrument performance in a more diverse population, including low-income parents. Additionally, the short period of time between administrations might explain in part the high ICCs (and lack of significant differences) observed during the reliability testing. However, only 4 of the 24 factors showed “substantial agreement”, thus illustrating some variability even during this short period of time. Finally, the study lacked evidence for criterion validity and sensitivity

to change. The HomeSTEAD study was designed to capture criterion validity evidence for characteristics of the physical environment using a comparison of the parent survey to an in-home assessment by trained data collectors. While the physical environment was measured (e.g. food availability), this 1-hour home visit provided insufficient opportunity to accurately assess parent food practices. Scheduling of these home visits often avoided meal times (which would have been necessary to observe food parenting practices), and the process of data collection tended to disrupt normal routines. Evidence for sensitivity to change is also lacking as it would require employment of the instrument in an intervention study. Several of the limitations noted were beyond the scope of the current project. These limitations are far outweighed by the numerous study strengths, one of which was the rigorous process used to develop and refine the item pool (incorporating a systematic review of existing measures¹⁵ and an emerging food parenting practices content map⁷). Additionally, identification of the final scales used a multi-step process that began with an EFA, but also worked to simplify and shorten scales to create an efficient assessment instrument. Furthermore, the study allowed for assessment of variation in responses across repeat administrations, test-retest reliability, and construct validity. Results of this extensive testing demonstrated sound psychometric properties and supports the usefulness of the HomeSTEAD family food practices survey in future research.

Conclusions

The HomeSTEAD instrument is intended to provide a comprehensive assessment of environmental factors in the home that influence children's food, physical activity, and screen time behaviors. HomeSTEAD's family food practices survey, the focus of this study, is an integral piece of this larger tool. The scales within this survey align closely with the current literature, integrating up-to-date terminology and definitions for food parenting practices. Psychometric testing demonstrated moderate to high levels of internal and external reliability and construct validity. While HomeSTEAD was designed to be comprehensive, it does not have to be used in its entirety. When used in future research, it will likely need to be customized based on what constructs are identified as most relevant for the study as well as the age of the children involved.

Acknowledgments

Funding Disclosure: Funding for this project was provided by the National Cancer Institute (NCI) at the National Institutes of Health (NIH), grant R21-CA134986. The project was conducted out of the University of North Carolina at Chapel Hill at the Center for Health Promotion and Disease Prevention, a Prevention Research Center funded through a cooperative agreement with the Centers for Disease Control and Prevention (CDC, U48-DP001944). Work was also conducted out of Washington University's Center for Diabetes Translation Research, funded by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK, P30-DK092950). Findings and conclusions in this article are those of the authors and do not necessarily represent the official position of the NIH, NCI, NIDDK, or CDC.

References

1. U.S. Department of Health & Human Services. [Accessed April 20, 2016] U.S. Department of Agriculture. 2015-2020 Dietary Guidelines for Americans. 8th2015. <http://health.gov/dietaryguidelines/2015/guidelines>

2. Golley RK, Hendrie GA, McNaughton SA. Scores on the dietary guideline index for children and adolescents are associated with nutrient intake and socio-economic position but not adiposity. *J Nutr*. 2011; 141:1340–1347. [PubMed: 21613454]
3. Diethelm K, Jankovic N, Moreno LA, et al. Food intake of European adolescents in the light of different food-based dietary guidelines: results of the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study. *Public Health Nutr*. 2012; 15:386–398. [PubMed: 21936969]
4. Kirkpatrick SI, Dodd KW, Reedy J, Krebs-Smith SM. Income and race/ethnicity are associated with adherence to food-based dietary guidance among US adults and children. *J Acad Nutr Diet*. 2012; 112:624–635. [PubMed: 22709767]
5. Reedy J, Krebs-Smith SM. Dietary sources of energy, solid fats, and added sugars among children and adolescents in the United States. *J Am Diet Assoc*. 2010; 110:1477–1484. [PubMed: 20869486]
6. Savage JS, Fisher JO, Birch LL. Parental influence on eating behavior: conception to adolescence. 2007; 35:22–34.
7. Vaughn AE, Ward DS, Fisher JO, et al. Fundamental constructs in food parenting practices: A content map to guide future research. *Nutr Rev*. 2016
8. Pearson N, Biddle SJ, Gorely T. Family correlates of fruit and vegetable consumption in children and adolescents: a systematic review. *Public Health Nutr*. 2009; 12:267–283. [PubMed: 18559129]
9. Ogden J, Cordey P, Cutler L, Thomas H. Parental restriction and children's diets: the chocolate coin and Easter egg experiments. *Appetite*. 2013; 61:36–44. [PubMed: 23142562]
10. Jansen E, Mulkens S, Emond Y, Jansen A. From the Garden of Eden to the land of plenty: restriction of fruit and sweets intake leads to increased fruit and sweets consumption in children. *Appetite*. 2008; 51:570–575. [PubMed: 18501474]
11. Jansen PW, Roza SJ, Jaddoe VW, et al. Children's eating behavior, feeding practices of parents and weight problems in early childhood: results from the population-based Generation R Study. *Int J Behav Nutr Phys Act*. 2012; 9:130. [PubMed: 23110748]
12. Batsell WR, Brown AS, Ansfield ME, Paschall GY. “You will eat all of that!” A retrospective analysis of forced consumption episodes. *Appetite*. 1998; 38:211–219.
13. Sleddens EF, Kremers SP, De Vries NK, Thijs C. Relationship between parental feeding styles and eating behaviours of Dutch children aged 6-7. *Appetite*. 2010; 54:30–36. [PubMed: 19747513]
14. Rodenburg G, Kremers SP, Oenema A, van de Mheen D. Associations of parental feeding styles with child snacking behaviour and weight in the context of general parenting. *Public Health Nutr*. 2014; 17:960–969. [PubMed: 23527513]
15. Vaughn AE, Tabak RG, Bryant MJ, Ward DS. Measuring parent food practices: a systematic review of existing measures and examination of instruments. *Int J Behav Nutr Phys Act*. 2013; 10:61. [PubMed: 23688157]
16. Hales D, Vaughn AE, Mazzucca S, et al. Development of HomeSTEAD's physical activity and screen time physical environment inventory. *Int J Behav Nutr Phys Act*. 2013; 10:132. [PubMed: 24313962]
17. Bryant MJ, Ward DS, Hales D, Vaughn A, Tabak RG, Stevens J. Reliability and validity of the Healthy Home Survey: a tool to measure factors within homes hypothesized to relate to overweight in children. *Int J Behav Nutr Phys Act*. 2008; 5:23. [PubMed: 18442392]
18. Swinburn B, Egger G, Raza F. Dissecting obesogenic environments: the development and application of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med*. 1999; 29:563–570. [PubMed: 10600438]
19. Rosenkranz RR, Dziewaltowski DA. Model of the home food environment pertaining to childhood obesity. *Nutr Rev*. 2008; 66:123–140. [PubMed: 18289177]
20. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977; 33:159–174. [PubMed: 843571]
21. Costello AB, Osborne J. Best practices in exploratory factor analysis: Four recommendations for getting the most from your analysis. *Practical Assessment, Research & Evaluation*. 2005; 10
22. Neuhouser ML, Lilley S, Lund A, Johnson DB. Development and validation of a beverage and snack questionnaire for use in evaluation of school nutrition policies. *J Am Diet Assoc*. 2009; 109:1587–1592. [PubMed: 19699839]
23. Muthén, LK., Muthén, BO. *Mplus User's Guide*. Los Angeles, CA: 1998-2011.

24. Suhr, DD. [Accessed on September 4, 2014] Exploratory or Confirmatory Factor Analysis?. Statistics and Data Analysis (paper 200-31). 2006. <http://www2.sas.com/proceedings/sugi31/200-31.pdf>
25. Bland JM, Altman DG. Cronbach's alpha. *Bmj*. 1997; 314:572. [PubMed: 9055718]
26. Shrout PE, Fleiss JL. Intraclass correlations: uses in assessing rater reliability. 1979; 86:420–428.
27. Shrout PE. Measurement reliability and agreement in psychiatry. 1998; 7:301–317.
28. Berge JM, Maclehose R, Loth KA, Eisenberg M, Bucchianeri MM, Neumark-Sztainer D. Parent conversations about healthful eating and weight: associations with adolescent disordered eating behaviors. *JAMA Pediatr*. 2013; 167:746–753. [PubMed: 23797808]
29. Neumark-Sztainer D, Bauer KW, Friend S, Hannan PJ, Story M, Berge JM. Family weight talk and dieting: how much do they matter for body dissatisfaction and disordered eating behaviors in adolescent girls? *J Adolesc Health*. 2010; 47:270–276. [PubMed: 20708566]
30. Koivisto UK, Sjoden PO. Reasons for rejection of food items in Swedish families with children aged 2-17. *Appetite*. 1996; 26:89–103. [PubMed: 8660035]
31. Vereecken CA, Van Damme W, Maes L. Measuring attitudes, self-efficacy, and social and environmental influences on fruit and vegetable consumption of 11-and 12-year-old children: Reliability and validity. *J Am Diet Assoc*. 2005; 105:257–261. [PubMed: 15668685]
32. Musher-Eizenman D, Holub S. Comprehensive Feeding Practices Questionnaire: validation of a new measure of parental feeding practices. *J Pediatr Psychol*. 2007; 32:960–972. [PubMed: 17535817]
33. Zeinstra GG, Koelen MA, Kok FJ, van der Laan N, de Graaf C. Parental child-feeding strategies in relation to Dutch children's fruit and vegetable intake. *Public Health Nutr*. 2010; 13:787–796. [PubMed: 19772690]
34. McGowan L, Croker H, Wardle J, Cooke LJ. Environmental and individual determinants of core and non-core food and drink intake in preschool-aged children in the United Kingdom. *Eur J Clin Nutr*. 2012; 66:322–328. [PubMed: 22252108]
35. Berge JM. A review of familial correlates of child and adolescent obesity: what has the 21st century taught us so far? *Int J Adolesc Med Health*. 2009; 21:457–483. [PubMed: 20306760]
36. Vereecken CA, Keukelier E, Maes L. Influence of mother's educational level on food parenting practices and food habits of young children. *Appetite*. 2004; 43:93–103. [PubMed: 15262022]
37. Satter EM. The feeding relationship. *J Am Diet Assoc*. 1986; 86:352–356. [PubMed: 3950279]
38. Satter E. The feeding relationship: problems and interventions. *J Pediatr*. 1990; 117:S181–189. [PubMed: 2199651]
39. Satter E. Feeding dynamics: helping children to eat well. *J Pediatr Health Care*. 1995; 9:178–184. [PubMed: 7629684]

Practice Implications

What is the current knowledge on this topic?

- There is great interest in how parent feeding practices impact child diet and weight; however, current measures assess only select practices and conceptualize these practices differently.

How does this research add to the knowledge on this topic?

- HomeSTEAD's family food practices survey provides a comprehensive assessment tool that can be used to evaluate parents' use of 24 different feeding practices.

How might this knowledge impact current dietetics practice?

- The family food practices survey is a valuable tool for research dietitians. Its further evaluation in larger and more diverse samples will help inform feeding practices to promote.





Figure 1. Timeline for HomeSTEAD data collection

Note: Reprinted from “Development of HomeSTEAD's physical activity and screen time physical environment inventory,” by Hales D et al., 2013, *International Journal of Behavioral Nutrition and Physical Activity*, volume 10, page 132. Reprinted with permission.



Black boxes with white text indicate where constructs were added or moved into a different section. Gray boxes indicate constructs that were not captured in this survey.

Figure 2.

Comparison of the food parenting practices content map to scales measured by HomeSTEAD's family food practices survey.

Note: Adapted from "Fundamental constructs in food parenting practices: A content map to guide future research," by Vaughn AE et al., 2016, *Nutrition Reviews*, in press. Adapted with permission.

Table 1
HomeSTEAD's^a Coercive Control scales with item wording, factor loadings, and internal consistency (Cronbach's α)

Scale Name and Items	Factor Loading	Internal consistency
Restriction		0.62
How often do you restrict (or try to restrict) your child's food intake so that s/he will not gain weight? ^b	0.74	
How often do you tell your child not to eat something because it will make him/her fat? ^b	0.76	
How many times have you told your child to eat less food or eat different foods in order to lose weight or to keep from gaining weight? ^b	0.78	
Soothing with food		0.81
I give my child something to eat or drink when s/he is bored or worried, even if I know s/he is not hungry. ^c	0.79	
Offering my child something to eat is one of the best ways to stop his/her temper tantrums. ^c	0.91	
How often do you use food as a way to distract your child (e.g., if they are preventing you from doing your chores)? ^b	0.56	
In order to get my child to behave him/herself I promise him/her something to eat. ^c	0.76	
Threats and bribes		0.72
If your child does not like something, how often do you tell him/her that s/he will get a dessert if s/he tries it? ^b	0.82	
How often do you promise your child something other than food (e.g., toy, favorite activity) in return for eating specific foods? ^b	0.77	
I have to punish or remove privileges to get my child to eat more. ^c	0.63	
"Clean plate" policy		0.78
This is a policy in our home...		
If my child puts food on his/her plate, s/he has to eat it. ^c	0.86	
My child must stay at the table until a specified amount of food has been eaten. ^c	0.71	
My child is expected to eat everything on his/her plate at dinner. ^c	0.88	
Pressure to eat		0.64
How often do you help feed your child (e.g., hold the spoon to put food into his/her mouth)? ^b	0.74	
How often do you beg your child to eat? ^b	0.66	
How often do you coax or sweet talk your child to get him/her to take a bite? ^b	0.80	

^aHomeSTEAD is an abbreviation for the Home Self-administered Tool for Environmental assessment of Activity and Diet

^bResponse options offer a 5-point scale: never, rarely, sometimes, often, and always.

^cResponse options offer a 5-point scale: strongly disagree, disagree, neutral, agree, and strongly agree.

Table 2
HomeSTEAD's^a Autonomy Support scales with item wording, factor loadings, and internal consistency (Cronbach's α)

Scale Name and Items ^b	Factor Loading	Internal consistency
Encouragement		0.69
I encourage my child to look forward to the meal. ^c	0.58	
I encourage my child to enjoy his/her food. ^c	0.75	
This is a policy in our home: My child has to at least try or taste new foods. ^c	0.60	
Reasoning		0.68
How often do you have to encourage your child to eat things s/he doesn't like (because those foods are good for him/her)? ^d	0.71	
How often do you encourage vegetable consumption by making a game of eating vegetables or telling a story around eating a vegetable? ^d	0.52	
I reason with my child to get him/her to eat. ^c	0.49	
I negotiate with my child how much s/he can leave on his/her plate. ^c	0.58	
Praise		0.85
I praise my child if s/he eats what I give him/her. ^c	0.91	
I praise my child if s/he eats a new food. ^c	0.61	
I praise my child for choosing a healthy snack. ^c	0.80	
Nutrition education		0.73
How often do you try to make foods more familiar to your child by telling him/her where it came from? ^d	0.44	
How often do you suggest your child have a fruit/vegetable at snack time? ^d	0.51	
I discuss with my child the nutritional value of foods. ^c	0.93	
Do you give your child reasons for the rules you make about food and eating? ^d	0.89	
Guided choices: when food is eaten		0.69
I let my child eat between meals whenever s/he wants. ^c	0.63	
I let my child decide when s/he would like to have his/her meal. ^c	0.83	
I decide the times when my child eats his/her meals. ^c (R)	-0.82	
Guided choices: what food is eaten		0.66
I allow my child to choose what s/he has for snacks. ^c	0.47	
I decide what my child eats between meals. ^c (R)	-0.77	
As a parent, I decide the kinds of food that my child eats. ^c (R)	-0.56	
Guided choices: amount of food eaten		0.74
During meals, I allow my child to decide when s/he has had enough to eat. ^c	0.69	
At snack time, I allow my child to decide when s/he has had enough to eat. ^c	0.53	
I know better than my child if s/he is hungry or full. ^c (R)	-0.85	
When your child says "I'm not hungry", how often do you reply "You need to eat anyway"? ^d (R)	-0.77	

^aHomeSTEAD is an abbreviation for the Home Self-administered Tool for Environmental assessment of Activity and Diet

^bR is used to designate items that require reverse coding.

^cResponse options offer a 5-point scale: strongly disagree, disagree, neutral, agree, and strongly agree.

^dResponse options offer a 5-point scale: never, rarely, sometimes, often, and always.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3
HomeSTEAD^a Structure scales with item wording, factor loadings, and internal consistency (Cronbach's α)

Scale Name and Items ^b	Factor Loading	Internal consistency
Monitoring of unhealthy foods		0.93
I keep track of (either in my head or written down)...		
the sweets (candy, ice cream, cake, pies, pastries) that my child eats. ^c	0.98	
the snack food (potato chips, Doritos, cheese puffs) that my child eats. ^c	0.98	
the sugary drinks (soda/pop, Kool-Aid) my child drinks. ^c	0.87	
Rules and limits around unhealthy foods		0.80
I place limits on the sweet or salty snacks (candy, ice cream, cake, potato chips, tortilla chips) that my child eats? ^c	0.71	
How often do you restrict (or try to restrict) your child's access to fruit juice? ^d	0.66	
How often do you restrict (or try to restrict) your child's access to sweetened beverages? ^d	0.99	
If my child asks for sweetened beverages (including juice drinks or soda), I will give it to him/her. ^c (R)	-0.64	
How often do you allow your child to help him/herself to snacks, including salty or sweet snacks, or candy when s/he is at home? ^d (R)	-0.67	
Meal setting		0.67
How often does your child eat in a bedroom? ^d (R)	-0.70	
Do you limit snacking to designated places in your home? ^d	0.60	
I insist my child eats meals at the table. ^d	0.69	
Family eating		0.71
What do family dinners on weekdays look like at your home? ^e	0.88	
What do family dinners on weekend days look like at your home? ^e	0.77	
When schedules allow, how often do you or another adult in your household sit and eat lunch at home with your child? ^d	0.86	
Atmosphere of meals		0.72
Dinner time is usually a pleasant time for the family. ^c	0.60	
How often would you say arguments about eating occur during dinner time? ^d (R)	-0.69	
How often do other arguments, not about eating, occur during dinner time? ^d (R)	-0.66	
How frequently is the evening meal an unpleasant or stressful time for your family? ^d (R)	-0.80	
Eating area/physical space		0.62
How often do you decorate your table with flowers? ^d	0.62	
How often do you decorate your table with tablecloth? ^d	0.52	
How often do you decorate your table with placemats? ^d	0.47	
How often do you decorate your table with candles? ^d	0.70	
Distractions		0.87
During a typical weekend day, how often is the TV on during your child's meals even if s/he isn't watching it?		

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Scale Name and Items ^b	Factor Loading	Internal consistency
a. breakfast ^d	0.89	
b. snack ^d	0.96	
c. dinner ^d	0.87	
Planning and preparation of healthy meals		0.72
How often do you plan your family's meals to provide a variety of food groups?	0.78	
How often do you serve packaged, canned or frozen dinners as the main dish of a meal? ^d (R)	-0.78	
How often do you bake, broil, barbeque or steam food? ^d	0.65	
How often do you try to cook colorful (dark green, red, orange, purple) vegetables instead of potatoes or corn? ^d	0.66	
When you serve potatoes, how often are they fried, like French fries or hash browns? ^d (R)	-0.43	
How often does your family eat fast food for the main meal each week? ^f (R)	-0.50	
Attractive presentation of healthy foods		0.75
How often do you encourage vegetable consumption by serving vegetables in an interesting or attractive way? ^d	0.92	
How often do you encourage vegetable consumption by preparing the vegetables in alternative ways? ^d	0.75	
How often do you encourage vegetable consumption by serving vegetables hidden in other foods? ^d	0.47	
Availability and accessibility of healthy foods		0.66
Do you have fruits and vegetables that your child likes available at home? ^d	0.32	
How often is there fresh fruit on the counter, table, or somewhere else where your child can easily get to it? ^d	0.70	
How often are there cut-up vegetables in the fridge for your child to eat? ^d	0.51	
Modeling		0.64
I try to eat healthy foods in front of my child, even if they are not my favorite. ^c	0.59	
My child learns to eat healthy snacks from me. ^c	0.87	
How I eat does not particularly influence my child's habits. ^c (R)	-0.59	
How often do you try not to eat unhealthy foods when your children are around? ^d	0.44	
How often do you drink soda (regular or diet) or other sweetened beverages at meals and snacks with your child? ^d (R)	-0.38	
I eat food I want my child to eat. ^d	0.54	
Weight talk		0.76
How often do you complain about your own weight where your child can hear you? ^d	0.72	
How often does your family comment on each other's weight? ^d	0.74	
How often does your family talk about weight or dieting? ^d	0.82	

^aHomeSTEAD is an abbreviation for the Home Self-administered Tool for Environmental assessment of Activity and Diet

^bR is used to designate items that require reverse coding.

^cResponse options offer a 5-point scale: strongly disagree, disagree, neutral, agree, and strongly agree.

^dResponse options offer a 5-point scale: never, rarely, sometimes, often, and always.

^eResponse options offer a 5-point scale: 1=do not have family dinners, 2=child eats by him/herself or with other children, 3=parent and children eat in different areas, 4=one parent sits and watches child eat, 5=family members sit down and eat together.

^fResponse options offer an 8-point scale ranging from seldom/never to 7 days per week

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4
Means (SD) for HomeSTEAD[®] Scale Composite Scores at Time 1, 2 and 3, Mean Differences, and Reliability Testing^b

Scale	Means (SD)			p-value ^c	Reliability ICC (95% CI) ^d
	Time 1	Time 2	Time 3		
Coercive Control					
Restriction	1.5 (0.6)	1.5 (0.7)	1.6 (0.6)	0.86	0.66 (0.58, 0.74)
Soothing with food	1.4 (0.5)	1.5 (0.5)	1.5 (0.5)	0.41	0.60 (0.51, 0.69)
Threats and bribes	1.8 (0.7)	1.8 (0.6)	1.7 (0.6)	0.86	0.70 (0.62, 0.77)
"Clean plate" policy	2.6 (1.0)	2.6 (1.0)	2.7 (1.0)	0.73	0.78 (0.72, 0.83)
Pressure to eat	1.7 (0.7)	1.6 (0.7)	1.7 (0.7)	0.72	0.73 (0.67, 0.80)
Autonomy Support					
Encouragement	4.0 (0.6)	4.0 (0.6)	4.0 (0.6)	0.92	0.70 (0.62, 0.77)
Reasoning in response to picky eating	2.6 (0.8)	2.6 (0.7)	2.7 (0.7)	0.97	0.70 (0.62, 0.77)
Praise	4.0 (0.8)	4.0 (0.7)	4.0 (0.7)	0.80	0.68 (0.60, 0.75)
Nutrition education	3.5 (0.7)	3.5 (0.8)	3.5 (0.7)	0.98	0.80 (0.75, 0.85)
Guided choices: when food is eaten ^e	2.3 (0.7)	2.3 (0.7)	2.3 (0.6)	0.72	0.71 (0.64, 0.78)
Guided choices: what food is eaten ^e	2.5 (0.7)	2.5 (0.6)	2.4 (0.6)	0.46	0.57 (0.47, 0.66)
Guided choices: amount of food eaten ^e	3.7 (0.7)	3.6 (0.8)	3.6 (0.8)	0.38	0.78 (0.72, 0.83)
Structure					
Monitoring of unhealthy foods	3.8 (1.0)	3.7 (1.0)	3.8 (1.0)	0.67	0.65 (0.56, 0.72)
Rules and limits around unhealthy foods	3.8 (0.8)	3.8 (0.8)	3.8 (0.8)	0.68	0.80 (0.74, 0.85)
Meal setting	4.4 (0.7)	4.4 (0.7)	4.3 (0.8)	0.64	0.84 (0.80, 0.88)
Family eating	4.3 (0.8)	4.4 (0.7)	4.4 (0.7)	0.53	0.85 (0.80, 0.88)
Atmosphere of meals	4.0 (0.6)	4.1 (0.6)	4.1 (0.6)	0.24	0.75 (0.67, 0.81)
Eating area/physical space	2.5 (0.9)	2.6 (0.8)	2.6 (0.9)	0.91	0.89 (0.85, 0.92)
Distractions	2.3 (1.2)	2.2 (1.1)	2.3 (1.2)	0.70	0.78 (0.72, 0.84)
Planning and preparation of healthy meals	3.9 (0.6)	3.9 (0.5)	4.0 (0.5)	0.91	0.86 (0.82, 0.90)
Attractive presentation of healthy foods	2.8 (0.9)	2.8 (0.8)	2.9 (0.8)	0.76	0.70 (0.62, 0.77)
Availability and accessibility of healthy foods	3.9 (0.8)	3.9 (0.7)	3.8 (0.7)	0.59	0.75 (0.69, 0.81)
Modeling	3.9 (0.6)	3.9 (0.6)	3.9 (0.6)	0.95	0.76 (0.70, 0.82)

Scale	Means (SD)			p-value ^c	Reliability ICC (95% CI) ^d
	Time 1	Time 2	Time 3		
Weight talk	2.1 (0.8)	2.0 (0.7)	2.0 (0.7)	0.29	0.67 (0.59, 0.74)

^aHomeSTEAD is an abbreviation for the Home Self-administered Tool for Environmental assessment of Activity and Diet

^bUnless otherwise indicated, composite scores range from 1 to 5, with 5 being “strongly agree” or “always” answer choice.

^cp-value from repeated measures ANOVA testing. Statistical significance was set at 0.05.

^dSingle-measure intraclass correlations (ICC).

^eComposite scores range from 1 to 5, with higher scores being more child versus parent choice.

Table 5
Correlations between HomeSTEAD^a Scale Composite Scores at Time 1 and Dietary Outcomes

Scale	Weekly consumption of food or drink by child ^{b,c,d}			
	Fruits and Vegetables	Sugar-sweetened beverages	Snacks	Sweets
Coercive Control				
Restriction	-0.27**	0.49**	0.11	0.02
Soothing with food	-0.15	0.05	0.02	0.02
Threats and bribes	-0.05	0.10	0.17	0.04
“Clean plate” policy	0.06	0.08	0.23*	0.09
Pressure to eat	0.08	0.08	0.22*	0.12
Autonomy Support				
Encouragement	0.38**	-0.15	0.06	-0.01
Reasoning	0.07	-0.04	0.12	0.03
Praise	0.03	0.03	0.18*	0.08
Nutrition education	0.36**	-0.18	-0.12	-0.20*
Guided choices: when food is eaten	-0.05	-0.06	-0.08	-0.05
Guided choices: what food is eaten	-0.16	0.12	0.23*	0.20*
Guided choices: amount of food eaten	-0.03	-0.28**	-0.37**	-0.17
Structure				
Monitoring of unhealthy foods	0.29*	-0.07	0.10	<0.01
Rules and limits around unhealthy foods	0.41**	-0.37**	-0.29**	-0.23**
Meal setting	0.33**	-0.18*	-0.06	-0.27
Family eating	0.33**	-0.33**	-0.14	-0.05
Atmosphere of meals	0.04	0.03	<0.01	-0.01
Eating area/physical space	0.23**	-0.07	0.06	0.18
Distractions	-0.21*	0.39**	0.25**	0.21*
Planning and preparation of healthy meals	0.50**	-0.42**	-0.26**	-0.21*
Attractive presentation of healthy foods	0.33**	-0.08	-0.04	-0.12
Availability and accessibility of healthy foods	0.37**	-0.20*	<0.01	<0.01
Modeling	0.47**	-0.39**	-0.25*	-0.31**
Weight talk	-0.15	0.16	0.06	0.10

^aHomeSTEAD is an abbreviation for the Home Self-administered Tool for Environmental assessment of Activity and Diet

^bSpearman rank correlations

^cWeekly consumption of food or drink was reported by parents and grouped into ordered categories (e.g., categories for sweets include never/<1/wk, 1/wk, 2-4/wk, 5-6/wk, 1/d, >1 to <3/d, and >3/d).

^dStatistical significance was set at 0.05;

* however, a single asterisk is used to indicate $p < 0.05$,

** and a double asterisk is used to indicate $p < 0.01$.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript