
Maps of US Zika Cases

Maps of Zika in the United States

Zika Cases Reported in the United States

Laboratory-confirmed Zika virus disease cases reported to ArboNET by state or territory (as of March 1, 2017)



Laboratory-confirmed symptomatic Zika virus disease cases and presumptive viremic blood donors reported to ArboNET by states and territories — United States, 2015–2017 (as of March 1, 2017)

States	Symptomatic disease cases* (N=5,074) No. (%)	* Pr	esumptive viremic blood donors† (N=38) No. (%)
Alabama	38 (1)	0	(0)
Arizona	55 (1)	1	(3)
Arkansas	15 (<1)	0	(0)
California	426 (9)	5	(13)
Colorado	55 (1)	0	(0)
Connecticut	58 (1)	0	(0)
Delaware	17 (<1)	0	(0)

District of Columbia	31 (1)	0	(0)
Florida‡	1,083 (21)	24	(63)
Georgia	109 (2)	0	(0)
Hawaii	16 (<1)	0	(0)
Idaho	5 (<1)	0	(0)
Illinois	94 (2)	0	(0)
Indiana	53 (1)	0	(0)
Iowa	26 (1)	1	(3)
Kansas	22 (<1)	0	(0)
Kentucky	32 (1)	0	(0)
Louisiana	39 (1)	0	(0)
Maine	14 (<1)	0	(0)
Maryland	131 (3)	0	(0)
Massachusetts	121 (2)	0	(0)
Michigan	68 (1)	0	(0)
Minnesota	64 (1)	0	(0)
Mississippi	25 (1)	0	(0)
Missouri	36 (1)	0	(0)
Montana	9 (<1)	0	(0)
Nebraska	13 (<1)	0	(0)
Nevada	22 (<1)	1	(3)
New Hampshire	12 (<1)	0	(0)
New Jersey	180 (4)	0	(0)
New Mexico	10 (<1)	0	(0)
New York	1,004 (21)	3	(8)
North Carolina	91 (2)	0	(0)
North Dakota	3 (<1)	0	(0)
Ohio	85 (2)	0	(0)
Oklahoma	29 (1)	0	(0)
Oregon	47 (1)	0	(0)
Pennsylvania	174 (4)	0	(0)
Rhode Island	54 (1)	0	(0)
South Carolina	54 (1)	0	(0)
South Dakota	2 (<1)	0	(0)
Tennessee	61 (1)	0	(0)
Texas‡	312 (6)	3	(8)
Utah	22 (<1)	0	(0)

	` '	-	1-1
Vermont	11 (<1)	0	(0)
Virginia	113 (2)	0	(0)
Washington	70 (1)	0	(0)
West Virginia	11 (<1)	0	(0)
Wisconsin	50 (1)	0	(0)
Wyoming	2 (<1)	0	(0)

^{**} Includes reported confirmed and probable Zika virus disease cases per the <u>CSTE case</u> <u>definitions.</u>

[†] Presumptive viremic blood donors are people who reported no symptoms at the time of donating blood, but whose blood tested positive when screened for the presence of Zika virus RNA by the blood collection agency. Some presumptive viremic blood donors develop symptoms after their donation or may have had symptoms in the past. These individuals may be reported as both Zika virus disease cases and presumptive viremic blood donors. ‡Includes 215 cases in Florida and 6 cases in Texas that were acquired through presumed local mosquito-borne transmission.